

US Patent & Trademark Office

Patent Public Search | Text View

| | |
|----------------------|----------------------|
| United States Patent | 12383503 |
| Kind Code | B2 |
| Date of Patent | August 12, 2025 |
| Inventor(s) | Nunes; Cletus et al. |

Adagrasib solid pharmaceutical compositions

Abstract

Pharmaceutical compositions in solid form comprising adagrasib, suitable for oral dosage to treat subjects having cancer; as well as methods of manufacturing the compositions, and methods of treating cancer.

Inventors: Nunes; Cletus (San Diego, CA), Kheiripour; Mehrdad (San Diego, CA), Gavireddi; Monika (San Diego, CA)

Applicant: MIRATI THERAPEUTICS, INC. (Princeton, NJ)

Family ID: 1000008748074

Assignee: MIRATI THERAPEUTICS, INC. (Princeton, NJ)

Appl. No.: 18/794390

Filed: August 05, 2024

Prior Publication Data

| Document Identifier | Publication Date |
|---------------------|------------------|
| US 20240398815 A1 | Dec. 05, 2024 |

Related U.S. Application Data

continuation parent-doc WO PCT/US2023/030689 20230821 PENDING child-doc US 18794390
us-provisional-application US 63400640 20220824
us-provisional-application US 63399619 20220819

Publication Classification

Int. Cl.: A61K9/20 (20060101); A61K9/00 (20060101); A61K9/28 (20060101); A61K31/519 (20060101)

U.S. Cl.:

CPC **A61K9/2054** (20130101); **A61K9/0053** (20130101); **A61K9/2009** (20130101);
A61K9/2013 (20130101); **A61K9/2018** (20130101); **A61K9/2027** (20130101);
A61K9/28 (20130101); **A61K31/519** (20130101);

Field of Classification Search

USPC: None

References Cited

U.S. PATENT DOCUMENTS

| Patent No. | Issued Date | Patentee Name | U.S. Cl. | CPC |
|--------------|-------------|------------------|----------|-----|
| 10689377 | 12/2019 | Blake et al. | N/A | N/A |
| 2005/0051922 | 12/2004 | Nangia et al. | N/A | N/A |
| 2010/0310651 | 12/2009 | Mittal | N/A | N/A |
| 2015/0196493 | 12/2014 | Szymczak et al. | N/A | N/A |
| 2019/0224175 | 12/2018 | Golden et al. | N/A | N/A |
| 2021/0113569 | 12/2020 | Goncalves et al. | N/A | N/A |

FOREIGN PATENT DOCUMENTS

| Patent No. | Application Date | Country | CPC |
|------------|------------------|---------|-----|
| 3201612 | 12/2021 | CA | N/A |
| 115521312 | 12/2021 | CN | N/A |
| 3628664 | 12/2019 | EP | N/A |
| 2019099524 | 12/2018 | WO | N/A |
| 2022056307 | 12/2021 | WO | N/A |

OTHER PUBLICATIONS

International Search Report and Written Opinion for PCT Application No. PCT/US2023/030689, mailed on Feb. 26, 2024, 12 pages. cited by applicant

P01116 · Rask_Human, UniProtKB/Swiss-Prot, Retrieved on Aug. 29, 2024, 14 pages. cited by applicant

Alamgeer et al. (2013) “Novel Therapeutic Targets in Non-small Cell Lung Cancer”, Curr Opin Pharmacol., 13(3):394-401. cited by applicant

Berge et al. (Jan. 1977) “Pharmaceutical Salts”, Journal of Pharmaceutical Sciences, 66(1):1-19. cited by applicant

Dogan et al. (2012) “Molecular Epidemiology of EGFR and KRAS Mutations in 3,026 Lung Adenocarcinomas: Higher Susceptibility of Women to Smoking-related Kras-mutant Cancers”, Clin Cancer Res., 18(22):6169-6177. cited by applicant

Fell et al. (2020) “Identification of the Clinical Development Candidate MRTX849, a Covalent KRASG12C Inhibitor for the Treatment of Cancer”, Journal of Medicinal Chemistry, 63(13): 6679-6693. cited by applicant

McCormick, Frank (2015) “KRAS as a Therapeutic Target”, Clin Cancer Res., 21(8):1797-1801. cited by applicant

Ostrem et al. (2013) “K-Ras(G12C) Inhibitors Allosterically Control GTP Affinity and Effector Interactions”, Nature, 503(7477):548-551 (14 pages). cited by applicant

Samatar et al. (2014) “Targeting RAS-ERK Signalling in Cancer: Promises and Challenges”, Nat Rev Drug Discov., 13(12):928-942. cited by applicant

Santos et al. (1984) “Malignant Activation of a K-Ras Oncogene in Lung Carcinoma but Not in Normal Tissue of the Same Patient”, Science, 223(4637):661-664. cited by applicant
International Preliminary Report on Patentability for PCT Application No. PCT/US2023/030689, mailed on March 6, 2025, 9 pages. cited by applicant

Primary Examiner: Tran; Susan T

Attorney, Agent or Firm: Mintz, Levin, Cohn, Ferris, Glovsky and Popeo, P.C.

Background/Summary

FIELD OF THE INVENTION

(1) The present invention generally relates to solid pharmaceutical compositions of adagrasib (2-[(2S)-4-[7-(8-chloro-1-naphthyl)-2-[(2S)-1-methylpyrrolidin-2-yl]methoxy]-6,8-dihydro-5H-pyrido[3,4-d]pyrimidin-4-yl]-1-(2-fluoroprop-2-enoyl)piperazin-2-yl]acetonitrile); methods for preparing these compositions, and methods of their use for the treatment of various diseases and disorders.

BACKGROUND OF THE INVENTION

(2) Kirsten Rat Sarcoma 2 Viral Oncogene Homolog (“KRas”) is a small GTPase and a member of the Ras family of oncogenes. KRas serves as a molecular switch cycling between inactive (GDP-bound) and active (GTP-bound) states to transduce upstream cellular signals received from multiple tyrosine kinases to downstream effectors regulating a wide variety of processes, including cellular proliferation (e.g., see Alamgeer et al., (2013) Current Opin Pharmacol. 13:394-401).

(3) The role of activated KRas in malignancy was observed over thirty years ago (e.g., see Santos et al., (1984) Science 223:661-664). Aberrant expression of KRas accounts for up to 20% of all cancers and oncogenic KRas mutations that stabilize GTP binding and lead to constitutive activation of KRas and downstream signaling have been reported in 25-30% of lung adenocarcinomas. (e.g., see Samatar and Poulidakos (2014) Nat Rev Drug Disc 13(12): 928-942 doi: 10.1038/nrd428). Single nucleotide substitutions that result in missense mutations at codons 12 and 13 of the KRas primary amino acid sequence comprise approximately 40% of these KRas driver mutations in lung adenocarcinoma, with a G12C transversion being the most common activating mutation (e.g., see Dogan et al., (2012) Clin Cancer Res. 18(22):6169-6177, published online 2012 Sep. 26. doi: 10.1158/1078-0432.CCR-11-3265).

(4) The well-known role of KRas in malignancy and the discovery of these frequent mutations in KRas in various tumor types made KRas a highly attractable target of the pharmaceutical industry for cancer therapy. Notwithstanding thirty years of large scale discovery efforts to develop inhibitors of KRas for treating cancer, no KRas inhibitor has demonstrated sufficient safety and/or efficacy to obtain regulatory approval (e.g., see McCormick (2015) Clin Cancer Res. 21 (8):1797-1801).

(5) Recently, irreversible, covalent inhibitors that target KRas G12C have been described (e.g., see Ostrem et al., (2013) Nature 503:548-551). For instance, commonly-owned and assigned U.S. Provisional Application Ser. No. 62/586,775 discloses potent, orally bioavailable compounds that irreversibly bind to KRas G12C for treating KRas G12C-mediated cancers.

(6) A covalent, irreversible inhibitor of KRas G12C is adagrasib (2-[(2S)-4-[7-(8-chloro-1-naphthyl)-2-[(2S)-1-methylpyrrolidin-2-yl]methoxy]-6,8-dihydro-5H-pyrido[3,4-d]pyrimidin-4-yl]-1-(2-fluoroprop-2-enoyl)piperazin-2-yl]acetonitrile), also known as MRTX849. An amorphous form of this compound was described in International Patent Application PCT/US2018/061060 filed Nov. 14, 2018, published as WO2019/099524A1 on May 23, 2019 at Example 478, and in

Fell et al., (2020) J.Med. Chem. 63, 6679-6693. A crystalline form of this compound was described in International Patent Application PCT/US2021/049940 filed Sep. 10, 2021, published as WO2022/056307 on Mar. 17, 2022.

(7) Furthermore, it is desirable to reduce adverse events (AE) (also known as side effects) such as nausea, vomiting, diarrhea when the formulations are administered to patients.

(8) A need therefore exists for a pharmaceutical composition of adagrasib which displays suitable bioavailability and shelf-life stability, which is not susceptible to liquid capsule leakage over time, and which has fewer side effects compared with capsule formulations.

SUMMARY OF THE INVENTION

(9) In one embodiment, the present invention provides a pharmaceutical composition, in solid form, suitable for oral administration to a subject, including but not limited to a human subject, which comprises adagrasib or a salt thereof, wherein the solid pharmaceutical composition, after administration to the subject, is capable of providing $AUC_{sub.0 \rightarrow \infty}$ (the area under the curve of a plot of plasma drug concentration versus time) for adagrasib of at least, or about, 12000 ng*hr/mL under fasted conditions and/or of at least, or about, 19000 ng*hr/mL under fed conditions.

(10) In another embodiment, the solid pharmaceutical composition, after administration to a subject, is capable of providing $AUC_{sub.0 \rightarrow last}$ for adagrasib of at least, or about, 12000 ng*hr/mL under fasted conditions and/or of at least, or about, 19000 ng*hr/mL under fed conditions.

(11) In another embodiment, the solid pharmaceutical composition, after administration to a subject, is capable of providing $C_{sub.max}$ for the compound of formula (I) of at least, or about, 480 ng/mL under fasted conditions and/or of at least, or about, 630 ng/mL under fed conditions.

(12) In another embodiment, the solid pharmaceutical composition, after administration to a subject, is capable of providing: a) $AUC_{sub.0 \rightarrow \infty}$, for adagrasib of at least, or about, 12000 ng*hr/mL under fasted conditions and/or of at least, or about, 19000 ng*hr/mL under fed conditions; b) $AUC_{sub.0 \rightarrow last}$ for adagrasib of at least, or about, 12000 ng*hr/mL under fasted conditions and/or of at least, or about, 19000 ng*hr/mL under fed conditions; and c) $C_{sub.max}$ for adagrasib of at least, or about, 480 ng/mL under fasted conditions and/or of at least, or about, 630 ng/mL under fed conditions.

(13) In another embodiment, the solid pharmaceutical composition provides a $T_{sub.max}$ of less than about 6.5 hours, and preferably between about 6.0-6.25 hours.

(14) In any of the embodiments, it is not required that the recited pharmacokinetic (PK) values, such as $AUC_{sub.0 \rightarrow \infty}$, $AUC_{sub.0 \rightarrow last}$, and $C_{sub.max}$, are achieved by administering a single pharmaceutical composition. The invention contemplates, and explicitly includes, embodiments where these PK values are achieved following administration of several solid pharmaceutical compositions as a single dose (e.g., if a solid pharmaceutical composition (e.g., a tablet) comprises 200 mg adagrasib, the single dose may include, for example, three of such pharmaceutical compositions, for the total administered amount of 600 mg adagrasib).

(15) In another embodiment, adagrasib is present as a salt thereof.

(16) In another embodiment, the solid pharmaceutical composition comprises at least one additional anticancer compound in addition to the adagrasib.

(17) In another embodiment, the solid pharmaceutical composition is in the form of a powder or a tablet, including an encapsulated powder.

(18) In another embodiment, the solid pharmaceutical composition is in the form of a tablet.

(19) In another embodiment, the tablet of the invention comprises a film coat.

(20) In another embodiment, the tablet of the invention comprises: adagrasib, one or more diluents, a disintegrant, a glidant, a lubricant, and a film coat.

(21) In a preferred embodiment, the diluent is selected from the group consisting of microcrystalline cellulose, mannitol, and combinations thereof.

- (22) In another preferred embodiment, the disintegrant comprises crospovidone.
- (23) In another preferred embodiment, the glidant comprises colloidal silicon dioxide.
- (24) In another preferred embodiment, the lubricant comprises magnesium stearate.
- (25) In a preferred embodiment, the tablet of the invention comprises adagrasib, microcrystalline cellulose, mannitol, crospovidone, colloidal silicon dioxide, magnesium stearate and a film coat.
- (26) In another embodiment, the solid pharmaceutical composition is provided as a unit dosage form.
- (27) In one embodiment, the amount of adagrasib in the solid pharmaceutical composition is at least, or about, 200 mg.
- (28) In another embodiment, the amount of adagrasib in the solid pharmaceutical composition is at least, or about, 300 mg.
- (29) In another embodiment, the amount of adagrasib in the solid pharmaceutical composition is at least, or about, 400 mg.
- (30) In another embodiment, the amount of adagrasib in the solid pharmaceutical composition is at least, or about, 600 mg.
- (31) In another embodiment, the composition is in the form of a tablet and comprises: (1) adagrasib constituting about 30-35% percent of the composition; (2) microcrystalline cellulose constituting about 50-55% of the composition; (3) mannitol constituting about 8-10% of the composition; (4) crospovidone constituting about 2% to about 5% of the composition; (5) colloidal silicon dioxide constituting up to about 0.5-1.5% of the composition; and (6) magnesium stearate constituting about 1-2% of the composition.

wherein all percentages are percentages by weight and wherein the total weight is 100%.

- (32) In another embodiment, the composition further comprises a film coat.
- (33) In another embodiment, the composition is in the form of a tablet and comprises: (1) adagrasib constituting about 30-67% percent of the composition; (2) microcrystalline cellulose constituting about 20-55% of the composition; (3) mannitol constituting about 0-10% of the composition; (4) crospovidone constituting about 2% to about 5% of the composition; (5) colloidal silicon dioxide constituting up to about 0.5-2% of the composition; and (6) magnesium stearate constituting about 1-3% of the composition.

wherein all percentages are percentages by weight and wherein the total weight is 100%.

- (34) In another embodiment, the composition is in the form of a tablet and comprises: (1) adagrasib constituting about 50-67% percent of the composition; (2) microcrystalline cellulose constituting about 20-45% of the composition; (3) mannitol constituting about 0-10% of the composition; (4) crospovidone constituting about 2% to about 5% of the composition; (5) colloidal silicon dioxide constituting up to about 0.5-2% of the composition; and (6) magnesium stearate constituting about 1-3% of the composition.

wherein all percentages are percentages by weight and wherein the total weight is 100%.

- (35) In a preferred embodiment, the composition of the invention comprises: (1) adagrasib at about 33.3% percent of the composition; (2) microcrystalline cellulose at about 51.2% of the composition; (3) mannitol at about 10.0% of the composition; (4) crospovidone at about 3.0% of the composition; (5) colloidal silicon dioxide at about 1.0% of the composition; and (6) magnesium stearate at about 1.5% of the composition, wherein all percentages are percentages by weight and wherein the total weight is 100%.

- (36) In another embodiment, the solid pharmaceutical composition is prepared by a process which comprises the steps of: (a) pre-blending, wherein diluents, a glidant, and adagrasib are blended together; (b) de-lumping the components of step (a) using a screening mill; (c) blending the components of step (b) with a disintegrant, and a glidant; (d) lubricating by adding a lubricant to the components of step (c); (e) performing a dry granulation of the components of step (d) using a roller compactor, producing a granulated blend; (f) blending a diluent, a disintegrant and a glidant with the granulated blend of step (e); (g) lubricating by adding a lubricant to the components of

step (f) to produce a lubricated blend; (h) performing compression by charging the lubricated blend of step (g) to a rotary tablet press and compressing the lubricated blend into core tablets; and (i) performing film coating by charging the core tablets of step (h) into a pan coater and adding a film coating agent.

(37) One or more pharmaceutically acceptable excipients (or combinations thereof) may be included as intra-granular material, i.e., included prior to conducting dry granulation; and/or such excipients may be added as extra-granular material to the dry granulation product, i.e., added after the dry granulation product has been made.

(38) In yet another embodiment, the invention is directed to a method of making a solid pharmaceutical composition comprising adagrasib or a salt thereof, wherein said solid pharmaceutical composition is made by dry granulation.

(39) In one embodiment, the method comprises the steps of: (a) pre-blending, wherein a diluent, a glidant, and adagrasib are blended together; (b) de-lumping the components of step (a) using a screening mill; (c) blending the components of step (b) with a disintegrant, and a glidant; (d) lubricating by adding a lubricant to the components of step (c); (e) performing a dry granulation of the components of step (d) using a roller compactor, producing a granulated blend; (f) blending a diluent, a disintegrant and a glidant with the granulated blend of step (e); (g) lubricating by adding a lubricant to the components of step (f) to produce a lubricated blend; (h) performing compression by charging the lubricated blend of step (g) to a rotary tablet press and compressing the lubricated blend into tablet cores; and (i) performing film coating by charging the core tablets of step (h) into a pan coater and adding a film coating agent.

(40) In one embodiment, the solid pharmaceutical composition (e.g., tablet) may comprise additional excipients selected from the group consisting of diluents, fillers, super-disintegrants, binders, glidants, lubricants, and combinations thereof.

(41) In another embodiment, the invention is directed to a method of treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of the solid pharmaceutical composition of the present invention. In one embodiment, the therapeutically effective amount is at least, or about, 600 mg of adagrasib. In one embodiment, the cancer is a KRas G12C-associated cancer. In one embodiment, the KRas G12C-associated cancer is lung cancer. In one embodiment, the solid pharmaceutical composition is a tablet.

(42) Also provided herein are methods for treating cancer in a subject in need thereof, the method comprising (a) determining that cancer is associated with a KRas G12C mutation (e.g., a KRas G12C-associated cancer) (e.g., as determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit); and (b) administering to the patient a therapeutically effective amount of the solid pharmaceutical composition of the present invention. In one embodiment, the solid pharmaceutical composition is a tablet.

(43) In one embodiment, the provided methods of treating cancer cause fewer side effects in the subject than side effects associated with administering a capsule composition of the same amount of adagrasib.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) FIG. 1 is a manufacturing process flow diagram depicting one way of making solid pharmaceutical compositions of the invention.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

(2) The term “adagrasib” as used herein refers to a compound with the following formula:

(3) ##STR00001##

as well as pharmaceutically acceptable salts of this compound. The compound has the following chemical name: (2-[(2S)-4-[7-(8-chloro-1-naphthyl)-2-[(2S)-1-methylpyrrolidin-2-yl]methoxy]-6,8-dihydro-5H-pyrido[3,4-d]pyrimidin-4-yl]-1-(2-fluoroprop-2-enoyl)piperazin-2-yl]acetonitrile). It is also known as MRTX849. The invention encompasses amorphous and crystalline forms of adagrasib. An amorphous form of this compound was described in International Patent Application PCT/US2018/061060 filed Nov. 14, 2018, published as WO2019/099524A1 on May 23, 2019 at Example 478, and in Fell et al., (2020) J. Med. Chem. 63, 6679-6693. A crystalline form of this compound was described in International Patent Application PCT/US2021/049940 filed Sep. 10, 2021, published as WO2022/056307 on Mar. 17, 2022. Form 1, as referred to in this application, corresponds to Form A in PCT/US2021/049940; Form 2, as referred to in this application, corresponds to Form B in PCT/US2021/049940. The tablets described herein preferably comprise crystalline form(s) of adagrasib. The contents of these patent applications and of the literature reference are hereby incorporated by reference in their entirety.

(4) As used herein, “KRas G12C” refers to a mutant form of a mammalian KRas protein that contains an amino acid substitution of a cysteine for a glycine at amino acid position 12. The assignment of amino acid codon and residue positions for human KRas is based on the amino acid sequence identified by UniProtKB/Swiss-Prot P01116: Variant p.Gly12Cys.

(5) The term “composition” as used herein is intended to encompass a product comprising the specified ingredients (and in the specified amounts, if indicated), as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. By “pharmaceutically acceptable” it is meant the diluent, excipient or carrier must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

(6) The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

(7) The term “solid forms” and related terms used herein, unless otherwise specified, refers to crystalline forms comprising adagrasib and its various salt forms.

(8) The term “crystalline” and related terms used herein, when used to describe a substance, component or product, means that the substance, component or product is crystalline as determined by X-ray diffraction. See, for example, *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton PA, p. 173 (1990); *The United States Pharmacopeia*, 23rd ed., pp. 1843-1844 (1995); the contents of which are hereby incorporated by reference in their entireties.

(9) The term “crystalline forms” and related terms herein refers to the various crystalline modifications of a given substance, including, but not limited to, polymorphs, solvates, hydrates, co-crystals and other molecular complexes, as well as salts, solvates of salts, hydrates of salts, other molecular complexes of salts, and polymorphs thereof.

(10) The term “pharmaceutically acceptable salts” is meant to include salts of active compounds which are prepared with relatively nontoxic acids. Acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic; propionic; isobutyric; maleic; malonic; benzoic; succinic; suberic; fumaric; mandelic; phthalic; benzenesulfonic; toluenesulfonic, including p-toluenesulfonic, m-toluenesulfonic, and o-toluenesulfonic; citric; tartaric; methanesulfonic; and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galacturonic acids and the like (see, for example, Berge et al. J. Pharm. Sci. 66:1-

19 (1977)).

(11) The term, “amorphous form,” as used herein, refers to a noncrystalline form of a substance.

(12) The terms, “polymorphs” and “polymorphic forms” and related terms herein refer to crystal forms of a molecule. Different polymorphs may have different physical properties such as, for example, melting temperatures, heats of fusion, solubilities, dissolution rates and/or vibrational spectra as a result of the arrangement or conformation of the molecules in the crystal lattice. The differences in physical properties exhibited by polymorphs affect pharmaceutical parameters such as storage stability, compressibility and density (important in formulation and product manufacturing), and dissolution rates (an important factor in bioavailability). Polymorphs of a molecule can be obtained by a number of methods, as known in the art. Such methods include, but are not limited to, melt recrystallization, melt cooling, solvent recrystallization, desolvation, rapid evaporation, rapid cooling, slow cooling, vapor diffusion and sublimation.

(13) Techniques for characterizing polymorphs include, but are not limited to, differential scanning calorimetry (DSC), X-ray powder diffractometry (XRPD), single crystal X-ray diffractometry, vibrational spectroscopy, e.g., IR and Raman spectroscopy, solid state NMR, hot stage optical microscopy, scanning electron microscopy (SEM), electron crystallography and quantitative analysis, particle size analysis (PSA), surface area analysis, solubility studies and dissolution studies.

(14) The term, “solvate,” as used herein, refers to a crystal form of a substance which contains solvent. The term “hydrate” refers to a solvate wherein the solvent is water.

(15) The term, “desolvated solvate,” as used herein, refers to a crystal form of a substance which can only be made by removing the solvent from a solvate.

(16) The term “excipient” refers to an inactive ingredient of the pharmaceutical compositions of the invention. It includes, but is not limited to, solvents, wetting agents, diluents, superdisintegrants, binders, glidants, and lubricants.

(17) The terms “treat”, “treating” or “treatment”, as used herein, refer to the reduction or amelioration of the progression, severity, and/or duration of a disorder or the eradication, reduction or amelioration of symptoms of a disorder, or the delay of the recurrence or onset of a disorder or one or more symptoms thereof in a subject that results from the administration of one or more compound.

(18) As used herein, the term “subject,” “individual,” or “patient,” used interchangeably, refers to any animal, including mammals such as mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, primates, and humans. In some embodiments, the patient is a human. In some embodiments, the subject has experienced and/or exhibited at least one symptom of the disease or disorder to be treated and/or prevented. In some embodiments, the subject has been identified or diagnosed as having a cancer having a KRas G12C mutation (e.g., as determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit). In some embodiments, the subject has a tumor that is positive for a KRas G12C mutation (e.g., as determined using a regulatory agency-approved assay or kit). The subject can be a subject with a tumor(s) that is positive for a KRas G12C mutation (e.g., identified as positive using a regulatory agency-approved, e.g., FDA-approved, assay or kit). The subject can be a subject whose tumors have a KRas G12C mutation (e.g., where the tumor is identified as such using a regulatory agency-approved, e.g., FDA-approved, kit or assay). In some embodiments, the subject is suspected of having a KRas G12C gene-associated cancer. In some embodiments, the subject has a clinical record indicating that the subject has a tumor that has a KRas G12C mutation (and optionally the clinical record indicates that the subject should be treated with any of the compositions provided herein).

(19) The term “pediatric patient” as used herein refers to a patient under the age of 16 years at the time of diagnosis or treatment. The term “pediatric” can be further be divided into various subpopulations including: neonates (from birth through the first month of life); infants (1 month up to two years of age); children (two years of age up to 12 years of age); and adolescents (12 years of

age through 21 years of age (up to, but not including, the twenty-second birthday)). Berhman R E, Kliegman R, Arvin A M, Nelson W E. Nelson Textbook of Pediatrics, 15th Ed. Philadelphia: W. B. Saunders Company, 1996; Rudolph A M, et al. Rudolph's Pediatrics, 21st Ed. New York: McGraw-Hill, 2002; and Avery M D, First L R. Pediatric Medicine, 2nd Ed. Baltimore: Williams & Wilkins; 1994.

(20) In some embodiments of any of the methods or uses described herein, an assay is used to determine whether the patient has KRas G12C mutation using a sample (e.g., a biological sample or a biopsy sample such as a paraffin-embedded biopsy sample) from a patient (e.g., a patient suspected of having a KRas G12C-associated cancer, a patient having one or more symptoms of a KRas G12C-associated cancer, and/or a patient that has an increased risk of developing a KRas G12C-associated cancer) can include, for example, next generation sequencing, immunohistochemistry, fluorescence microscopy, break apart FISH analysis, Southern blotting, Western blotting, FACS analysis, Northern blotting, and PCR-based amplification (e.g., RT-PCR, quantitative real-time RT-PCR, allele-specific genotyping or ddPCR). As is well-known in the art, the assays are typically performed, e.g., with at least one labelled nucleic acid probe or at least one labelled antibody or antigen-binding fragment thereof.

(21) The term “regulatory agency” is a country's agency for the approval of the medical use of pharmaceutical agents with the country. For example, a non-limiting example of a regulatory agency is the U.S. Food and Drug Administration (FDA).

(22) As used herein, a “therapeutically effective amount” is an amount that is sufficient to ameliorate, or in some manner reduce a symptom or stop or reverse progression of a condition, or negatively modulate or inhibit the activity of KRas G12C. Such amount may be administered as a single dosage or may be administered according to a regimen, whereby it is effective.

(23) As used herein, treatment means any manner in which the symptoms or pathology of a condition, disorder or disease are ameliorated or otherwise beneficially altered. Treatment also encompasses any pharmaceutical use of the compositions herein.

(24) As used herein, amelioration of the symptoms of a particular disorder by administration of a particular pharmaceutical composition refers to any lessening, whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the composition.

(25) As used herein, the term “about” or “approximately” means an acceptable error for a particular value as determined by one of ordinary skill in the art, which depends in part on how the value is measured or determined. In certain embodiments, the term “about” or “approximately” means within 1, 2, 3, or 4 standard deviations. In certain embodiments, the term “about” or “approximately” means within 50%, 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, or 0.05% of a given value or range.

(26) The term “fasted conditions” refers to a subject fasting at least 10 hours before being administered compositions of the invention.

(27) The term “fed conditions” refers to a subject being administered compositions of the inventions no later than 30-45 minutes after consuming food.

(28) The term, “AUC,” as used herein, refers to the area under the curve of a plot of plasma drug concentration versus time.

(29) The term, “T.sub.max,” as used herein, refers to the time after administration of a drug when the maximum plasma concentration is reached.

(30) A “KRas G12C-associated disease or disorder” as used herein refers to diseases or disorders associated with or mediated by or having a KRas G12C mutation. A non-limiting example of a KRas G12C-associated disease or disorder is a KRas G12C-associated cancer.

DETAILED DESCRIPTION OF COMPOSITIONS AND METHODS

(31) The present invention is based on a surprising discovery that solid pharmaceutical compositions (e.g., tablets) comprising adagrasib result in fewer adverse events (AEs, also referred to as “side effects”), such as nausea, vomiting, diarrhea, when administered to patients, than

capsule formulations comprising the same amount of adagrasib. Without wishing to be bound to a specific theory, it is hypothesized that the lower number of the upper GI related AEs could be due to lower maximum concentration (C_{max}) compared to the capsules. Emerging data from the ongoing Phase 3 study indicate that the tablet administration also results in fewer upper GI AEs. Another possibility is that, in the case of the tablet, the disintegration mechanism and greater density may lead to more rapid clearance of the drug from the stomach. In contrast, the capsule may float or adhere to the stomach lining and rapidly release the drug in the stomach, leading to higher local drug levels. If the drug is irritating to stomach this could lead to higher AEs.

(32) Thus, in one embodiment, the present invention provides a pharmaceutical composition, in solid form, suitable for oral administration to a subject, including but not limited to a human subject, which comprises adagrasib or a salt thereof, wherein the solid pharmaceutical composition, after administration to the subject, is capable of providing AUC_{sub.0.fwdarw.∞} (the area under the curve of a plot of plasma drug concentration versus time) for adagrasib of at least, or about, 12000 ng*hr/mL under fasted conditions and/or of at least, or about, 19000 ng*hr/mL under fed conditions.

(33) In another embodiment, the solid pharmaceutical composition, after administration to a subject, is capable of providing AUC_{sub.0.fwdarw.last} for adagrasib of at least, or about, 12000 ng*hr/mL under fasted conditions and/or of at least, or about, 19000 ng*hr/mL under fed conditions.

(34) In another embodiment, the solid pharmaceutical composition, after administration to a subject, is capable of providing C_{sub.max} for the compound of formula (I) of at least, or about, 480 ng/mL under fasted conditions and/or of at least, or about, 630 ng/mL under fed conditions.

(35) In another embodiment, the solid pharmaceutical composition, after administration to a subject, is capable of providing: a) AUC_{sub.0.fwdarw.∞} for adagrasib of at least, or about, 12000 ng*hr/mL under fasted conditions and/or of at least, or about, 19000 ng*hr/mL under fed conditions; b) AUC_{sub.0.fwdarw.last} for adagrasib of at least, or about, 12000 ng*hr/mL under fasted conditions and/or of at least, or about, 19000 ng*hr/mL under fed conditions; and c) C_{sub.max} for adagrasib of at least, or about, 480 ng/mL under fasted conditions and/or of at least, or about, 630 ng/mL under fed conditions.

(36) In another embodiment, the solid pharmaceutical composition provides a T_{sub.max} of less than about 6.5 hours, and preferably between about 6.0-6.25 hours.

(37) In any of the embodiments, it is not required that the recited pharmacokinetic (PK) values, such as AUC_{sub.0.fwdarw.∞}, AUC_{sub.0.fwdarw.last}, and C_{sub.max}, are achieved by administering a single pharmaceutical composition. The invention contemplates, and explicitly includes, embodiments where these PK values are achieved following administration of several solid pharmaceutical compositions as a single dose (e.g., if a solid pharmaceutical composition (e.g., a tablet) comprises 200 mg adagrasib, the single dose may include, for example, three of such pharmaceutical compositions, for the total administered amount of 600 mg adagrasib).

(38) In another embodiment, adagrasib is present as a salt thereof.

(39) In another embodiment, the solid pharmaceutical composition comprises at least one additional anticancer compound in addition to the adagrasib.

(40) In another embodiment, the solid pharmaceutical composition is in the form of a powder or a tablet, including an encapsulated powder.

(41) In another embodiment, the solid pharmaceutical composition is in the form of a tablet.

(42) In one embodiment, the solid pharmaceutical composition (e.g., tablet) may comprise various excipients selected from the group consisting of solvents, wetting agents, diluents, fillers, super-disintegrants, binders, glidants, lubricants, and combinations thereof. It should be understood that the invention contemplates use of other excipients that serve substantially the same functions in substantially the same manner as those described above.

(43) Tablets may contain the solid active ingredient in admixture with at least one pharmaceutically

acceptable excipient which is suitable for the manufacture of tablets. The excipient may be, for example; a disintegrating agent, such as a super-disintegrant; a binder; a diluent; a glidant; a lubricant; an emulsifier; or any other excipient known to one of skill in the art.

(44) Preferred tablets are those which provide good potency, content uniformity, hardness, friability and dissolution, and which contribute to the chemical and physical stability of the pharmaceutical compositions.

(45) In another embodiment, the tablet of the invention comprises a film coat.

(46) In another embodiment, the tablet of the invention comprises: adagrasib, one or more diluents, a disintegrant, a glidant, a lubricant, and a film coat.

(47) In a preferred embodiment, the diluent is selected from the group consisting of microcrystalline cellulose, mannitol, and combinations thereof.

(48) In another preferred embodiment, the disintegrant comprises crospovidone.

(49) In another preferred embodiment, the glidant comprises colloidal silicon dioxide.

(50) In another preferred embodiment, the lubricant comprises magnesium stearate.

(51) In a preferred embodiment, the tablet of the invention comprises adagrasib, microcrystalline cellulose, mannitol, crospovidone, colloidal silicon dioxide, magnesium stearate and a film coat.

(52) In another embodiment, the solid pharmaceutical composition is provided as a unit dosage form.

(53) In one embodiment, the amount of adagrasib in the solid pharmaceutical composition is at least, or about, 200 mg.

(54) In another embodiment, the amount of adagrasib in the solid pharmaceutical composition is at least, or about, 300 mg.

(55) In another embodiment, the amount of adagrasib in the solid pharmaceutical composition is at least, or about, 400 mg.

(56) In another embodiment, the amount of adagrasib in the solid pharmaceutical composition is at least, or about, 600 mg.

(57) In another embodiment, the composition is in the form of a tablet and comprises: (1) adagrasib constituting about 30-35% percent of the composition; (2) microcrystalline cellulose constituting about 50-55% of the composition; (3) mannitol constituting about 8-10% of the composition; (4) crospovidone constituting about 2% to about 5% of the composition; (5) colloidal silicon dioxide constituting up to about 0.5-1.5% of the composition; and (6) magnesium stearate constituting about 1-2% of the composition.

wherein all percentages are percentages by weight and wherein the total weight is 100%.

(58) In another embodiment, the composition is in the form of a tablet and comprises: (1) adagrasib constituting about 30-67% percent of the composition; (2) microcrystalline cellulose constituting about 20-45% of the composition; (3) mannitol constituting about 0-10% of the composition; (4) crospovidone constituting about 2% to about 5% of the composition; (5) colloidal silicon dioxide constituting up to about 0.5-2% of the composition; and (6) magnesium stearate constituting about 1-3% of the composition.

wherein all percentages are percentages by weight and wherein the total weight is 100%.

(59) In another embodiment, the composition is in the form of a tablet and comprises: (1) adagrasib constituting about 50-67% percent of the composition; (2) microcrystalline cellulose constituting about 20-45% of the composition; (3) mannitol constituting about 0-10% of the composition; (4) crospovidone constituting about 2% to about 5% of the composition; (5) colloidal silicon dioxide constituting up to about 0.5-2% of the composition; and (6) magnesium stearate constituting about 1-3% of the composition.

wherein all percentages are percentages by weight and wherein the total weight is 100%.

(60) In another embodiment, the composition further comprises a film coat.

(61) In a preferred embodiment, the composition of the invention comprises: (1) adagrasib at about 33.3% percent of the composition; (2) microcrystalline cellulose at about 51.2% of the

composition; (3) mannitol at about 10.0% of the composition; (4) crospovidone at about 3.0% of the composition; (5) colloidal silicon dioxide at about 1.0% of the composition; and (6) magnesium stearate at about 1.5% of the composition, wherein all percentages are percentages by weight and wherein the total weight is 100%.

(62) In another embodiment, the solid pharmaceutical composition is prepared by a process which comprises the steps of: (a) pre-blending, wherein diluents, a glidant, and adagrasib are blended together; (b) de-lumping the components of step (a) using a screening mill; (c) blending the components of step (b) with a disintegrant, and a glidant; (d) lubricating by adding a lubricant to the components of step (c); (e) performing a dry granulation of the components of step (d) using a roller compactor, producing a granulated blend; (f) blending a diluent, a disintegrant and a glidant with the granulated blend of step (e); (g) lubricating by adding a lubricant to the components of step (f) to produce a lubricated blend; (h) performing compression by charging the lubricated blend of step (g) to a rotary tablet press and compressing the lubricated blend into tablet cores; and (i) performing film coating by charging the core tablets of step (h) into a pan coater and adding a film coating agent.

(63) One or more pharmaceutically acceptable excipients (or combinations thereof) may be included as intragranular material, i.e., included prior to conducting dry granulation; and/or such excipients may be added as extra-granular material to the dry granulation product, i.e., added after the dry granulation product has been dried.

(64) In yet another embodiment, the invention is directed to a method of making a solid pharmaceutical composition comprising adagrasib or a salt thereof, wherein said solid pharmaceutical composition is made by dry granulation.

(65) In one embodiment, the method comprises the steps of: (a) pre-blending, wherein diluents, a glidant, and adagrasib are blended together; (b) de-lumping the components of step (a) using a screening mill; (c) blending the components of step (b) with a disintegrant, and a glidant; (d) performing an intra-granular lubrication by adding a lubricant to the components of step (c); (e) performing a dry granulation of the components of step (d) using a roller compactor, producing a granulated blend; (f) blending a diluent, a disintegrant and a glidant with the granulated blend of step (e); (g) performing an extra-granular lubrication by adding a lubricant to the components of step (f) to produce a lubricated blend; (h) performing compression by charging the lubricated blend of step (g) to a rotary tablet press and compressing the lubricated blend into tablet cores; and (i) adding a film coating to the tablet cores of step (h).

(66) The pharmaceutical compositions provided herein can further include one or more pharmaceutically acceptable additives such as a suspending agent, a flavoring agent, a sweetening agent, a dispersing agent, a surfactant, a colorant, a solubilizer, a moistening agent, a plasticizer, a stabilizer, a penetration enhancer, an anti-foaming agent, an antioxidant, an preservative, or a mixture thereof.

(67) In another embodiment, the invention is directed to a method of treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of the solid pharmaceutical composition of the present invention. In one embodiment, the therapeutically effective amount is at least, or about, 600 mg of adagrasib. In one embodiment, the cancer is a KRas G12C-associated cancer. In one embodiment, the KRas G12C-associated cancer is lung cancer. In one embodiment, the solid pharmaceutical composition is a tablet.

(68) Also provided herein are methods for treating cancer in a subject in need thereof, the method comprising (a) determining that cancer is associated with a KRas G12C mutation (e.g., a KRas G12C-associated cancer) (e.g., as determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit); and (b) administering to the patient a therapeutically effective amount of the solid pharmaceutical composition of the present invention. In one embodiment, the solid pharmaceutical composition is a tablet.

(69) In one embodiment, the provided methods of treating cancer cause fewer side effects in the

subject than side effects associated with administering a capsule composition of the same amount of adagrasib.

(70) It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including, for example, the activity of the specific polymorph employed, the metabolic stability and length of action of that polymorph, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, and the severity of the patient's condition.

(71) The pharmaceutical compositions provided herein can be combined with other compounds having related utilities to treat or prevent cancer. In many instances, administration of the subject pharmaceutical compositions in conjunction with these alternative agents enhances the efficacy of such agents. Accordingly, in some instances, the present pharmaceutical compositions, when combined or administered in combination with, e.g., anti-cancer agents, can be used in dosages which are less than the expected amounts when used alone, or less than the calculated amounts for combination therapy.

(72) High drug load formulations containing greater than 50% drug loading with and without Mannitol were successfully evaluated for their processability, in vitro dissolution and bioavailability in preclinical species (dog model). Formulations up to 67% drug loading were successfully evaluated and it was observed that the amount of Mannitol impacts both processability and bioavailability. It was found that reduction of mannitol w/w % from 10% w/w to 5% w/w and 0% w/w increases bioavailability of the drug and improves processability.

EXAMPLES OF THE INVENTION

(73) The following Examples illustrate the invention.

Example 1: Adagrasib Formulations

(74) This example illustrates an adagrasib tablet formulation, which was made and tested for bioavailability and food effect.

(75) The tablet is an immediate release (IR) film-coated tablet for oral administration. The dosage strength of each tablet is 200 mg or 300 mg, or 400 mg or 600 mg active (adagrasib). The qualitative and quantitative composition per unit of drug product is illustrated in Tables 1-9. No overages are used in the product.

(76) TABLE-US-00001 TABLE 1 Drug Product, 200 mg (33% (w/w)), Qualitative and Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
Adagrasib Drug 200 33.33 Drug Substance^a Substance Microcrystalline 307 51.17 Diluent
Cellulose.sup.b Mannitol 60 10 Diluent Crospovidone 18 3 Disintegrant Colloidal Silicon Dioxide
6 1 Glidant Magnesium Stearate.sup.c 9 1.5 Lubricant Core Tablet Weight 600 100 Film Coat
Agent 18-30 3-5 Film Coat Total Tablet Weight 618-630 103-105

(77) TABLE-US-00002 TABLE 2 Drug Product, 300-600 mg (50-67% (w/w)), Qualitative and Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
Adagrasib Drug Substance 300-600 50-67 Drug Substance Microcrystalline Cellulose 120-360 20-
45 Diluent Mannitol 0-100 0-10 Diluent Crospovidone 10-45 2-5 Disintegrant Colloidal Silicon
Dioxide 3-18 0.5-2 Glidant Magnesium Stearate.sup.d 6-27 1-3 Lubricant Core Tablet Weight
450-900 100 Film Coat Agent 10-54 2-6 Film Coat Total Tablet Weight 460-954 102-106

(78) TABLE-US-00003 TABLE 3 Drug Product, 300 mg (50% (w/w)), Qualitative and Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
Adagrasib Drug Substance 300 50 Drug Substance Microcrystalline Cellulose 192 32 Diluent
Mannitol 60 10 Diluent Crospovidone 30 5 Disintegrant Colloidal Silicon Dioxide 6 1 Glidant
Magnesium Stearate.sup.e 12 2 Lubricant Core Tablet Weight 600 100 Film Coat Agent 18 3 Film
Coat Total Tablet Weight 618 103

(79) TABLE-US-00004 TABLE 4 Drug Product, 400 mg (50% (w/w)), Qualitative and Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
Adagrasib Drug Substance 400 50 Drug Substance Microcrystalline Cellulose 256 32 Diluent

Mannitol 80 10 Diluent Crospovidone 40 5 Disintegrant Colloidal Silicon Dioxide 8 1 Glidant
 Magnesium Stearate.sup.f 16 2 Lubricant Core Tablet Weight 800 100 Film Coat Agent 24 3 Film
 Coat Total Tablet Weight 824 103

(80) TABLE-US-00005 TABLE 5 Drug Product, 300 mg (50% (w/w)), Qualitative and
 Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
 Adagrasib Drug Substance 300 50 Drug Substance Microcrystalline Cellulose 222 37 Diluent
 Mannitol 30 5 Diluent Crospovidone 30 5 Disintegrant Colloidal Silicon Dioxide 6 1 Glidant
 Magnesium Stearate.sup.g 12 2 Lubricant Core Tablet Weight 600 100 Film Coat Agent 18 3 Film
 Coat Total Tablet Weight 618 103

(81) TABLE-US-00006 TABLE 6 Drug Product, 400 mg (50% (w/w)), Qualitative and
 Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
 Adagrasib Drug Substance 400 50 Drug Substance Microcrystalline Cellulose 296 37 Diluent
 Mannitol 40 5 Diluent Crospovidone 40 5 Disintegrant Colloidal Silicon Dioxide 8 1 Glidant
 Magnesium Stearate.sup.h 16 2 Lubricant Core Tablet Weight 800 100 Film Coat Agent 24 3 Film
 Coat Total Tablet Weight 824 103

(82) TABLE-US-00007 TABLE 7 Drug Product, 300 mg (57.14% (w/w)), Qualitative and
 Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
 Adagrasib Drug Substance 300 57.14 Drug Substance Microcrystalline Cellulose 190.75 36.36
 Diluent Mannitol 0 0 Diluent Crospovidone 15.75 3 Disintegrant Colloidal Silicon Dioxide 10.5 2
 Glidant Magnesium Stearate.sup.i 8 1.5 Lubricant Core Tablet Weight 525 100 Film Coat Agent
 15.75 3 Film Coat Total Tablet Weight 540.75 103

(83) TABLE-US-00008 TABLE 8 Drug Product, 400 mg (57.14% (w/w)), Qualitative and
 Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
 Adagrasib Drug Substance 400 57.14 Drug Substance Microcrystalline Cellulose 254.5 36.36
 Diluent Mannitol 0 0 Diluent Crospovidone 21 3 Disintegrant Colloidal Silicon Dioxide 14 2
 Glidant Magnesium Stearate.sup.j 10.5 1.5 Lubricant Core Tablet Weight 700 100 Film Coat Agent
 21 3 Film Coat Total Tablet Weight 721 103

(84) TABLE-US-00009 TABLE 9 Drug Product, 600 mg (66.67% (w/w)), Qualitative and
 Quantitative Unit Composition Quantity per Tablet Weight % Component (mg) per Tablet Function
 Adagrasib Drug Substance 600 66.67 Drug Substance Microcrystalline Cellulose 241.47 26.83
 Diluent Mannitol 0 0 Diluent Crospovidone 27 3 Disintegrant Colloidal Silicon Dioxide 18 2
 Glidant Magnesium Stearate.sup.k 13.5 1.5 Lubricant Core Tablet Weight 900 100 Film Coat Agent
 27 3 Film Coat Total Tablet Weight 927 103

Example 2: Pharmacokinetics

(85) A first objective of the study was to investigate the relative bioavailability (BA) and
 pharmacokinetic (PK) profiles of MRTX849 (i.e., adagrasib) in capsule and tablet formulations
 following single-dose oral administration in healthy subjects under fasted conditions, to evaluate
 the effect of food on the PK profile of MRTX849 in tablet formulation following single-dose oral
 administration in healthy subjects, and to evaluate side effects.

(86) The secondary objective of the study was to investigate the safety and tolerability, including
 incidence of adverse events (side effects), of a single-dose oral administration of capsule and tablet
 formulations of MRTX849 in healthy subjects under fasted and fed conditions.

(87) Finally, the exploratory objectives of the study were to investigate the PK profiles of potential
 MRTX849 metabolites following single-dose oral administration of MRTX849 capsule and tablet
 formulations in healthy subjects under fasted and fed conditions.

(88) Methodology

(89) This study was a phase 1, randomized, open-label, 2×2 crossover trial in healthy subjects and
 consisted of 2 parts. In part 1, 38 subjects were enrolled and received a single oral dose of
 adagrasib 600 mg administered as capsule (Reference [R]) or tablet (Test [T]) formulations under
 fasting conditions in 1 of the 2 randomized treatment sequences (RT or TR).

(90) In part 2, 20 subjects were enrolled and received a 600 mg dose of adagrasib as three tablets of 200 mg adagrasib under fasting (R) or fed (a high-fat, high-calorie meal; T) conditions in 1 of the 2 randomized treatment sequences (RT or TR). Treatments within the sequence were separated by a washout period of 9 to 12 days. Serial pharmacokinetic (PK) blood samples were collected from pre-dose through 168 hours post-dose. Noncompartmental analysis was conducted to derive the primary PK parameters (C.sub.max, AUC.sub.last, and AUC.sub.∞). Log-transformed PK parameters were analyzed by a mixed effect model which included treatment, period, and sequence as fixed effects and subject nested within sequence as a random effect. For each PK parameter, a point estimate and its associated 90% confidence interval (CI) were constructed for the treatment difference between T and R treatments (ie, tablet vs capsule, fed vs fasted) and this difference and its 90% CI were exponentiated to obtain the ratio of geometric least-squares mean (GLSM) and its 90% CI.

Results

(91) In part 1, 35 subjects provided evaluable PK data for statistical analysis. The relative BA of the tablet formulation was approximately 89% compared to the capsule formulation. The GLSM ratios (90% CIs) for C.sub.max, AUC.sub.last, and AUC.sub.∞ were 86.47% (77.53%-96.43%), 88.82% (79.87%-98.78%), and 88.90% (79.98%-98.81%), respectively.

(92) In part 2, 15 subjects provided evaluable PK data for statistical analysis. Food increased C.sub.max and AUC of the tablet formulation by approximately 20% and 38%, respectively. The GLSM ratios (90% CIs) for C.sub.max, AUC.sub.last, and AUC.sub.∞ were 120.33% (95.27%-151.99%), 137.64% (113.86%-166.37%), and 137.53% (113.85%-166.12%), respectively. Adagrasib was generally well tolerated, with gastrointestinal (GI) disorders most frequently reported. The frequency of diarrhea, nausea, and vomiting was lower for the tablet vs capsule formulations (19.4% vs 35.1%, 13.9% vs 32.4%, and 0% vs 2.7%, respectively) and for the tablet fed vs fasted conditions (12.5% vs 47.4%, 6.3% vs 36.8%, and 0% vs 21.1%, respectively).

(93) Table 10 shows results of a bioavailability study of a single dose of 600 mg adagrasib mixed form capsules (reference) and mixed form tablets (Test 1) under Fasted Conditions.

(94) TABLE-US-00010 TABLE 10 Statistical Analysis of MRTX849 Primary Pharmacokinetic Endpoints to Assess Relative Bioavailability of a Single Dose of 600 mg MRTX849 Mixed Form Capsules (Reference) and Mixed Form Tablets (Test 1) under Fasted Conditions Within-Com-90% for subject Para- Treat- par- Ration % CV meter ment n GLSM ison Ratio Lower Upper (%)

| Parameter | R | T1 | Ratio (%) | Lower (%) | Upper (%) |
|--------------|--------------------|--------------------|-----------|-----------|-----------|
| AUC.sub.∞ | 12927.42 (h*ng/ml) | 12315.56 (h*ng/ml) | 95.27 | 84.93 | 106.86 |
| AUC.sub.last | 12886.28 (h*ng/ml) | 12272.80 (h*ng/ml) | 95.24 | 84.87 | 106.87 |
| C.sub.max | 514.13 (ng/ml) | 495.02 (ng/ml) | 96.28 | 86.59 | 107.06 |

22.9 ml) R
23.0 ml) R
21.1 R

AUC.sub.∞ = area under the plasma concentration-time curve from time zero to infinity; AUC.sub.last = area under the curve from time zero to time of last measurable concentration; CI = confidence interval; C.sub.max = maximum (peak) plasma drug concentration; GLSM = geometric least squares mean; ln = natural logarithm; n = number of subject with valid observations; R: 600 mg MRTX849 Mixed Form capsules (reference) T1: 600 mg MRTX849 Mixed Form tablets (intermediate test1) Model: ln(PK Parameter) = Sequence + Treatment(R and T1) + Random Error, with Subject fitted as a random effect. Degrees of Freedom method = KR For each model, data for the reference and test drugs were included exclusively. The ratio and corresponding confidence limits (expressed as percentages) were back-transformed from the difference and confidence limits calculated on the log scale. Only subjects with valid PK parameters for both treatments were included for the statistical analysis.

(95) Table 10 shows that the relative BA of the Mixed Form tablet formulation was approximately 5% lower compared to the Mixed Form capsule formulation. Mixed Form capsules and tablets include a mixture of Form 1 and Form 2 crystalline forms of adagrasib. Bioequivalence was demonstrated for Mixed Form tablets versus Mixed Form capsules and Form 2 capsules versus Mixed Form capsules as the 90% CI for the GLSM ratios of MRTX849 primary PK parameters

(C.sub.max, AUC.sub.last, and AUC.sub.∞) were within the regulatory BE limits of 80.00% to 125.00%. Within-subject variability was less than 25%.

(96) Table 11 shows summary PK parameters for adagrasib in plasma.

(97) TABLE-US-00011 TABLE 11 Summary of MRTX849 Pharmacokinetic Parameters After Administration of a Single Dose of 600 mg MRTX849 Mixed Form Capsules (Reference), Mixed Form Tablets (Test 1), and Form 2 Capsules (Test 2) under Fasted Conditions 600 mg MRTX849 600 mg MRTX849 600 mg MRTX849 Mixed Form Capsules, Mixed Form Tablets, Form 2 Capsules, Fasted Fasted Fasted (Reference) (Intermediate Test 1) (Intermediate Test 2) Parameter (N = 23) (N = 24) (N = 23) AUC.sub.last (h*ng/mL) 12920 (30.7) [23] 12550 (42.0) [24] 12680 (40.9) [22] AUC.sub.∞ (h*ng/mL) 12960 (30.7) [23] 12590 (41.9) [24] 12730 (40.8) [22] C.sub.max (ng/mL) 516 (28.5) [23] 502 (40.3) [24] 533 (40.5) [22] CL/F (L/h) 46.3 (30.7) [23] 47.7 (41.9) [24] 47.1 (40.8) [22] t.sub.max (h) 6.07 (6.00-12.02) [23] 6.07 (4.02-12.00) [24] 6.02 (6.00-8.02) [22] t.sub.1/2 (h) 16.9 (18.0) [23] 16.8 (18.8) [24] 16.7 (17.5) [22] V.sub.z/F (L) 1113 (33.3) [23] 1136 (39.9) [24] 1117 (44.3) [22] AUC.sub.∞ = area under the plasma concentration-time curve from time zero to infinity; AUC.sub.last = area under the concentration-time curve from time 0 to the time of the last quantifiable concentration (t.sub.last); CL/F = apparent total clearance of the drug from plasma after oral administration; C.sub.max = maximum (peak) plasma drug concentration; CV = coefficient of variation (%); n = number of subjects with valid observations; N = number of subjects; NC = not calculated; t.sub.1/2 = elimination half-life; t.sub.max = time to reach maximum (peak) plasma concentration following drug administration; V.sub.z/F = apparent volume of distribution during terminal phase after non-intravenous administration; Geometric mean (CV) [n] statistics presented; for t.sub.max, median (min-max) [n] statistics presented; for t.sub.1/2, arithmetic mean (arithmetic CV) statistics presented.

(98) Table 12 also shows summary PK parameters for adagrasib in plasma.

(99) TABLE-US-00012 TABLE 12 Summary of MRTX849 Pharmacokinetic Parameters After Administration of a Single Dose of 600 mg MRTX849 Mixed Form Capsules (Reference) and Form 2 Tablets (Test) under Fasted Conditions 600 mg MRTX849 600 mg MRTX849 Mixed Form Capsules, Form 2 Tablets, Fasted Fasted (Reference) (Test) Parameter (N = 37) (N = 36) AUC.sub.last (h*ng/mL) 14020 (58.5) [36] 12530 (62.9) [36] AUC.sub.∞ (h*ng/mL) 14070 (58.3) [36] 12590 (62.6) [36] C.sub.max (ng/mL) 556 (49.0) [36] 486 (54.6) [36] CL/F (L/h) 42.7 (58.3) [36] 47.7 (62.6) [36] t.sub.max (h) 7.50 (4.05-12.00) [36] 6.07 (4.03-12.00) [36] t.sub.1/2 (h) 17.0 (17.7) [36] 17.2 (16.9) [36] V.sub.z/F (L) 1030 (52.0) [36] 1166 (55.7) [36] AUC.sub.∞ = area under the plasma concentration-time curve from time zero to infinity; AUC.sub.last = area under the concentration-time curve from time 0 to the time of the last quantifiable concentration (t.sub.last); CL/F = apparent total clearance of the drug from plasma after oral administration; C.sub.max = maximum (peak) plasma drug concentration; CV = coefficient of variation (%); n = number of subjects with valid observations; N = number of subjects; NC = not calculated; t.sub.1/2 = elimination half-life; t.sub.max = time to reach maximum (peak) plasma concentration following drug administration; V.sub.z/F = apparent volume of distribution during terminal phase after non-intravenous administration; Geometric mean (CV) [n] statistics presented; for t.sub.max, median (min-max) [n] statistics presented; for t.sub.1/2, arithmetic mean (arithmetic CV) statistics presented.

(100) Table 13 presents statistical analysis for the assessment of the relative BA of 600 mg MRTX849 Mixed Form capsules (reference) and Form 2 tablets (test) based on the primary PK parameters of MRTX849.

(101) TABLE-US-00013 TABLE 13 Statistical Analysis of MRTX849 Primary Pharmacokinetic Endpoints to Assess Relative Bioavailability of a Single Dose of 600 mg MRTX849 Mixed Form Capsules (Reference) and Form 2 Tablets (Test) under Fasted Conditions Ratio (%) 90% CI For Within- Para- Treat- GLSM the Ratio (%) subject meter ment n GLSM (T vs R) Lower Upper CV (%) AUC.sub.∞ R 35 14154.12 (h*ng/ T 35 12582.72 88.90 79.98 98.81 26.6 ml) AUC.sub.last R

35 14102.37 (h*ng/ T 35 12526.36 88.82 79.87 98.78 26.7 ml) C.sub.max R 35 558.67 (ng/ml) T 35 483.07 86.47 77.53 96.43 27.4 AUC.sub.∞ = area under the plasma concentration-time curve from time zero to infinity; AUC.sub.last = area under the plasma concentration-time curve from time zero to time of last measurable concentration; CI = confidence interval; C.sub.max = maximum (peak) plasma drug concentration; GLSM = geometric least squares mean; ln = natural logarithm; n = number of subjects with valid observations; R: 600 mg MRTX849 Mixed Form capsules (reference) T: 600 mg MRTX849 Form 2 tablets (test) Model: ln(PK Parameter) = Sequence + Period + Treatment + Random Error, with Subject fitted as a random effect. Degrees of Freedom method = KR The ratio and corresponding confidence limits (expressed as percentages) were back-transformed from the difference and confidence limits calculated on the log scale.

(102) Results showed that relative BA of Form 2 tablets was approximately 89% compared to the Mixed Form capsules based on AUCs (Table 11). However, BE between Form 2 tablets and Mixed Form capsules was not met as the 90% CIs for the GLSM ratios for C.sub.max, AUC.sub.last, and AUC.sub.∞ were not contained within the regulatory BE limits of 80.00% to 125.00%. While the upper boundary of the 90% CI for the primary PK parameters was within the BE limit of 125.00%, the lower boundary of the 90% CI for C.sub.max, AUC.sub.last, and AUC.sub.∞ (77.53%, 79.87%, and 79.98%, respectively) was slightly below the BE limit of 80.00%.

(103) Food effect was investigated as part of the study.

(104) Eligible subjects were randomized to 1 of 2 treatment sequences. For Periods 1 and 2, eligible subjects were admitted into the clinical research unit (CRU) on Day -1 and confined to the CRU until discharge on Day 8. A single 600-mg dose of MRTX849 (Form 2 tablets) was to be administered either under fasting conditions or with a high-fat, high-calorie breakfast on Day 1 of each of Periods 1 and 2. The dose of MRTX849 was determined based on the preliminary results from Part 1. Each period included serial blood collections obtained from predose through 168 hours postdose for analysis of plasma concentrations of MRTX849 (and metabolites, if applicable). A washout period of 9 to 12 days after dose administration in Period 1 was required. A follow-up phone call was scheduled between 2 and 5 days, inclusive, after discharge from Period 2.

(105) Results showed that the high-fat, high-calorie meal increased MRTX849 Form 2 tablets' C.sub.max and AUC by approximately 20% and 38%, respectively (Table 14). The efficacy and safety profiles of MRTX849 in patients are being characterized using Mixed Form capsules under fasting conditions. Accounting for the relative BA of Form 2 tablets compared to Mixed Form capsules, Form 2 tablets under fed conditions are expected to result in approximately 4% higher C.sub.max and 22% higher AUC compared to the Mixed Form capsules under fasted conditions. Furthermore, Form 2 tablets under fed conditions showed lower between-subject variability than under fasted conditions which may result in less variability in clinical response in patients.

Therefore, the effect of food on MRTX849 Form 2 tablet is not considered clinically meaningful.

(106) TABLE-US-00014 TABLE 14 Statistical Analysis of MRTX849 Primary Pharmacokinetic Endpoints to Assess the Effect of Food on a Single Dose of 600 mg MRTX849 Form 2 Tablets

| Ratio (%) | 90% CI For Within- Treat- GLSM the Ratio (%) | subject Parameter | ment n | GLSM (T:R) |
|--------------|---|-------------------|--|--|
| Lower | Upper | CV (%) | AUC.sub.∞ | R 15 14113.65 (h*ng/ml) T 15 19409.83 137.53 113.85 166.12 29.8 |
| AUC.sub.last | R 15 14055.46 (h*ng/ml) T 15 19345.25 137.64 113.86 166.37 29.9 | C.sub.max | R 15 527.43 (ng/ml) T 15 634.67 120.33 95.27 151.99 37.2 | AUC.sub.∞ = area under the plasma concentration-time curve from time zero to infinity; AUC.sub.last = area under the plasma concentration-time curve from time zero to time of last measurable concentration; CI = confidence interval; C.sub.max = maximum (peak) plasma drug concentration; GLSM = geometric least squares mean; ln = natural logarithm; n = number of subjects with valid observations; R: 600 mg MRTX849 Form 2 tablets, fasted condition (reference) T: 600 mg MRTX849 Form 2 tablets, fed condition (test) Model: ln(PK Parameter) = Sequence + Period + Treatment + Random Error, with Subject fitted as a random effect The ratio and corresponding confidence limits (expressed as percentages) were back-transformed from the difference and confidence limits calculated on the log |

scale. Only subjects with data for both treatments (fasted and fed) are included in the analysis.

(107) The frequency of gastrointestinal disorders, in particular diarrhea, nausea, and vomiting, appeared to be lower during tablet dosing compared to during capsule dosing. Also, the frequency of gastrointestinal disorders, particularly diarrhea, nausea, abdominal pain, and vomiting, appeared to be lower when tablets were taken with food compared to fasted conditions. A summary of side effects is displayed in Table 15.

(108) TABLE-US-00015 TABLE 15 Summary of Treatment-emergent Adverse Events by System Organ Class and Preferred Term

| Part 1 | Part 2 | Part 3 | Overall MF capsules | MF tablets | F2 capsules | MF capsules | F2 tablets | F2 tablets (R) | (T1) | (T2) | (R) | (T) | (fasted; R) | (fed; T) | System Organ Class (N = 23) | (N = 24) | (N = 23) | (N = 37) | (N = 36) | (N = 19) | (N = 16) | (N = 82) | Preferred Term nS (%) | nS (%) | nS (%) | nS (%) | nS (%) | nS (%) | nS (%) | nS (%) | Overall |
|--------|--------|--------|---------------------|------------|-------------|-------------|------------|----------------|------|------|-----|-----|-------------|----------|-----------------------------|----------|----------|----------|----------|----------|----------|----------|-----------------------|--------|--------|--------|--------|--------|--------|--------|---------|
| 11 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | 7 | 5 | 13 | 7 | 9 | 2 | 37 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 |
| 15 | 9 | 15 | 21 | 8 | 10 | 4 | 53 | 11 | 8 | 12 | 20 | 8 | 10 | 3 | 48 | 10 | | | | | | | | | | | | | | | |

components.

Compression 9. Charge the final lubricated blend to a rotary tablet press and compress the blend into tablet cores. 10. Perform dedusting and metal detection of core tablet products.

In-Process Controls:

(114) Tablet appearance, weight, and hardness are monitored and controlled during the compression step.

(115) Film Coating

(116) 11. Prepare an aqueous suspension of film coat agent and purified water using a mixer and appropriate size vessel. 12. Charge the core tablets into a perforated pan coater, and coat the tablet by spraying the suspension from previous step to 3-4% weight gain.

(117) All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of the specification that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

Claims

1. A tablet formulation for oral administration in a human comprising (1) adagrasib; (2) one or more diluents selected from the group consisting of microcrystalline cellulose, mannitol, and a combination thereof; (3) crospovidone; (4) colloidal silicon dioxide; and (5) magnesium stearate; wherein a single oral dose of the tablet formulation provides a C.sub.max for adagrasib in the human that is lower as compared to a C.sub.max for adagrasib in the human provided by a single oral dose of a capsule formulation comprising the same amount of adagrasib.
2. The tablet formulation of claim 1, wherein the C.sub.max for adagrasib in the tablet formulation is about 13% lower than the C.sub.max of the capsule formulation when the human is under fasted conditions.
3. The tablet formulation of claim 1, wherein a side effect of nausea in the human is lower as compared to the side effect of nausea in humans administered the capsule formulation.
4. The tablet formulation of claim 1, wherein a side effect of vomiting in the human is lower as compared to the side effect of vomiting in humans administered the capsule formulation.
5. The tablet formulation of claim 1, wherein a side effect of diarrhea in the human is less as compared to the side effect of diarrhea in humans administered the capsule formulation.
6. The tablet formulation of claim 1, wherein a side effect of nausea in the human occurs in less than 17% of the total subject population.
7. The tablet formulation of claim 1, wherein a side effect of vomiting in the human occurs in less than 5% of the total subject population.
8. The tablet formulation of claim 1, wherein a side effect of diarrhea in the human occurs in less than 20% of the total subject population.
9. The tablet formulation of claim 1, wherein adagrasib constitutes about 30-67% of the tablet formulation; crospovidone constitutes about 2% to about 5% of the tablet formulation; colloidal silicon dioxide constitutes 0.5-2% of the tablet formulation; and magnesium stearate constitutes about 1-3% of the tablet formulation, wherein all percentages are percentages by weight.
10. The tablet formulation of claim 1, wherein adagrasib constitutes about 30-35% of the tablet formulation; the crospovidone constitutes about 2% to about 5% of the tablet formulation; the colloidal silicon dioxide constitutes up to about 0.5-1.5% of the tablet formulation; and the magnesium stearate constitutes about 1-2% of the tablet formulation, wherein all percentages are percentages by weight.

11. A tablet formulation for oral administration in a human comprising (1) adagrasib; (2) one or more diluents selected from the group consisting of microcrystalline cellulose, mannitol, and a combination thereof; (3) crospovidone; (4) colloidal silicon dioxide; and (5) magnesium stearate; wherein a single oral dose of the tablet formulation provides an $AUC_{sub.0-fwdarw.\infty}$ for adagrasib in the human that is lower as compared to the $AUC_{sub.0-fwdarw.\infty}$ for adagrasib in the human provided by a single oral dose of a capsule formulation comprising the same amount of adagrasib.
 12. The tablet formulation of claim 11, wherein the $AUC_{sub.0-fwdarw.\infty}$ for adagrasib in the tablet formulation is about 10% lower than the $AUC_{sub.0-fwdarw.\infty}$ of the capsule formulation when the human is under fasted conditions.
 13. The tablet formulation of claim 11, wherein a side effect of nausea in the human is lower as compared to the side effect of nausea in humans administered the capsule formulation.
 14. The tablet formulation of claim 11, wherein a side effect of vomiting in the human is lower as compared to the side effect of vomiting in humans administered the capsule formulation.
 15. The tablet formulation of claim 11, wherein a side effect of diarrhea in the human is less as compared to the side effect of diarrhea in humans administered the capsule formulation.
 16. The tablet formulation of claim 11, wherein a side effect of nausea in the human occurs in less than 17% of the total subject population.
 17. The tablet formulation of claim 11, wherein a side effect of vomiting in the human occurs in less than 5% of the total subject population.
 18. The tablet formulation of claim 11, wherein a side effect of diarrhea in the human occurs in less than 20% of the total subject population.
 19. The tablet formulation of claim 11, wherein adagrasib constitutes about 30-67% of the tablet formulation; the crospovidone constitutes about 2% to about 5% of the tablet formulation; the colloidal silicon dioxide constitutes 0.5-2% of the tablet formulation; and the magnesium stearate constitutes about 1-3% of the tablet formulation, wherein all percentages are percentages by weight.
 20. The tablet formulation of claim 11, wherein adagrasib constitutes about 30-35% of the tablet formulation; the crospovidone constitutes about 2% to about 5% of the tablet formulation; the colloidal silicon dioxide constitutes up to about 0.5-1.5% of the tablet formulation; and the magnesium stearate constitutes about 1-2% of the tablet formulation, wherein all percentages are percentages by weight.
 21. The tablet formulation of claim 1, wherein the one or more diluents comprise mannitol.
 22. The tablet formulation of claim 1 wherein the mannitol comprises about 0-10% of the tablet formulation by weight percentage.
 23. The tablet formulation of claim 1, wherein the one or more diluents comprise microcrystalline cellulose.
 24. The tablet formulation of claim 23, wherein the microcrystalline cellulose comprises about 20-55% of the tablet formulation by weight percentage.
 25. The tablet formulation of claim 1, further comprising a film coat.
 26. The tablet formulation of claim 11, wherein the one or more diluents comprise mannitol.
 27. The tablet formulation of claim 11, wherein the mannitol comprises about 0-10% of the tablet formulation by weight percentage.
 28. The tablet formulation of claim 11, wherein the one or more diluents comprise microcrystalline cellulose.
 29. The tablet formulation of claim 28, wherein the microcrystalline cellulose comprises about 20-55% of the tablet formulation by weight percentage.
 30. The tablet formulation of claim 11, further comprising a film coat.
-