

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent Application Publication

20250248988

Kind Code

A1

Publication Date

August 07, 2025

Inventor(s)

Pan; Ke et al.

COMBINATION USES OF NITROXOLINE AND ITS ANALOGUES WITH CHEMOTHERAPIES AND IMMUNOTHERAPIES IN THE TREATMENT OF CANCERS

Abstract

Novel combination therapies involving nitroxoline, its analogue or pharmaceutically acceptable salt thereof with at least one additional anti-cancer chemotherapy or immunotherapy agent are described. Related kits, pharmaceutical compositions and methods of production are also described.

Inventors: Pan; Ke (Chadds Ford, PA), Huang; Peng (Okayama, JP), Li; Qiang (China Medical City, CN)

Applicant: Jiangsu Yahong Meditech Co., Ltd. (Taizhou Jiangsu, CN)

Family ID: 1000008557849

Appl. No.: 19/189410

Filed: April 25, 2025

Related U.S. Application Data

parent US continuation 17656903 20220329 parent-grant-document US 12310960 child US 19189410

Publication Classification

Int. Cl.: A61K31/47 (20060101); A61K9/00 (20060101); A61K31/282 (20060101); A61K31/337 (20060101); A61K31/407 (20060101); A61K31/704 (20060101); A61K33/243 (20190101); A61K39/00 (20060101); A61K39/04 (20060101); A61K39/395 (20060101); A61P35/00 (20060101); C07K16/28 (20060101)

U.S. Cl.:

CPC **A61K31/47** (20130101); **A61K9/0019** (20130101); **A61K9/0053** (20130101);
A61K31/282 (20130101); **A61K31/337** (20130101); **A61K31/407** (20130101);
A61K31/704 (20130101); **A61K33/243** (20190101); **A61K39/04** (20130101);
A61K39/3955 (20130101); **A61P35/00** (20180101); **C07K16/2818** (20130101);
A61K2039/505 (20130101); A61K2039/54 (20130101); C07K2317/76 (20130101)

Background/Summary

CROSS REFERENCE TO RELATED APPLICATIONS [0001] This application is a Section 371 of International Application No. PCT/US2017/025388, filed Mar. 31, 2017, which published in the English Language on Oct. 5, 2017, under International Publication No. WO 2017/173278, which claims priority to U.S. Provisional Patent Application No. 62/315,774, filed on Mar. 31, 2016, the disclosures of which are hereby incorporated by reference herein in their entirety.

FIELD OF THE INVENTION

[0002] The present invention relates to a combination therapy for the treatment of cancer, particularly to combinations of nitroxoline, or its analogues or pharmaceutically acceptable salts thereof, with at least one additional anti-cancer chemotherapy or immunotherapy agent, wherein the combination therapy shows synergistic anti-cancer effects.

BACKGROUND OF THE INVENTION

[0003] Combination therapies for the treatment of cancers serve to minimize the chances of the cancer developing resistance to any one anti-cancer therapy, and they allow for the use of lower doses of the individual anti-cancer therapies, resulting in reduced toxicities. Depending on the nature of drug-drug interactions, the combination of two active ingredients may result in a synergistic effect, additive effect or antagonistic effect.

[0004] Nitroxoline (NIT) is an antimicrobial agent which has been commercially marketed for a long time for the treatment of urinary tract infections. It was recently discovered that nitroxoline is also active in inhibiting angiogenesis (4) and inhibiting the growth and invasion of cancer (5, 6). NIT and its analogues, such as oxyquinoline, clioquinol and iodoquinol, share a common clinical use in treating infectious diseases. NIT analogues were also reported to have anti-cancer cytotoxicities (7). However, to the knowledge of the inventors, none of these drugs have been investigated in combination with other anti-cancer drugs, and drug-drug interaction of NIT or its analogues with any of the other cancer therapies has not been reported.

[0005] There exists a need in the art for novel combination therapies to treat cancers with enhanced anti-cancer effects or reduced toxicities.

BRIEF SUMMARY OF THE INVENTION

[0006] The present invention satisfies these needs by providing novel combination therapies involving nitroxoline, its analogues or pharmaceutically acceptable salts thereof with at least one additional anti-cancer chemotherapy or immunotherapy agent. The combination therapies demonstrate synergistic effects on the inhibition of cancer cell growth.

[0007] In one general aspect, the present invention provides a method for treating cancer by administering to a subject in need thereof nitroxoline, its analogue, or a pharmaceutically acceptable salt thereof, in combination with a chemotherapy. The chemotherapy is preferably selected from the group consisting of a microtubule disassembly inhibitor, an agent that cross-links DNA, and a platinum compound.

[0008] In another general aspect, the present invention provides a method for treating cancer by administering to a subject in need thereof nitroxoline, its analogue, or a pharmaceutically acceptable salt thereof, in combination with an immunotherapy agent. The immunotherapy agent is preferably an agent that can stimulate an effective immune response and inhibit immune-

suppression. More preferably, the immunotherapy agent is an inhibitor or modulator of regulatory T cells or myeloid-derived suppressor cells.

[0009] Other aspects, features and advantages of the invention will be apparent from the following disclosure, including the detailed description of the invention and its preferred embodiments and the appended claims.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] The foregoing summary, as well as the following detailed description of the invention, will be better understood when read in conjunction with the appended drawings. For the purpose of illustrating the invention, they are shown in the drawings embodiments which are presently preferred. It should be understood, however, that the invention is not limited to the precise arrangements and instrumentalities shown. In the drawings:

[0011] FIG. 1 shows the structures of nitroxoline (NIT) and examples of its analogues;

[0012] FIG. 2 shows combination index (CI) plots of the inhibition of the growth of human bladder cancer cell line 5637 by the combination of NIT and carboplatin;

[0013] FIG. 3 shows CI plots of the inhibition of the growth of human bladder cancer cell line 5637 by the combination of NIT and paclitaxel;

[0014] FIG. 4 shows CI plots of the inhibition of the growth of human liver cancer cell line HepG2 by the combination of NIT and carboplatin;

[0015] FIG. 5 shows CI plots of the inhibition of the growth of human liver cancer cell line HepG2 by the combination of NIT and paclitaxel;

[0016] FIG. 6 shows CI plots of the inhibition of the growth of human bladder cancer cell line 5637 by the combination of NIT and mitomycin C;

[0017] FIG. 7 shows CI plots of the inhibition of the growth of human bladder cancer cell line 5637 by the combination of NIT and epirubicin;

[0018] FIG. 8 shows CI plots of the inhibition of the growth of human bladder cancer cell line 5637 by the combination of NIT and pirarubicin;

[0019] FIG. 9 shows the inhibition of tumor growth by the combination of oral NIT and mitomycin C in an orthotopic mouse MBT-2-Luc bladder cancer model;

[0020] FIG. 10 shows the inhibition of tumor growth by the combination of oral NIT and intravesical BCG in an orthotopic mouse MBT-2-Luc bladder cancer model;

[0021] FIG. 11 shows the inhibition of tumor growth by the combination of oral NIT and anti-PD-1 antibody in an orthotopic mouse MBT-2-Luc bladder cancer model; and

[0022] FIG. 12 shows the inhibition of tumor growth by the combination of oral NIT and anti-PD-1 antibody in an orthotopic mouse RM-9-Luc prostate cancer model.

DETAILED DESCRIPTION OF THE INVENTION

[0023] Various publications, articles and patents are cited or described in the background and throughout the specification; each of these references is herein incorporated by reference in its entirety. Discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is for the purpose of providing context for the invention. Such discussion is not an admission that any or all of these matters form part of the prior art with respect to any inventions disclosed or claimed.

Definitions

[0024] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention pertains. Otherwise, certain terms used herein have the meanings as set in the specification. All patents, published patent applications and publications cited herein are incorporated by reference as

if set forth fully herein. It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise.

[0025] Unless otherwise stated, any numerical value, such as a concentration or a concentration range described herein, are to be understood as being modified in all instances by the term “about.” Thus, a numerical value typically includes $\pm 10\%$ of the recited value. For example, a concentration of 1 mg/mL includes 0.9 mg/mL to 1.1 mg/mL. Likewise, a concentration range of 1% to 10% (w/v) includes 0.9% (w/v) to 11% (w/v). As used herein, the use of a numerical range expressly includes all possible subranges, all individual numerical values within that range, including integers within such ranges and fractions of the values unless the context clearly indicates otherwise.

[0026] As used herein, the terms “treat,” “treating,” and “treatment” are all intended to refer to an amelioration or reversal of at least one measurable physical parameter related to a cancer which is not necessarily discernible in the subject, but can be discernible in the subject. The terms “treat,” “treating,” and “treatment,” can also refer to causing regression, preventing the progression, or at least slowing down the progression of the cancer. In a particular embodiment, “treat,” “treating,” and “treatment” refer to an alleviation, prevention of the development or onset, or reduction in the duration of one or more symptoms associated with the cancer. In a particular embodiment, “treat,” “treating,” and “treatment” refer to prevention of the recurrence of the cancer. In a particular embodiment, “treat,” “treating,” and “treatment” refer to an increase in the survival of a subject having the cancer. In a particular embodiment, “treat,” “treating,” and “treatment” refer to elimination of the cancer in the subject.

[0027] As used herein, the term “subject” refers to an animal, and preferably a mammal. According to particular embodiments, the subject is a mammal including a non-primate (e.g., a camel, donkey, zebra, cow, pig, horse, goat, sheep, cat, dog, rat, rabbit, guinea pig or mouse) or a primate (e.g., a monkey, chimpanzee, or human). In particular embodiments, the subject is a human.

[0028] As used herein, the term “effective amount” or “therapeutically effective amount” refers to an amount of an active ingredient or component that elicits the desired biological or medicinal response in a subject. In particular embodiments, the effective amount is the amount of an active ingredient or compound that is effective to achieve a synergistic effect with another active ingredient or compound. As used herein, a “synergistic effect” refers to an effect which is greater than the additive effect of the two individual active ingredients or compounds on the treatment of cancer. A therapeutically effective amount can be determined empirically and in a routine manner, in relation to the stated purpose. For example, in vitro assays can optionally be employed to help identify optimal dosage ranges. Selection of a particular effective dose can be determined (e.g., via clinical trials) by those skilled in the art based upon the consideration of several factors, including the disease to be treated or prevented, the symptoms involved, the patient's body mass, the patient's immune status and other factors known by the skilled artisan. The precise dose to be employed in the formulation will also depend on the route of administration, and the severity of disease, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses can be extrapolated from dose-response curves derived from in vitro or animal model test systems. For example, drug combinations can be evaluated using a mathematical algorithm based on Loewe's model of additivity. In this model, combination index (CI) values are calculated. $CI < 1$, $CI = 1$, and $CI > 1$ indicate synergistic, additive and antagonistic interactions, respectively.

[0029] As used herein, the term “pharmaceutically acceptable salt” refers to a salt form of nitroxoline or one of its analogues that is safe and effective. Non-limiting examples of pharmaceutically acceptable salts include: hydrochloride, hydrobromide, hydroiodide, sulfate, bisulfate, citrate, acetate, succinate, ascorbate, oxalate, nitrate, pears salts, hydrogen phosphate, dihydrogen phosphate, salicylate, hydrogen citrate, tartrate, maleate, fumarate, formate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, and p-toluenesulfonate. Preferably, a

pharmaceutically acceptable salt of nitroxoline is a base addition salt of nitroxoline, such as those described in International Patent Application No. PCT/US14/30532, the entire content of which is incorporated herein by reference.

[0030] As used herein, the term “chemotherapy agent” refers to any chemical substance that is an anti-cancer drug.

[0031] As used herein, the term “immunologic agent” or “immunologic modulator” refers to any agent that is capable of stimulating an immune response and/or inhibiting immune-suppression.

[0032] As used herein, the term “in combination,” in the context of the administration of two or more therapies to a subject, refers to the use of more than one therapy. The use of the term “in combination” does not restrict the order in which therapies are administered to a subject. For example, a first therapy (e.g., an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof described herein) can be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 16 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 16 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy (e.g., an effective amount of an immunotherapy agent or an effective amount of a chemotherapeutic agent) to a subject.

[0033] As used herein, the term “carrier” refers to any excipient, diluent, filler, salt, buffer, stabilizer, solubilizer, oil, lipid, lipid containing vesicle, microsphere, liposomal encapsulation, or other material well known in the art for use in pharmaceutical formulations. It will be understood that the characteristics of the carrier, excipient or diluent will depend on the route of administration for a particular application. As used herein, the term “pharmaceutically acceptable carrier” refers to a non-toxic material that does not interfere with the effectiveness of a composition according to the invention or the biological activity of a composition according to the invention. According to particular embodiments, in view of the present disclosure, any pharmaceutically acceptable carrier suitable for use in a nitroxoline-based pharmaceutical composition can be used in the invention. Non-limiting examples of carriers include saline and water.

[0034] Nitroxoline (NIT) and its analogues

[0035] As shown in FIG. 1, NIT and its analogues, such as oxyquinoline, clioquinol and iodoquinol, share a common clinical use in treating infectious diseases. NIT is a broad-spectrum antibiotic, and it has been marketed to treat urinary tract infections since the 1960s. Its mechanism of action in inhibiting bacterial growth remains uncharacterized. NIT was found to inhibit adherence of uropathogenic *E. coli* to uroepithelial cells and urinary catheters at sub-MICs of NIT, and it is thought that NIT promotes disorganization of the bacterial outer membrane by chelating the divalent ions Mg^{2+} and Mn^{2+} (1). Oxyquinoline (Oxine) is an antiseptic with mild fungistatic, bacteriostatic, anthelmintic, and amebicidal activities. Oxine is capable of inhibiting isolated *E. coli* RNA polymerase without directly contacting the enzyme by essentially depriving the enzyme of Mn^{2+} and Mg^{2+} (2). Oxine is also used as a metal chelating agent, and so as a carrier for radio-indium for diagnostic purposes. Clioquinol has long been used as a topical anti-infective, intestinal antiamebic, and vaginal trichomonacide. The oral preparation of clioquinol has been shown to cause subacute myelo-optic neuropathy and has thus been banned worldwide. Clioquinol was found to halt cognitive decline in Alzheimer's disease in animal models, possibly due to its ability to act as a chelator of Cu^{2+} and Zn^{2+} (3). Iodoquinol is another halogenated 8-quinolinol widely used as an intestinal antiseptic, especially as an antiamebic agent. It is also used topically for other infections and may cause CNS and eye damage. Its mechanism of action remains unknown.

[0036] NIT and its analogues have been shown to have anti-cancer activities. It was shown that NIT is a MetAP2 inhibitor and can therefore block cancer angiogenesis by inhibiting proliferation

of human umbilical vein endothelial cells (HUVECs) (4). NIT was also shown to be able to induce apoptosis of prostate cancer cells (5), and its cytotoxicity against cancer cells was more pronounced when administered in combination with Cu.sup.2+ (6). NIT analogues, such as clioquinol, oxine and iodoquinol, were also reported to have anti-cancer cytotoxicities (7).

Chemotherapies

[0037] Cancer is the uncontrolled growth of cells coupled with malignant behavior including invasion and metastasis, among other features. It is caused by the interaction between genetic susceptibility and environmental factors. These factors lead to accumulations of genetic mutations in oncogenes, genes that control the growth rate of cells, and tumor suppressor genes, genes that help to prevent cancer, giving cancer cells their malignant characteristics, such as uncontrolled growth. Chemotherapy is a category of cancer treatment that uses chemical substances, especially one or more anti-cancer drugs. In the broad sense, most chemotherapeutic drugs work by impairing mitosis (cell division), effectively targeting fast-dividing cells. As these drugs cause damage to cells, they are termed cytotoxic. Anti-cancer drugs prevent mitosis by various mechanisms including damaging DNA and inhibition of the cellular machinery involved in cell division. One theory as to why these drugs kill cancer cells is that they induce a programmed form of cell death known as apoptosis. There are many types of cytotoxic chemotherapies in clinical use for treating cancer, and they are grouped into different classes according to the chemical structural characteristics. See, for example, Table 1 below.

TABLE-US-00001 TABLE 1 List of anti-cancer chemotherapies

Class	Mechanism of Action	Drug Names
Nucleoside	DNA methyltransferase inhibitor and incorporates decitabine, floxuridine itself into RNA, hence inhibiting gene expression thymidylate synthase	capecitabine, carmofur, inhibitor fluorouracil, tegafur
ribonucleotide reductase and		clofarabine
DNA polymerase inhibitor	DNA polymerase inhibitor, cytarabine, fludarabine	S-phase specific; incorporates its metabolites into DNA
DNA synthesis inhibitor,	gemcitabine induces apoptosis specifically in S-phase	purine synthesis inhibitor mercaptopurine, nelarabine, tioguanine
adenosine deaminase	pentostatin inhibitor	Anti- dihydrofolate reductase methotrexate, pemetrexed, metabolites inhibitors pralatrexate, raltitrexed
Inhibits DNA synthesis by	hydroxycarbamide inhibiting the enzyme	ribonucleotide reductase
Topoisomerase	topoisomerase inhibitors	irinotecan, topotecan
I inhibitors Anthracyclines	inhibits DNA and RNA synthesis by intercalating	epirubicin, idarubicin, DNA base pairs; inhibits mitoxantrone, valrubicin
DNA repair by inhibiting	topoisomerase II	Podophyllo- topoisomerase II inhibitor etoposide, teniposide
toxins Taxanes	Microtubule disassembly	cabazitaxel, docetaxel, inhibitor. Arrests cells in paclitaxel, and taxanes
late G2 phase and M phase.	mixed or conjugated with proteins or antibodies	Vinca microtubule assembly
vinblastine, vincristine,	alkaloids inhibitor, arrests cells	vindesine, vinflunine, in M phase
vinorelbine	Alkylating alkylates DNA	bendamustine, busulfan, agents
carmustine, chlorambucil,	chlormethine, cyclophosphamide, dacarbazine, Fotemustine, Ifosfamide, lomustine, melphalan, streptozotocin, temozolomide, trabectedin	Platinum reacts with DNA, inducing carboplatin, cisplatin, compounds
apoptosis, non-cell cycle	nedaplatin, oxaliplatin	specific
Miscellaneous	unclear, reactive	altretamine others intermediates covalently bind to microsomal proteins and DNA, possibly causing DNA damage
inhibits DNA and to a lesser	bleomycin extent	RNA synthesis, produces single and double strand breaks in DNA possibly by free radical formation
proteasome inhibitor	bortezomib, carfilzomib	complexes with DNA
dactinomycin	interfering with DNA- dependent	RNA synthesis
antimicrotubule and	estramustine	estrogenic actions promotes tubulin
ixabepilone	polymerisation and stabilises microtubular function, causing cell cycle arrest at G2/M phase and subsequently induces apoptosis	cross-links DNA
mitomycin C	inhibits DNA, RNA and	procarbazine protein synthesis

[0038] Traditional chemotherapeutic agents are cytotoxic, and they act by killing cells that divide rapidly, which is one of the fundamental properties of most cancer cells. This targeting of rapidly-dividing cells means that chemotherapy also harms cells that divide rapidly under normal

circumstances, such as cells in the bone marrow, the digestive tract, and in hair follicles, resulting in the most common side-effects of chemotherapy, including myelosuppression (decreased production of blood cells, hence also immunosuppression), mucositis (inflammation of the lining of the digestive tract), and alopecia (hair loss). Combination therapy involves treating a patient with a number of different drugs simultaneously. Because the drugs differ in their mechanism and side-effects, a combination therapy may minimize the chances of resistance developing to any one agent, thus can be used to treat resistant cancers or at lower doses to reduce toxicity. However, because of the unpredictable nature of drug-drug interactions, a combination of known treatments may also result in reduced efficacy or increased side effects.

Cancer Immunotherapies

[0039] Cancer immunotherapy attempts to stimulate the immune system to reject and destroy tumors. Tumors have been found to be able to escape host immunity by manipulating the tumor microenvironment and driving immunosuppression, such that although the components necessary for mounting an effective anti-tumor immune response are present in patients with cancer, the host usually fails to arrest tumor progression (8). Numerous molecular and cellular mechanisms have been proposed in the past to explain this counterintuitive scenario, largely focusing on the “immune escape” of tumors and suggesting that tumors develop the capability to avoid tumor-specific immune responses generated by the host, or they altogether disable the host anti-tumor immunity. This process, referred to as “tumor-induced immune suppression” has been recognized in recent years and is being intensively investigated. It appears that tumors can interfere with all components of the immune system, affecting all stages of the anti-tumor immune response.

[0040] Molecular alterations that occur in tumor cells as the tumor progresses from the pre-malignant to metastatic phenotype are a result of genetic instability, now recognized as a principal characteristic of all tumors. Genetic changes are detectable during early stages of tumorigenesis, become more pronounced as the tumor progresses, are the greatest in metastatic cells and are responsible for tumor heterogeneity and alterations in the antigenic epitope profile of tumor cells. It has been suggested that the immune system might drive these antigenic changes via “immune editing” by eliminating those malignant cells that are sensitive to immune intervention and allowing for the selection and survival of immune-resistant variants. The net result of immune editing is that the tumor escapes from the host immune system. Regulatory T cells (Treg) and myeloid-derived suppressor cells (MDSC) are two types of cells used by the tumor to execute an immune escape. Protection of immune cells from the adverse effects of Treg, MDSC or inhibitory factors, and thus enhancement of their effector functions, can restore effective anti-tumor immunity in patients with cancer.

[0041] CD4^{sup}.+CD25^{sup}.highFOXP3^{sup}.+Treg cells accumulate in human tumors and in the peripheral circulation of patients with cancer. It is unclear whether these cells migrate to tumors or expand in situ. Because tumor-associated antigens (TAA) are self-antigens, it is possible that Treg accumulation is a response to enforce immune tolerance. Treg cells contribute to the down-regulation of immune activity of effector T cells by a variety of mechanisms including IL-10 and TGF- β 1 production, enzymatic degradation of ATP to immunosuppressive adenosine, or the engagement of the Fas/FasL and granzyme/perforin pathways. Tumors benefit from immunosuppressive effects mediated by Treg.

[0042] Bone marrow-derived immature myeloid cells MDSC

(CD34^{sup}.+CD33^{sup}.+CD13^{sup}.+CD15^{sup}.-) are present at an elevated frequency in the peripheral circulation and tumors of nearly all cancer patients. They are recruited by tumor-derived soluble factors such as TGF- β 1, IL-10, VEGF, GM CSF, IL-6, PGE2. They promote tumor growth by suppressing T-cell responses via several mechanisms, including production of arginase-1, an enzyme involved in L-arginine metabolism, as well as activation of inducible nitric oxide synthase (iNOS). They also control the tumor's production of indoleamine-2,2-dioxygenase (IDO), which is involved in the catabolism of tryptophan, an amino acid essential for T-cell differentiation.

[0043] Bacillus Calmette Guerin (BCG) immunotherapy for early stage (non-invasive) bladder cancer utilizes instillation of attenuated live bacteria into the bladder, and is effective in preventing recurrence in up to two thirds of cases. The immune response to BCG can be summarized as follows (9): infection of urothelial and bladder tumor cells by BCG results in internalization of BCG, which increases the expression of antigen-presenting molecules. This induces an immune response via the release of cytokines, such as Th1 cytokines (IL-2, tumor necrosis factor, IL-12, and IFN- γ) and Th2 cytokines (IL-4, IL-5, IL-6, and IL-10) along with IL-8 and IL-17. This complex immune cascade induces antitumor activity mediated by cytotoxic T lymphocytes, natural killer cells, neutrophils, and macrophages. In another study of BCG in an orthotopic mouse bladder tumor model, the population of MDSCs was significantly downregulated following the high-dose BCG therapy compared with the low-dose therapy. In the same study, CD4^{sup.}/Foxp3^{sup.}+Tregs also exhibited the same change. Following the BCG treatment, the population of CD4^{sup.}/Foxp3^{sup.}+ Tregs was decreased in the blood. These inhibitory effects on the immune-suppressive factors may explain the robust antitumor therapeutic effects of the BCG therapy (10).

[0044] Immune checkpoint regulators, which can be both costimulatory and coinhibitory molecules, regulate the immune system. The balance between these checkpoints signals regulates lymphocyte activation and consequently the immune response. Tumors can use these checkpoint regulators to protect themselves from the immune system. Immune checkpoint therapies can enhance the proliferation, migration, persistence, and/or cytotoxic activity of T cells in a subject and, in particular, by increasing the numbers of tumor infiltrating T cells (11). The best characterized immune checkpoint receptors are cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4; also known as CD152), programmed cell death protein 1 (PD-1; also known as CD 279), and indoleamine 2,3-dioxygenase (IDO), and agents targeting these molecules are either approved or being extensively tested in clinical trials for treatment of multiple solid or hematological cancers.

[0045] CTLA-4, an inhibitory receptor, is a global immune checkpoint regulator engaged in priming immune responses via down-regulating the initial stages of T-cell activation. CTLA-4 was the first clinically validated checkpoint pathway target. CTLA-4 is homologous to the T-cell costimulatory protein CD28, and both molecules bind to CD80 and CD86 on antigen-presenting cells. CTLA-4 binds CD80 and CD86 with a markedly higher affinity and avidity than CD28 does, enabling CTLA-4 to outcompete CD 28 for its ligands, resulting in an effective inhibition of T cell activation (12). Blocking CTLA-4 activity by using an antagonistic antibody interferes with this mechanism and thus preserves the activity of the T cells. Currently, Bristol-Myers Squibb's anti-CTLA-4 mAb, ipilimumab, has received US Food and Drug Administration (FDA) approval for patients with metastatic melanoma. Additionally, AstraZeneca's anti-CTLA-4 mAb, tremelimumab, has been granted orphan drug designation by the FDA for the treatment of patients with malignant mesothelioma.

[0046] PD-1 is another inhibitory receptor expressed on activated T and B cells, and it functions to dampen the immune response (13). PD-1 acts as an immune checkpoint regulator, and upon binding of one of its ligands, PD-L1 (B7-H1, CD274) or PD-L2 (B7-DC, CD273), PD-1 inhibits proliferation and cytokine production of T cells. Overexpression of PD-L1 or PD-L2 in the tumor microenvironment leads to the inhibition of the intratumoral immune responses (14). Inhibition of the interaction between PD-1 and PD-L1 by anti-PD-1/PD-L1 antibodies can inhibit the deactivation of T cells, and thus enhance anti-tumor responses, delay tumor growth, and facilitate tumor rejection (15). Two anti-PD-1 mAbs, Bristol-Myers Squibb's nivolumab, and Merck's pembrolizumab, have received US FDA approval for patients with metastatic melanoma and non-small-cell lung cancer. Recently, the FDA approved nivolumab as a treatment for patients with metastatic renal cell carcinoma. In 2016, Roche's anti-PD-L1 mAb, atezolizumab, was approved by the FDA for treating metastatic urothelial cancer and non-small cell lung cancer. Other anti-PD-L1

mAbs, such as AstraZeneca's durvalumab and Pfizer's avelumab, are in late-stage clinical trials. [0047] Indoleamine 2,3-dioxygenase (IDO) is an enzyme that catalyzes the oxidative cleavage of tryptophan (16). IDO plays a role in suppressing the immune system because T cells undergoing antigen-dependent activation require tryptophan for cell proliferation and survival (17). IDO is overexpressed in most tumors and/or tumor-draining lymph nodes and it plays a significant role in helping tumors to evade attack from the immune system. IDO inhibitors block the IDO enzyme which reduces the depletion of tryptophan and ultimately may help to promote an enhanced immune response against the tumor (18). Currently, a number of clinical trials are underway to evaluate IDO inhibitors for both monotherapy and combination cancer therapies.

[0048] In addition to PD-1, CTLA-4, and IDO, other immune checkpoints are also involved in the occurrence and development of malignant tumors, including T cell membrane protein-3 (TIM-3), LAG3, T-cell immunoreceptor with Ig and immunoreceptor tyrosine-based inhibitory motif (ITIM) domains (TIGIT), BTLA, inducible T-cell costimulator (ICOS), killer inhibitory receptors (KIR), and V-domain Ig-containing suppressor of T cell activation (VISTA). Similar to PD-1, CTLA-4 and IDO, these immune checkpoints inhibit lymphocyte activity and/or induce lymphocyte anergy, and thus are ideal targets for cancer immunotherapy. Blocking antibodies for these immune checkpoints have shown specific anti-tumor activities in animal models, and some are being tested in clinical trials.

Methods of Treatment

[0049] In some aspects, the invention involves methods of combination therapy with an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of at least one second agent including any chemotherapeutic agents or immunotherapy agents for treating various types of cancer. The present invention provides methods for combining an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, with an effective amount of an immunotherapy agent or a chemotherapeutic agent to produce significant enhancement of the anti-cancer effect, which is preferably synergistic.

[0050] The cancer can be any cancer. In some embodiments, the cancer is melanoma, cervical cancer, breast cancer, ovarian cancer, prostate cancer, testicular cancer, urothelial carcinoma, bladder cancer, non-small cell lung cancer, small cell lung cancer, sarcoma, colorectal adenocarcinoma, gastrointestinal stromal tumors, gastroesophageal carcinoma, colorectal cancer, pancreatic cancer, kidney cancer, hepatocellular cancer, malignant mesothelioma, leukemia, lymphoma, myelodysplastic syndrome, multiple myeloma, transitional cell carcinoma, neuroblastoma, plasma cell neoplasms, Wilm's tumor, glioblastoma, retinoblastoma, or hepatocellular carcinoma. In particular embodiments, the cancer is bladder cancer, prostate cancer, kidney cancer, urothelial carcinoma, testicular cancer, non-small cell lung cancer, breast cancer, or hepatocellular carcinoma. In more particular embodiments, the cancer is bladder cancer, prostate cancer, kidney cancer or urothelial carcinoma. In more particular embodiments, the cancer is bladder cancer or prostate cancer.

[0051] The second agent can be any chemotherapeutic agent that enhances the anti-cancer effect of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof. In some embodiments, the second agent is a microtubule disassembly inhibitor, an agent that cross-links DNA, an agent that intercalates DNA base pairs, or a platinum compound. In some embodiments, the second agent is a microtubule disassembly inhibitor, such as a taxane, such as paclitaxel, docetaxel or cabazitaxel, or a taxane mixed or conjugated with proteins or antibodies. In some embodiments, the second agent is an agent that cross-links DNA, such as mitomycin C. In some embodiments, the second agent is an agent that intercalates DNA base pairs, such as an anthracycline, such as daunorubicin, doxorubicin, epirubicin, pirarubicin, idarubicin, mitoxantrone, or valrubicin. In some embodiments, the second agent is a platinum compound, such as carboplatin, cisplatin, nedaplatin or oxaliplatin. In particular embodiments, the second agent is selected from the group consisting of paclitaxel, mitomycin C, epirubicin, pirarubicin, cisplatin and carboplatin.

[0052] The second agent can also be any immunotherapy agent that enhances the anti-cancer effect of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof. In some embodiments, the second agent is an immunologic agent that can stimulate an effective immune response and/or inhibit immune-suppression. In some embodiments, the second inhibitor is an agent that modulates, particularly inhibits and downregulates immune-suppressive factors such as regulatory T cells and M DSCs, including any inhibitors or antibodies of the programmed cell death 1 ligand 1 (PD-L1)/programmed cell death protein 1 (PD-1) pathway. In some embodiments, the second inhibitors are any effective inhibitors/antibodies capable of modulating immunocytes activities including but not limited to targeting cytotoxic T lymphocyte-associated antigen 4 (CTLA-4), CD20, CD19, IL-17a, CD25, arginase 1 (ARG1), indoleamine-2,3-dioxygenase (IDO), or tryptophan 2,3-dioxygenase (TDO2). In some embodiments, the second inhibitor is BCG. In particular embodiments, the second agent is an antibody against PD-1 or an antigen binding fragment thereof. In other particular embodiments, the second agent is an antibody against PD-L1 or an antigen binding fragment thereof. In other particular embodiments, the second agent is an antibody against CTLA-4 or an antigen binding fragment thereof. In other particular embodiments, the second agent is a BCG therapy, preferably an attenuated BCG therapy, more preferably mycobacterial cell wall fragments, and most preferably mycobacterial cell wall fragments with biologically active nucleic acids derived from *Mycobacterium phlei*.

[0053] The dosage of the compounds or agents are selected, e.g., based on required dose, fluid volumes, viscosities, etc., according to the particular mode of administration selected, to preferably achieve a synergistic effect. In some embodiments, the nitroxoline or its analogue or pharmaceutically acceptable salt thereof, is administered, preferably orally, in a dosage of about 100 mg/day to about 1600 mg/day, such as about 100 mg/day, 200 mg/day, 300 mg/day, 400 mg/day, 500 mg/day, 600 mg/day, 700 mg/day, 800 mg/day, 900 mg/day, 1000 mg/day, 1100 mg/day, 1200 mg/day, 1300 mg/day, 1400 mg/day, 1500 mg/day and 1600 mg/day. Preferably, the nitroxoline or its analogue or pharmaceutically acceptable salt thereof, is administered, preferably orally, in a dosage of about 600 mg/day to about 1600 mg/day.

[0054] In some embodiments, the second agent is an anti-PD-1 antibody or an antigen binding fragment thereof, preferably in a dosage of about 0.1 mg/kg to about 20 mg/kg, such as about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 mg/kg, administered, preferably by intravenous or intramuscular injection, once every 2, 3 or 4 weeks. More preferably, the anti-PD-1 antibody or antigen binding fragment thereof is administered, preferably by intravenous or intramuscular injection, in a total dosage of about 2 mg/kg to about 15 mg/kg in a period of about 3 weeks, and the treatment is optionally repeated one or more times.

[0055] In some embodiments, the second agent is an anti-PD-L1 antibody or an antigen binding fragment thereof, preferably in a dosage of about 1 mg/kg to about 40 mg/kg, such as about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 mg/kg, administered, preferably by intravenous or intramuscular injection, once every 2, 3 or 4 weeks. More preferably, the anti-PD-L1 antibody or antigen binding fragment thereof is administered, preferably by intravenous or intramuscular injection, in a total dosage of about 10 mg/kg to about 30 mg/kg in a period of 4 weeks, and the treatment is optionally repeated one or more times.

[0056] In some embodiments, the second agent is an anti-CTLA-4 antibody or an antigen binding fragment thereof, preferably in a dosage of about 0.1 mg/kg to about 6mg/kg, such as about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5 or 6 mg/kg administered, preferably by intravenous or intramuscular injection, once every 3, 4 or 5 weeks. More preferably, the anti-CTLA-4 antibody or antigen binding fragment thereof is administered, preferably by intravenous or intramuscular injection, in a total dosage of about 1 mg/kg to about 4 mg/kg in a period of about 4 weeks, and the treatment is optionally repeated one or more times.

[0057] In some embodiments, the second agent is a BCG therapy, preferably in a dosage of about 0.5×10^8 to about 50×10^8 colony forming units (CFUs), such as about $0.5 \times$, $1 \times$, $2 \times$, $4 \times$, $6 \times$, $8 \times$, $10 \times$, $12 \times$, $14 \times$, $16 \times$, $18 \times$, $20 \times$, $22 \times$, $24 \times$, $26 \times$, $28 \times$, $30 \times$, $32 \times$, $34 \times$, $36 \times$, $38 \times$, $40 \times$, $42 \times$, $44 \times$, $46 \times$, $48 \times$ or 50×10^8 CFUs, administered, preferably intravesically, once every 1, 2, 3, 4, or 5 weeks. More preferably, the BCG therapy is administered, preferably intravesically, in a dose of about 1×10^8 to about 8×10^8 CFUs once a week, and the treatment is optionally repeated one or more times.

[0058] In some embodiments, the second agent is a chemotherapeutic agent administered, preferably intravesically, at a concentration of about 0.1 mg/ml or more, up to the agent's highest solubility in water or saline. In some embodiments, the chemotherapeutic agent is a DNA cross-linking or intercalating agent, such as mitomycin C, an anthracycline (e.g., epirubicin, pirarubicin, etc.), a platinum-based antineoplastic agents (e.g., carboplatin, cisplatin, oxaliplatin, nedaplatin), at a concentration of about 0.1 mg/ml to about 5 mg/ml, such as about 0.1, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5 or 5 mg/ml, administered, preferably intravesically, in a bladder instillation fluid, once every 1, 2, 3, 4 or 5 weeks. More preferably, the chemotherapeutic agent is administered, preferably intravesically, once a week at a concentration of about 0.5 mg/mL to 2 mg/mL in a bladder instillation fluid, and the treatment is optionally repeated one or more times.

[0059] In some embodiments, the chemotherapeutic agent is a taxane drug that targets tubulin, such as paclitaxel, docetaxel, cabazitaxel, a conjugate thereof with a protein or antibody, etc., at a concentration of about 0.5 mg/ml to about 10 mg/ml of the taxane drug, such as about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/ml, administered, preferably intravesically, in a bladder instillation fluid, once every 1, 2, 3, 4 or 5 weeks. More preferably, the taxane drug is administered, preferably intravesically, once a week at a concentration of about 1 mg/mL to 5 mg/mL in a bladder instillation fluid, and the treatment is optionally repeated one or more times.

[0060] The compounds or agents can be administered by any acceptable route. In some embodiments, the compounds are administered orally, intraadiposally, intraarterially, intraarticularly, intracranially, intradermally, intralesionally, intramuscularly, intranasally, intraocularly, intrapericardially, intraperitoneally, intrapleurally, intraprostatically, intrarectally, intrathecally, intratracheally, intratumorally, intraumbilically, intravaginally, intravenously, intravesically, intravitreally, liposomally, locally, mucosally, orally, parenterally, rectally, subconjunctival, subcutaneously, sublingually, topically, transbuccally, transdermally, vaginally, in creams, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof. In some embodiments, the nitroxoline or its analogue or pharmaceutically acceptable salt thereof, is administered orally or by injections, and is preferably administered orally. In some embodiments, the immunotherapy agent is an antibody that is administered by injections. In some embodiments, the immunotherapy agent is a BCG therapy that is administered intravesically or by injections, and preferably is administered intravesically. In some embodiments, the chemotherapeutic agent is administered intravesically or by injections, and preferably is administered intravesically.

[0061] In some embodiments, the compounds or agents are administered once daily. In other embodiments, the compounds or agents are administered twice daily. In other embodiments, the compounds are administered multiple times a day, once every two days, once every three days, once every four days, once every five days, once every six days, once every seven days, once every two weeks, once every three weeks, once every four weeks, once every two months, once every three months, once every four months, once every five months, once every six months, or once per year. The dosage regimen of the nitroxoline or its analogue, or a pharmaceutically acceptable salt thereof can be different from that of the immunotherapy agent or of the chemotherapeutic agent.

[0062] The compounds or agents can be administered for one day, two days, three days, four days,

five days, six days, seven days, two weeks, three weeks, four weeks, two months, three months, four months, five months, six months, one year, two years three years, four years, five years, ten years, of fifteen years.

[0063] Methods of the present invention can be used in combination with additional cancer therapies. In some embodiments, the additional cancer therapy comprises surgery, radiotherapy, chemotherapy, toxin therapy, immunotherapy, cryotherapy or gene therapy. In some embodiments, the cancer is a chemotherapy-resistant, immunotherapy-resistant or radio-resistant cancer.

[0064] A combination treatment of the present invention according to embodiments of the invention can be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. A method of treatment in the present invention can be applied as the combination of an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof with an effective amount of an immunotherapy agent or an effective amount of a chemotherapeutic agent, in any animal model or in the clinic.

Compositions/Kits

[0065] Aspects of the invention also relate to a pharmaceutical composition or kit comprising an effective amount of nitroxoline or its analogue, or a pharmaceutically acceptable salt thereof, an effective amount of an immunotherapy agent or a chemotherapeutic agent, and a pharmaceutically acceptable carrier, as well as methods of manufacturing the pharmaceutical composition or kit by combining the nitroxoline or its analogues, or the pharmaceutically acceptable salts thereof, the at least one additional anti-cancer immunotherapy or chemotherapy agent, and a pharmaceutically acceptable carrier using methods known in the art in view of the present disclosure.

[0066] According to embodiments of the invention, a kit comprises one or more pharmaceutical compositions having an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of an immunotherapy agent. According to other embodiments of the invention, a kit comprises one or more pharmaceutical compositions having an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of a chemotherapeutic agent. The effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and the effective amount of an immunotherapy agent or chemotherapeutic agent can be present in one pharmaceutical composition. They can also be present in separate pharmaceutical compositions. The kit further contains instructions on using the combination of the effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and the effective amount of an immunotherapy agent or chemotherapeutic agent for treating a cancer.

[0067] Compositions or kits according to embodiments of the invention can be prepared using methods known in the art in view of the present disclosure.

[0068] According to particular embodiments, an effective amount refers to the amount of each active ingredient or compound that is effective to achieve a synergistic effect on one, two, three, four, or more of the following effects: (i) reduce or ameliorate the severity of the cancer to be treated or a symptom associated therewith; (ii) reduce the duration of the cancer to be treated, or a symptom associated therewith; (iii) prevent the progression of the cancer to be treated, or a symptom associated therewith; (iv) cause regression of the cancer to be treated, or a symptom associated therewith; (v) prevent the development or onset of the cancer to be treated, or a symptom associated therewith; (vi) prevent the recurrence of the cancer to be treated, or a symptom associated therewith; (vii) reduce hospitalization of a subject having the cancer to be treated, or a symptom associated therewith; (viii) reduce hospitalization length of a subject having the cancer to be treated, or a symptom associated therewith; (ix) increase the survival of a subject with the cancer to be treated, or a symptom associated therewith; (xi) inhibit or reduce the cancer to be treated, or a symptom associated therewith in a subject; and/or (xii) enhance or improve the prophylactic or therapeutic effect(s) of another therapy.

EMBODIMENTS

[0069] Embodiment 1 is a method of treating a cancer comprising administering to a subject in need thereof an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of an immunotherapy agent, preferably the immunotherapy agent is an inhibitor or modulator of regulatory T cells or myeloid-derived suppressor cells, and the combination of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the immunotherapy agent results in a synergistic effect.

[0070] Embodiment 2 is the method of Embodiment 1, wherein the immunotherapy agent is an inhibitor of the PD-L1/PD-1 pathway or an inhibitor of the CTLA-4 pathway, preferably the immunotherapy agent is an anti-PD-1 antibody, an anti-PD-L1 antibody, an anti-CTLA-4 antibody, or an antigen binding fragment thereof, more preferably, the antibody or fragment thereof is administered by intravenous or intramuscular injection.

[0071] Embodiment 3 is the method of Embodiment 1, wherein the immunotherapy agent is a Bacillus Calmette-Guerin (BCG) therapy, preferably an attenuated BCG therapy, more preferably mycobacterial cell wall fragments, and most preferably mycobacterial cell wall fragments with biologically active nucleic acids derived from Mycobacterium phlei, more preferably, the BCG therapy is administered intravesically.

[0072] Embodiment 4 is the method of Embodiment 3, wherein nitroxoline, the analogue or the pharmaceutically acceptable salt thereof is not administered within 24 hours of the intravesical instillation of the BCG therapy.

[0073] Embodiment 5 is the method of any one of Embodiments 1 to 4, wherein the cancer is selected from the group consisting of melanoma, cervical cancer, breast cancer, ovarian cancer, prostate cancer, testicular cancer, urothelial carcinoma, bladder cancer, non-small cell lung cancer, small cell lung cancer, sarcoma, colorectal adenocarcinoma, gastrointestinal stromal tumors, gastroesophageal carcinoma, colorectal cancer, pancreatic cancer, kidney cancer, hepatocellular cancer, malignant mesothelioma, leukemia, lymphoma, myelodysplastic syndrome, multiple myeloma, transitional cell carcinoma, neuroblastoma, plasma cell neoplasms, Wilm's tumor, glioblastoma, retinoblastoma, and hepatocellular carcinoma.

[0074] Embodiment 6 is the method of Embodiment 5, wherein the cancer is bladder cancer.

[0075] Embodiment 7 is the method of Embodiment 5, wherein the cancer is prostate cancer.

[0076] Embodiment 8 is the method of any one of Embodiments 1 to 7, wherein the effective amount of nitroxoline, or the analogue or pharmaceutically acceptable salt thereof is administered orally or by intravenous or intramuscular injection.

[0077] Embodiment 9 is a method of treating a cancer, preferably a bladder cancer, a liver cancer, or prostate cancer, comprising administering to a subject in need thereof an effective amount of nitroxoline or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of a chemotherapeutic agent, wherein the combination of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the chemotherapeutic agent results in a synergistic effect.

[0078] Embodiment 10 is the method of Embodiment 9, wherein the chemotherapeutic agent is a taxane drug that targets tubulin, preferably paclitaxel, docetaxel, cabazitaxel; or the chemotherapeutic agent is DNA cross-linking or intercalating agent, preferably mitomycin C, an anthracycline that is preferably epirubicin, pirarubicin, or a platinum-based antineoplastic agent that is preferably carboplatin, cisplatin, oxaliplatin or nedaplatin.

[0079] Embodiment 11 is the method of Embodiment 9 or 10, wherein the chemotherapeutic agent is administered intravesically once every week, or once every 2, 3, 4 or 5 weeks.

[0080] Embodiment 12 is the method of any one of Embodiments 1 to 11, wherein the nitroxoline analogue is selected from the group consisting of oxyquinoline, clioquinol and iodoquinol.

[0081] Embodiment 13 is the method of any one of Embodiments 1 to 12, wherein the effective amount of nitroxoline or the analogue or pharmaceutically acceptable salt thereof is 100 mg to 1600 mg orally administered per day, preferably 600 mg to 1600 mg orally administered per day.

[0082] Embodiment 14 is a method of treating a cancer, preferably a bladder cancer or a prostate cancer, comprising administering to a subject in need thereof an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of an anti-PD-1 antibody or an antigen binding fragment thereof, wherein nitroxoline or the analogue or pharmaceutically acceptable salt thereof is orally administered 100 mg to 1600 mg per day, preferably 600 mg to 1600 mg per day, and the anti-PD-1 antibody or antigen binding fragment thereof is administered by intravenous or intramuscular injection 0.1 mg/kg to 20 mg/kg once every 1 to 4 weeks, preferably 2 mg/kg to 15 mg/kg in total in a period of 3 weeks, and the combination of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the anti-PD-1 antibody or antigen binding fragment thereof results in a synergistic effect.

[0083] Embodiment 15 is a method of treating a cancer, preferably a bladder cancer or a prostate cancer, comprising administering to a subject in need thereof an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of an anti-PD-L1 antibody or an antigen binding fragment thereof, wherein nitroxoline or the analogue or pharmaceutically acceptable salt thereof is orally administered 100 mg to 1600 mg per day, preferably 600 mg to 1600 mg per day, and the anti-PD-L1 antibody or antigen binding fragment thereof is administered by intravenous or intramuscular injection 1 mg/kg to 40 mg/kg once every 2 to 4 weeks, preferably 10 mg/kg to 30 mg/kg in total in a period of 4 weeks, and the combination of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the anti-PD-L1 antibody or antigen binding fragment thereof results in a synergistic effect.

[0084] Embodiment 16 is a method of treating a cancer, preferably a bladder cancer or a prostate cancer, comprising administering to a subject in need thereof an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of an anti-CTLA-4 antibody or an antigen binding fragment thereof, wherein nitroxoline or the analogue or pharmaceutically acceptable salt thereof is orally administered 100 mg to 1600 mg per day, preferably 600 mg to 1600 mg per day, and the anti-CTLA-4 antibody or antigen binding fragment thereof is administered by intravenous or intramuscular injection 0.1 mg/kg to about 6 mg/kg once every 3 to 5 weeks, preferably 1 mg/kg to 4 mg/kg in total in a period of 4 weeks, and the combination of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the anti-CTLA-4 antibody or antigen binding fragment thereof results in a synergistic effect.

[0085] Embodiment 17 is a method of treating a cancer, preferably a bladder cancer or a prostate cancer, comprising administering to a subject in need thereof an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of a BCG therapy, wherein nitroxoline or the analogue or pharmaceutically acceptable salt thereof is orally administered 100 mg to 1600 mg per day, preferably 600 mg to 1600 mg per day, and the BCG therapy is intravesically administered in a dosage of 0.5×10^8 to 50×10^8 colony forming units (CFUs) once every 1 to 5 weeks, preferably 1×10^8 to 8×10^8 CFUs once a week, and the combination of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the BCG therapy results in a synergistic effect.

[0086] Embodiment 18 is a method of treating a cancer, preferably a bladder cancer, a liver cancer or a prostate cancer, comprising administering to a subject in need thereof an effective amount of nitroxoline, or an analogue or pharmaceutically acceptable salt thereof, and an effective amount of a chemotherapeutic agent, wherein nitroxoline or the analogue or pharmaceutically acceptable salt thereof is orally administered 100 mg to 1600 mg per day, preferably 600 mg to 1600 mg per day, the effective amount of the chemotherapeutic agent is administered intravesically or by intravenous or intramuscular injection, and the combination of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the chemotherapeutic agent results in a synergistic effect.

[0087] Embodiment 19 is the method of Embodiment 18, wherein the chemotherapeutic agent is a DNA cross-linking or intercalating agent selected from the group consisting of mitomycin C, epirubicin, pirarubicin, carboplatin, cisplatin, oxaliplatin and nedaplatin, and the chemotherapeutic

agent is administered intravesically once every 1-5 weeks in a bladder instillation fluid at a concentration of 0.1 mg/ml to 5 mg/ml, preferably 0.5 mg/ml to 2 mg/mL.

[0088] Embodiment 20 is the method of Embodiment 18, wherein the chemotherapeutic agent is a taxane drug selected from the group consisting of paclitaxel docetaxel and cabazitaxel, and the chemotherapeutic agent is administered intravesically in a bladder instillation fluid at a concentration of 0.5 to 10 mg/mL, preferably 1 to 5 mg/mL.

[0089] Embodiment 21 is the method of any one of Embodiments 1 to 20, wherein nitroxoline or a pharmaceutically acceptable salt thereof is administered orally or by injection to the subject in need thereof.

[0090] Embodiment 22 is the method of any one of Embodiments 1 to 20, wherein a nitroxoline analogue selected from the group consisting of oxyquinoline, clioquinol and iodoquinol or a pharmaceutically acceptable salt thereof is administered orally or by injection to the subject in need thereof.

[0091] Embodiment 23 is a kit comprising an effective amount of nitroxoline or its analogue, or a pharmaceutically acceptable salt thereof, an effective amount of an immunotherapy agent or chemotherapeutic agent, and one or more pharmaceutically acceptable carriers, wherein the effective amount of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the effective amount of the immunotherapy agent or chemotherapeutic agent are present in the same pharmaceutical composition or separate pharmaceutical compositions.

[0092] Embodiment 24 is the kit of Embodiment 23, wherein the kit comprises a pharmaceutical composition containing 100 mg and 1600 mg, preferably 600 mg to 1600 mg, per dosage form of nitroxoline or its analogue, or a pharmaceutically acceptable salt thereof.

[0093] Embodiment 25 is the kit of Embodiment 23 or 24, wherein the kit comprises a pharmaceutical composition containing the immunotherapy agent selected from the group consisting of an anti-PD-1 antibody or an antigen binding fragment thereof; an anti-PD-L1 antibody or an antigen binding fragment thereof; an anti-CTLA-4 antibody or an antigen binding fragment thereof; and a BCG therapy.

[0094] Embodiment 26 is the kit of Embodiment 23 or 24, wherein the kit comprises a pharmaceutical composition containing the chemotherapeutic agent selected from the group consisting of mitomycin C, epirubicin, pirarubicin, carboplatin, cisplatin, oxaliplatin, nedaplatin, paclitaxel docetaxel and cabazitaxel.

[0095] Embodiment 27 is the kit of any one of Embodiments 23 to 26, wherein the kit comprises a pharmaceutical composition containing nitroxoline or a pharmaceutically acceptable salt thereof.

[0096] Embodiment 28 is the kit of any one of Embodiments 23 to 26, wherein the kit comprises a pharmaceutical composition containing a nitroxoline analogue selected from the group consisting of oxyquinoline, clioquinol and iodoquinol, or a pharmaceutically acceptable salt thereof.

[0097] Embodiment 29 is a method of manufacturing the kit of any one of Embodiments 23 to 28, comprising combining the nitroxoline, its analogue, or the pharmaceutically acceptable salt thereof, the immunotherapy agent or the chemotherapeutic agent, and the one or more pharmaceutically acceptable carriers in the kit.

[0098] Embodiment 30 is a method of manufacturing the kit of any one of Embodiments 23 to 28, comprising obtaining a first pharmaceutical composition comprising the effective amount of nitroxoline, its analogue, or the pharmaceutically acceptable salt thereof, obtaining a second pharmaceutical composition comprising the effective amount of the immunotherapy agent or the chemotherapeutic agent, and combining the first and second pharmaceutical compositions in the kit.

[0099] Embodiment 31 is a use of a kit of any one of Embodiments 23 to 28 in the manufacture of a medicament for the treatment of cancer, and the combined use of nitroxoline or the analogue or pharmaceutically acceptable salt thereof and the immunotherapy agent or chemotherapeutic agent results in a synergistic effect.

[0100] Embodiment 32 is the use of Embodiment 31, wherein the cancer is selected from the group consisting of melanoma, cervical cancer, breast cancer, ovarian cancer, prostate cancer, testicular cancer, urothelial carcinoma, bladder cancer, non-small cell lung cancer, small cell lung cancer, sarcoma, colorectal adenocarcinoma, gastrointestinal stromal tumors, gastroesophageal carcinoma, colorectal cancer, pancreatic cancer, kidney cancer, hepatocellular cancer, malignant mesothelioma, leukemia, lymphoma, myelodysplastic syndrome, multiple myeloma, transitional cell carcinoma, neuroblastoma, plasma cell neoplasms, Wilm's tumor, glioblastoma, retinoblastoma, and hepatocellular carcinoma.

[0101] Embodiment 33 is the use of Embodiment 32, wherein the cancer is bladder cancer.

[0102] Embodiment 34 is the use of Embodiment 32, wherein the cancer is prostate cancer.

[0103] Embodiment 35 is the use of Embodiment 32, wherein the cancer is liver cancer.

EXAMPLES

[0104] The following examples of the invention are to further illustrate the nature of the invention. It should be understood that the following examples do not limit the invention and that the scope of the invention is to be determined by the appended claims.

Example 1: Synergistic Inhibition of Tumor Growth by a Combination of NIT and One or More Chemotherapies

[0105] NIT was tested with a group of chemotherapies including carboplatin, paclitaxel, mitomycin C, epirubicin and pirarubicin in the growth inhibition of two human cancer cell lines. As shown in Tables 2 and 3, NIT inhibited the growth of the human bladder cancer cell line 5637 and of the human liver cancer cell line HepG2.

TABLE-US-00002 TABLE 2 Incubation of NIT with the human bladder cancer cell line 5637 for 120 hours resulted in inhibition of cell growth. Inhibition NIT (μg/ml) OD Values Mean SD Rates

0	1.3	1.26	1.29	1.316	1.292	0.024	0.25	1.148	1.126	0.798	1.164	1.059	0.175	18.0%	0.5	0.606	0.624
0.632	0.618	0.620	0.011	52.0%	1	0.254	0.268	0.25	0.264	0.259	0.008	79.9%	2	0.097	0.104	0.098	
0.097	0.099	0.003	92.3%	4	0.056	0.06	0.058	0.056	0.058	0.002	95.5%						

TABLE-US-00003 TABLE 3 Incubation of NIT with the human liver cancer cell line HepG2 for 120 hours resulted in inhibition of cell growth. Inhibition NIT(μg/ml) OD Values Mean SD Rates

0	1.255	1.243	1.241	1.289	1.257	0.022	0.5	1.264	1.195	1.243	1.164	1.217	0.045	3.2%	1	0.975	1.046
0.977	0.939	0.984	0.045	21.7%	2	0.273	0.255	0.245	0.217	0.248	0.023	80.3%	4	0.113	0.124	0.124	
0.118	0.120	0.005	90.5%	8	0.086	0.082	0.08	0.081	0.082	0.003	93.5%	16	0.057	0.06	0.061	0.06	
0.060	0.002	95.3%															

[0106] As shown in Tables 4 and 5, carboplatin inhibited the growth of the human bladder cancer cell line 5637 and of the human liver cancer cell line HepG2.

TABLE-US-00004 TABLE 4 Incubation of carboplatin with the human bladder cancer cell line 5637 for 120 hours resulted in inhibition of cell growth. Inhibition Carbo(μg/ml) OD Values Mean SD Rates

0	1.3	1.26	1.29	1.316	1.292	0.024	0.5	0.836	0.854	0.892	0.956	0.885	0.053	31.5%	1		
0.576	0.616	0.574	0.628	0.599	0.028	53.7%	2	0.396	0.403	0.386	0.369	0.389	0.015	69.9%	4	0.163	
0.149	0.153	0.151	0.154	0.006	88.1%	8	0.11	0.113	0.105	0.112	0.110	0.004	91.5%				

TABLE-US-00005 TABLE 5 Incubation of carboplatin with the human liver cancer cell line HepG2 for 120 hours resulted in inhibition of cell growth. Carbo Inhibition (μg/ml) OD Values Mean SD Rates

0	1.255	1.243	1.241	1.289	1.257	0.022	4	0.761	0.747	0.767	0.796	0.768	0.021				
38.9%	8	0.361	0.308	0.346	0.328	0.336	0.023	70.7%	16	0.148	0.177	0.18	0.175	0.170	0.015	85.2%	
32	0.104	0.125	0.133	0.136	0.125	0.014	89.1%	64	0.08	0.083	0.084	0.076	0.081	0.004	93.0%	128	
0.074	0.068	0.068	0.068	0.070	0.003	93.9%											

[0107] As shown in Tables 6 and 7, paclitaxel inhibited the growth of the human bladder cancer cell line 5637 and of the human liver cancer cell line HepG2.

TABLE-US-00006 TABLE 6 Incubation of paclitaxel with the human bladder cancer cell line 5637 for 112 hours resulted in inhibition of cell growth. Inhibition Taxol(ng/ml) OD Values Mean SD Rates

0	1.028	1.023	0.987	0.988	1.007	0.022	0.125	0.884	0.899	0.885	0.879	0.887	0.009	11.9%			
---	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	--	--	--

0.25 0.531 0.523 0.549 0.506 0.527 0.018 47.6% 0.5 0.149 0.159 0.161 0.163 0.158 0.006 84.3% 1
0.093 0.097 0.099 0.101 0.098 0.003 90.3% 2 0.03 0.031 0.034 0.035 0.033 0.002 96.8%
TABLE-US-00007 TABLE 7 Incubation of paclitaxel with the human liver cancer cell line HepG2
for 112 hours resulted in inhibition of cell growth. Inhibition Taxol(ng/ml) OD Values Mean SD
Rates 0 1.79 1.764 1.73 1.845 1.782 0.049 0.5 1.642 1.718 1.69 1.664 1.679 0.033 5.8% 1 1.544
1.548 1.519 1.523 1.534 0.015 14.0% 2 0.974 0.914 0.95 0.904 0.936 0.032 47.5% 4 0.428 0.466
0.41 0.423 0.432 0.024 75.8% 8 0.177 0.153 0.173 0.182 0.171 0.013 90.4% 16 0.128 0.134 0.123
0.138 0.131 0.007 92.7%

[0108] The IC.sub.50 values of these compounds' inhibition of cell growth are summarized in Table 8.

TABLE-US-00008 TABLE 8 IC.sub.50 values of NIT, carboplatin and paclitaxel for inhibition of
cell growth of human bladder cancer cell line 5637 and of human liver cancer cell line HepG2. NIT
Carboplatin Paclitaxel Cell Line (μg/ml) (μg/ml) (ng/ml) 5637 0.517 0.917 0.289 HepG2 1.86
3.687 2.502

[0109] Mitomycin C, epirubicin and pirarubicin were similarly tested for inhibition of the growth
of bladder cancer cell line 5637, and their IC.sub.25, IC.sub.50 and IC.sub.75 values are
summarized in Table 9.

TABLE-US-00009 TABLE 9 IC.sub.25, IC.sub.50 and IC.sub.75 values of mitomycin C,
epirubicin and pirarubicin for inhibition of cell growth of human bladder cancer cell line 5637.
Drug IC.sub.25(ng/ml) IC.sub.50(ng/ml) IC.sub.75(ng/ml) Mitomycin C 48.36 152.53 536.62
Epirubicin 34.97 45.21 63.93 Pirarubicin 96.58 142.57 219.58

[0110] The combinational studies of NIT and chemotherapies were conducted by mixing NIT with
individual chemotherapies at five to seven concentration levels, with the medium concentration
levels close to their IC.sub.50 values. The combinational indices (CIs) were calculated and plotted
according to Ting-Chao Chou's method (19) to estimate the combinational effects, including
synergism, addition and antagonism. As shown in Tables 10 and 11, NIT inhibited the growth of
human bladder cancer 5637 in combination with carboplatin and paclitaxel, respectively, and the CI
values of the studies were calculated and plotted in FIGS. 2 and 3, respectively.

TABLE-US-00010 TABLE 10 Incubation of a combination of NIT and carboplatin with the human
bladder cancer cell line 5637 for 120 hours resulted in inhibition of cell growth. NIT Carbo
Inhibition (μg/ml) (μg/ml) OD Values Mean SD Rates 0 0 1.324 1.364 1.296 1.348 1.333 0.030
0.25 0.5 0.36 0.394 0.41 0.362 0.382 0.025 71.4% 0.5 1 0.134 0.144 0.132 0.15 0.140 0.008
89.5% 1 2 0.072 0.08 0.07 0.07 0.073 0.005 94.5% 2 4 0.038 0.046 0.038 0.042 0.041 0.004
96.9% 4 8 0.02 0.019 0.017 0.017 0.018 0.002 98.6%

TABLE-US-00011 TABLE 11 Incubation of a combination of NIT and paclitaxel with the human
bladder cancer cell line 5637 for 96 hours resulted in inhibition of cell growth. NIT Taxol
Inhibition (μg/ml) (ng/ml) OD Values Mean SD Rates 0 0 0.793 0.786 0.766 0.784 0.782 0.012
0.125 0.125 0.567 0.536 0.542 0.516 0.540 0.021 30.9% 0.25 0.25 0.375 0.42 0.385 0.424 0.401
0.025 48.7% 0.5 0.5 0.082 0.083 0.081 0.086 0.083 0.002 89.4% 1 1 0.027 0.03 0.026 0.027
0.028 0.002 96.5% 2 2 0.015 0.014 0.013 0.015 0.014 0.001 98.2%

[0111] As shown in Tables 12 and 13, NIT inhibited the growth of human liver cancer cell line
HepG2 in combination with carboplatin and paclitaxel, respectively, the CI values of the studies are
plotted in FIGS. 4 and 5, respectively.

TABLE-US-00012 TABLE 12 Incubation of a combination of NIT and carboplatin with the human
liver cancer cell line HepG2 for 120 hours resulted in inhibition of cell growth. NIT Carbo
Inhibition (μg/ml) (μg/ml) OD Values Mean SD Rates 0 0 1.261 1.243 1.235 1.257 1.249 0.012 0.5
4 0.469 0.496 0.472 0.454 0.473 0.017 62.1% 1 8 0.133 0.134 0.138 0.146 0.138 0.006 89.0% 2 16
0.098 0.094 0.093 0.097 0.096 0.002 92.4% 4 32 0.077 0.08 0.079 0.081 0.079 0.002 93.7% 8 64
0.051 0.05 0.049 0.052 0.051 0.001 96.0% 16 128 0.047 0.043 0.046 0.043 0.045 0.002 96.4%

TABLE-US-00013 TABLE 13 Incubation of a combination of NIT and paclitaxel with the human

liver cancer cell line HepG2 for 120 hours resulted in inhibition of cell growth. NIT Taxol
 Inhibition (µg/ml) (ng/ml) OD Values Mean SD Rates 0 0 1.393 1.397 1.451 1.414 1.414 0.026 0.5
 0.5 1.115 1.134 1.164 1.129 1.136 0.021 19.7% 1 1 0.928 0.938 1.009 0.963 0.960 0.036 32.1% 2 2
 0.516 0.552 0.556 0.474 0.525 0.038 62.9% 4 4 0.199 0.214 0.199 0.182 0.199 0.013 86.0% 8 8
 0.116 0.116 0.111 0.097 0.110 0.009 92.2% 16 16 0.059 0.067 0.068 0.066 0.065 0.004 95.4%
 [0112] Similarly, NIT was combined with mitomycin C, epirubicin and pirarubicin to inhibit the
 growth of bladder cancer cell line 5637, and their combinational inhibitory results and CI plots are
 shown in Tables 14-17 and FIGS. 6-8.

TABLE-US-00014 TABLE 14 Incubation of a combination of NIT and mitomycin C with the
 human bladder cancer cell line 5637 for 96 hours resulted in inhibition of cell growth. NIT 5637
 Average Fa 10000 5000 2500 1250 625 0 Mitomycin C 200 94.1% 85.9% 82.1% 78.6% 80.3%
 79.4% 50 93.5% 79.5% 71.2% 75.9% 77.2% 77.9% 12.5 93.5% 75.0% 51.4% 29.7% 24.9% 27.7%
 3.13 94.2% 75.3% 36.4% 0.5% 4.8% 1.6% 0 94.0% 74.3% 23.3% -3.8% -7.2% 0.0%

TABLE-US-00015 TABLE 15 CI values of the combination of NIT and mitomycin C when
 incubated with the human bladder cancer cell line 5637 for 96 hours. NIT CI 10000 5000 2500
 1250 625 0 Mitomycin 200 1.764 2.032 2.011 2.122 1.874 C 50 1.289 1.263 1.094 0.749 0.613
 12.5 1.194 0.996 0.861 0.952 0.905 3.13 1.126 0.888 0.772 5.148 1.033 0

TABLE-US-00016 TABLE 16 CI values of the combination of NIT and epirubicin when incubated
 with the human bladder cancer cell line 5637 for 96 hours. NIT CI 10000 5000 2500 1250 625 0
 Epirubicin 50 1.826 1.728 1.51 1.288 1.154 16.67 1.339 1.158 1.009 0.885 0.718 5.56 1.223 1.015
 1.035 4.16 3.337 1.85 1.166 0.914 0.857 2.483 1.66 0

TABLE-US-00017 TABLE 17 CI values of the combination of NIT and pirarubicin when
 incubated with the human bladder cancer cell line 5637 for 96 hours. NIT CI 10000 5000 2500
 1250 625 0 pirarubicin 200 2.266 1.885 1.536 1.39 1.276 66.67 1.405 1.915 1.664 1.333 1.219
 22.22 1.185 0.969 0.596 0.409 0.315 7.41 1.067 0.965 1.437 5.521 11.846 0

[0113] According to Chou (19), a CI value below 0.9 represents a synergistic combination of the
 two drugs, and values of 0.1-0.3, 0.3-0.7, 0.7-0.85, and 0.85-0.9 represent strong synergism,
 synergism, moderate synergism and slight synergism, respectively. Therefore, the CI plots in FIGS.
 2 and 4 support the conclusion that NIT and carboplatin inhibited the growth of the bladder cancer
 cell line 5637 and of the liver cancer cell line HepG2 in a manner that is classified as synergistic or
 strongly synergistic. The CI plot in FIG. 3 shows that the combination of NIT and paclitaxel
 demonstrated slight synergism in inhibiting the growth of the bladder cancer cell line 5637, and
 FIG. 5 shows that the inhibition of the growth of the liver cancer cell line HepG2 with this
 combination is partially synergistic when the drug concentrations' Fa (fraction absorbed) is lower
 than 0.3. This partial synergism at concentrations below the drugs' IC50 values was also found for
 the combination of NIT and mitomycin C for inhibition of the bladder cancer cell line 5637, as
 shown in Table 15 and FIG. 6. Tables 16 and 17 and FIGS. 7 and 8 showed that the combination of
 NIT with epirubicin or pirarubicin are partially synergistic at some concentrations for inhibition of
 the growth of the bladder cancer cell line 5637.

[0114] The synergistic tumor growth inhibition exhibited by NIT in combination with mitomycin C
 (MMC) was further evaluated in a mouse bladder cancer orthotopic xenograft. FIG. 9 shows the
 effect of combining orally dosed NIT and intravesically administered MMC on the inhibition of
 tumor development in a MBT-2-luc orthotopic mouse bladder cancer model. The tumor volume was
 analyzed by the Xenogen IVIS200 system following the indicated treatment. FIG. 9A shows a
 typical IVIS image for each treatment group. The tumor volume of each mouse was determined by
 region-of-interest analysis of total photons per second. Six mice were analyzed in each group. As
 shown in FIG. 9B, the mean tumor volume for the group of 15 mg/kg Bid oral NIT in combination
 with Q7d 1 mg/ml intravesical mitomycin C (NIT+MMC) was significantly different from that for
 single NIT or single MMC treatment groups (p<0.0001 and p=0.0001 respectively), indicating a
 remarkable enhancement in tumor growth inhibition by the combination treatment of NIT with

MMC. The Kaplan-Meier analysis was also performed to assess mouse survival for each treatment and is summarized in FIG. 9C. Consistently, the survival status for the group of NIT plus MMC was significantly improved, in contrast to the control and single drug treatment groups.

[0115] The results of the tumor growth inhibition and survival proportions, as well as the Bliss independence model calculations of additivity are summarized in Table 18.

TABLE-US-00018 TABLE 18 Bliss independence model calculations of additivity for oral NIT in combination with MMC in an orthotopic mouse MBT-2-Luc bladder cancer model. NIT + MMC Expected Bliss Actual Single drug Additive Observed NIT MMC Value Value Interaction Tumor 0.33 0.56 0.7 0.71 Synergy Growth Inhibition Survival 0.33 0.57 0.71 1 Synergy Proportion

[0116] NIT administered orally at a dose of 15 mg/kg (Bid) inhibited tumor growth at a rate of 0.33, and the proportion of surviving mice at the end of the experiment (45 days) was 0.33. The tumor inhibition rate and the proportion of surviving mice for intravesically administered MMC treatment was 0.56 and 0.57 respectively. When oral NIT was dosed with MMC, the observed tumor inhibition rate and the proportion of surviving mice was significantly increased to 0.71 and 1, respectively, which were greater than their Expected Bliss Additive Values, indicating a synergy for tumor inhibition by the combination of NIT and MMC.

[0117] The discovery that NIT has different degrees of synergism with different classes of chemotherapies can be used for the combinational treatment of a variety of solid and blood tumors. Example 2: Synergistic Inhibition of Tumor Growth by combination of NIT with Immunotherapies

[0118] NIT was tested with a group of immunotherapies including BCG and anti-PD-1 antibody for tumor growth inhibition of mouse bladder and prostate cancer orthotopic xenograft models.

[0119] FIG. 10 shows the effect of combining orally dosed NIT and intravesically instilled BCG on inhibiting tumor development in a MBT-2-luc orthotopic mouse bladder cancer model. The tumor volume was analyzed by the Xenogen IVIS200 system following the indicated treatment, and a typical IVIS image for each treatment group of indicated treatment is shown in FIG. 10A. The tumor volume of each mouse was determined by region-of-interest analysis of total photons per second. Eight mice were analyzed in each group. As shown in FIG. 10B, 30 mg/kg NIT administered orally inhibited the cancer growth, either continuously or with a 48-hour break, at a slightly higher or similar activity compared to that caused by BCG administered intravesically. When oral NIT (administered continuously or with a 48-hour break) was dosed with BCG, the inhibition rates were significantly increased, compared to single NIT or BCG treatments ($p < 0.0001$ for all comparisons). The Kaplan-Meier analysis was also performed to evaluate mouse survival for each treatment and is summarized in FIG. 10C. The survival status for the group of oral NIT (administered continuously or with a 48-hour break) in combination with BCG was significantly improved, compared to the single drug treatment groups.

[0120] The results of the Bliss independence model calculations of additivity for tumor growth inhibition and survival proportions are summarized in Table 19 (continuous NIT) and Table 20 (NIT with 48 h break).

TABLE-US-00019 TABLE 19 Bliss independence model calculations of additivity for oral NIT (continuous) in combination with intravesical BCG in an orthotopic mouse MBT-2-Luc bladder cancer model. NIT (continuous) + BCG Expected Single Drug Bliss Actual NIT Additive Observed (continuous) BCG Value Value Interaction Tumor 0.52 0.35 0.69 0.75 Synergy Growth Inhibition Survival 0.4 0.2 0.52 0.60 Synergy Proportion

TABLE-US-00020 TABLE 20 Bliss independence model calculations of additivity for oral NIT (48 h-break) in combination with intravesical BCG in an orthotopic mouse MBT-2-Luc bladder cancer model. NIT (48 h-break) + BCG Expected Single Drug Bliss Actual NIT Additive Observed (48 h-break) BCG Value Value Interaction Tumor 0.35 0.35 0.58 0.85 Synergy Growth Inhibition Survival 0.4 0.2 0.52 0.60 Synergy Proportion

[0121] When oral NIT was administered in combination with intravesical BCG, either continuously or with a 48-hour break, the observed tumor inhibition rates and survival proportions were

significantly increased, compared to that of single NIT and BCG treatments. The actual tumor inhibition rates and survival proportions of NIT in combination with BCG were much greater than their Expected Bliss Additive Value, suggesting a robust synergy in tumor inhibition by the combination of NIT and BCG.

[0122] To assess a synergistic tumor inhibition effect of NIT in combination with anti-PD-1, the combination was tested for tumor growth inhibition of mouse bladder and prostate cancer orthotopic xenograft models.

[0123] FIG. 11 shows the effect of orally dosed NIT combined with intraperitoneal anti-PD-1 antibody on inhibiting tumor growth in a MBT-2-luc orthotopic mouse bladder cancer model. Typical IVIS images for each indicated treatment group are shown in FIG. 11A. The tumor volume of each mouse was determined by region-of-interest analysis of total photons per second. Five mice were analyzed in each group. As shown in FIG. 11B, 10 mg/kg intraperitoneal anti-PD-1 intensively inhibited the tumor growth at a rate of 80%, while 15 mg/kg NIT administered orally showed a tumor inhibition rate of 31%. When oral NIT was combined with anti-PD-1, the inhibition rates were significantly enhanced (90%), compared to single NIT or anti-PD-1 treatments ($p<0.0001$ and $p=0.039$ respectively). The Kaplan-Meier analysis was also performed and is summarized in FIG. 11C. The survival status for the groups of NIT plus anti-PD-1 or the single anti-PD-1 was significantly higher than the single NIT group.

[0124] The results of the Bliss independence model calculations of additivity for tumor growth inhibition are summarized in Table 21. When NIT was administered in combination with anti-PD-1, its actual observed tumor inhibition rate was greater than the Expected Bliss Additive Value, suggesting a synergy in tumor inhibition.

TABLE-US-00021 TABLE 21 Bliss independence model calculations of additivity for oral NIT in combination with anti-PD-1 antibody in an orthotopic mouse MBT-2-Luc bladder cancer model.

	NIT	anti-PD-1	Expected Bliss	Actual	Single Drug	Additive	Observed	Value	Value
Interaction	Tumor	0.31	0.8	0.86	0.9	Synergy	Growth Inhibition		

[0125] FIG. 12 shows the effect of orally dosed NIT combined with intraperitoneal anti-PD-1 antibody on inhibiting tumor growth in a RM-9-luc orthotopic mouse prostate cancer model. Typical IVIS images for each indicated treatment group are shown in FIG. 12A. The tumor volume of each mouse was determined by region-of-interest analysis of total photons per second. Eight mice were analyzed for each group. As shown in FIG. 12B, 10 mg/kg intraperitoneal anti-PD-1 showed robust tumor growth inhibition (76%), while 15 mg/kg NIT administered orally showed a tumor inhibition rate of 52%. When NIT was combined with anti-PD-1, the tumor inhibition rate was significantly enhanced to 96%, with p values of 0.017 and 0.027 compared to single NIT and anti-PD-1 treatments, respectively. FIG. 12C shows images of tumors collected from each group. The tumor weight of each mouse was determined and is summarized in FIG. 12D. Analysis of tumor weight data revealed that the combination of NIT and anti-PD-1 resulted in significant enhancement of tumor inhibition (74%), compared to single NIT and anti-PD-1 treatments (37% and 56% respectively), with p values of 0.0006 and 0.0111, respectively.

[0126] The results of the Bliss independence model calculations of additivity for tumor volume and weight are summarized in Table 22. When NIT was administered in combination with anti-PD-1, its actual observed tumor volume and weight inhibition rate were 0.96 and 0.74, respectively, greater than their Expected Bliss Additive Values (0.89 and 0.72, respectively), suggesting a synergy in tumor inhibition.

TABLE-US-00022 TABLE 22 Bliss independence model calculations of additivity for oral NIT in combination with anti-PD-1 antibody in an orthotopic mouse RM-9-Luc prostate cancer model.

	NIT	anti-PD-1	Expected Bliss	Actual	Single drug	Additive	Observed	NIT	anti-PD-1	Value	Value	
Interaction	Tumor	0.52	0.76	0.89	0.96	Synergy	Growth Inhibition	Tumor	0.37	0.56	0.72	0.74
								Synergy	Weight Proportion			

[0127] The discovery that NIT has different degrees of synergism with different immunotherapies

can be used for the combinational treatment of a variety of solid and blood tumors.

Example 3: Experimental Procedures

3.1 Study Materials

[0128] Compounds: NIT, carboplatin, paclitaxel, epirubicin and pirarubicin were purchased from suppliers, and were dissolved in the DMSO to stock solution and store in -20°C . Before use, the stock solutions were diluted to different concentrations of working solutions. The DMSO concentration in working solutions was less than 1%. All cancer cell lines were purchased from vendors.

[0129] Cell lines and reagents: The murine bladder cancer cell lines MBT-2 and RM-9 were provided by the American Type Culture Collection (Rockville, MD, USA). All cells were maintained in RPM I-1640 medium supplemented with 10% fetal bovine serum. The cells were cultured at 37°C in a 5% CO_2 atmosphere and routinely passaged by trypsin-EDTA treatment in 100-cm² flasks containing BCG (81 mg; Connaught substrain, ImmuCyst, Nihou Kayaku, Inc., Tokyo, Japan), and phosphate-buffered saline (PBS) for in vivo studies.

[0130] MBT-2 and RM-9 cells stably expressing luciferase (MBT-2-Luc and RM-9-Luc; luciferase L 4899 obtained from Sigma-Aldrich Japan G.K.) were generated by transfecting MBT-2 and RM-9 cells with the pGL 3-Luc plasmid using a TransIT®-3T3 transfection kit (Mirus Bio LLC, Madison, WI, USA). Cells that stably expressed luciferase were obtained by selection with 500 $\mu\text{g}/\text{mL}$ of G418 for two weeks. Following G418 selection, growth medium from MBT-2-Luc and RM-9-Luc was tested for luciferase activity to confirm the expression and secretion of luciferase into the cell medium.

[0131] R at anti-mouse PD-1 mAb (RM P1-14; IgG2a) was purchased from BioX Cell (West Lebanon, NH, USA).

[0132] Animals: Eight-week-old female C3H/HeN and C57/BL 6 mice were obtained from vendors. The mice were maintained at the animal facility in a specific pathogen-free environment with food and water provided ad libitum.

3.2 In vitro Assays

[0133] An MTT assay was used as the test method. An appropriate amount of cells ($2 \times 10^3/100 \mu\text{L}$) was seeded into 96-well plate and incubated at 37°C in a CO_2 incubator overnight for cell adhesion and adaptation. Cancer cells were also seeded in a separate 96-well plate (plate T0), 12 wells/cell line, to measure the OD value at time zero (T0). After overnight adaptation, 20 μL MTT reagent were added per well (final concentration 0.5 mg/mL) to plate T0 and incubated at 37°C for 4 hours. The supernatant medium was pipetted out, and about 150 μL DMSO were added per well. The plate was read on a plate reader using 550 nm as the test wavelength and 630 nm as the reference wavelength to obtain the OD value at time zero (T0). After overnight adaptation, the test compounds and vehicle were added into the plate to treat the cells. The test compounds were tested at 6 concentrations in triplicate wells. The plate was incubated at 37°C in a CO_2 incubator for 48 h. A longer incubation time can be used. 20 μL MTT reagent were added per well and incubated at 37°C for 4 h. The supernatant medium was pipetted out, and about 150 μL DMSO were added per well. The plate was read on a plate reader (TECAN, Infinite M200) using about 550 nm as the test wavelength and about 630 nm as the reference wavelength. Cell viability (%) was calculated in two ways: [0134] % Viability = $[(T-B)/(C-B)] \times 100\%$ [0135] % Viability = $[(T-C_{\text{sub.0}})/(C-C_{\text{sub.0}})] \times 100\%$ [0136] T: Mean Absorbance of Treatment at various hours; [0137] C: Mean Absorbance of negative Control at various hours; [0138] B: Mean Absorbance of blank well (only culture medium) at various hours; [0139] $C_{\text{sub.0}}$: Mean Absorbance of negative Control at 0 h.

[0140] The results are expressed as mean \pm SD. The IC_{50} was calculated by XL fit software. The test for each compound was independently repeated once.

3.3 In Vivo Effects on the Murine Bladder and Prostate Cancer Models.

[0141] To establish the orthotopic bladder cancer tumors, the mice were anesthetized by the

administration, .circle-solid.: MMC intravesical administration, Δ: CMC p.o, .box-tangle-solidup.: NIT p.o

[0144] For the NIT and anti-PD-1 combination, the mice carrying orthotopic bladder cancer tumors were randomly divided into four groups: control (PBS), NIT, anti-PD-1, and the combination of NIT and anti-PD-1. Detailed descriptions of the drug dosing are shown in Table 25.

TABLE-US-00025 TABLE 25 Drug dosing schematic. Day 1, Day 2, Day 3, Day 4, Day 5, Groups 6, 11 7, 12 8, 13 9, 14 10, 15 Group1: vehicle ΔΔ ΔΔ ΔΔ ΔΔ ○ Group 2: NIT .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. (bid,p.o15 mg/kg) ○ Group 3: anti-PD-1 ΔΔ ΔΔ ΔΔ ΔΔ (q5d, .circle-solid. intraperitoneal administration, 10 mg/kg) Group 4: .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. NIT + anti-PD-1 .circle-solid. ○: PBS intraperitoneal administration, .circle-solid.: anti-PD-1 intraperitoneal administration, Δ: CMC p.o, .box-tangle-solidup.: NIT p.o

[0145] To establish the orthotopic prostate cancer tumors, C57/BL 6 mice were intraperitoneally anesthetized with a ketamine/xylazine solution at a dose of 0.1 ml/10 g body weight (K 113; Sigma-Aldrich Japan G.K., Tokyo, Japan). A low abdominal transverse incision was made, and the bilateral dorsal lobes of prostate were exposed. Following the trypsinization of RM-9-Luc cells, 5.0×10^3 cells in 10 μ l of Hanks' balanced salt solution were injected using a new 24-gauge catheter directly into the right dorsal lobe of the prostate. A clearly recognizable bleb within the injected prostatic lobe was considered as a sign of a technically satisfactory injection. The abdominal wound was closed with stainless steel clips (Autoclip; Becton Dickinson Co., Sparks, MD). One week after the RM-9-Luc cell injection, when the tumor diameter reached 5 mm, the mice carrying orthotopic prostate cancer tumors were randomly divided into four groups: control (PBS), NIT, anti-PD-1, and the combination of NIT and anti-PD-1. Detailed descriptions of the drug dosing are shown in Table 26.

TABLE-US-00026 TABLE 26 Drug dosing schematic. Day Day Day Day 1, 5, 2, 6, 3, 7, 4, 8, Groups 9, 13 10, 14 11, 15 12, 16 Group1: vehicle ΔΔ ΔΔ ΔΔ ΔΔ ○ Group 2: NIT .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. (bid,p.o15 mg/kg) ○ Group 3: anti-PD-1 ΔΔ ΔΔ ΔΔ ΔΔ (q4d, .circle-solid. intraperitoneal administration, 10 mg/kg) SGroup 4: .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. .box-tangle-solidup..box-tangle-solidup. NIT + anti-PD-1 .circle-solid. ○: PBS intraperitoneal administration, .circle-solid.: anti-PD-1 intraperitoneal administration, Δ: CMC p.o, .box-tangle-solidup.: NIT p.o

[0146] The mice received i.p. injections of luciferin, and the luciferase expression in the tumors was measured by the Xenogen IVIS200 System. The mice were killed using CO.sub.2 for euthanasia and according to the guidelines for the euthanasia of animals (Edition, 2013).

3.4 Statistical Analysis.

[0147] Data from all quantitative assays are expressed as the mean±standard deviation and were analyzed statistically using a one-way analysis of variance (ANOVA) and the independent-samples t test. Statistical calculations were performed using GraphPad Prism 5. P values of less than 0.05 were considered statistically significant.

[0148] Additivity was determined using the fractional product concept or Bliss Independence model: $E_{sub.xy} = E_{sub.x} + E_{sub.y} - (E_{sub.x}E_{sub.y})$, where $E_{sub.xy}$ is the additive effect of the 2 compounds x and y as calculated by the product of the individual effect of the 2 compounds, $E_{sub.x}$ and $E_{sub.y}$. Synergy was established when the actual observed tumor inhibition value was greater than the expected tumor inhibition value determined by the Bliss Independence model (20).

REFERENCES

[0149] 1. Pelletier C, Prognon P, Bourlioux P Roles of divalent cations and pH in mechanism of action of nitroxoline against *Escherichia coli* strains. *Antimicrob Agents Chemother*. 1995;39(3):707-13. [0150] 2. Fraser RS, Creanor J. The mechanism of inhibition of ribonucleic acid synthesis by 8-hydroxyquinoline and the antibiotic lomofungin *Biochem J*. 1975;147(3):401-10. [0151] 3. Adlard PA et al. Rapid restoration of cognition in Alzheimer's transgenic mice with 8-hydroxy quinoline analogs is associated with decreased interstitial A-beta. *Neuron*. 2008;59(1):43-55. [0152] 4. Shim JS, et al. Effect of nitroxoline on angiogenesis and growth of human bladder cancer. *J Natl Cancer Inst*. 2010;102(24):1855-73. [0153] 5. Chang WL et al. Repurposing of nitroxoline as a potential anticancer agent against human prostate cancer: a crucial role on AM PK/mTOR signaling pathway and the interplay with Chk2 activation. *Oncotarget*. 2015; 6(37):39806-20. [0154] 6. Jiang H. et al. Nitroxoline (8-hydroxy-5-nitroquinoline) is more a potent anti-cancer agent than clioquinol (5-chloro-7-iodo-8-quinoline) *Cancer Letters* 2011; 312:11-17. [0155] 7. Ding WQ, Liu B, Vaught JL, Yamauchi H, and Lind SE. Anticancer activity of the antibiotic clioquinol. *Cancer Res* 2005; 65:3389-95. [0156] 8. Whiteside TL, Inhibiting the Inhibitors: Evaluating Agents Targeting Cancer Immuno-suppression. *Expert Opin Biol Ther*. 2010; 10(7): 1019-1035. [0157] 9. Fuge O, Vasdev N, Allchorne P, Green J S, Immunotherapy for bladder cancer. *Res Rep Urol*. 2015; 7:65-79. [0158] 10. Huang P, Ma C, Xu P, Guo K, Xu A, Liu C, Efficacy of intravesical *Bacillus Calmette-Guérin* therapy against tumor immune escape in an orthotopic model of bladder cancer. *Exp Ther Med*. 2015;9(1):162-166. [0159] 11. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer*. 2012; 12(4):252-64. [0160] 12. Krummel MF, Allison J P. CD28 and CTLA-4 have opposing effects on the response of T cells to stimulation. *J Exp Med*. 1995; 182(2):459-65. [0161] 13. Hamid O, Robert C, Daud A, Hodi FS, Hwu WJ, Kefford R. et al. Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma. *N Engl J Med*. 2013;369(2):134-44. [0162] 14. Freeman GJ, Long AJ, Iwai Y et al. Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med*. 2000; 192:1027-1034. [0163] 15. Hawkes EA, Grigg A, Chong G. Programmed cell death-1 inhibition in lymphoma. *Lancet Oncol*. 2015;16(5):e234-45. [0164] 16. Liu X, Shin N, Koblisch HK, et al. Selective inhibition of IDO1 effectively regulates mediators of antitumor immunity. *Blood*. 2010;115:3520-3530. [0165] 17. Opitz CA, Litzenburger UM, Opitz U, et al. The indoleamine-2,3-dioxygenase (IDO) inhibitor 1-methyl-D-tryptophan upregulates IDO1 in human cancer cells. *PloS One*. 2011; 6:e19823. [0166] 18. Muller AJ, Prendergast GC. Marrying immunotherapy with chemotherapy: why say IDO? *Cancer Res*. 2005; 65:8065-8068. [0167] 19. Ting-Chao Chou, Theoretical Basis, Experimental Design, and Computerized Simulation of Synergism and Antagonism in Drug Combination Studies. *Pharmacol Rev* 2006; 58:621-681. [0168] 20. Yan H, Zhang B, Li S, Zhao Q. A formal model for analyzing drug combination effects and its application in TNF-alpha-induced NFkappaB pathway. *BMC Syst Biol*. 2010;4:50.

Claims

1.-15. (canceled)

16. A method of treating urothelial carcinoma comprising administering to a subject in need thereof an effective amount of nitroxoline, an analogue or pharmaceutically acceptable salt thereof, and an effective amount of a second agent, wherein the combination of nitroxoline, the analogue or pharmaceutically acceptable salt thereof and the second agent results in a synergistic effect, wherein the analogue is selected from the group consisting of oxyquinoline, clioquinol, and iodoquinol, and wherein the second agent is an immunotherapy agent or a chemotherapeutic agent.

17. The method of claim 16, wherein the second agent is an immunotherapy agent.

18. The method of claim 17, wherein the immunotherapy agent is an anti-PD-1 (programmed cell death protein 1) antibody or an antigen binding fragment thereof, an anti-PD-L1 (programmed cell

death ligand 1) antibody or an antigen binding fragment thereof, an anti-CTLA-4 (targeting cytotoxic T lymphocyte-associated antigen 4) antibody or an antigen binding fragment thereof, or Bacillus Calmette-Guerin (BCG).

19. The method of claim 18, wherein the immunotherapy agent is an anti-PD-1 antibody or an antigen binding fragment thereof.

20. The method of claim 19, wherein the effective amount of nitroxoline, the analogue or pharmaceutically acceptable salt thereof is 100 mg to 1600 mg administered orally per day, and the effective amount of the anti-PD-1 antibody or antigen binding fragment thereof is 1 mg/kg to 20 mg/kg administered intravenously every 1 to 4 weeks.

21. The method of claim 18, wherein the immunotherapy agent is an anti-PD-L1 antibody or an antigen binding fragment thereof.

22. The method of claim 21, wherein the effective amount of nitroxoline, the analogue or pharmaceutically acceptable salt thereof is 100 mg to 1600 mg administered orally per day, and the effective amount of the anti-PD-L1 antibody or antigen binding fragment thereof is 4 mg/kg to 40 mg/kg administered intravenously every 1 to 4 weeks.

23. The method of claim 18, wherein the immunotherapy agent is an anti-CTLA-4 antibody or an antigen binding fragment thereof.

24. The method of claim 23, wherein the effective amount of nitroxoline, the analogue or pharmaceutically acceptable salt thereof is 100 mg to 1600 mg administered orally per day, and the effective amount of the anti-CTLA-4 antibody or antigen binding fragment thereof is 0.6 mg/kg to 6 mg/kg administered intravenously every 1 to 4 weeks.

25. The method of claim 18, wherein the immunotherapy agent comprises Bacillus Calmette-Guerin (BCG) therapy.

26. The method of claim 25, wherein the effective amount of nitroxoline, the analogue or pharmaceutically acceptable salt thereof is 100 mg to 1600 mg administered orally per day, and the effective amount of the BCG therapy is 0.5×10^8 to 50×10^8 colony forming units (CFUs) administered intravesically every 1 to 4 weeks.

27. The method of claim 16 wherein the second agent is a chemotherapeutic agent.

28. The method of claim 27, wherein the chemotherapeutic agent is selected from the group consisting of paclitaxel, docetaxel, cabazitaxel, mitomycin C, epirubicin, pirarubicin, carboplatin, cisplatin, oxaliplatin and nedaplatin.

29. The method of claim 28, wherein the chemotherapeutic agent is administered intravesically.

30. The method of claim 16, wherein the nitroxoline, analogue, or pharmaceutically acceptable salt thereof and the chemotherapeutic agent are administered by way of the simultaneous, sequential or separate administration.

31. The method of claim 28, wherein the effective amount of nitroxoline, the analogue or pharmaceutically acceptable salt thereof is 100 mg to 1600 mg administered orally per day, and mitomycin C, epirubicin, pirarubicin, carboplatin, cisplatin, oxaliplatin or nedaplatin is administered intravesically in a bladder instillation fluid at a concentration of 0.1 mg/mL to 5 mg/mL.

32. The method of claim 28, wherein the effective amount of nitroxoline, the analogue or pharmaceutically acceptable salt thereof is 100 mg to 1600 mg administered orally per day, and paclitaxel, docetaxel or cabazitaxel is administered intravesically in a bladder instillation fluid at a concentration of 0.5 mg/mL to 10 mg/mL.

33. A composition comprising an effective amount of nitroxoline, an analogue, or a pharmaceutically acceptable salt thereof, an effective amount of a chemotherapeutic agent, and one or more pharmaceutically acceptable carriers, wherein the analogue is selected from the group consisting of oxyquinoline, clioquinol, and iodoquinol, and wherein the chemotherapeutic agent is selected from the group consisting of mitomycin C, epirubicin, pirarubicin, carboplatin, cisplatin, oxaliplatin and nedaplatin.

