

# initial exploration

Stuart Barnum

8/29/2022

```
knitr::opts_chunk$set(echo = TRUE)
```

```
library(tidyverse)
```

```
## -- Attaching packages ----- tidyverse 1.3.2 --
## v ggplot2 3.3.6      v purrr  0.3.4
## v tibble  3.1.8      v dplyr  1.0.9
## v tidyr   1.2.0      v stringr 1.4.1
## v readr   2.1.2      v forcats 0.5.2
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
```

```
library(lubridate)
```

```
##
## Attaching package: 'lubridate'
##
## The following objects are masked from 'package:base':
##
##     date, intersect, setdiff, union
```

```
library(funtimes)
```

```
library(boot)
```

```
historic <- read_csv("NYPD_Arrests_Data__Historic.csv",
                     guess_max = Inf)
```

```
## Rows: 5308876 Columns: 19
## -- Column specification -----
## Delimiter: ","
## chr (10): ARREST_DATE, PD_DESC, OFNS_DESC, LAW_CODE, LAW_CAT_CD, ARREST_BORO...
## dbl (9): ARREST_KEY, PD_CD, KY_CD, ARREST_PRECINCT, JURISDICTION_CODE, X_CO...
##
## i Use `spec()` to retrieve the full column specification for this data.
## i Specify the column types or set `show_col_types = FALSE` to quiet this message.
```

```
summary(historic)
```

```
##   ARREST_KEY      ARREST_DATE      PD_CD      PD_DESC
## Min.   : 9926901 Length:5308876 Min.   : 0.0 Length:5308876
## 1st Qu.: 61436632 Class :character 1st Qu.:269.0 Class :character
## Median : 85671028 Mode  :character Median :511.0 Mode  :character
## Mean   :102879939          Mean   :505.8
## 3rd Qu.:150090000          3rd Qu.:748.0
```

```
## Max. :238513928 Max. :997.0
## NA's :313
## KY_CD OFNS_DESC LAW_CODE LAW_CAT_CD
## Min. :101.0 Length:5308876 Length:5308876 Length:5308876
## 1st Qu.:126.0 Class :character Class :character Class :character
## Median :341.0 Mode :character Mode :character Mode :character
## Mean :298.4
## 3rd Qu.:348.0
## Max. :995.0
## NA's :9169
## ARREST_BORO ARREST_PRECINCT JURISDICTION_CODE AGE_GROUP
## Length:5308876 Min. : 1.00 Min. : 0.000 Length:5308876
## Class :character 1st Qu.: 33.00 1st Qu.: 0.000 Class :character
## Mode :character Median : 60.00 Median : 0.000 Mode :character
## Mean : 60.76 Mean : 1.296
## 3rd Qu.: 84.00 3rd Qu.: 0.000
## Max. :123.00 Max. :97.000
## NA's :10
## PERP_SEX PERP_RACE X_COORD_CD Y_COORD_CD
## Length:5308876 Length:5308876 Min. : 913357 Min. : 121131
## Class :character Class :character 1st Qu.: 993280 1st Qu.: 186857
## Mode :character Mode :character Median :1004892 Median : 209285
## Mean :1005355 Mean : 214587
## 3rd Qu.:1015924 3rd Qu.: 236614
## Max. :1067302 Max. :8202360
## NA's :1 NA's :1
## Latitude Longitude Lon_Lat
## Min. :40.50 Min. : -74.25 Length:5308876
## 1st Qu.:40.68 1st Qu.: -73.97 Class :character
## Median :40.74 Median : -73.93 Mode :character
## Mean :40.76 Mean : -73.92
## 3rd Qu.:40.82 3rd Qu.: -73.89
## Max. :62.08 Max. : -73.68
## NA's :1 NA's :1
```

```
historic_w_date <- historic %>%
  mutate(Date = as.Date(ARREST_DATE, "%m/%d/%Y"))
summary(historic_w_date)
```

```
## ARREST_KEY ARREST_DATE PD_CD PD_DESC
## Min. : 9926901 Length:5308876 Min. : 0.0 Length:5308876
## 1st Qu.: 61436632 Class :character 1st Qu.:269.0 Class :character
## Median : 85671028 Mode :character Median :511.0 Mode :character
## Mean :102879939 Mean :505.8
## 3rd Qu.:150090000 3rd Qu.:748.0
## Max. :238513928 Max. :997.0
## NA's :313
## KY_CD OFNS_DESC LAW_CODE LAW_CAT_CD
## Min. :101.0 Length:5308876 Length:5308876 Length:5308876
## 1st Qu.:126.0 Class :character Class :character Class :character
## Median :341.0 Mode :character Mode :character Mode :character
## Mean :298.4
## 3rd Qu.:348.0
## Max. :995.0
## NA's :9169
```

```
## ARREST_BORO      ARREST_PRECINCT  JURISDICTION_CODE  AGE_GROUP
## Length:5308876   Min.      : 1.00    Min.      : 0.000   Length:5308876
## Class :character  1st Qu.: 33.00    1st Qu.: 0.000   Class :character
## Mode  :character  Median   : 60.00    Median   : 0.000   Mode   :character
##                  Mean    : 60.76    Mean    : 1.296
##                  3rd Qu.: 84.00    3rd Qu.: 0.000
##                  Max.    :123.00    Max.    :97.000
##                  NA's    :10
## PERP_SEX          PERP_RACE          X_COORD_CD          Y_COORD_CD
## Length:5308876    Length:5308876    Min.      : 913357   Min.      : 121131
## Class :character  Class :character  1st Qu.: 993280    1st Qu.: 186857
## Mode  :character  Mode  :character  Median :1004892    Median : 209285
##                  Mean    :1005355    Mean    : 214587
##                  3rd Qu.:1015924    3rd Qu.: 236614
##                  Max.    :1067302    Max.    :8202360
##                  NA's    :1          NA's    :1
## Latitude          Longitude          Lon_Lat          Date
## Min.      :40.50   Min.      : -74.25   Length:5308876    Min.      :2006-01-01
## 1st Qu.:40.68   1st Qu.: -73.97   Class :character  1st Qu.:2009-05-05
## Median :40.74   Median : -73.93   Mode  :character  Median :2012-07-10
## Mean    :40.76   Mean    : -73.92                    Mean    :2012-11-10
## 3rd Qu.:40.82   3rd Qu.: -73.89                    3rd Qu.:2016-02-04
## Max.    :62.08   Max.    : -73.68                    Max.    :2021-12-31
## NA's    :1      NA's    :1
```

```
year_to_date <- read_csv("NYPD_Arrest_Data_Year_to_Date_.csv") %>%
  mutate(Date = as.Date(ARREST_DATE, "%m/%d/%Y"))
```

```
## Rows: 93238 Columns: 19
## -- Column specification -----
## Delimiter: ","
## chr (10): ARREST_DATE, PD_DESC, OFNS_DESC, LAW_CODE, LAW_CAT_CD, ARREST_BORO...
## dbl (9): ARREST_KEY, PD_CD, KY_CD, ARREST_PRECINCT, JURISDICTION_CODE, X_CO...
##
## i Use `spec()` to retrieve the full column specification for this data.
## i Specify the column types or set `show_col_types = FALSE` to quiet this message.
```

```
all_arrests <- bind_rows(historic_w_date, year_to_date)
```

```
summary(all_arrests)
```

```
## ARREST_KEY      ARREST_DATE      PD_CD      PD_DESC
## Min.      : 9926901   Length:5402114   Min.      : 0.0   Length:5402114
## 1st Qu.: 62053042   Class :character  1st Qu.:268.0   Class :character
## Median : 86350562   Mode  :character  Median :510.0   Mode  :character
## Mean    :105300015                    Mean    :504.1
## 3rd Qu.:152357173                    3rd Qu.:748.0
## Max.    :247417454                    Max.    :997.0
##                  NA's    :546
## KY_CD          OFNS_DESC          LAW_CODE          LAW_CAT_CD
## Min.      :101.0   Length:5402114   Length:5402114   Length:5402114
## 1st Qu.:126.0   Class :character  Class :character  Class :character
## Median :341.0   Mode  :character  Mode  :character  Mode  :character
## Mean    :297.5
## 3rd Qu.:348.0
```

```
## Max.      :995.0
## NA's      :9473
## ARREST_BORO      ARREST_PRECINCT JURISDICTION_CODE AGE_GROUP
## Length:5402114   Min.      : 1.0   Min.      : 0.00   Length:5402114
## Class :character 1st Qu.: 33.0   1st Qu.: 0.00   Class :character
## Mode  :character Median : 60.0   Median : 0.00   Mode  :character
##                Mean  : 60.8   Mean  : 1.29
##                3rd Qu.: 84.0   3rd Qu.: 0.00
##                Max.   :123.0   Max.   :97.00
##                NA's   :10
## PERP_SEX          PERP_RACE          X_COORD_CD          Y_COORD_CD
## Length:5402114   Length:5402114   Min.      : 913357   Min.      : 121131
## Class :character Class :character 1st Qu.: 993212   1st Qu.: 186857
## Mode  :character Mode  :character Median :1004892   Median : 209223
##                Mean  :1005347   Mean  : 214481
##                3rd Qu.:1015947   3rd Qu.: 236608
##                Max.   :1067302   Max.   :8202360
##                NA's   :1        NA's   :1
## Latitude          Longitude          Lon_Lat          Date
## Min.      :40.50   Min.      : -74.25   Length:5402114   Min.      :2006-01-01
## 1st Qu.:40.68   1st Qu.: -73.97   Class :character 1st Qu.:2009-05-22
## Median :40.74   Median : -73.93   Mode  :character Median :2012-08-20
## Mean  :40.76   Mean  : -73.92
## 3rd Qu.:40.82   3rd Qu.: -73.89
## Max.   :62.08   Max.   : -73.68
## NA's   :1        NA's   :1
## New Georeferenced Column
## Length:5402114
## Class :character
## Mode  :character
##
##
##
##
```

## Question 1


Has the arrest rate been decreasing from 2018-2022? Describe the trend and

defend any statistical tests used to support this conclusion.

The arrest record has not decreased from 2018. It decreased from 2018 through mid 2020 and then increased thereafter.

I begin by having quick look at counts for 30 day periods for the entire period from 2006. I use all of the years, for now, to perhaps uncover cyclical affects (e.g. over each year) that may be distorted by COVID.

```
ggplot(all_arrests) + geom_histogram(aes(Date), binwidth=30) +
  labs(y = "Number of arrests", title="Count totals from 2006 by 30-day periods") +
  theme_bw()
```




Find arrest counts for each day, to facilitate possible modeling (in the end, I used counts by week in my analysis). Have a quick look, noting that the seeming cyclical effects, though somewhat evident for earlier years, are now more difficult to discern.

```
arrest_day_counts <-
  all_arrests %>%
  group_by(Date) %>%
  summarize(daily_arrest_count = n())

#plot the daily counts together with a loess-smoothed curve
ggplot(arrest_day_counts, aes(x=Date, y = daily_arrest_count)) +
  geom_smooth(span = .2) + geom_point(alpha = .15)+
  theme_bw()

## `geom_smooth()` using method = 'gam' and formula 'y ~ s(x, bs = "cs")'
```




I might expect to see differences in week days, with e.g. more arrests of certain types on weekends. I will bin the weeks, obtaining the count for each week and assigning these counts to the median day of the week (Wednesday). With this, we might better visualize trends over time periods greater than a few weeks.

```
week_counts <- all_arrests %>%
  mutate(week_floor = floor_date(Date, unit="weeks"),
         week_median_date = week_floor + 3) %>%
  group_by(week_median_date) %>%
  summarize(week_count = n())

ggplot(week_counts, aes(week_median_date, week_count)) +
  geom_point(alpha = .25) +
  geom_smooth(method = "loess", span = .1) +
  scale_x_date(breaks = '1 year') +
  theme(axis.text.x = element_text(angle = 90)) +
  labs(title = "Weekly counts from 2006, with smoothing showing seasonal\n variation in earlier years")

## `geom_smooth()` using formula 'y ~ x'
```

Weekly counts from 2006, with smoothing showing seasonal variation in earlier years




```
week_counts_fm_2018 <- week_counts %>%
  filter(week_median_date >= ymd("2018-01-01"))

ggplot(week_counts_fm_2018, aes(week_median_date, week_count)) +
  geom_point(alpha = .3) +
  geom_smooth(method = "loess", span = .15) +
  theme_bw() +
  scale_x_date(breaks = '1 year') +
  theme(axis.text.x = element_text(angle = 90)) +
  labs(title = "Weekly arrests from 2018, with smoothing showing seasonal\n variation, perhaps distorted",
       x = "Median week date (the Wednesday)",
       y = "Weekly number of arrests")

## `geom_smooth()` using formula 'y ~ x'
```

Weekly arrests from 2018, with smoothing showing seasonal variation, perhaps distorted due to COVID and other aspects




It appears that arrests decreased until the middle of 2020 and then started to increase. I will divide the data into these two time periods and apply a Mann-Kendall test with sieve bootstrap to the two time series resulting from separation of weeks before middle 2020 and weeks after middle 2020. This test allows that data be autocorrelated and that there be periodicity, which seems to exist, even if it is rather irregular.

As a first step, I put the relevant data into a time series R object and create a quick plot of the results


```
#these two results might be used with daily counts
first_obs <- week_counts_fm_2018 %>%
  filter(week_median_date==min(week_median_date))
last_obs <- week_counts_fm_2018 %>%
  filter(week_median_date==max(week_median_date))

#the times series object for weekly counts
timeseries_both <- ts(week_counts_fm_2018$week_count,
  start = c(2018, 1),
  frequency = 52)
plot(timeseries_both)
```






```
timeseries_earlier <- window(timeseries_both,  
                             end = c(2020, 26))  
plot(timeseries_earlier)
```



```
timeseries_later <- window(timeseries_both,  
                           start = c(2020, 26))  
plot(timeseries_later)
```



Now do the statistical significance tests on upward and downward trends for the respective time period. The test is for no trend (the null hypothesis versus some trend)

```
notrend_test(timeseries_earlier, B=1000, test='MK')
```

```
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: timeseries_earlier
## Mann--Kendall's tau = -0.6877, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 1
##
## $AR_coefficients
## phi_1
## 0.6165258
```

```
notrend_test(timeseries_later, B=1000, test='MK')
```

```
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: timeseries_later
## Mann--Kendall's tau = 0.54822, p-value = 0.002
## alternative hypothesis: monotonic trend.
## sample estimates:
```

```
## $AR_order
## [1] 4
##
## $AR_coefficients
##      phi_1      phi_2      phi_3      phi_4
## 0.66380318 -0.24681672  0.09182166  0.16597143
```

## Question 2

What are the top 5 most frequent arrests as described in the column 'pd\_desc' in 2018-2022? Compare & describe the overall trends of these arrests across time.

Moving on the Question 2, we count numbers of arrests by 'pd\_desc' for all weeks since 2018, and then subset the arrests dataset to list only the arrests within pd\_desc categories with the top 5 counts.

```
pd_desc_counts_top5 <- all_arrests %>%
  filter(Date >= ymd('2018-01-01')) %>%
  count(PD_DESC) %>%
  arrange(desc(n)) %>%
  top_n(5)
```

```
## Selecting by n
```

```
print(pd_desc_counts_top5)
```

```
## # A tibble: 5 x 2
##   PD_DESC                                n
##   <chr>                                <int>
## 1 ASSAULT 3                            99757
## 2 LARCENY,PETIT FROM OPEN AREAS, 56001
## 3 ASSAULT 2,1,UNCLASSIFIED            51694
## 4 TRAFFIC,UNCLASSIFIED MISDEMEAN 40527
## 5 ROBBERY,OPEN AREA UNCLASSIFIED 29773
```

```
pd_desc_subset <- all_arrests %>%
  filter(Date >= ymd('2018-01-01')) %>%
  inner_join(pd_desc_counts_top5, by="PD_DESC")
```

```
pd_desc_counts <- pd_desc_subset %>%
  mutate(week_floor = floor_date(Date, unit="weeks"),
         week_median_date = week_floor + 3) %>%
  group_by(PD_DESC, week_median_date) %>%
  summarize(week_count = n())
```


```
## `summarise()` has grouped output by 'PD_DESC'. You can override using the
## `.groups` argument.
```

Visualizing the apparent trend, for the 5 categories with the highest count, we see similar trends as with the overall counts—decreasing until about middle 2020 and then increasing. Except that at least one of the crime categories shows week trends or perhaps no statistically significant trend at all. We check the statistical significance using the same test as with Question 1. Note that two of the categories show no 2018 data. These may be new or renamed categories (I have not explored this).

```
ggplot(pd_desc_counts, aes(x=week_median_date, y=week_count, color=PD_DESC)) +
  geom_point(alpha = .25) +
  geom_smooth(method = "loess", span = .2) +
  theme_bw() +
```

```
scale_x_date(breaks = '1 year') +
theme(axis.text.x = element_text(angle = 45, hjust = 1)) +
labs(title = "Top 5 crime categories from 2018, with smoothing\n showing trends similar to the overall.",
x = "Median date of week",
y = "Number of arrests")
```

```
## `geom_smooth()` using formula 'y ~ x'
```



Put the counts for the 5 categories into separate variables and then do the statistical tests for the downward trend before mid 2020 and the upward trend after mid 2020

```
top5wide <- pd_desc_counts %>%
  pivot_wider(names_from = PD_DESC,
              values_from = week_count)
```

```
top5wide %>% head()
```

```
## # A tibble: 6 x 6
##   week_median_date `ASSAULT 2,1,UNCLASSIFIED` ASSAULT ~1 LARCE~2 ROBBE~3 TRAFF~4
##   <date>          <int>          <int>    <int>    <int>    <int>
## 1 2018-01-03      166          376      NA      NA      215
## 2 2018-01-10      209          496      NA      NA      386
## 3 2018-01-17      221          474      NA      NA      458
## 4 2018-01-24      211          556      NA      NA      496
## 5 2018-01-31      237          497      NA      NA      394
## 6 2018-02-07      210          504      NA      NA      418
## # ... with abbreviated variable names 1: `ASSAULT 3`,
```

```
## # 2: `LARCENY,PETIT FROM OPEN AREAS`,` 3: `ROBBERY,OPEN AREA UNCLASSIFIED`,
## # 4: `TRAFFIC,UNCLASSIFIED MISDEMEAN`
```

```
for (i in 2:6){

  print(str_c("Column ", i))

  ts <- ts(top5wide[,i],
           start = c(2018, 1),
           frequency = 52)

  if (i %in% 4:5) {
    ts_early <- window(ts,
                       start = c(2019, 1),
                       end = c(2020, 26))
  } else {
    ts_early <- window(ts,
                       end = c(2020, 26))
  }

  print("Earlier Period")
  print(notrend_test(ts_early, B=1000, test='MK'))

  print("Later Period")
  ts_late <- window(ts,
                    start = c(2020, 26))
  print(notrend_test(ts_late, B=1000, test='MK'))
}
```

```
## [1] "Column 2"
## [1] "Earlier Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_early
## Mann--Kendall's tau = -0.20504, p-value = 0.022
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 1
##
## $AR_coefficients
## phi_1
## 0.4007805
##
##
## [1] "Later Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_late
## Mann--Kendall's tau = 0.47345, p-value = 0.002
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
```

```

## [1] 4
##
## $AR_coefficients
##      phi_1      phi_2      phi_3      phi_4
## 0.29067871 0.08557285 -0.13506349 0.30474540
##
##
## [1] "Column 3"
## [1] "Earlier Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_early
## Mann--Kendall's tau = -0.51921, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 2
##
## $AR_coefficients
##      phi_1      phi_2
## 0.3195167 0.2159243
##
##
## [1] "Later Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_late
## Mann--Kendall's tau = 0.58399, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 4
##
## $AR_coefficients
##      phi_1      phi_2      phi_3      phi_4
## 0.287532895 0.005090214 0.080753378 0.115167169
##
##
## [1] "Column 4"
## [1] "Earlier Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_early
## Mann--Kendall's tau = -0.57969, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 2
##
## $AR_coefficients
##      phi_1      phi_2

```

```

## 0.56170486 0.04507668
##
##
## [1] "Later Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_late
## Mann--Kendall's tau = 0.7002, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 4
##
## $AR_coefficients
##      phi_1      phi_2      phi_3      phi_4
## 0.52260456 0.04355166 0.02173111 0.03941950
##
##
## [1] "Column 5"
## [1] "Earlier Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_early
## Mann--Kendall's tau = -0.0096958, p-value = 0.959
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 1
##
## $AR_coefficients
##      phi_1
## 0.5378248
##
##
## [1] "Later Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_late
## Mann--Kendall's tau = 0.41491, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 1
##
## $AR_coefficients
##      phi_1
## 0.1379164
##
##
## [1] "Column 6"
## [1] "Earlier Period"

```



```
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_early
## Mann--Kendall's tau = -0.5449, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 4
##
## $AR_coefficients
##      phi_1      phi_2      phi_3      phi_4
## 0.65124765 0.12845373 -0.13398382 0.07014356
##
##
## [1] "Later Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_late
## Mann--Kendall's tau = 0.43043, p-value = 0.001
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 3
##
## $AR_coefficients
##      phi_1      phi_2      phi_3
## 0.520254352 0.002511877 0.060611831
```

### Question 3

If we think of arrests as a sample of total crime, is there more crime in precinct 19 (Upper East Side) than precinct 73 (Brownsville)? Describe the trend, variability and justify any statistical tests used to support this conclusion.

This question asks us to compare crime rates, not just trends in crime rates. It can be misleading to say that one area has more crime than another area, without consideration of the population of those areas. I therefore found some population numbers and compared the per capital crime rates.

My sources:

<https://johnkeefe.net/nyc-police-precinct-and-census-data>

[https://github.com/jkeefe/census-by-precincts/blob/master/data/nyc/DECENNIALPL2020.P1\\_metadata\\_2022-02-06T162722.csv](https://github.com/jkeefe/census-by-precincts/blob/master/data/nyc/DECENNIALPL2020.P1_metadata_2022-02-06T162722.csv)

Another tricky aspect with Question 3 is the seeming autocorrelation (over time) in the weekly crime rates. Most statistical tests that would compare, say, the average weekly crime rates of these areas require independence of the data points. This condition fails. I use bootstrapping to partially ameliorate this issue. I am not certain that my solution is ideal.

```
precinct_counts <- all_arrests %>%
  filter(ARREST_PRECINCT %in% c(19, 73)) %>%
  mutate(precinct = case_when(ARREST_PRECINCT == 19 ~ "Upper East Side",
                              ARREST_PRECINCT == 73 ~ "Brownsville")) %>%
```


```
mutate(week_floor = floor_date(Date, unit="weeks"),
       week_median_date = week_floor + 3) %>%
group_by(precinct, week_median_date) %>%
summarize(week_count = n())
```

## `summarise()` has grouped output by 'precinct'. You can override using the  
## `.groups` argument.

```
ggplot(precinct_counts, aes(x=week_median_date, y=week_count, color=precinct)) +
  geom_point(alpha = .25) +
  geom_smooth(method = "loess", span = .2) +
  scale_x_date(breaks = '1 year') +
  theme(axis.text.x = element_text(angle = 45, hjust = 1)) +
  labs(title = "Two precinct from 2006, with smoothing showing trends\n similar to the overall arrest t
```

## `geom\_smooth()` using formula 'y ~ x'

Two precinct from 2006, with smoothing showing trends  
similar to the overall arrest trend




```
precinct_counts_fm_2018 <- precinct_counts %>%
  filter(week_median_date >= ymd("2018-01-01"))

ggplot(precinct_counts_fm_2018, aes(x=week_median_date, y=week_count, color=precinct)) +
  geom_point(alpha = .25) +
  geom_smooth(method = "loess", span = .2) +
  theme_bw() +
  scale_x_date(breaks = '1 year') +
  theme(axis.text.x = element_text(angle = 45, hjust = 1)) +
  labs(title = "Two precincts from 2018, with smoothing showing trends\n similar to the overall arrest t
```

```
x = "Median date of week",
y = "Number of arrests")
```

```
## `geom_smooth()` using formula 'y ~ x'
```

Two precincts from 2018, with smoothing showing trends similar to the overall arrest trend



```
precinct_wide <- precinct_counts_fm_2018 %>%
  pivot_wider(names_from = precinct,
              values_from = week_count)
```

```
precinct_wide %>% head()
```

```
## # A tibble: 6 x 3
##   week_median_date Brownsville `Upper East Side`
##   <date>           <int>           <int>
## 1 2018-01-03         89             47
## 2 2018-01-10        111             58
## 3 2018-01-17        153             57
## 4 2018-01-24        151             50
## 5 2018-01-31        120             50
## 6 2018-02-07        115             55
```

```
for (i in 2:3){

  print(str_c("Column ", i))

  ts <- ts(precinct_wide[,i],
          start = c(2018, 1),
```

```

        frequency = 52)

ts_early <- window(ts,
                    end = c(2020, 26))

print("Earlier Period")
print(notrend_test(ts_early, B=1000, test='MK'))

print("Later Period")
ts_late <- window(ts,
                  start = c(2020, 26))
print(notrend_test(ts_late, B=1000, test='MK'))
}

## [1] "Column 2"
## [1] "Earlier Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data:  ts_early
## Mann--Kendall's tau = -0.60437, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 4
##
## $AR_coefficients
##      phi_1      phi_2      phi_3      phi_4
## 0.27209432 0.02793462 -0.13415858 0.17738374
##
## [1] "Later Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data:  ts_late
## Mann--Kendall's tau = 0.35488, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 1
##
## $AR_coefficients
##      phi_1
## 0.2825638
##
## [1] "Column 3"
## [1] "Earlier Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data:  ts_early
## Mann--Kendall's tau = -0.17689, p-value = 0.04

```

```
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 2
##
## $AR_coefficients
##      phi_1      phi_2
## 0.2056698 0.1416509
##
##
## [1] "Later Period"
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_ate
## Mann--Kendall's tau = 0.33838, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 3
##
## $AR_coefficients
##      phi_1      phi_2      phi_3
## 0.08875647 -0.07324782 0.23561994

john_keefe_data = read_csv("nyc_precinct_2020pop (1).txt")

## Rows: 77 Columns: 145
## -- Column specification -----
## Delimiter: ","
## dbl (145): precinct, P1_001N, P1_002N, P1_003N, P1_004N, P1_005N, P1_006N, P...
##
## i Use `spec()` to retrieve the full column specification for this data.
## i Specify the column types or set `show_col_types = FALSE` to quiet this message.
```

Look at the arrests per capita for the two precincts. From the Keefe data, per the 2020 census, Precinct 19 has population 220,261, while Precinct 73 has population 98,506. I will assume that these numbers are roughly accurate from 2018 to the present.

```
summary(precinct_wide)
```

```
## week_median_date      Brownsville      Upper East Side
## Min. :2018-01-03 Min. : 21.00 Min. : 9.00
## 1st Qu.:2019-02-16 1st Qu.: 51.00 1st Qu.:28.00
## Median :2020-04-01 Median : 71.00 Median :37.00
## Mean :2020-04-01 Mean : 72.63 Mean :37.77
## 3rd Qu.:2021-05-15 3rd Qu.: 91.00 3rd Qu.:47.00
## Max. :2022-06-29 Max. :166.00 Max. :76.00
```

```
#arrests per capita as PERCENT
```

```
precinct_wide_adjusted <- precinct_wide %>%
  mutate(`Upper East Side per capita` = 100*`Upper East Side` / 220261,
         `Brownsville per capita` = 100 * Brownsville / 98506)
```

```
#make long to facilitate visualization with ggplot
```

```
precinct_long_adusted <- precinct_wide_adjusted %>%
```

```

pivot_longer(cols = c(`Upper East Side per capita`,
                      `Brownsville per capita`),
             names_to = "Precinct",
             values_to = "Arrests per capita")


precinct_long_adusted %>% head()

## # A tibble: 6 x 5
##   week_median_date Brownsville `Upper East Side` Precinct      Arrests
##   <date>           <int>           <int> <chr>           <dbl>
## 1 2018-01-03         89             47 Upper East Side per ca~ 0.0213
## 2 2018-01-03         89             47 Brownsville per capita 0.0903
## 3 2018-01-10        111             58 Upper East Side per ca~ 0.0263
## 4 2018-01-10        111             58 Brownsville per capita 0.113
## 5 2018-01-17        153             57 Upper East Side per ca~ 0.0259
## 6 2018-01-17        153             57 Brownsville per capita 0.155
## # ... with abbreviated variable name 1: `Arrests per capita`

ggplot(precinct_long_adusted, aes(x=week_median_date, y=`Arrests per capita`, color=Precinct)) +
  geom_point(alpha = .25) +
  geom_smooth(method = "loess", span = .2) +
  theme_bw() +
  scale_x_date(breaks = '1 year') +
  theme(axis.text.x = element_text(angle = 45, hjust = 1)) +
  labs(title = "Arrests per capita for two precincts",
       x = "Median date of week",
       y = "Arrests per capita (%)") +
  scale_color_discrete(labels = c("Brownsville", "Upper East Side"))

## `geom_smooth()` using formula 'y ~ x'

```




Some analysis of the differences between the per capita arrest rates

```
percapita_difference <- precinct_wide_adjusted %>%
  mutate(difference = `Brownsville per capita` -
    `Upper East Side per capita`)

ggplot(percapita_difference, aes(x=week_median_date, y=difference)) +
  geom_point(alpha = .25) +
  geom_smooth(method = "loess", span = .2) +
  theme_bw() +
  scale_x_date(breaks = '1 year') +
  theme(axis.text.x = element_text(angle = 45, hjust = 1)) +
  labs(title = "Difference in weekly arrests per capita",
    subtitle = "Brownsville and Upper East Side",
    x = "Median date of week",
    y = "Arrests per capita difference (% points)") +
  scale_color_discrete(labels = c("Brownsville", "Upper East Side"))

## `geom_smooth()` using formula 'y ~ x'
```



As with my previous analysis, there are notable differences between the time period before mid 2020 and the period after mid 2020. I will separate the data as before and then look for both trends and for overall differences between the precincts. The analysis of the overall differences, such as whether the differences in weekly arrest rates, between the two precincts, is statistically significant is complicated by the fact that the observations for each week are not independent. Most statistical tests, including non-parametric tests, assume independence. I ameliorate this issue by using bootstrap methods—resampling data to be used in statistical tests. All of my analysis will use population-adjusted arrest rates and will focus on the differences between the two precincts.

The trend analysis, again using sieve bootstrap.

```
# analyze the earlier and later bits within R timeseries objects
```

```
ts <- ts(percapi_difference$difference,
        start = c(2018, 1),
        frequency = 52)
```


```
ts_early <- window(ts,
                  end = c(2020, 26))
```

```
print("Earlier Period")
```

```
## [1] "Earlier Period"
```

```
plot(ts_early)
```





```
print(notrend_test(ts_early, B=1000, test='MK'))
```


```
##
## Sieve-bootstrap Mann--Kendall's trend test
##
## data: ts_early
## Mann--Kendall's tau = -0.56858, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 4
##
## $AR_coefficients
##      phi_1      phi_2      phi_3      phi_4
## 0.16126208 0.07275539 -0.10805570 0.12628703
```

```
print("Later Period")
```

```
## [1] "Later Period"
```

```
ts_late <- window(ts,
                  start = c(2020, 26))
plot(ts_late, main="Difference in weekly per capita arrest rate",
      xlab = "Midpoint of week", ylab="Percentage point difference")
```

## Difference in weekly per capita arrest rate



```
print(notrend_test(ts_early, B=1000, test='MK'))
```

```
##
## Sieve-bootstrap Mann-Kendall's trend test
##
## data: ts_early
## Mann-Kendall's tau = 0.2671, p-value < 2.2e-16
## alternative hypothesis: monotonic trend.
## sample estimates:
## $AR_order
## [1] 1
##
## $AR_coefficients
## phi_1
## 0.1556195
```

We now look at the overall difference in the per capita arrest rates between the precincts, examining whether the differences in the weekly means are statistically significant.

```
early_differences <- percapita_difference[percapita_difference$week_median_date < ymd('2020-07-01'),]
early_differences %>% head()
```

```
## # A tibble: 6 x 6
##   week_median_date Brownsville `Upper East Side` Upper East Si-1 Brown-2 diffe-3
##   <date>           <int>           <int>           <dbl>   <dbl>   <dbl>
## 1 2018-01-03         89             47         0.0213  0.0903  0.0690
## 2 2018-01-10        111             58         0.0263  0.113   0.0864
```

```

## 3 2018-01-17          153          57          0.0259 0.155 0.129
## 4 2018-01-24          151          50          0.0227 0.153 0.131
## 5 2018-01-31          120          50          0.0227 0.122 0.0991
## 6 2018-02-07          115          55          0.0250 0.117 0.0918
## # ... with abbreviated variable names 1: `Upper East Side per capita`,
## # 2: `Brownsville per capita`, 3: difference

fc <- function(d, i){
  d2 <- d[i,]
  return(mean(d2$difference))
}

bootmean_early <- boot(early_differences, fc, R=1000)
boot.ci(bootmean_early, type = c('perc', 'bca'))

## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
## Based on 1000 bootstrap replicates
##
## CALL :
## boot.ci(boot.out = bootmean_early, type = c("perc", "bca"))
##
## Intervals :
## Level      Percentile      BCa
## 95%      ( 0.0656, 0.0738 ) ( 0.0658, 0.0741 )
## Calculations and Intervals on Original Scale

later_differences <- percapita_difference[percapita_difference$week_median_date >= ymd('2020-07-01'),]

later_differences %>% head()

## # A tibble: 6 x 6
##   week_median_date Brownsville `Upper East Side` Upper East Si~1 Brown~2 diffe~3
##   <date>          <int>          <int>          <dbl>    <dbl>    <dbl>
## 1 2020-07-01          30           9          0.00409 0.0305 0.0264
## 2 2020-07-08          30          11          0.00499 0.0305 0.0255
## 3 2020-07-15          30          14          0.00636 0.0305 0.0241
## 4 2020-07-22          21           9          0.00409 0.0213 0.0172
## 5 2020-07-29          36          21          0.00953 0.0365 0.0270
## 6 2020-08-05          45          27          0.0123 0.0457 0.0334
## # ... with abbreviated variable names 1: `Upper East Side per capita`,
## # 2: `Brownsville per capita`, 3: difference

bootmean_later <- boot(later_differences, fc, R=1000)
boot.ci(bootmean_later, type = c('perc', 'bca'))

## BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
## Based on 1000 bootstrap replicates
##
## CALL :
## boot.ci(boot.out = bootmean_later, type = c("perc", "bca"))
##
## Intervals :
## Level      Percentile      BCa
## 95%      ( 0.0379, 0.0432 ) ( 0.0380, 0.0433 )
## Calculations and Intervals on Original Scale

```

## Question 4

**Given the available data, what model would you build to predict crime to better allocate NYPD resources?**

I would try a cluster analysis, or a number of cluster analyses, using as independent variables geographic coordinates,  $(x, y)$  pairs indicating specific points in the city, together with variables such as time of day, day of the week, the month, type of crime or zoning status (e.g. residential), etc. The type of model could be a simple k-means analysis or (preferably) a mixture model, with statistical distributions assigned to the clusters (the task of the model fit is to estimate the parameters of these distributions). Either type of model (k-means or mixture model) may produce centers where crimes and arrests of certain types tend to be concentrated. These centers in  $n$ -dimensional space, e.g. in 3 dimensions with the  $x$  and  $y$  coordinates and time since the start of the week. It might be useful to create separate models for certain, different, crime categories.

In a mixture model, the value of the dependent variable we are predicting (e.g. a  $y$  on the left-hand side of the model equation) would be the value of the probability distribution function (for the probability of, say, an arrest). However, in addition to independent variables that have values in the data (e.g. the geographic coordinates), a model such as this will have latent traits or variables such as the cluster centers or the variances of the clusters. Variables or traits of particular interest will include (e.g.) cluster centers and (for mixture models) other parameters describing the clusters (e.g. variance of a normal distribution). In a similar fashion, in k-means clustering, the cluster centers will be of particular interest.

A model such as this might enhance efficient placement of police resources in certain locations at certain times, based on the properties of the clusters this model might discern. It might also, depending on the data that is used, allow allocation of police officers with certain specialties (with focus on certainly types of crimes).

As for model evaluation, the AIC or BIC criteria could be used to select one model over another. However, these measures will not tell us whether the final result is a good one (only that it is the best among possibly poorly-performing models). Cross validation could also be used for selecting one model over another. Two ways of evaluating a final model would be validation with data that is held out (a test set) or building a simulated dataset using the model. The actual data distributions and the simulated data distributions might then be compared using QQ plots (quantiles of one distribution plotted against quantiles of the other).

Challenges I might face include lack of direct access to what we want to measure. For example, we may know the arrest rate and need to use the arrest rate as proxy for the crime rate. The arrest rates might somewhat reflect greater amounts of policing in certain areas rather than greater amounts of crimes in those areas. Further analysis might be required to disentangle numbers of arrests due to actual crime numbers and numbers or arrests due to greater presence of policing. Another challenge, as always with large amounts of data and sophisticated models, will be limitations on computing resources.