Health Interests and Practices Survey – Tobacco Summary Data

Item	2006	2004	2002	2001	1999	1997	1995	1993	19
30 day	29.8	24.4	25.3	28.3	38.8	41.1	38.1	29.3	24
prevalence of	95% CI	95% CI	95% CI	95% CI	95% CI	95% CI	95% CI	95% CI	95
cigarettes	(27-33%)	(21-28%)	(22-29%)	(24-33%)	(34-44%)	(38-44%)	(35-41%)	(26-32%)	(2)
Lifetime prevalence	,	41.5	46.8	44.8	58.8				
Average age of starting smoking	17	16.5	16.5	17.5	17.9				
% of "heavy" smokers – ½ pack daily or more	12.5	7.4	11.1	22.5	20.6				
% of smokers smoking less than 1 cigarette per day	74.3	70.4	65.9	23.0	45.0				
Smokeless tobacco – 30 day prevalence	8.8	8.2	20.2	15.5					
% at least one quit attempt in past 12 months	46.0	48.0	53	46.0	48.2				
Average yearly quit attempts	2.2	1.59	1.81	1.71	1.66				
% daily smokers	5.9	8.9	9.5	12.3	10.4				
% using Hookah in last 12 months	46.2								
N and Response Rate	N=875	N=752, Response rate: 97%	N=656; Response rate: 97%	N=424; Response rate: 96%	N=428; Response rate: 98%	N=962; Resp. rate: 98%	N=978: Resp. rate: 97%	N=860; Resp. rate: 96%	N= Re rat

Results weighted by gender and grade level of participants. Sarah L. Hansen, M.A., CHES and Tanya Villhauer, M.A., Principal Investigators

Survey is anonymous and conducted in classrooms at the eight-week point in the fall semester.