Assignment 2 - Causal inference

RF

2/5/2020

Assignment 2 - Exploring causal inference issues

In this assignment we explore some issues related to multiple regressions (regressions with more than one predictor), and inferred (causal) relations between variables. N.B. the data is simulated (to make sure I know the actual mechanism generating it), but it's based on a real study. So bear with a longish introduction to get into the details of what we are doing and why it is important.

Altercentric intrusion in schizophrenia

People with schizophrenia often report altered control and distinction of self-other representations: intrusive thoughts, hearing of voices, delusions of mind reading, paranoia, etc (a substantial portion of the psychotic symptoms experienced in schizophrenia). These have been variously attributed to hypermentalizing (over attribution of mental states to others), social impairment (over preoccupation with own thought processes), hyper socialization (inability to inhibit information from others), etc.

The current study investigates 1) whether schizophrenia is indeed related to altered control and distinction of self-other representations, in particular altercentric intrusions (inability to inhibit social information), and 2) whether these are related to the relevant psychotic symptoms. N.B. the actual study also investigates egocentric intrusion, do check the papers below if interested.

The task is a slightly modified version of this: https://www.ncbi.nlm.nih.gov/pubmed/20731512 (also what Nicole and Louise described in their guest talk) You look at a picture with some dots visible to you, as well as with a different person with a different set of dots visible to them. The number of dots you see and that the other sees can be the same (congruent condition) or not (incongruent condition). You are tasked to indicate whether a given number (e.g. 3) matches the number of dots you see (and the dots visible to the other person are irrelevant to the task).

The tasks investigates altercentric intrusion: will your reaction time change according to whether the other person is seeing the same amount of dots as you, or not? The idea is that if you correctly inhibit social information, your reaction time should not change, as the information about the other person is not relevant. On the contrary, if you nevertheless use task irrelevant social information, you'll be slower at indicating whether 3 is the right number of dots when the other person sees a different amount of dots than you (conflicting information). The bigger the difference between RTs in the congruent and incongruent condition the bigger the altercentric intrusion effect.

For each participant you have 6 variables: 1) ID, 2) AltercentricIntrusion (continuous score), 3) Diagnosis (schizophrenia vs. control), 4) VoiceHearing (severity of voice hearing symptoms, continuous score of the severity of the symptom as measured by a clinician), 5) MindReading (severity of delusions of mind reading, continuous score of the severity of the symptom as measured by a clinician); 6) Apathy (severity of lack of motivation in taking care of oneself, from washing to showing up at work, continuous score of the severity of the symptom as measured by a clinician). N.B. Real clinical scores of symptoms would be on a likert scale, and altercentric intrusions would be on a slightly complex continuous scale. But life is too short for

that shit, and we want to focus on multivariate models and causal inference, so all those variables in the assignment (but ID and Diagnosis) have been simulated as Gaussian distributions.

The research questions you have to answer are the following:

```
#SIGRID
# Loading packages
pacman::p_load(tidyverse, brms, here, patchwork, bayesplot, ggdag)

# Reading data
df <- read_csv("Ass2.csv") %>%
  # Standardizing all the predictors
  mutate(
    across(c(VoiceHearing:AltercentricIntrusion), ~ scale(.x))
    )
```

```
##
## -- Column specification --------------------------------------------------------
## cols(
##   VoiceHearing = col_double(),
##   MindReading = col_double(),
##   Apathy = col_double(),
##   AltercentricIntrusion = col_double(),
##   ID = col_double(),
##   Diagnosis = col_double()
## )
```

First part

Q1.1) Does schizophrenia involve altercentric intrusion? Define model and priors. Test the implications of your priors (prior predictive checks) and if needed adjust them. Run the model. Test the quality of the fitted model (posterior predictive checks, prior-posterior updates). Assess the evidence in favor of an increased altercentric intrusion in schizophrenia. Report the model and the results, including plots.

```
############################ Inspecting the predictor ################################
# ANDERS
ggplot(df, aes(AltercentricIntrusion))+
  geom_density()
```




```
ggplot(df, aes(x = as.factor(Diagnosis), y = AltercentricIntrusion))+
  geom_boxplot()
```




```
############################ CHECKING THE PRIORS OF THE NULL MODEL ################################
# GUSTAV
# Defining formula
AI_f0 <- bf(AltercentricIntrusion ~ 1)

# Defining priors
AI_p0 <- c(
  prior(normal(0,1), class = Intercept),
  prior(normal(1,0.5), class = sigma)
)

# Making the model and only checking the priors
AI_m0_prior <- brm(
  formula = AI_f0,
  prior = AI_p0,
  family = gaussian,
  data = df,
  sample_prior = "only",
  file = here("Assignment_2","models_gustav", "AI_m0_prior")
)

pp_check(AI_m0_prior, nsamples = 100)
```




```
############################## CHECKING THE PRIORS WITH A PREDICTOR ##############################
# SIGRID
# Defining formula
AI_f1 <- bf(AltercentricIntrusion ~ 1 + Diagnosis)


# Defining priors
AI_p1 <- c(
  prior(normal(0,1), class = Intercept),
  prior(normal(1,0.5), class = sigma),
  prior(normal(0,0.2), class = b, coef = Diagnosis)
)

# Making the model and only checking the priors
AI_m1_prior <- brm(
  formula = AI_f1,
  prior = AI_p1,
  family = gaussian,
  data = df,
  sample_prior = "only" ,
  file = here("Assignment_2","models_gustav", "AI_m1_prior")
)

pp_check(AI_m1_prior, nsamples = 100)
```




```
##################### FITTING THE MODELS #####################
# ANDERS
AI_m0 <- brm(
  formula = AI_f0,
  prior = AI_p0,
  family = gaussian,
  data = df,
  sample_prior = T ,
  file = here("Assignment_2","models_gustav", "AI_m0")
)

AI_m1 <- brm(
  formula = AI_f1,
  prior = AI_p1,
  family = gaussian,
  data = df,
  sample_prior = T ,
  file = here("Assignment_2","models_gustav", "AI_m1")
)

p1<- pp_check(AI_m0, nsamples=100)
p2<- pp_check(AI_m1, nsamples=100)
p1 + p2
```




```
################## SUMMARY ##############
# ANDERS
posterior_summary(AI_m1) %>% round(digits = 2)
```

```
##                    Estimate Est.Error    Q2.5   Q97.5
## b_Intercept           -0.07      0.06   -0.19    0.06
## b_Diagnosis            0.27      0.11    0.05    0.50
## sigma                  0.99      0.04    0.92    1.08
## prior_Intercept       -0.03      1.00   -2.03    1.84
## prior_b_Diagnosis      0.00      0.20   -0.37    0.38
## prior_sigma            1.03      0.47    0.17    2.00
## lp__                -424.04      1.28 -427.30 -422.60
```

```
######################## POSTERIOR UPDATE FROM PRIORS ##########################################
# GUSTAV

posterior <- posterior_samples(AI_m1)

posterior %>%
  select(-lp__) %>%
  rename(
    b_sigma = sigma,
    prior_Diagnosis = prior_b_Diagnosis) %>%
  pivot_longer(cols = everything(),
               names_to = "Parameter",
```

```
               values_to = "Value") %>%
  separate(Parameter, into = c("Distribution", "Parameter"), sep = "_") %>%
  mutate(Distribution = ifelse(Distribution == "b", "Posterior", "Prior")) %>%
  # Plotting -----------------------------------------
  ggplot(aes(Value, fill = Distribution))+
  geom_density( alpha = 0.8)+
  scale_fill_manual(values = wesanderson::wes_palette("GrandBudapest2"))+
  facet_wrap(~Parameter, nrow = 1, )+
  labs(title = "Posterior update from priors")+
  theme_bw()+
  theme(strip.text = element_text(size=15),
        axis.title=element_text(size=15),
        plot.title = element_text(size = 20, face = "bold", hjust = 0.5))
```




```
# SIGRID
color_scheme_set("orange")

mcmc_plot(AI_m1,
          type = "intervals",
          prob = .5,
          prob_outer = .95,
          point_est = "mean") +
  labs(title = "The coefficient plot for the model with Diagnosis as predictor") +
  theme(axis.text.y = element_text(hjust = 0),
        panel.grid.minor = element_blank(),
        strip.text = element_text(hjust = 0))
```



Q1.2) Is altercentric intrusion related to specific symptoms *in the patients*? Identify which of the symptoms could be relevant (given their description above). Should you include more than one symptom? Build models, priors, predictive checks. Assess the evidence and report models and results, including plots. Discuss whether the results make sense.

We would predict that both mindreading and voicehearing are symptoms that are related to altercentric intrusion. Therefore

---

```
################################## UNIVARIATE ANALYSIS OF SYMPTOMS ########################
# ANDERS
# Subsetting the patients
patients <- df %>%
  filter(Diagnosis == 1)
```

```
############ CHECKING PRIORS #############
# GUSTAV

######### VOICEHEARING #########
VH_f <- bf(AltercentricIntrusion ~ VoiceHearing)

VH_prior <- c(
  prior(normal(0,1), class = Intercept),
  prior(normal(0,0.2), class = b),
  prior(normal(1,0.5), class = sigma)
)

AI_VH_prior <- brm(
  formula = VH_f,
  prior = VH_prior,
  family = gaussian,
  data = patients,
  sample_prior = 'only',
  file = here('Assignment_2', 'models_sigrid', 'AI_VH_prior')
)

pp_check(AI_VH_prior, nsamples = 100)
```




```
############ MINDREADING ##############
MR_f <- bf(AltercentricIntrusion ~ MindReading)

MR_prior <- c(
  prior(normal(0,1), class = Intercept),
  prior(normal(0,0.2), class = b),
  prior(normal(1,0.5), class = sigma)
)

AI_MR_prior <- brm(
  formula = MR_f,
  prior = MR_prior,
  family = gaussian,
  data = patients,
  sample_prior = 'only',
  file = here('Assignment_2', 'models_sigrid', 'AI_MR_prior')
)

pp_check(AI_MR_prior, nsamples = 100)
```




```
# Fitting the models ------------------------------------------------------------------------
# SIGRID

# Voicehearing
VH_m0 <- brm(
  formula = VH_f,
  data = patients,
  family = gaussian,
  prior = VH_prior,
  sample_prior = T,
  file = here("Assignment_2","models_gustav","VH_m0")
)

# Mindreading
MR_m0 <- brm(
  formula = MR_f,
  data = patients,
  family = gaussian,
  prior = MR_prior,
  sample_prior = T,
  file = here("Assignment_2","models_gustav","MR_m0")
)

##### SUMMARY

pp_check(VH_m0, nsamples = 100)
```




```
pp_check(MR_m0, nsamples = 100)
```




```
posterior_summary(VH_m0) %>% round(digits = 2)
```

```
##                      Estimate Est.Error    Q2.5  Q97.5
## b_Intercept              0.24      0.16   -0.07   0.55
## b_VoiceHearing           0.05      0.12   -0.19   0.29
## sigma                    0.91      0.08    0.77   1.07
## prior_Intercept          0.03      1.00   -1.89   2.00
## prior_b_VoiceHearing     0.00      0.20   -0.40   0.39
## prior_sigma              1.03      0.47    0.17   1.97
## lp__                   -98.76      1.25 -102.06 -97.35
```

```
posterior_summary(MR_m0) %>% round(digits = 2)
```

```
##                     Estimate Est.Error    Q2.5  Q97.5
## b_Intercept             0.25      0.12    0.01   0.49
## b_MindReading           0.06      0.10   -0.14   0.25
## sigma                   0.90      0.07    0.77   1.06
## prior_Intercept         0.00      1.00   -2.00   1.99
## prior_b_MindReading     0.00      0.20   -0.40   0.40
## prior_sigma             1.02      0.47    0.16   1.96
## lp__                  -98.59      1.19 -101.69 -97.25
```

```
################################### PLOTTING ###################################
# ANDERS
```

```r
# Voicehearing ------------------------------------------------------------------------------
post_VH <- posterior_samples(VH_m0)

VH_plot <- post_VH %>%
  select(-lp__) %>%
  rename(
    b_sigma = sigma,
    prior_VoiceHearing = prior_b_VoiceHearing) %>%
  pivot_longer(cols = everything(),
               names_to = "Parameter",
               values_to = "Value") %>%
  separate(Parameter, into = c("Distribution", "Parameter"), sep = "_") %>%
  mutate(Distribution = ifelse(Distribution == "b", "Posterior", "Prior"),
         Parameter = fct_relevel(Parameter, "VoiceHearing")) %>%
  # Plotting -------------------------------------------
  ggplot(aes(Value, fill = Distribution))+
  geom_density( alpha = 0.8)+
  scale_fill_manual(values = wesanderson::wes_palette("GrandBudapest2"))+
  facet_wrap(~Parameter, nrow = 1, )+
  labs(title = "Posterior update from priors - Voicehearing")+
  theme_bw()+
  theme(strip.text = element_text(size=15),
        axis.title=element_text(size=15),
        plot.title = element_text(size = 20, face = "bold", hjust = 0.5))

# Mindreading -------------------------------------------------------------------------------
post_MR <- posterior_samples(MR_m0)

MR_plot <- post_MR %>%
  select(-lp__) %>%
  rename(
    b_sigma = sigma,
    prior_MindReading = prior_b_MindReading) %>%
  pivot_longer(cols = everything(),
               names_to = "Parameter",
               values_to = "Value") %>%
  separate(Parameter, into = c("Distribution", "Parameter"), sep = "_") %>%
  mutate(Distribution = ifelse(Distribution == "b", "Posterior", "Prior"),
         Parameter = fct_relevel(Parameter, "MindReading")) %>%
  # Plotting -------------------------------------------
  ggplot(aes(Value, fill = Distribution))+
  geom_density( alpha = 0.8)+
  scale_fill_manual(values = wesanderson::wes_palette("GrandBudapest2"))+
  facet_wrap(~Parameter, nrow = 1, )+
  labs(title = "Posterior update from priors - Mindreading")+
  theme_bw()+
  theme(strip.text = element_text(size=15),
        axis.title=element_text(size=15),
        plot.title = element_text(size = 20, face = "bold", hjust = 0.5))

VH_plot / MR_plot
```






```
########### PLOTTING #############
#GUSTAV

color_scheme_set("orange")

# MindReading
MR_m0_p <- mcmc_plot(MR_m0,
         type = "intervals",
         prob = .5,
         prob_outer = .95,
         point_est = "mean") +
  labs(title = "MindReading") +
  theme(axis.text.y = element_text(hjust = 0),
        panel.grid.minor = element_blank(),
        strip.text = element_text(hjust = 0))

# VoiceHearing
VH_m0_p <- mcmc_plot(VH_m0,
         type = "intervals",
         prob = .5,
         prob_outer = .95,
```

```
         point_est = "mean") +
  labs(title = "VoiceHearing") +
  theme(axis.text.y = element_text(hjust = 0),
        panel.grid.minor = element_blank(),
        strip.text = element_text(hjust = 0))

MR_m0_p / VH_m0_p
```



---

* After inspecting the symptoms' individual effect with a conservative prior, it seems that neither mindreading or voicehearing have a convincing effect on altercentric intrusion. The symptoms did not deviate from 0 effect. Next we will try to combine the symptoms in a multiple linear model*

```
########################### MULTIPLE LINEAR MODEL ##################################
######################### VoiceHearing + MindReading ##################################
# SIGRID

multi_f0 <- bf(AltercentricIntrusion ~ 1 + VoiceHearing + MindReading)


# We will use the same priors as for diagnosis. We expect 0 effect, but accept values on up to 2 SD, th

multi_prior0 <- c(
  prior(normal(0,1), class = Intercept),
```

```r
  prior(normal(1,0.5), class = sigma),
  prior(normal(0,0.2), class = b, coef = VoiceHearing),
  prior(normal(0,0.2), class = b, coef = MindReading)
)

# Fitting the models ------------------------------------------------------------------------------

# Voicehearing
multi_m0 <- brm(
  formula = multi_f0,
  data = patients,
  family = gaussian,
  prior = multi_prior0,
  sample_prior = T,
  file = here("Assignment_2","models_gustav","multi_m0")
)


# Inspecting models -------------------------------------------------------------------------------
posterior_summary(multi_m0) %>% round(digits = 2)
```

```
##                       Estimate Est.Error    Q2.5  Q97.5
## b_Intercept               0.16      0.19   -0.21   0.53
## b_VoiceHearing            0.08      0.12   -0.16   0.32
## b_MindReading             0.08      0.11   -0.12   0.29
## sigma                     0.90      0.08    0.77   1.07
## prior_Intercept           0.02      1.00   -1.92   1.93
## prior_b_VoiceHearing      0.00      0.20   -0.41   0.40
## prior_b_MindReading       0.00      0.20   -0.40   0.40
## prior_sigma               1.02      0.48    0.15   2.02
## lp__                    -98.28      1.48 -102.08 -96.45
```

```r
################## PLOTTING ###############
#ANDERS

color_scheme_set("orange")

mcmc_plot(multi_m0,
         type = "intervals",
         prob = .5,
         prob_outer = .95,
         point_est = "mean") +
  labs(title = "The coefficient plot for the model with both predictors") +
  theme(axis.text.y = element_text(hjust = 0),
        panel.grid.minor = element_blank(),
        strip.text = element_text(hjust = 0))
```




```
# MORE PLOTTING
# GUSTAV
post_multi0 <- posterior_samples(multi_m0)

post_multi0 %>%
  select(-lp__) %>%
  rename(
    b_sigma = sigma,
    prior_VoiceHearing = prior_b_VoiceHearing,
    prior_MindReading = prior_b_MindReading) %>%
  pivot_longer(cols = everything(),
               names_to = "Parameter",
               values_to = "Value") %>%
  separate(Parameter, into = c("Distribution", "Parameter"), sep = "_") %>%
  mutate(Distribution = ifelse(Distribution == "b", "Posterior", "Prior"),
         Parameter = fct_relevel(Parameter, "VoiceHearing")) %>%
  # Plotting -----------------------------------------
  ggplot(aes(Value, fill = Distribution))+
  geom_density( alpha = 0.8)+
  scale_fill_manual(values = wesanderson::wes_palette("GrandBudapest2"))+
  facet_wrap(~Parameter, nrow = 2, )+
  labs(title = "VoiceHearing + MindReading")+
  theme_bw()+
  theme(strip.text = element_text(size=15),
        axis.title=element_text(size=15),
        plot.title = element_text(size = 20, face = "bold", hjust = 0.5))
```




```
########## INSPECTING CORRELATIONS ########3
# SIGRID
patients %>%
  select(AltercentricIntrusion, MindReading, VoiceHearing) %>%
  pairs(col = "forestgreen")
```



Second part

Q2.1) However, we know that the diagnosis is based on symptom assessment: if the overall sum of symptoms is severe enough, the participant gets a diagnosis. In other words, by selecting the patients, and including the symptoms in the model we might have inadvertently introduced an issue in our inference. Do try to draw a causal graph (Directed Acyclical Graph) of the variables and compare it with the types of causal graphs presented in the slides. Discuss which biases you might have introduced.

```
############################ CREATING A DAG ################################
# ANDERS

# AI = AltercentricIntrusion
# VH = VoiceHearing
# MR = MindReading
# AP = Apathy
# D = Diagnosis

dag_coords <-
  tibble(name = c("AP","AI", "VH", "MR", "D"),
         x    = c(0, -1.2, 0, 0, 1.2),
         y    = c(0, 1, 2, 1, 1))

# save our DAG
dag <-
  dagify(D ~ VH + MR + AP,
```

```
         AI ~ VH + MR ,
         coords = dag_coords)

# plot
dag %>%
  ggplot(aes(x = x, y = y, xend = xend, yend = yend)) +
  geom_dag_point(color = "steelblue", alpha = 1/2, size = 6.5) +
  geom_dag_text(color = "black") +
  geom_dag_edges() +
  theme_dag()
```



Q2.2.) Redesign your analysis following the graph and report how the results change

```
################### CREATING A NEW MULTIPLE MODEL WITH ALL PREDICTORS #######################
#################### VoiceHearing + MindReading ##############################
# GUSTAV

# Fitting the models ------------------------------------------------------------------------

# All predictors
updated_m0 <- brm(
  formula = multi_f0,
  data = df,
  family = gaussian,
  prior = multi_prior0,
  sample_prior = T,
```

```r
  file = here("Assignment_2","models_gustav","updated_m0")
)


# Expecting models ------------------------------------------------------------------------
summary(updated_m0)
```

```
##  Family: gaussian
##   Links: mu = identity; sigma = identity
## Formula: AltercentricIntrusion ~ 1 + VoiceHearing + MindReading
##    Data: df (Number of observations: 300)
## Samples: 4 chains, each with iter = 2000; warmup = 1000; thin = 1;
##          total post-warmup samples = 4000
##
## Population-Level Effects:
##              Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
## Intercept       -0.00      0.06    -0.11     0.11 1.00     4289     2927
## VoiceHearing     0.16      0.06     0.05     0.27 1.00     3960     3034
## MindReading      0.16      0.05     0.05     0.26 1.00     3865     2862
##
## Family Specific Parameters:
##       Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
## sigma     0.97      0.04     0.90     1.06 1.00     4081     3235
##
## Samples were drawn using sampling(NUTS). For each parameter, Bulk_ESS
## and Tail_ESS are effective sample size measures, and Rhat is the potential
## scale reduction factor on split chains (at convergence, Rhat = 1).
```

```r
posterior_summary(updated_m0) %>% round(digits = 2)
```

```
##                      Estimate Est.Error    Q2.5   Q97.5
## b_Intercept              0.00      0.06   -0.11    0.11
## b_VoiceHearing           0.16      0.06    0.05    0.27
## b_MindReading            0.16      0.05    0.05    0.26
## sigma                    0.97      0.04    0.90    1.06
## prior_Intercept          0.00      0.98   -1.89    1.90
## prior_b_VoiceHearing     0.00      0.20   -0.39    0.39
## prior_b_MindReading      0.00      0.20   -0.39    0.38
## prior_sigma              1.03      0.47    0.18    1.95
## lp__                  -417.45      1.42 -421.00 -415.69
```

```r
################## PLOTTING ###############
# SIGRID

color_scheme_set("orange")

mcmc_plot(updated_m0,
         type = "intervals",
         prob = .5,
         prob_outer = .95,
         point_est = "mean") +
  labs(title = "The coefficient plot for the model with both predictors") +
```

```
  theme(axis.text.y = element_text(hjust = 0),
        panel.grid.minor = element_blank(),
        strip.text = element_text(hjust = 0))
```




```
# MORE PLOTTING
# ANDERS

post_updated_m0 <- posterior_samples(updated_m0)

post_updated_m0 %>%
  select(-lp__) %>%
  rename(
    b_sigma = sigma,
    prior_VoiceHearing = prior_b_VoiceHearing,
    prior_MindReading = prior_b_MindReading) %>%
  pivot_longer(cols = everything(),
               names_to = "Parameter",
               values_to = "Value") %>%
  separate(Parameter, into = c("Distribution", "Parameter"), sep = "_") %>%
  mutate(Distribution = ifelse(Distribution == "b", "Posterior", "Prior"),
         Parameter = fct_relevel(Parameter, "VoiceHearing")) %>%
  # Plotting -------------------------------------------
  ggplot(aes(Value, fill = Distribution))+
  geom_density( alpha = 0.8)+
  scale_fill_manual(values = wesanderson::wes_palette("GrandBudapest2"))+
  facet_wrap(~Parameter, nrow = 2, )+
  labs(title = "VoiceHearing + MindReading")+
  theme_bw()+
  theme(strip.text = element_text(size=15),
        axis.title=element_text(size=15),
        plot.title = element_text(size = 20, face = "bold", hjust = 0.5))
```



Third part

These issues are very difficult to think through, and not knowing the causal mechanisms generating the data in advance makes our inferences even more unreliable. To explore these issues, I recommend using simulations. In other words, defining a "true" model, generating data from it and assessing what different analyses would lead you to infer (and therefore which biases they might introduce). You can find the code I used to simulate your data below.

Q3.1) Look through the code and identify whether the results you have match the underlying truth. Discuss what you have learned.

Q3.2) OPTIONAL: is this a general pattern? Try varying the parameters (e.g. correlation values) and assess whether the new dataset(s) leads to the same biases in your analysis.

```
pacman::p_load(MASS, tidyverse, psych)

seed <- 1981 # Defining a seed so the results are always the same
n <- 300 # Defining the amount of participants

SymptomCorr <- .2 # Defining the correlation of symptoms (as they tend to co-occur)
EffectCorrRel <- .2 # Defining the correlation between relevant symptoms and effect (Some symptoms are
EffectCorrIrrel <- 0 # Defining the correlation between irrelevant symptoms and effect (none)

# Creating the variance-covariance matrix for the variables we want to generate (3 symptoms, 1 effect)
Sigma <- matrix(data=c(1,SymptomCorr,SymptomCorr,EffectCorrRel,
                       SymptomCorr,1,SymptomCorr,EffectCorrRel,
```

```r
                        SymptomCorr,SymptomCorr,1,EffectCorrIrrel,
                        EffectCorrRel,EffectCorrRel,EffectCorrIrrel,1),
                        nrow=4,ncol=4)

## Generate data from a multivariate (mvr) normal (n) distribution
d <- mvrnorm(n = n, # number of participant
        mu = c(1.2, 1.2, 1.2, 4), # mean of each variable
        Sigma) # variance co-variance matrix

# Giving meaningful names to variables and add ID
d <- data.frame(
  VoiceHearing = d[,1],
  MindReading =  d[,2],
  Apathy =  d[,3],
  AltercentricIntrusion = d[,4],
  ID = seq(nrow(d)))

# Assessing whether the participant has schizophrenia (high enough sum of symptoms)
# Here we choose participants scoring above 75% percentile (the most severe ones)
d$Diagnosis <- 0
d$Diagnosis[(d$VoiceHearing + d$MindReading + d$Apathy) >
              quantile(d$VoiceHearing + d$MindReading + d$Apathy, .75)] <- 1

## Plotting the relation between variables all participants
pairs.panels(dplyr::select(d,-Diagnosis, -ID))
```



```r
## Plotting the relation between variables in schizophrenia
d1 <- d %>% subset(Diagnosis==1) %>% dplyr::select(-Diagnosis, -ID)
pairs.panels(d1)
```




```r
#write_csv(d, "data/Ass2.csv")
```