

Effects of climate change on four New England groundfish species

Emily S. Klein · Sarah L. Smith · Jacob P. Kritzer

Received: 23 March 2016 / Accepted: 19 July 2016
© Springer International Publishing Switzerland 2016

Abstract Multiple groundfish stocks in New England remain depleted despite management measures that have been effective elsewhere. A growing body of research suggests that environmental change driven by increasing concentrations of carbon dioxide in the atmosphere and ocean is unfolding more rapidly in New England than elsewhere, and is an important factor in the failure of these stocks to respond to management. We reviewed research on effects of changes in temperature, salinity, dissolved oxygen, pH, and ocean currents on pelagic life stages, post-settlement life stages, and reproduction of four species in the New England groundfish fishery: Atlantic cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), winter flounder (*Pseudopleuronectes americanus*), and yellowtail flounder (*Limanda ferruginea*). The volume of research on cod was nearly equal to that on the other

three species combined. Similarly, many more studies examined effects of temperature than other factors. The majority of studies suggest adverse outcomes, with less evidence for mixed or positive effects. However, for all of the factors other than temperature, there are more knowledge gaps than known effects. Importantly, most work to date examines impacts in isolation, but effects might combine in nonlinear ways and cause stronger reductions in stock productivity than expected. Management strategies will need to account for known effects, nonlinear interactions, and uncertainties if fisheries in New England are to adapt to environmental change.

Keywords Climate change · Groundfish · New England · Temperature · Productivity · Fisheries management

E. S. Klein
Department of Ecology and Evolutionary Biology,
Princeton University, 106A Guyot Hall, Princeton,
NJ 08544, USA

Present Address:
E. S. Klein
Southwest Fisheries Science Center, National Oceanographic
and Atmospheric Administration, 8901 La Jolla Shores Drive,
La Jolla, CA 92037, USA

S. L. Smith · J. P. Kritzer (✉)
Fishery Solutions Center, Environmental Defense Fund,
18 Tremont Street, Boston, MA 02108, USA
e-mail: jkritzter@edf.org

Introduction

Fisheries management in the United States has evolved in important ways over the past few decades. Rebuilding plans for overfished species managed at the federal level have been required since statutory revisions to the Magnuson Stevens Act in 1996, a strategy aimed at deterring management inaction and accelerating recovery (Safina et al. 2005). The effectiveness of rebuilding plans initially proved disappointing because fishing mortality could not be kept to

sufficiently low levels in practice (Rosenberg et al. 2006). However, further revisions to federal law in 2007 required that risk-averse buffers be included within catch limits, a strategy that can account for the often considerable scientific uncertainties underlying management (Restrepo and Powers 1999). The 2007 revisions also required that management plans include accountability measures for quota overages. This has motivated increased use of both output controls and incentive-based allocation systems, also known as ‘catch shares’, across US fisheries, approaches that are more effective at meeting fishing mortality targets (Worm et al. 2009; Melnychuk et al. 2012), reducing risk of collapse (Costello et al. 2008), dampening inter-annual variability (Essington 2010; Essington et al. 2012), and improving socio-economic outcomes (Gutierrez et al. 2011; Grimm et al. 2012). As a result of these accumulated changes, the economic and biological performance of fisheries in the United States has continually improved over the past decade (NOAA 2015).

A notable exception to the generally positive trends in US fisheries is the New England groundfish fishery (NOAA 2015), the oldest commercial fishery in the nation. Management of the groundfish fishery includes all of the elements outlined above that have contributed to success in many other US fisheries: rebuilding plans, risk-averse buffers in catch limits, output controls, and catch shares. Yet, many stocks not only remain severely depleted but are also trending downward (NEFSC 2015). Stocks at very low levels of biomass can be the least responsive to management measures and are most vulnerable to environmental variability (Vert-pre et al. 2013), which might explain the disappointing outcomes. Furthermore, environmental changes linked to rising concentrations of CO₂ in the atmosphere and ocean are especially pronounced in the waters off New England (see “Ecosystem context” below). The magnitude of these changes recently motivated a comprehensive assessment of the vulnerability of 82 species in the waters off New England to the effects of climate change (Hare et al. 2016).

Climate change can affect stock productivity, potential yield and management targets (Pörtner and Peck 2010; Sumaila et al. 2011; Doney et al. 2012; Pinsky and Mantua 2014). Fogarty et al. (2008) re-estimated biological reference points for Atlantic cod (*Gadus morhua*) accounting for predicted effects of


increased temperature on growth and recruitment. Their results suggest that maximum sustainability yield (MSY) and the fishing mortality rate that will produce MSY (F_{MSY}) will be lower with a 2 °C increase relative to baseline conditions. Pershing et al. (2015) corroborated these results and illustrated that management strategies that have not accounted for these changes are partly to blame for the continued decline of the Gulf of Maine cod stock. In contrast, the vulnerability assessment by Hare et al. (2016) suggests that groundfish species might be affected less than other functional groups, notably benthic invertebrates and diadromous species. However, the authors also noted that more detailed empirical and process-oriented studies focused on individual species might yield different outcomes. Such studies are especially needed for groundfish in light of their poor status and high economic, cultural, and ecological importance.

To help guide future research and management strategies for New England groundfish that account for ongoing environmental change, we conducted a structured literature review of evidence for the effects of climatic and oceanographic factors on four of the most economically important species in the New England groundfish complex. We considered two gadoids, Atlantic cod and haddock (*Melanogrammus aeglefinus*), and two pleuronectoids, winter flounder (*Pseudopleuronectes americanus*) and yellowtail flounder (*Limanda ferruginea*). Our objectives were to identify and synthesize what is known about the positive and negative effects of climatic and oceanographic changes on each species, trends that are consistent across species or unique to each, and knowledge gaps that require future research.

Ecosystem context

The marine waters off New England are defined by two prominent geographic features: the Gulf of Maine and Georges Bank (Fig. 1). The Gulf of Maine is bound by the coastlines of the U.S. states of Massachusetts, New Hampshire and Maine to the west and north, the Scotian Peninsula and Scotian Shelf in Canada to the north and east, and Georges Bank to the south. Canada’s Bay of Fundy opens into the Gulf of Maine, which itself extends into the wider Atlantic Ocean by the Northeast Channel. The Gulf is strongly influenced by freshwater input from more than a dozen

Fig. 1 Geography and topography of the coastal and marine waters off New England, USA



large rivers and many smaller tributaries. Its seafloor includes underwater banks, ledges, shoals, basins and other features, providing considerable topographic complexity. In contrast, Georges Bank is a considerably more uniform sandy underwater plateau, albeit with important variation in substrate composition and diversity.

As the North Atlantic as a whole is a comparatively young marine system, the region exhibits relatively low marine biodiversity (Witman et al. 2004). However, the historical fisheries productivity of the region was remarkably high (e.g., Rosenberg et al. 2005), bolstered by the strong linkages with freshwater ecosystems (Hall et al. 2012; Ames and Lichter 2013). Multiple anthropogenic stressors have altered ecosystem structure and function in the centuries since European colonization of the region, including over-fishing, coastal development, pollution, and other watershed manipulations. For example, kelp forests in

the Gulf of Maine that were once dominated by large predatory fish have become increasingly dominated by lower trophic level crustaceans (Steneck et al. 2004).

More recently, climatic and oceanographic effects of rising concentrations of greenhouse gases are increasingly altering the marine ecosystems off New England. In the Gulf of Maine, sea surface temperatures have increased at a rate of $0.026\text{ }^{\circ}\text{C year}^{-1}$ over the last 30 years, accelerating to $0.26\text{ }^{\circ}\text{C year}^{-1}$ between 2004 and 2012 (Mills et al. 2013), a rate faster than almost anywhere else on the globe. Average annual precipitation has increased about 5 % in the Northeast since 1991 (Walsh et al. 2014), and is expected to increase by 5–10 % by 2050 according to IPCC forecast models (Fernandez et al. 2015). Precipitation is also becoming more variable, resulting in more flood and drought years (Balch et al. 2012). Stronger precipitation events have already caused ocean freshening (Balch et al. 2012), and

continued decreases in salinity are expected due to melting of the Arctic ice cap and freshening of the Labrador Current (Nummelin et al. 2016). Oceans have become about 0.1 pH units more acidic worldwide and continued increases in CO₂ concentration and decreases in salinity are expected to cause a further reduction of 0.2–0.3 pH units (Feeley et al. 2009; also see Caldeira and Wickett 2003, 2005).

Climate change can also result in lower levels of dissolved oxygen due to the reduced solubility of warmer waters. Increased nutrient runoff from increased precipitation can exacerbate hypoxic conditions in coastal waters, while alterations to water column stratification resulting from temperature and salinity changes can expand deeper hypoxic zones into shallower depths (Rabalais et al. 2010; Doney et al. 2012). Increased prevalence of hypoxia in coastal or offshore waters has not been widely reported for the New England region, but those changes might unfold as other climatic and oceanographic changes progress. Likewise, increased warming of the upper water column combined with changes in wind stress and salinity has the potential to one day alter the velocity and direction of ocean currents (Hoegh-Guldberg and Bruno 2010).

These changes in the physical environment have already resulted in measurable ecological alterations, such as decline in abundance of zooplankton species that are important prey for larval fish (Friedland et al. 2013). Many fish species are also exhibiting substantial shifts in their distribution (Nye et al. 2009; Kleisner et al. 2016), and these shifts are happening faster in the waters off New England than anywhere else in the United States (Pinsky et al. 2013). These changes are altering species composition and trophic guild structure (Garrison and Link 2000; Kleisner et al. 2016), and likely altering the system state to one dominated by lower trophic level pelagic species over higher trophic level demersal groundfish (Choi et al. 2004).

Literature review

We searched Web of Science using combinations of the scientific and common names of our four focal species with a geographic constraint (Gulf of Maine, Georges Bank, New England), and specific (temperature, temp*, salinity, ocean acidification, hypoxia,

dissolv* oxygen) and general (environment* effects, environ*, climate change) topical search terms. Using references lists from the resulting literature base, we identified additional sources for topics with few studies in the initial search, allowing a wider geographic range across the North Atlantic. The waters off New England are toward the southern extent of the range of all four species we considered, especially Atlantic cod. However, cold water delivered south toward New England by the Labrador Current, combined with warm water moved away from the region by the Gulf Stream and North Atlantic Drift, mean that the New England region has historically maintained temperature profiles similar to many locales at higher latitudes across the North Atlantic. Therefore, studies from those locales can likely provide useful insights. Finally, in addition to published papers, we also reviewed relevant grey literature, such as Essential Fish Habitat source documents and stock assessment reports.

The search resulted in 136 core references, the topics of which are summarized in Table 1. The majority of these focused on Atlantic cod, with winter flounder a distant second. Many were empirical studies addressing effects of environmental attributes more generally, rather than the outcomes of climate change or other directional environmental change specifically, but we utilized these empirical findings to evaluate the likely effects of ongoing and forthcoming environmental change. Fewer studies used projections or otherwise worked within the context of climate change explicitly. Effects of temperature received the most research attention, with a dearth of literature addressing effects of hypoxia and especially ocean acidification. Additional research on the effects of these factors exists for other components of the ecosystem (species, habitats, processes) that can affect our species of interest, but we concentrated on studies that addressed the focal species directly.


Temporal trends in the studies we considered (Fig. 2) show that the considerable attention given to temperature effects over others has generally been maintained through time. However, the volume of studies increased dramatically from the 1980s to the 1990s. More importantly, the number of studies examining interacting effects of different environmental factors also increased substantially more recently. This is a promising trend in light of the multiple changes unfolding in the ecosystem.

Table 1 Number of studies addressing effects of changing environmental conditions on four species of New England groundfish

Species	Environmental factors								
	Temperature	Salinity	Acidification	Hypoxia	Currents	Multiple or interacting effects	Climate change projections	Other effects	Total
Atlantic cod	22 (15)	13 (4)	1	2 (2)	2	13 (7)	3	2	65
Haddock	10 (4)	3	1	1	6	7 (1)	0	0	24
Winter flounder	11 (9)	4	0	3 (3)	1	10 (1)	0	0 (2)	30
Yellowtail flounder	11	2 (1)	0	1	0	7 (3)	0	2 (1)	17
Total	74	25	2	11	9	41	3	7	

Numbers in parentheses are studies focused on areas other than the North American continental shelf. Some references provided information on more than one effect or species, and therefore the total number of references per species or factor is not a direct sum of the cells in a row or column in all cases

Fig. 2 Temporal trends in the number of studies considered in the literature review of effects of environmental change on four species of New England groundfish, summed across the four species. “Multiple effects” include those references addressing more than one environmental factor, and these references are reflected in the totals for each of those factors as well



Patterns in the literature

We summarize the empirical findings reported in the literature in three broad categories: Pelagic life stages (i.e., growth, development, condition, survival, and behavior of eggs and larvae), post-settlement life stages (growth, development, condition, survival, and behavior of juveniles and adults), and reproduction (maturation, fecundity, spawning and recruitment; the linkage between pelagic and post-settlement life stages). These categories are not distinct, of course. For example, growth and survival during pelagic life stages impacts recruitment, and post-settlement growth is a key determinant of fecundity. Within each of these categories, we consider temperature

separately from the other environmental factors as it was focus for the majority of studies considered.

Results of the literature review are summarized in Table 2. There is considerable complexity inherent in the environmental-biological interactions we are examining. In some instances, the effects of directional change in a given environmental attribute depend upon the magnitude of the changes relative to a species' tolerances, the rate of change, the location in question along various gradients (especially latitudinal, inshore-offshore, and depth) that determine localized adaptations, and interactions among these factors. This complexity is difficult to summarize, and the content in Table 2 is therefore generalized to a degree by necessity, although some mixed effects are

Table 2 Summary of impacts on four species of New England groundfish due to expected changes in environmental conditions based on literature review

Species	Environmental change	Pelagic stages (eggs & larvae)			Post-settlement (juveniles & adults)			Reproduction			
		Growth & development	Survival	Dispersal & settlement	Growth & condition	Survival	Behavior	Maturation	Fecundity	Spawning	Recruitment
Atlantic cod	↑ Temperature	Green	Red	Grey	Green	Red	Grey	Green	Red	Red	Green
	↓ Salinity	Red	Red	Grey	Red	Red	Grey	Grey	Grey	Red	Red
	↓ pH	Red	Red	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey
	↓ DO	Grey	Grey	Grey	Red	Red	Grey	Grey	Red	Grey	Red
	Δ Currents	Grey	Grey	Red	Red	Red	Grey	Grey	Grey	Grey	Red
Haddock	↑ Temperature	Green	Red	Grey	Green	Red	Grey	Green	Red	Red	Red
	↓ Salinity	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey
	↓ pH	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey
	↓ DO	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey
	Δ Currents	Grey	Grey	Red	Red	Red	Grey	Grey	Grey	Grey	Red
Yellowtail flounder	↑ Temperature	Green	Red	Red	Green	Red	Red	Grey	Green	Red	Green
	↓ Salinity	Grey	Red	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Green
	↓ pH	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey
	↓ DO	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey
	Δ Currents	Grey	Grey	Red	Grey	Grey	Grey	Grey	Grey	Grey	Red
Winter flounder	↑ Temperature	Green	Red	Red	Green	Red	Red	Grey	Red	Red	Red
	↓ Salinity	Red	Red	Grey	Red	Red	Grey	Grey	Grey	Grey	Grey
	↓ pH	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey
	↓ DO	Grey	Grey	Grey	Red	Red	Red	Grey	Grey	Grey	Grey
	Δ Currents	Grey	Grey	Red	Grey	Grey	Grey	Grey	Grey	Grey	Red

Mixed effects are conveyed in some instances, but a number of factors can result in very different and even opposing outcomes (e.g., rate and magnitude of change; latitude). Therefore, these results are generalized to a degree by necessity. Green = positive effects, red = negative effects, light green or light red = indirect positive or negative effect (e.g., due to a change in another life history trait), multiple colors = evidence for different types of effects, DO = dissolved oxygen

captured. The following text attempts to delineate different outcomes expected due to the nature of environmental change and the position along environmental gradients in more detail.

Pelagic life stages

Temperature effects

Elevated ocean temperatures can increase early growth and development rates of cod (Laurence 1978; Brander 1995; Kristiansen et al. 2011), haddock (Laurence 1978; Cargnelli et al. 1999), yellowtail flounder (Colton 1959; Laurence and Howell 1981), and winter flounder (Laurence 1975; Williams 1975; Rogers 1976; Buckley et al. 1990), suggesting potential benefits of climate change. Reduced incubation and faster development reduces time in life stages when mortality and predation rates are highest, and hastens the transition to larger, free-swimming juveniles that are better at foraging and predator avoidance. Length of incubation time can be especially crucial for species such as winter flounder that are

dependent on nearshore and estuarine areas where they are more susceptible to other anthropogenic impacts, such as pollution, during critical developmental periods (Pereira et al. 1999).

If warmer waters do increase growth rates, larvae will still need enough appropriately sized prey to meet size-specific metabolic needs (O'Connor et al. 2007; Kristiansen et al. 2014). In early life, failure to find food is one of the most important causes of mortality in fish larvae (Blaxter 1986). The availability of appropriately sized forage is crucial for larval feeding and growth, and therefore recruitment and year class strength, of cod (Buckley et al. 2010) and winter flounder (Laurence 1977; Stoner et al. 2001). Prey mismatches, due to temperature impacts on prey themselves, earlier spawning or rapid larval growth, can lead to larval starvation. In the North Sea, there is evidence that warming waters have caused shifts in plankton communities and in turn reduced survival of larval cod (Beaugrand et al. 2003), and may be a major driver in recent recruitment failure of cod (Nicolas et al. 2014). Similarly, changes in environmental conditions may be causing biogeographical and

potentially regime shifts in copepod communities in the North Atlantic (Beaugrand et al. 2002; Beaugrand 2004) and limiting the overall production of planktonic prey (Boyce et al. 2010), with potential impacts on cod recruitment (Beaugrand et al. 2003). Kristiansen et al. (2011) determined warm years can increase prey availability and overlap with larval cod, but Friedland et al. (2013) found a recent reduction in the distribution and abundance of certain plankton in the waters off New England, resulting in localized feeding mismatches for larval cod. Changing conditions may also be limiting food availability for larval flatfish, especially in southern areas (Gibson 2013). In addition, as planktonic prey vary in caloric content, increasing temperature may also impact early life stages via the quality of prey (Nicolas et al. 2014). Furthermore, warmer temperatures may limit primary productivity (Steinacher et al. 2010; Capotondi et al. 2012), also reducing prey availability (Kristiansen et al. 2014). These impacts may vary by location and may be stronger in southern waters where temperate groundfish species are already at the limit of their environmental tolerances.

The potential benefits of warmer waters on early growth and development may be limited for other reasons as well. First, more rapid incubation, hatching and growth might occur only within a certain range of temperature increase, beyond which adverse effects unfold, as is evident for haddock (Laurence and Rogers 1976; Laurence 1978). Second, the rate at which waters warm may also be critical. For example, sudden and drastic increases in temperature can increase mortality in yellowtail flounder larvae (Colton 1959). Third, warmer waters may have other, deleterious effects. For cod, high temperatures can adversely affect fertilization, egg development (Nicolas et al. 2014), and egg size (Miller et al. 1995). Increased temperature can reduce the growth and survival of haddock (Peck et al. 2003; Eriksen et al. 2012), and hatching success (Williams 1975; Rogers 1976; Keller and Klein-MacPhee 2000), initial larval size (Buckley et al. 1990; Keller and Klein-MacPhee 2000), and larval survival (Rogers 1976; Taylor and Danila 2005; Manderson 2008) in winter flounder. Fourth, changing temperature may have indirect effects. Projected rates of warming may cause earlier water column stratification in summer, potentially reducing settlement rates and increasing mortality in juvenile yellowtail flounder (Sullivan et al. 2005).

Again, these possible consequences of warming waters will likely vary by location, given spatial comparisons of temperature effects on the early life history of cod (Planque and Fredou 1999; Pörtner et al. 2001, 2008) and yellowtail flounder (Steves et al. 2000).

Finally, temperature increases may alter predation rates on larvae, which has important effects on survival of larval winter flounder (Dickie and McCracken 1955). The spawning cycle of winter flounder is timed to align with favorable environmental conditions and reduced predation (Jeffries and Johnson 1974; Jeffries et al. 1989; Collette and Klein-MacPhee 2002; Taylor and Danila 2005), which can be disrupted by changes in temperature regimes and predator dynamics. With warmer winters and earlier springs, predators move into critical estuarine nursery grounds earlier and are more active (Taylor and Danila 2005; Bell et al. 2014), resulting in lower survival and abundance of eggs and larvae (Taylor and Collie 2003; Taylor and Danila 2005). Population modeling suggests that the increased predator abundance can result in consumption of up to 50 % of flounder spawn in a single season (Taylor and Danila 2005). These effects might offset advantages of increased temperature (Rajasilta et al. 1993; Taylor and Danila 2005; Keller and Klein-MacPhee 2000).

Other environmental effects

In addition to temperature, changes in salinity may also impact the early life stages of groundfish. For cod, low salinity results in a decline in fertilization and normal development (Davenport et al. 1981; Westin and Nissling 1991), delayed hatching and increased egg mortality (Laurence and Rogers 1976; Koester et al. 2001), and caused eggs to sink more readily (Lough 2004). Winter flounder appear relatively tolerant to changes in salinity, likely due to their reliance on inshore and estuarine waters, but extreme salinities were found to cause abnormal development (Rogers 1976) and lower egg and larval survival and hatching success (Buckley 1989). Salinity and temperature may also interact to affect early life history stages. Laurence and Howell (1981) found that survival of yellowtail flounder eggs is greatly reduced in low salinities at both low and high temperatures, and Rogers (1976) discovered salinity had a stronger effect on hatching success and larval survival in winter flounder under higher temperatures.

Ocean acidification has the potential to influence calcification, acid-based regulation, blood circulation, respiration, and the nervous system in marine organisms (Frommel et al. 2012). Fish are particularly vulnerable to these impacts in early life when they lack the ability to regulate internal pH levels (Sayer et al. 1993). Frommel et al. (2012) found severe to lethal tissue damage for many internal organs in cod raised under three CO₂ levels projected for future scenarios, although adverse effects were remedied once the physiological ability to regulate pH developed within a larva. Cod in this study were from lower-latitude stocks in the northeast Atlantic, and the authors note impacts may be greater in northern populations, as rates of ocean acidification are projected to be higher there (Steinacher et al. 2009).

Finally, changes in the processes that retain larvae on or transport larvae to critical nursery grounds, such as the speed or direction of ocean currents, can also influence larval survival and ultimately recruitment success. Hjort (1914) identified transport and retention of larval fish on nursery areas as critical for recruitment, a conclusion backed by patterns of recruitment in cod (Churchill et al. 2011), haddock (Boucher et al. 2013), and winter flounder (Pearcy 1962; Crawford and Carey 1985). Adaptive responses to patterns in ocean currents allow young cod to settle in particular areas at the correct time to both avoid predators and gain access to abundant food (Lough et al. 1989), and spawning by winter flounder may be timed to utilize the hydrodynamics of small narrow estuaries to maximize egg and larval retention (Pearcy 1962; Pereira et al. 1999). Therefore, changes in currents and circulation as well as conditions in river and estuarine systems may have significant impacts on larval survival and settlement.

Post-settlement life stages

Temperature

Research assessing the impact of climate change on marine fishes often focuses on distributional shifts and behavioral responses to changing conditions (e.g., Murawski 1993; Perry et al. 2005; Harley et al. 2006; Dulvy et al. 2008; Nye et al. 2009; Harley 2011; Simpson et al. 2011; Pinsky and Mantua 2014; Kleisner et al. 2016). Groundfish often move away from even slight temperature changes in the water

column, as observed for cod (Rose et al. 1994; Herbert and Steffensen 2005) and haddock (Eriksen et al. 2012). High temperatures can also alter adult winter flounder migration and distribution (McCracken 1963), and the movement of yellowtail flounder (Walsh and Morgan 2004). Additionally, winter flounder juveniles and adults respond to extreme cold or heat by moving or burying themselves (Topp 1968; Olla et al. 1969; Pereira et al. 1999).

Northward movement to avoid warming waters may come with unexpected consequences, as temperature increases will not impact all populations equally. Populations can exhibit different temperature preferences, as seen in yellowtail flounder (Johnson et al. 1999; Collette and Klein-MacPhee 2002; Simpson and Walsh 2004; Sullivan et al. 2005), and may have adapted differently to local thermal regimes, as observed for cod (Dutil and Brander 2003; Mieszewska et al. 2009). Purchase and Brown (2000) found that, if southern cod stocks move into higher latitudes, they may be less productive and less able to cope with cold periods and thermal anomalies than stocks adapted to those areas. Temperature-driven impacts might also be exacerbated by fishing pressure (Jeffries and Terceiro 1985; Collie et al. 2008) and increased predation (Bell et al. 2014), interactions that can vary with location. Outcomes may not always be detrimental, however. Simpson and Walsh (2004) determined warming waters might have helped some northern stocks of yellowtail flounder rebound from heavy fishing pressure.

Increasing temperatures will impact more than distribution and behavior (Harley et al. 2006). Warmer waters may initially result in increased growth rates, but this may only be to a threshold temperature, after which body size declines (Huey and Berrigan 2001; Baudron et al. 2011; Eriksen et al. 2012). Consequently, adults may ultimately be smaller and less productive in warmer waters. Furthermore, as temperatures increase, other metabolic and physiological rates increase as well (Fonds et al. 1992; MacIsaac et al. 1997), and rising temperatures can therefore affect the fitness and physiology of adult cod (Sartoris et al. 2003), haddock (Baudron et al. 2011), and yellowtail flounder (MacIsaac et al. 1997), and may increase adult mortality in yellowtail flounder (Sullivan et al. 2005) and winter flounder (Nichols 1918; McCracken 1963). Jeffries and Johnson (1974) found that increased mortality of adult winter flounder with

rising temperature in the mid-1960s and early 1970s explained depressed catch rates at that time, a result confirmed more recently by Bell et al. (2014).

Other environmental effects

In addition to temperature, salinity can affect juvenile and adult groundfish. Although generally tolerant to a range of salinities, very low salinity can increase adult mortality in winter flounder (McCracken 1963). Cod can endure low salinity waters (Odense et al. 1966; Dutil et al. 1992; Provencher et al. 1993), but may require sufficient acclimation time, as sudden decreases in salinity can increase mortality (Provencher et al. 1993). Even seemingly tolerable salinities can still have physiological effects, such as altering ionic and osmotic regulation in adult cod (Fletcher 1978a, b), which in turn influences the energy available for growth (Lambert et al. 1994).

Changing oxygen levels may also have consequences for groundfish. Claireaux and Dutil (1992) found nonlinear and detrimental responses in cod from the Gulf of St. Lawrence and Newfoundland waters with increasing hypoxia, including hyperventilation and metabolic acidosis. They determined that effects would be further exacerbated with the greater oxygen requirements needed for digestion, pursuit of prey, or escape from predators. In Long Island Sound, winter flounder exhibited decreased abundance and mean length during low oxygen events, as larger fish may have moved to more oxygen-rich areas (Howell and Simpson 1994). Depressed oxygen can also reduce growth rates (Bejda et al. 1992; Howell and Simpson 1994) and juvenile production (Meng et al. 2001; Stierhoff et al. 2006) in winter flounder. These problems might be exacerbated with rising temperatures if species seek out cooler and deeper waters that are more likely to be hypoxic (Claireaux and Dutil 1992).

Effects of different stressors can interact in complex and deleterious ways. For example, Deutsch et al. (2015) found the combined impact of increased temperatures and decreased dissolved oxygen levels will reduce metabolic indices in the ocean, especially in higher latitudes. Hypoxia might interact with changes in salinity and temperature to influence physiology, and the combination of conditions can exacerbate effects that would otherwise be modest in isolation. Low oxygen conditions in conjunction with

unfavorable changes in salinity place added stress on important physiological processes in cod, including respiration and metabolism (Claireaux and Dutil 1992). The connection between limited oxygen and temperature also influences aerobic capacity and therefore energy budgets (Pörtner et al. 1998), which in turn affect growth, fecundity, and recruitment (Pörtner et al. 2001). Collectively, this research indicates that interactions among environmental stressors can have sub-lethal impacts that manifest as depressed stock productivity rather than direct mortality.

Finally, changes in ocean currents can also affect juvenile groundfish. Lough et al. (1989) argued that young cod and haddock expend more energy swimming in strong currents to stay over preferred bottom. This expenditure could become increasingly significant with escalating stress due to higher temperatures or lack of prey, with consequences for development, growth, condition, and survival.

Reproduction

Temperature effects

Environmental change also affects maturation, fecundity, spawning, and ultimately recruitment. Temperature is inversely correlated with age and length at maturity for cod stocks across the Atlantic (e.g. Jorgensen 1990; O'Brien 1999). When cod mature earlier, the resulting spawning stock is on average younger, smaller, less fecund, and spawns for a shorter period of time (Kjesbu et al. 1996). Similarly, in North Sea haddock, warmer waters resulted in earlier maturation and larger young haddock, but they reached a smaller maximum size overall, resulting in lower spawning stock biomass and yield (Baudron et al. 2011). Deleterious impacts of temperature on adult groundfish may also have indirect effects on reproduction and recruitment via adult condition, especially for females, as maternal contributions can drive recruitment (Brodziak and O'Brien 2005). For cod, condition of females prior to the spawning season directly impacts fecundity (Kjesbu et al. 1991), energy use during spawning (Lambert and Dutil 2000; Ratz and Lloret 2003), and post-spawning health and mortality (Krivobok and Tokareva 1972). In haddock, female fecundity is dependent upon individual body mass (Palmer and Clark 1979; Smith and Morse 1985;

Page and Frank 1989), and in yellowtail flounder there is a non-linear relationship between size and maternal contribution (egg size and yolk quantity), which can be altered by rising temperatures (Benoit and Pepin 1999).

After maturation, changing temperatures can alter the timing of spawning for cod (Hutchings and Myers 1994; Fahay et al. 1999) and haddock (Page and Frank 1989; Cargnelli et al. 1999), which can create mismatches with optimal conditions for larval dispersal, feeding, survival and growth (Rijnsdorp et al. 2009). The timing of haddock spawning is affected by the duration of fall and winter in the previous year (Smith et al. 1981). For winter flounder, abundance is negatively correlated with warmer winter temperatures during spawning (Jeffries and Johnson 1974; Keller and Klein-MacPhee 2000), and winter flounder spawning is also timed to take advantage of both cold winters and spring warming to maximize initial larval size and early growth rates (Buckley et al. 1990; Wilber et al. 2013), which would be disrupted by warmer winters and earlier springs.

Temperature effects will vary by location. Brodziak and O'Brien (2005) found the relative importance of temperature, among other environmental variables, differed across stocks for several groundfish species in the Northwest Atlantic. Warmer temperatures result in earlier recruitment of cod, especially for northern stocks in colder waters (Brander 1994; Drinkwater 2002), and this earlier recruitment may result in smaller, less fecund adults (Kjesbu et al. 1996). However, in southern waters, temperature is negatively correlated with cod recruitment (Ottersen et al. 1994; Planque and Fredou 1999; Ottersen et al. 2006). Differences in temperature effects among populations have also been found in yellowtail flounder (Sissenwine 1974; Sutcliffe et al. 1977). In southern New England winter flounder, Manderson (2008) found warmer spring temperatures correlated with increased synchrony among age-0 fish, and that such synchrony can destabilize regional age-class structure and population dynamics.

Other environmental effects

Reproductive success is affected by environmental factors other than temperature. Sutcliffe et al. (1983) and Myers et al. (1993) found cod recruitment to be positively correlated with salinity, whereas Walsh

et al. (2004) found a negative correlation between salinity and yellowtail flounder recruitment. Larger scale and interacting conditions are also likely critical, as is evident for yellowtail flounder. In the Mid-Atlantic Bight, smoothed recruitment of yellowtail flounder was highly correlated with the North Atlantic Oscillation (NAO) and air temperature (Sullivan et al. 2005). Prior to the collapse of the Grand Bank yellowtail flounder stock, the NAO was strongly positive, resulting in colder air and water, extensive ice cover, and low surface salinity, impacts that were likely exacerbated by fishing pressure (Brodie et al. 2010). Collectively, this suggests the potential importance of salinity and NAO, as well as the potential for differential impacts across space. Finally, changes in currents and circulation may alter the transport to and retention of larvae on critical nursery grounds, which will ultimately impact recruitment success and stock structure of cod and haddock, which have already been altered by human impacts (Ames 1997).

Discussion

Patterns across species

Our review revealed both important insights and gaps in knowledge of the potential response of cod, haddock, winter flounder, and yellowtail flounder to ongoing climatic and oceanographic change. Temperature is one of the most important and widely studied environmental parameters influencing the biology and ecology of marine fish, affecting their physiology (e.g., Fry 1971), migration (e.g., Hoar 1953), and distribution (e.g., Nye et al. 2009). Our review suggests that rising temperatures may have some positive effects on all of the species considered by causing faster growth at multiple life stages, which can move fish through smaller and more vulnerable size classes more quickly, and achieve earlier maturation. Such outcomes can accelerate recovery from depletion through enhanced population growth. However, increased growth also entails metabolic costs, and might come at the expense of future growth, fecundity and egg quality. More importantly, research also highlights many adverse effects of warmer waters that would offset these benefits, especially as temperatures become too high and approach species' upper thermal tolerances. Other impacts driven by

temperature increases might further undermine advantages, such as higher predation on larval winter flounder.

The outcomes of other climactic and oceanographic changes are less well studied than temperature, and our review found virtually no positive effects of expected decreases in pH, salinity, and dissolved oxygen, or potential changes in ocean currents. Alteration of these other environmental parameters appears to be especially important for pelagic life stages of our focal species. This is perhaps not surprising given that eggs and larvae lack well developed physiological regulation mechanisms and the ability to seek out favorable environmental conditions. Also, pelagic life stages are highly responsive to changes in the planktonic community that is similarly affected by these factors.

Collectively, our review did not reveal fundamentally different responses among the focal species to the environmental changes considered. This may be expected, as all of these species co-occur and have co-evolved within the ecosystem. Consideration of species historically rare on Georges Bank and in the Gulf of Maine but expected to become more abundant (e.g., Atlantic croaker; Hare et al. 2010) might reveal very different responses.

That is not to say that certain environmental changes will not affect some species more than others. Cod and haddock in particular exhibit narrower thermal tolerance than the flatfishes (Cargnelli et al. 1999; Fahay et al. 1999; Johnson et al. 1999; Pereira et al. 1999), and might be more vulnerable to adverse effects of warming waters. However, the flatfishes show behaviors and habitat use patterns that might increase their vulnerability to certain changes, such as spawning site fidelity and dependence on localized spawning, nursery, and feeding grounds, demonstrated by winter flounder (McCracken 1963; Howell et al. 1999; Fairchild et al. 2013), or the strong habitat-dependence (e.g. Gibson and Robb 1992; Gibson 1994), limited movements (Morgan and Walsh 1996), and concentration on preferred habitats with abundance declines (Brodie et al. 1998; Simpson and Walsh 2004; Pereira et al. 2012) of yellowtail flounder. These characteristics suggests both flounders may have limited ability to shift as conditions change, and may experience greater stress if there are mismatches between preferred thermal and physical habitat. Furthermore, yellowtail flounder show density-dependent effects on demographic traits (Myers

and Cadigan 1993; Beverton 1995), which will exacerbate environmental stresses. However, other aspects of the biology and ecology of the species here might confer resilience (see “Mechanisms for adaptation” below).

Our results generally align with the conclusions of Fogarty et al. (2008), Pershing et al. (2015), and Hare et al. (2016). The latter concluded winter flounder is the most vulnerable among the four species we considered, followed by cod, with haddock and yellowtail flounder being the least vulnerable. This conclusion is not necessarily evident in the overall summary of our review (Table 2). However, winter flounder and to a lesser extent cod will be affected by changes in a greater diversity of aquatic habitats, in particular shallower nearshore areas that are being more profoundly affected by climate change and other anthropogenic impacts. Our review corroborates some of the more detailed components of the assessment by Hare et al. (2016). For example, they conclude that winter flounder will be more susceptible to changes in oceans currents than the other three species. Our review agrees with this conclusion, although the literature suggests that Hare et al. may be underestimating the effect given the more precise and limited settlement habitats used by winter flounder, and the greater influence of changes in freshwater flow on dispersal and retention.

Interactions among effects

Another critical finding of our review is the importance of understanding how changes in various environmental effects will interact. Studies examining such interactions and their collective consequences for biological attributes of our species of interest are rare. Those that do reveal nonlinear outcomes, and the number of impacts we identify suggests potential for significant cumulative effects. Understanding these combined impacts is critical, as they can result in cohort- and population-level effects that compromise resilience, ecosystem function, and economic value (e.g., Harley et al. 2006; Friedland et al. 2013; Kristiansen et al. 2014).

Figure 3 illustrates how interactions among and accumulated effects of many of the individual outcomes summarized in Table 2 might unfold to shape a population of cod. Increasing temperatures can have both positive and negative effects on the growth of




Fig. 3 Schematic diagram of many of the ways that environmental change can have positive and negative impacts over the life cycle of Atlantic cod (*Gadus morhua*), illustrating the cumulative outcomes and potentially nonlinear interactions among individual effects

larval cod, depending upon the magnitude of the increase, while acidification and especially decreasing salinity are expected to have largely negative effects. The combination of these effects is likely to be greater than that estimated for any in isolation. Higher temperatures and lower salinities are also expected to have adverse effects on survival of eggs and larvae, effects that again might be greater in combination than estimated when considering only one type of environmental change. There are also indirect consequences of interacting environmental stressors both pre- and post-settlement. For the former, the importance of body size for survivorship means that the combined growth effects are likely to reduce survivorship further than that expected due to the direct lethal effects of environmental change. Collectively, this means fewer and smaller fish available to make the pelagic-benthic

transition at the end of the larval stage. The resulting reduced larval pool might further exhibit lower settlement rates due to changes in both ocean currents and water column stratification, as well as fewer or lower quality prey.

After settlement, growth rates and body size are likely to be lower if the incoming cohort of settling larvae is comprised of smaller fish. Increased temperature might offset or exacerbate this effect, again depending upon the magnitude of the increase and other external effects, such as the abundance and quality of prey. Hypoxia can further compromise growth, which can in turn compromise survivorship given size-dependence. Survivorship can also be affected directly by other changes, especially sudden decreases in salinity. On the other hand, higher temperatures can hasten maturation, minimizing the

number of fish that do not reach spawning age due to lower survivorship and increasing near-term productivity. This might have short-term benefits, but at a longer term cost due to the consequences of earlier maturation for future growth, survivorship, fecundity and egg quality. In other words, an apparently counterbalancing environmental effect might mean a fundamental shift in the structure and reproductive success of the spawning population, with implications for the ability of the population to grow and withstand periods of recruitment failure over time. Also, depressed adult fitness may mean lower quality eggs and larvae over time, placing those individuals at a disadvantage from the outset as they confront future environmental stressors.

Mechanisms for adaptation

Although we identify generally adverse effects of environmental change, which may be even greater in combination, our focal species all have attributes that may help them adapt. Metapopulation structure can provide inherent resilience for a species because not all environmental changes will be experienced homogeneously across a species' range (Planque et al. 2010). Subpopulations that are affected less can buffer against depletion or even localized extinction elsewhere (Kritzer and Sale 2004), although spatial population structure will not be advantageous in the face of adverse environmental changes that are experienced at the stock or metapopulation scale (Koenig 2002). Further, serial depletion of subpopulations can erode the potential for bet-hedging and recolonization benefits. Among our focal species, the spatial structure of cod populations has been especially well documented (Ames 2004; Lage et al. 2004; Wirgin et al. 2007; Kovach et al. 2010; Runge et al. 2010), which has illuminated the costs of serial depletion at the metapopulation scale (Smedbol and Wroblewski 2002; Steneck and Wilson 2010).

Spatial structure also creates the potential for behavioral and life history traits that are tailored to different conditions through localized adaptation and based upon genetic diversity (Sale et al. 2006). Localized adaptation can be a disadvantage at the individual or population scale if new conditions deviate too much from those that shaped the adaptation. However, at a larger scale across a metapopulation, a higher degree of genetic diversity increases the

ability to adapt to new conditions. Spatial patterns in genetic diversity have been described for both cod (Hauser and Carvalho 2008; Kovach et al. 2010) and winter flounder (Howell et al. 1992). These differences are maintained in cod by spawning patterns (Kovach et al. 2010) and larval retention strategies (Mieszowska et al. 2009; Runge et al. 2010), and in winter flounder by a high degree of site fidelity (Howe and Coates 1975; Phelan 1992; Gibson 2013). Unfortunately, our review suggests that both of these strategies might be compromised by changing conditions.

Although spatial structure and genetic, behavioral, and life history diversity can confer resilience and allow for adaptation at the metapopulation scale, certain behavioral and life history attributes can also contribute to resilience and adaptation at smaller scales. Winter flounder are generally viewed as an inshore spawning species, but exhibit a greater degree of spawning in deeper offshore waters than is generally appreciated (DeCelles and Cadrin 2010). This plasticity over relatively small scales might allow for localized adaptation in the future. Mobility can allow fish to seek out a better location if local conditions become unfavorable, provided of course that suitable habitat in new locations has not been unduly affected by climate change or other impacts (Kritzer et al. 2016). Cod generally exhibit the longest range movements among our focal species, especially in the Northwest Atlantic (Robichaud and Rose 2004), which should allow stronger positive responses to environmental change.

One potential downside of greater mobility is that fewer benefits of marine protected areas (MPAs) will be realized (Sherwood and Grabowski 2010). MPAs can also allow recovery of complete age and size structure (Berkeley et al. 2004), and this demographic diversity can contribute to resilience given the robustness and high reproductive value of older and larger fish (Planque et al. 2010). Fortunately, the behavioral diversity exhibited by cod has allowed some fish to benefit from spatial management in New England, resulting in more complete age and size structure within MPAs (Sherwood and Grabowski 2015). Haddock, winter flounder and yellowtail flounder all exhibit larger body size in MPAs as well (Link et al. 2005). Because impacts of environmental changes should be experienced across MPA boundaries, these outcomes underscore the continued importance of fishing pressure as a demographic driver. A

recent analysis of long-term trends in cod populations across the North Atlantic reveals that effects of fishing mortality are comparable to environmental effects (Frank et al. 2016). Therefore, the observed differences in population structure across MPA boundaries suggest that spatial protection could become especially important in buffering the effects of fishing pressure in the face of rising and compounding effects of environmental change.

Implications for research and management

Our review revealed that cod is the most widely studied among the species we considered, and that effects of temperature are the most widely studied among the environmental factors. This suggests that more research on the other species and environmental factors is needed. However, the disparity in research between cod and the other species is less than that between temperature and the other factors (Tables 1, 2). Therefore, future research should focus on acidification, salinity, hypoxia, ocean currents, and their interactions. The vulnerability assessment by Hare et al. (2016) supports prioritization of research on ocean acidification in particular, given that its magnitude is expected to increase substantially in the coming decades. Furthermore, Hare et al. conclude that the exposure of all four species to acidification will be very high, but their sensitivity will be relatively low. These divergent outcomes raise important questions about whether the effects will be significant or not, and our review reveals that the existing literature offers little direct insight (Tables 1, 2).

Future studies should examine as broad a taxonomic spectrum as possible. We considered four of the 15 groundfish species managed in New England, and among many more managed and unmanaged species in the region. We found generally consistent trends in the effects examined, at least in terms of directionality if not magnitude, but within a single functional group (c.f., Hare et al. 2016; Kleisner et al. 2016). Given the high dimensionality of a complete research matrix for the region (i.e., [environmental factors \times species \times life stages \times life history traits]), or an expanded version of Table 2), the most efficient and effective strategy will aim to extract as many general trends as possible. This would allow robust assumptions to be made where knowledge gaps inevitably remain. Furthermore, considerably more work is needed on the interactions

among effects, including accumulated and synergistic impacts within and across life stages. In fact, this should arguably be the highest priority for future research, given that predictions for individual effects alone will likely underestimate aggregate outcomes (Fig. 3).

Building new science that increases understanding of the effects of climatic and oceanographic changes will take time, but management strategies will need to adapt in the near-term both to what we know and to the many uncertainties. Fishing will remain a dominant driver of groundfish population dynamics, and will interact with climate change (Harley et al. 2006; Araujo and Bundy 2012; Bundy et al. 2012) and increase the sensitivity of fish stocks to climate perturbations (Anderson et al. 2008; Hsieh et al. 2008; Pinsky and Mantua 2014). Therefore, harvest rates will need to be adjusted to contemporary productivity resulting from environmental change (Brander 2010; Planque et al. 2010; Nye et al. 2013). Fogarty et al. (2008) and Pershing et al. (2015) have re-estimated key biological parameters and reference points for cod in light of the potential impacts of climate change, illustrating the lower biomass and fishing mortality targets to which we should be managing.

In addition, the diversity of physical, chemical and biological changes unfolding in the environment means that new analytical tools will be needed to account for the impact of changes and their interactions in determining management targets. This may mean greater use of multispecies production models (e.g., Fogarty et al. 2012), especially in the face of changing species composition within management areas, supplemented by models of intermediate complexity for ecosystem assessment (MICE) to inform those models on environmental effects (Collie et al. 2014; Plaganyi et al. 2014). However, catch limits derived from either single- or multi-species models alone are unlikely to account for the full suite of changes, complexities, and uncertainties we face. Therefore, spatial management measures that protect ecosystem attributes such as habitat, biodiversity, and trophic structure (McCleod et al. 2009), and rebuild spatial structure, age structure and size structure (Kritzer and Sale 2004; Planque et al. 2010) will also be critical.

Better integration among fisheries management entities spanning municipal, state and national boundaries is already needed in light of the fact that many

species do not respect these boundaries over the course of their life history (Kritzer and Liu 2013). Beyond governance of fisheries, more integrated management of different human uses that contribute to climate change and other environmental stressors will be necessary to minimize impacts on and maximize the value derived from a variety of ecosystem services (Crowder et al. 2006). In general, managers, fishermen and scientists will need to be more responsive, adaptive and risk-averse in light of ongoing environmental change and the uncertainties inherent in how it will impact groundfish and other species.

Acknowledgments This review was prepared through support from the Gordon and Betty Moore Foundation to the Environmental Defense Fund (EDF). The authors benefited from discussions with the Northeast Shelf Fishery Ecosystem Work Group convened by EDF. JPK is especially grateful to his colleagues on the New England Fishery Management Council's Scientific and Statistical Committee, with whom he grapples on a regular basis with the practical implications of changes in the region's marine ecosystem. However, the content of this review and any errors are the responsibility of the authors alone.

References

- Ames EP (1997) Cod and haddock spawning grounds in the Gulf of Maine: from Grand Manan Channel to Ipswich Bay. The Island Institute, Rockland
- Ames EP (2004) Atlantic cod stock structure in the Gulf of Maine. *Fisheries* 29:10–28
- Ames EP, Lichter J (2013) Gadids and alewives: structure within complexity in the Gulf of Maine. *Fish Res* 141:70–78
- Anderson CNK, Hsieh CH, Sandin SA, Hewitt R, Hollowed A, Beddington J, May RM, Sugihara G (2008) Why fishing magnifies fluctuations in fish abundance. *Nature* 452: 835–839
- Araujo JN, Bundy A (2012) Effects of environmental change, fisheries and trophodynamics on the ecosystem of the western Scotian Shelf, Canada. *Mar Ecol Prog Ser* 464: 51–67. doi:10.3354/meps09792
- Balch WM, Drapeau DT, Bowler BC, Huntington TG (2012) Step-changes in the physical, chemical and biological characteristics of the Gulf of Maine, as documented by the GNATS time series. *Mar Ecol Prog Ser* 450:11–35
- Baudron AR, Needle CL, Marshall CT (2011) Implications of a warming North Sea for the growth of haddock *Melanogrammus aeglefinus*. *J Fish Biol* 78:1874–1889. doi:10.1111/j.1095-8649.2011.02940.x
- Beaugrand G (2004) The North Sea regime shift: evidence, causes, mechanisms and consequences. *Prog Oceanogr* 60:245–262. doi:10.1016/j.pocean.2004.02.018
- Beaugrand G, Reid PC, Ibanez F, Lindley JA, Edwards M (2002) Reorganization of North Atlantic marine copepod biodiversity and climate. *Science* 296:1692–1694. doi:10.1126/science.1071329
- Beaugrand G, Brander KM, Lindley JA, Souissi S, Reid PC (2003) Plankton effect on cod recruitment in the North Sea. *Nature* 426:661–664. doi:10.1038/nature02164
- Bejda AJ, Phelan BA, Studholme AL (1992) The effect of dissolved-oxygen on the growth of young-of-the-year winter flounder, *Pseudopleuronectes americanus*. *Environ Biol Fish* 34:321–327. doi:10.1007/bf00004780
- Bell RJ, Hare JA, Manderson JP, Richardson DE (2014) Externally driven changes in the abundance of summer and winter flounder. *ICES J Mar Sci*. doi:10.1093/icesjms/fsu069
- Benoit HP, Pepin P (1999) Interaction of rearing temperature and maternal influence on egg development rates and larval size at hatch in yellowtail flounder (*Pleuronectes ferrugineus*). *Can J Fish Aquat Sci* 56:785–794. doi:10.1139/cjfas-56-5-785
- Berkeley SA, Hixon MA, Larson RJ, Love MS (2004) Fisheries sustainability via protection of age structure and spatial distribution of fish populations. *Fisheries* 29:23–32
- Beverton RJH (1995) Spatial limitation of population size; the concentration hypothesis. *Neth J Sea Res* 34:1–6. doi:10.1016/0077-7579(95)90010-1
- Blaxter JHS (1986) Development of sense-organs and behavior of teleost larvae with special reference to feeding and predator avoidance. *Trans Am Fish Soc* 115:98–114
- Boucher JM, Chen CS, Sun YF, Beardsley RC (2013) Effects of interannual environmental variability on the transport-retention dynamics in haddock *Melanogrammus aeglefinus* larvae on Georges Bank. *Mar Ecol Prog Ser* 487:201–215. doi:10.3354/meps10462
- Boyce DG, Lewis MR, Worm B (2010) Global phytoplankton decline over the past century. *Nature* 466:591–596. doi:10.1038/nature09268
- Brander KM (1994) Patterns of distribution, spawning, and growth in Northern Atlantic cod—the utility of inter-regional comparisons. In: Jakobsson J, Atthorsson OS, Beverton RJH et al (eds) Cod and climate change—proceedings of a symposium. *ICES Mar Sci Symp* 198: 406–413
- Brander KM (1995) The effect of temperature on growth of Atlantic cod (*Gadus morhua* L.). *ICES J Mar Sci* 52:1–10. doi:10.1016/1054-3139(95)80010-7
- Brander K (2010) Impacts of climate change on fisheries. *J Mar Syst* 79:389–402. doi:10.1016/j.jmarsys.2008.12.015
- Brodie WB, Walsh SJ, Atkinson DB (1998) The effect of stock abundance on range contraction of yellowtail flounder (*Pleuronectes ferrugineus*) on the Grand Bank of Newfoundland in the Northwest Atlantic from 1975 to 1995. *J Sea Res* 39:139–152. doi:10.1016/s1385-1101(97)00056-7
- Brodie WB, Walsh SJ, Parsons DM (2010) An evaluation of the collapse and recovery of the yellowtail flounder (*Limanda ferruginea*) stock on the Grand Bank. *ICES J Mar Sci* 67:1887–1895. doi:10.1093/icesjms/fsq121
- Brodziak J, O'Brien L (2005) Do environmental factors affect recruits per spawner anomalies of New England groundfish? *ICES J Mar Sci* 62:1394–1407. doi:10.1016/j.icesjms.2005.04.019

- Buckley J (1989) Species profiles: life histories and environmental requirements of coastal fish and invertebrates (North Atlantic)—winter flounder. US Fish and Wildlife Service Biological Reports, vol 82(11.87)/TR EL-82-4
- Buckley LJ, Smigielski AS, Halavik TA, Laurence GC (1990) Effects of water temperature on size and biochemical composition of winter flounder *Pseudopleuronectes americanus* at hatching and feeding initiation. Fish Bull 88:419–428
- Buckley LJ, Lough RG, Mountain D (2010) Seasonal trends in mortality and growth of cod and haddock larvae result in an optimal window for survival. Mar Ecol Prog Ser 405:57–69. doi:[10.3354/meps08503](https://doi.org/10.3354/meps08503)
- Bundy A, Coll M, Shannon LJ, Shin Y (2012) Global assessments of the status of marine exploited ecosystems and their management: what more is needed? Curr Opin Environ Sustain 4:292–299
- Caldeira K, Wickett ME (2003) Anthropogenic carbon and ocean pH. Nature 425:365. doi:[10.1038/425365a](https://doi.org/10.1038/425365a)
- Caldeira K, Wickett ME (2005) Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean. J Geophys Res C Oceans. doi:[10.1029/2004jc002671](https://doi.org/10.1029/2004jc002671)
- Capotondi A, Alexander MA, Bond NA, Curchitser EN, Scott JD (2012) Enhanced upper ocean stratification with climate change in the CMIP3 models. J Geophys Res C Oceans. doi:[10.1029/2011jc007409](https://doi.org/10.1029/2011jc007409)
- Cargnelli LM, Griesbach SJ, Berrien PL, Morse WM, Johnson DL (1999) Essential fish habitat source document: haddock, *Melanogrammus aeglefinus*, life history and habitat characteristics. NOAA Technical Memorandum NMFS-NE-128
- Choi JS, Frank KT, Leggett WC, Drinkwater K (2004) Transition to an alternate state in a continental shelf ecosystem. Can J Fish Aquat Sci 61:505–510. doi:[10.1139/f04-079](https://doi.org/10.1139/f04-079)
- Churchill JH, Runge J, Chen CS (2011) Processes controlling retention of spring-spawned Atlantic cod (*Gadus morhua*) in the western Gulf of Maine and their relationship to an index of recruitment success. Fish Oceanogr 20:32–46. doi:[10.1111/j.1365-2419.2010.00563.x](https://doi.org/10.1111/j.1365-2419.2010.00563.x)
- Claireaux G, Dutil JD (1992) Physiological response of the Atlantic cod (*Gadus morhua*) to hypoxia at various environmental salinities. J Exp Biol 163:97–118
- Collette BB, Klein-MacPhee G (2002) Fishes of the Gulf of Maine, 3rd edn. Smithsonian Institution Press, Washington, DC
- Collie JS, Wood AD, Jeffries HP (2008) Long-term shifts in the species composition of a coastal fish community. Can J Fish Aquat Sci 65:1352–1365
- Collie JS, Botsford LW, Hastings A, Kaplan IC, Largier JL, Livingston PA, Plaganyi E, Rose KA, Wells BK, Werner FE (2014) Ecosystem models for fisheries management: finding the sweet spot. Fish Fish 17:101–125. doi:[10.1111/faf.12093](https://doi.org/10.1111/faf.12093)
- Colton JB (1959) A field observation of mortality of marine fish larvae due to warming. Limnol Oceanogr 4:219–222. doi:[10.4319/lo.1959.4.2.0219](https://doi.org/10.4319/lo.1959.4.2.0219)
- Costello C, Gaines SD, Lynham J (2008) Can catch shares prevent fisheries collapse? Science 321:1678–1681. doi:[10.1126/science.1159478](https://doi.org/10.1126/science.1159478)
- Crawford RE, Carey CG (1985) Retention of winter flounder larvae within a Rhode Island salt pond. Estuaries 8:217–227. doi:[10.2307/1352202](https://doi.org/10.2307/1352202)
- Crowder LB, Osherenko G, Young OR, Airame S, Norse EA, Baron N, Day JC, Bouvere F, Ehler CN, Halpern BS, Langdon SJ, McLeod KL, Ogden JC, Peach RE, Rosenberg AA, Wilson JA (2006) Resolving mismatches in US ocean governance. Science 313:617–618
- Davenport J, Lonning S, Kjorsvik E (1981) Osmotic and structural changes during early development of eggs and larvae of the cod, *Gadus morhua* L. J Fish Biol 19:317–331. doi:[10.1111/j.1095-8649.1981.tb05835.x](https://doi.org/10.1111/j.1095-8649.1981.tb05835.x)
- DeCelles GR, Cadrin SX (2010) Movement patterns of winter flounder (*Pseudopleuronectes americanus*) in the southern Gulf of Maine: observations with the use of passive acoustic telemetry. Fish Bull 108:408–419
- Deutsch C, Ferrel A, Seibel B, Pörtner HO, Huey RB (2015) Climate change tightens a metabolic constraint on marine habitats. Science 348:1132–1135. doi:[10.1126/science.1251605](https://doi.org/10.1126/science.1251605)
- Dickie LM, McCracken FD (1955) Isoleth diagrams to predict equilibrium yields of a small flounder fishery. J Fish Res Board Can 12:187–209
- Doney SC, Ruckelshaus M, Duffy JE, Barry JP, Chan F, English CA, Galindo HM, Grebmeier JM, Hollowed AB, Knowlton N, Polovina J, Rabalais NN, Sydeman WJ, Talley LD (2012) Climate change impacts on marine ecosystems. Ann Rev Mar Sci 4:11–37. doi:[10.1146/annurev-marine-041911-111611](https://doi.org/10.1146/annurev-marine-041911-111611)
- Drinkwater KF (2002) A review of the role of climate variability in the decline of northern cod. In: McGinn NA (ed) Fisheries in a changing climate. Am Fish Soc Symp 32:113–129
- Dulvy NK, Rogers SI, Jennings S, Stelzenmuller V, Dye SR, Skjoldal HR (2008) Climate change and deepening of the North Sea fish assemblage: a biotic indicator of warming seas. J Appl Ecol 45:1029–1039. doi:[10.1111/j.1365-2664.2008.01488.x](https://doi.org/10.1111/j.1365-2664.2008.01488.x)
- Dutil JD, Brander K (2003) Comparing productivity of North Atlantic cod (*Gadus morhua*) stocks and limits to growth production. Fish Oceanogr 12:502–512. doi:[10.1046/j.1365-2419.2003.00243.x](https://doi.org/10.1046/j.1365-2419.2003.00243.x)
- Dutil JD, Munro J, Audet C, Besner M (1992) Seasonal variation in the physiological response of Atlantic cod (*Gadus morhua*) to low salinity. Can J Fish Aquat Sci 49:1149–1156. doi:[10.1139/f92-128](https://doi.org/10.1139/f92-128)
- Eriksen E, Ingvaldsen R, Stiansen JE, Johansen GO (2012) Thermal habitat for 0-group fish in the Barents Sea; how climate variability impacts their density, length, and geographic distribution. ICES J Mar Sci 69:870–879. doi:[10.1093/icesjms/fsr210](https://doi.org/10.1093/icesjms/fsr210)
- Essington TE (2010) Ecological indicators display reduced variation in North American catch share fisheries. Proc Natl Acad Sci USA 107:754–759. doi:[10.1073/pnas.0907252107](https://doi.org/10.1073/pnas.0907252107)
- Essington TE, Melnychuk MC, Branch TA, Heppell SS, Jensen OP, Link JS, Martell SJD, Parma AM, Pope JG, Smith ADM (2012) Catch shares, fisheries, and ecological stewardship: a comparative analysis of resource responses to a rights-based policy instrument. Conserv Lett 5:186–195. doi:[10.1111/j.1755-263X.2012.00226.x](https://doi.org/10.1111/j.1755-263X.2012.00226.x)

- Fahay MP, Berrien PL, Johnson DL, Morse WM (1999) Essential fish habitat source document: Atlantic cod, *Gadus morhua*, life history and habitat characteristics. NOAA Technical Memorandum NMFS-NE-124
- Fairchild EA, Siceloff L, Howell WH, Hoffman B, Armstrong MP (2013) Coastal spawning by winter flounder and a reassessment of essential fish habitat in the Gulf of Maine. *Fish Res* 141:118–129
- Feeley RA, Doney SC, Cooley SR (2009) Ocean acidification: present conditions and future changes in a high-CO₂ world. *Oceanography* 22:36–47
- Fernandez IJ, Schmitt CV, Birkel SD, Stancioff E, Pershing AJ, Kelley JT, Runge JA, Jacobson GL, Mayewski PA (2015) Maine's climate future: 2015 update. University of Maine, Orono, p 24
- Fletcher CR (1978a) Osmotic and ionic regulation in cod (*Gadus callarias* L): 1. Water balance. *J Comp Physiol* 124:149–155
- Fletcher CR (1978b) Osmotic and ionic regulation in cod (*Gadus callarias* L): 2. Salt balance. *J Comp Physiol* 124:157–168
- Fogarty M, Incze L, Hayhoe K, Mountain D, Manning J (2008) Potential climate change impacts on Atlantic cod (*Gadus morhua*) off the northeastern USA. *Mitigation Adapt Strateg Glob Chang* 13:453–466. doi:10.1007/s11027-007-9131-4
- Fogarty MJ, Overholtz WJ, Link JS (2012) Aggregate surplus production models for demersal fishery resources of the Gulf of Maine. *Mar Ecol Prog Ser* 459:247–258
- Fonds M, Cronie R, Vethaak AD, Vanderpuyl P (1992) Metabolism, food consumption and growth of plaice (*Pleuronectes platessa*) and flounder (*Platichthys flesus*) in relation to fish size and temperature. *Neth J Sea Res* 29:127–143. doi:10.1016/0077-7579(92)90014-6
- Frank KT, Petrie B, Leggett WC, Boyce DG (2016) Large scale, synchronuous variability of marine fish populations driven by commercial exploitation. *Proc Nat Acad Sci USA*. doi:10.1073/pnas.1602325113
- Friedland KD, Kane J, Hare JA, Lough RG, Fratantoni PS, Fogarty MJ, Nye JA (2013) Thermal habitat constraints on zooplankton species associated with Atlantic cod (*Gadus morhua*) on the US Northeast Continental Shelf. *Prog Oceanogr* 116:1–13. doi:10.1016/j.pocean.2013.05.011
- Frommel AY, Maneja R, Lowe D, Malzahn AM, Geffen AJ, Folkvord A, Piatkowski U, Reusch TBH, Clemmesen C (2012) Severe tissue damage in Atlantic cod larvae under increasing ocean acidification. *Nate Clim Change* 2:42–46
- Fry FEJ (1971) The effect of environmental factors on the physiology of fish. In: Hoar WS, Randall DJ (eds) *Fish physiology*, vol 6. Academic Press, pp 1–98. doi:10.1016/S1546-5098(08)60146-6
- Garrison LP, Link JS (2000) Fishing effects on spatial distribution and trophic guild structure of the fish community in the Georges Bank region. *ICES J Mar Sci* 57:723–730
- Gibson RN (1994) Impact of habitat quality and quantity on the recruitment of juvenile flatfishes. *Neth J Sea Res* 32:191–206. doi:10.1016/0077-7579(94)90040-x
- Gibson MR (2013) Assessing the local population of winter flounder with a two-era biomass dynamic model: a narrower view of Southern New England. Rhode Island Division of Fish and Wildlife, Jamestown
- Gibson RN, Robb L (1992) The relationship between body size, sediment grain size and the burying ability of juvenile plaice, *Pleuronectes platessa* L. *J Fish Biol* 40:771–778. doi:10.1111/j.1095-8649.1992.tb02623.x
- Grimm D, Barkhorn I, Festa D, Bonzon K, Boomhower J, Hovland V, Blau J (2012) Assessing catch shares' effects evidence from federal United States and associated British Columbian fisheries. *Mar Policy* 36:644–657. doi:10.1016/j.marpol.2011.10.014
- Gutierrez NL, Hilborn R, Defeo O (2011) Leadership, social capital and incentives promote successful fisheries. *Nature* 470:386–389. doi:10.1038/nature09689
- Hall CJ, Jordaan A, Frisk MG (2012) Centuries of anadromous forage fish loss: consequences for ecosystem connectivity and productivity. *Bioscience* 62:723–731
- Hare JA, Alexander MA, Fogarty MJ, Williams EH, Scott JD (2010) Forecasting the dynamics of a coastal fishery species using a coupled climate-population model. *Ecol App* 20:452–464. doi:10.1890/08-1863.1
- Hare JA, Morrison WE, Nelson MW, Stachura MM, Teeters EJ, Griffis RB, Alexander MA, Scott JD, Alade L, Bell RJ, Chute AS, Curti KL, Curtis TH, Kircheis D, Kocik JF, Lucey SM, McCandless CT, Milke LM, Richardson DE, Robillard E, Walsh HJ, McManus MC, Marancik KE, Griswold CA (2016) A vulnerability assessment of fish and invertebrates to climate change on the Northeast U.S. continental shelf. *PLoS ONE* 11:e0146756. doi:10.1371/journal.pone.0146756
- Harley CDG (2011) Climate change, keystone predation, and biodiversity loss. *Science* 334:1124–1127. doi:10.1126/science.1210199
- Harley CDG, Hughes AR, Hultgren KM, Miner BG, Sorte CJB, Thornber CS, Rodriguez LF, Tomanek L, Williams SL (2006) The impacts of climate change in coastal marine systems. *Ecol Lett* 9:228–241. doi:10.1111/j.1461-0248.2005.00871.x
- Hauser L, Carvalho GR (2008) Paradigm shifts in marine fisheries genetics: ugly hypotheses slain by beautiful facts. *Fish Fish* 9:333–362
- Herbert NA, Steffensen JF (2005) The response of Atlantic cod, *Gadus morhua*, to progressive hypoxia: fish swimming speed and physiological stress. *Mar Biol* 147:1403–1412. doi:10.1007/s00227-005-0003-8
- Hjort J (1914) Fluctuations in the great fisheries of Northern Europe viewed in the light of biological research. Copenhagen conseil permanent international pour l'exploration de la mer. Rapports et Proces-Verbaux des Reunions 20:234
- Hoar WS (1953) Control and timing of fish migration. *Biol Rev* 28:437–452. doi:10.1111/j.1469-185X.1953.tb01387.x
- Hoegh-Guldberg O, Bruno JF (2010) The impact of climate change on the world's marine ecosystems. *Science* 328:1523–1528. doi:10.1126/science.1189930
- Howe AB, Coates PG (1975) Winter flounder movements, growth, and mortality off Massachusetts. *Trans Am Fish Soc* 104:13–29. doi:10.1577/1548-8659(1975)104<13:wfmam>2.0.co;2
- Howell P, Simpson D (1994) Abundance of marine resources in relation to dissolved oxygen in Long Island Sound. *Estuaries* 17:394–402. doi:10.2307/1352672

- Howell P, Howe A, Gibson M, Ayvazian S (1992) Fishery management plan for inshore stocks of winter flounder, *Pleuronectes americanus*. ASMFC fisheries management report 21
- Howell PT, Molnar DR, Harris RB (1999) Juvenile winter flounder distribution by habitat type. *Estuaries* 22:1090–1095. doi:[10.2307/1353086](https://doi.org/10.2307/1353086)
- Hsieh CH, Reiss CS, Hewitt RP, Sugihara G (2008) Spatial analysis shows that fishing enhances the climatic sensitivity of marine fishes. *Can J Fish Aquat Sci* 65:947–961
- Huey RB, Berrigan D (2001) Temperature, demography, and ectotherm fitness. *Am Nat* 158:204–210
- Hutchings JA, Myers RA (1994) Timing of cod reproduction—interannual variability and the influence of temperature. *Mar Ecol Prog Ser* 108:21–31. doi:[10.3354/meps108021](https://doi.org/10.3354/meps108021)
- Jeffries HP, Johnson WC (1974) Seasonal distributions of bottom fishes in Narragansett Bay area—7 year variations of abundance of winter flounder (*Pseudopleuronectes americanus*). *J Fish Res Board Can* 31:1057–1066
- Jeffries HP, Terceiro M (1985) Cycle of changing abundance in the fishes of the Narragansett Bay area. *Mar Ecol Prog Ser* 25:239–244. doi:[10.3354/meps025239](https://doi.org/10.3354/meps025239)
- Jeffries P, Keller A, Hale S (1989) Predicting winter flounder (*Pseudopleuronectes americanus*) catches by time-series analysis. *Can J Fish Aquat Sci* 46:650–659
- Johnson DL, Morse WM, Berrien PL, Vitaliano JJ (1999) Essential fish habitat source document: yellowtail flounder, *Limanda ferruginea*, life history and habitat characteristics. NOAA Technical Memorandum NMFS-NE-140
- Jorgensen T (1990) Long-term changes in age at sexual maturity of Northeast Arctic cod (*Gadus morhua* L.). *ICES J Mar Sci* 46:235–248
- Keller AA, Klein-MacPhee G (2000) Impact of elevated temperature on the growth, survival, and trophic dynamics of winter flounder larvae: a mesocosm study. *Can J Fish Aquat Sci* 57:2382–2392. doi:[10.1139/cjfas-57-12-2382](https://doi.org/10.1139/cjfas-57-12-2382)
- Kjesbu OS, Klungsoyr J, Kryvi H, Witthames PR, Walker MG (1991) Fecundity, atresia, and egg size of captive Atlantic cod (*Gadus morhua*) in relation to proximate body composition. *Can J Fish Aquat Sci* 48:2333–2343. doi:[10.1139/f91-274](https://doi.org/10.1139/f91-274)
- Kjesbu OS, Solemdal P, Bratland P, Fonn M (1996) Variation in annual egg production in individual captive Atlantic cod (*Gadus morhua*). *Can J Fish Aquat Sci* 53:610–620. doi:[10.1139/cjfas-53-3-610](https://doi.org/10.1139/cjfas-53-3-610)
- Kleisner KM, Fogarty MJ, McGee S, Barnett A, Fratantoni P, Greene J, Hare JA, Lucey SM, McGuire C, Odell J, Saba VS, Smith L, Weaver KJ, Pinsky ML (2016) The effects of sub-regional climate velocity on the distribution and spatial extent of marine species assemblages. *PLoS ONE* 11:e0149220. doi:[10.1371/journal.pone.0149220](https://doi.org/10.1371/journal.pone.0149220)
- Koenig WD (2002) Global patterns of environmental synchrony and the Moran effect. *Ecography* 25:283–288
- Koester FW, Hinrichsen H-H, St. John MA, Schnack D, MacKenzie BR, Tomkiewicz J, Plikshs M (2001) Developing Baltic cod recruitment models. II. Incorporation of environmental variability and species interactions. *Can J Fish Aquat Sci* 58:1534–1556
- Kovach AI, Breton TS, Berlinsky DL, Maceda L, Wirgin I (2010) Fine-scale spatial and temporal genetic structure of Atlantic cod off the Atlantic coast of the USA. *Mar Ecol Prog Ser* 410:177–195. doi:[10.3354/meps08612](https://doi.org/10.3354/meps08612)
- Kristiansen T, Drinkwater KF, Lough RG, Sundby S (2011) Recruitment variability in North Atlantic cod and match-mismatch dynamics. *PLoS ONE* 6:e17456. doi:[10.1371/journal.pone.0017456](https://doi.org/10.1371/journal.pone.0017456)
- Kristiansen T, Stock C, Drinkwater KF, Curchitser EN (2014) Mechanistic insights into the effects of climate change on larval cod. *Glob Change Biol* 20:1559–1584. doi:[10.1111/gcb.12489](https://doi.org/10.1111/gcb.12489)
- Kritzer JP, Liu OR (2013) Fishery management strategies for addressing complex spatial structure in marine fish stocks. In: Cadrin SX, Kerr LA, Mariani S (eds) Stock identification methods, 2nd edn. Elsevier, Boston, pp 29–58
- Kritzer JP, Sale PF (2004) Metapopulation ecology in the sea: from Levins' model to marine ecology and fisheries science. *Fish Fish* 5:131–140
- Kritzer JP, Delucia M, Greene E, Shumway C, Topolski MF, Thomas-Blate J, Chiarella L, Davy KB, Smith K (2016) The importance of benthic habitats for coastal fisheries. *Bioscience*. doi:[10.1093/biosci/biw014](https://doi.org/10.1093/biosci/biw014)
- Krivobok MN, Tokareva GI (1972) Dynamics of weight variations of the body and separate organs of Baltic cod during the maturation of gonads. *Trudy vses. nauchno-issled. Inst Morsk Ryb Khoz Okeanogr* 85:46–55
- Lage C, Kuhn K, Kornfield I (2004) Genetic differentiation among Atlantic cod (*Gadus morhua*) from Browns Bank, Georges Bank, and Nantucket Shoals. *Fish Bull* 102:289–297
- Lambert Y, Dutil JD (2000) Energetic consequences of reproduction in Atlantic cod (*Gadus morhua*) in relation to spawning level of somatic energy reserves. *Can J Fish Aquat Sci* 57:815–825. doi:[10.1139/cjfas-57-4-815](https://doi.org/10.1139/cjfas-57-4-815)
- Lambert Y, Dutil JD, Munro J (1994) Effects of intermediate and low-salinity conditions on growth rate and food conversion of Atlantic cod (*Gadus morhua*). *Can J Fish Aquat Sci* 51:1569–1576. doi:[10.1139/f94-155](https://doi.org/10.1139/f94-155)
- Laurence GC (1975) Laboratory growth and metabolism of winter flounder *Pseudopleuronectes americanus* from hatching through metamorphosis at 3 temperatures. *Mar Biol* 32:223–229. doi:[10.1007/bf00399202](https://doi.org/10.1007/bf00399202)
- Laurence GC (1977) Bioenergetic model for analysis of feeding and survival potential of winter flounder, *Pseudopleuronectes americanus*, larvae during period from hatching to metamorphosis. *Fish Bull* 75:529–546
- Laurence GC (1978) Comparative growth, respiration and delayed feeding abilities of larval cod (*Gadus morhua*) and haddock (*Melanogrammus aeglefinus*) as influenced by temperature during laboratory studies. *Mar Biol* 50:1–7. doi:[10.1007/bf00390536](https://doi.org/10.1007/bf00390536)
- Laurence GC, Howell WH (1981) Embryology and influence of temperature and salinity on early development and survival of yellowtail flounder *Limanda ferruginea*. *Mar Ecol Prog Ser* 6:11–18. doi:[10.3354/meps006011](https://doi.org/10.3354/meps006011)
- Laurence GC, Rogers CA (1976) Effects of temperature and salinity on comparative embryo development and mortality of Atlantic cod (*Gadus morhua* L.) and haddock (*Melanogrammus aeglefinus* L.). *ICES J Mar Sci* 36:220–228
- Link J, Almeida F, Valentine P, Auster P, Reid R, Vitaliano J (2005) The effects of closed areas on Georges Bank. *Am Fish Soc Symp* 41:345–368

- Lough RG (2004) Essential fish habitat source document: Atlantic cod, *Gadus morhua*, life history and habitat characteristics. 2nd edn. NOAA Technical Memorandum NMFS-NE 190:i-vi, 1–94
- Lough RG, Valentine PC, Potter DC, Audire PJ, Bolz GR, Neilson JD, Perry RI (1989) Ecology and distribution of juvenile cod and haddock in relation to sediment type and bottom currents on eastern Georges Bank. *Mar Ecol Prog Ser* 56:1–12. doi:[10.3354/meps056001](https://doi.org/10.3354/meps056001)
- MacIsaac PF, Goff GP, Speare DJ (1997) Comparison of routine oxygen consumption rates of three species of pleuronectids at three temperatures. *J Appl Ichthyol* 13:171–176. doi:[10.1111/j.1439-0426.1997.tb00117.x](https://doi.org/10.1111/j.1439-0426.1997.tb00117.x)
- Manderson JP (2008) The spatial scale of phase synchrony in winter flounder (*Pseudopleuronectes americanus*) production increased among southern New England nurseries in the 1990s. *Can J Fish Aquat Sci* 65:340–351. doi:[10.1139/f07-169](https://doi.org/10.1139/f07-169)
- McCleod E, Salm R, Green A, Almany J (2009) Designing marine protected area networks to address the impacts of climate change. *Front Ecol Environ*. doi:[10.1890/070211](https://doi.org/10.1890/070211)
- McCracken FD (1963) Seasonal movements of the winter flounder, *Pseudopleuronectes americanus* (Walbaum), on the Atlantic coast. *J Fish Res Board Can* 20:551–586
- Melnichuk MC, Essington TE, Branch TA, Heppell SS, Jensen OP, Link JS, Martell SJD, Parma AM, Pope JG, Smith ADM (2012) Can catch share fisheries better track management targets? *Fish Fish* 13:267–290. doi:[10.1111/j.1467-2979.2011.00429.x](https://doi.org/10.1111/j.1467-2979.2011.00429.x)
- Meng L, Powell JC, Taplin B (2001) Using winter flounder growth rates to assess habitat quality across an anthropogenic gradient in Narragansett Bay, Rhode Island. *Estuaries* 24:576–584. doi:[10.2307/1353258](https://doi.org/10.2307/1353258)
- Mieszkowska N, Genner MJ, Hawkins SJ, Sims DW (2009) Effects of climate change and commercial fishing on Atlantic cod *Gadus morhua*. *Adv Mar Biol* 56:213–273. doi:[10.1016/S0065-2881\(09\)56003-8](https://doi.org/10.1016/S0065-2881(09)56003-8)
- Miller TJ, Herra T, Leggett WC (1995) An individual based analysis of the variability of eggs and their newly-hatching larvae of Atlantic cod (*Gadus morhua*) on the Scotian Shelf. *Can J Fish Aquat Sci* 52:1083–1093. doi:[10.1139/f95-106](https://doi.org/10.1139/f95-106)
- Mills KE, Pershing AJ, Brown CJ, Chen Y, Chiang F-S, Holland DS, Lehuta S, Nye JA, Sun JC, Thomas AC, Wahle RA (2013) Fisheries management in a changing climate lessons from the 2012 ocean heat wave in the Northwest Atlantic. *Oceanography* 26:191–195
- Morgan MJ, Walsh SJ (1996) Tracking movements of juvenile yellowtail flounder in the nursery area on the Southern Grand Bank, NAFO Division 3LNO. NAFO SCR
- Murawski SA (1993) Climate-change and marine fish distributions—forecasting from historical analogy. *Trans Am Fish Soc* 122:647–658. doi:[10.1577/1548-8659\(1993\)122<0647:ccamfd>2.3.co;2](https://doi.org/10.1577/1548-8659(1993)122<0647:ccamfd>2.3.co;2)
- Myers RA, Cadigan NG (1993) Density-dependent juvenile mortality in marine demersal fish. *Can J Fish Aquat Sci* 50:1576–1590. doi:[10.1139/f93-179](https://doi.org/10.1139/f93-179)
- Myers RA, Drinkwater KF, Barrowman NJ, Baird JW (1993) Salinity and recruitment of Atlantic cod (*Gadus morhua*) in the Newfoundland region. *Can J Fish Aquat Sci* 50:1599–1609. doi:[10.1139/f93-181](https://doi.org/10.1139/f93-181)
- National Oceanographic and Atmospheric Administration [NOAA] (2015) Status of stocks 2014. Annual report to congress on the status of U.S. fisheries. http://www.fisheries.noaa.gov/sfa/publications/feature_stories/2015/status_of_stocks_2014.html. Accessed 24 Feb 2016
- Nichols JT (1918) An abnormal winter flounder and others. *Copeia* 55:37–39. doi:[10.2307/1436085](https://doi.org/10.2307/1436085)
- Nicolas D, Rochette S, Llope M, Licandro P (2014) Spatio-temporal variability of the North Sea cod recruitment in relation to temperature and zooplankton. *PLoS ONE*. doi:[10.1371/journal.pone.0088447](https://doi.org/10.1371/journal.pone.0088447)
- Northeast Fisheries Science Center [NEFSC] (2015) Stock assessment update of 20 Northeast groundfish stocks through 2014. <http://www.nefsc.noaa.gov/publications/crd/crd1524/>. Accessed 13 March 2015
- Nummelin A, Ilıcak M, Li C, Smedsrud LH (2016) Consequences of future increased Arctic runoff on Arctic Ocean stratification, circulation and sea ice cover. *J Geophys Res Oceans*. doi:[10.1002/2015JC011156](https://doi.org/10.1002/2015JC011156)
- Nye JA, Link JS, Hare JA, Overholtz WJ (2009) Changing spatial distribution of fish stocks in relation to climate and population size on the Northeast United States continental shelf. *Mar Ecol Prog Ser* 393:111–129
- Nye JA, Gamble RJ, Link JS (2013) The relative impact of warming and removing top predators on the Northeast US large marine biotic community. *Ecol Modell* 264:157–168. doi:[10.1016/j.ecolmodel.2012.08.019](https://doi.org/10.1016/j.ecolmodel.2012.08.019)
- O'Brien L (1999) Factors influencing the rate of sexual maturity and the effect on spawning stock for Georges Bank and Gulf of Maine Atlantic cod *Gadus morhua* stocks. *J Northwest Atl Fish Sci* 25:179–203
- O'Connor MI, Bruno JF, Gaines SD, Halpern BS, Lester SE, Kinlan BP, Weiss JM (2007) Temperature control of larval dispersal and the implications for marine ecology, evolution, and conservation. *Proc Natl Acad Sci USA* 104:1266–1271. doi:[10.1073/pnas.0603422104](https://doi.org/10.1073/pnas.0603422104)
- Odense P, Bordeleau A, Guilbault R (1966) Tolerance levels of cod (*Gadus morhua*) to low salinity. *J Fish Res Board Can* 23:1465–1467
- Olla BL, Wicklund R, Wilk S (1969) Behavior of winter flounder in a natural habitat. *Trans Am Fish Soc* 98:717–720. doi:[10.1577/1548-8659\(1969\)98<5B717%3ABOWFIA%5D2.0.CO%3B2](https://doi.org/10.1577/1548-8659(1969)98<5B717%3ABOWFIA%5D2.0.CO%3B2)
- Ottersen G, Loeng H, Raknes A (1994) Influence of temperature variability on recruitment of cod in the Barents Sea. In: Jakobsson J, Atthorsson OS, Beverton RJH et al (eds) Cod and climate change—proceedings of a symposium. ICES Mar Sci Symp 198:471–481
- Ottersen G, Hjermann DO, Stenseth NC (2006) Changes in spawning stock structure strengthen the link between climate and recruitment in a heavily fished cod (*Gadus morhua*) stock. *Fish Oceanogr* 15:230–243. doi:[10.1111/j.1365-2419.2006.00404.x](https://doi.org/10.1111/j.1365-2419.2006.00404.x)
- Page FH, Frank KT (1989) Spawning time and egg stage duration in Northwest Atlantic haddock (*Melanogrammus aeglefinus*) stocks with emphasis on Georges and Browns Bank. *Can J Fish Aquat Sci* 46:68–81
- Palmer JE, Clark SH (1979) Haddock: traditional New England foodfish. Marine resources of the Atlantic Coast, 2 edn. Atlantic State Marine Fisheries Commission, Washington, DC

- Pearcy WG (1962) Ecology of an estuarine population of winter flounder, *Pseudopleuronectes americanus* (Walbaum). Bull Bingham Oceanogr Collect 17:78
- Peck MA, Buckley LJ, Caldarone EM, Bengtson DA (2003) Effects of food consumption and temperature on growth rate and biochemical-based indicators of growth in early juvenile Atlantic cod *Gadus morhua* and haddock *Melanogrammus aeglefinus*. Mar Ecol Prog Ser 251:233–243. doi:[10.3354/meps251233](https://doi.org/10.3354/meps251233)
- Pereira JJ, Goldberg R, Ziskowski JJ, Berrien P, Morse WM, Johnson DL (1999) Essential fish habitat source document: winter flounder, *Pseudopleuronectes americanus*, life history and habitat characteristics. NOAA Technical Memorandum NMFS-NE-138
- Pereira JJ, Schultz ET, Auster PJ (2012) Geospatial analysis of habitat use in yellowtail flounder *Limanda ferruginea* on Georges Bank. Mar Ecol Prog Ser 468:279–290. doi:[10.3354/meps10035](https://doi.org/10.3354/meps10035)
- Perry AL, Low PJ, Ellis JR, Reynolds JD (2005) Climate change and distribution shifts in marine fishes. Science 308:1912–1915. doi:[10.1126/science.1111322](https://doi.org/10.1126/science.1111322)
- Pershing AJ, Alexander MA, Hernandez CM, Kerr LA, Le Bris A, Mills KE, Nye JA, Record NR, Scannell HA, Scott JD, Sherwood GD, Thomas AC (2015) Slow adaptation in the face of rapid warming leads to collapse of the Gulf of Maine cod fishery. Science 350:809–812. doi:[10.1126/science.aac9819](https://doi.org/10.1126/science.aac9819)
- Phelan BA (1992) Winter flounder movements in the inner New York Bight. Trans Am Fish Soc 121:777–784. doi:[10.1577/1548-8659\(1992\)121<0777:wfmitti>2.3.co;2](https://doi.org/10.1577/1548-8659(1992)121<0777:wfmitti>2.3.co;2)
- Pinsky ML, Mantua NJ (2014) Emerging adaptation approaches for climate-ready fisheries management. Oceanography 27:146–159. doi:[10.5670/oceanog.2014.93](https://doi.org/10.5670/oceanog.2014.93)
- Pinsky ML, Worm B, Fogarty MJ, Samiento JL, Levin SA (2013) Marine taxa track local climate velocities. Science 341:1239–1242. doi:[10.1126/science.1239352](https://doi.org/10.1126/science.1239352)
- Plaganyi EE, Punt AE, Hillary R, Morello EB, Thebaud O, Hutton T, Pillans RD, Thorson JT, Fulton EA, Smith ADM, Smith F, Bayliss P, Haywood M, Lyne V, Rothlisberg PC (2014) Multispecies fisheries management and conservation: tactical applications using models of intermediate complexity. Fish Fish 15:1–22. doi:[10.1111/j.1467-2979.2012.00488.x](https://doi.org/10.1111/j.1467-2979.2012.00488.x)
- Planque B, Fredou T (1999) Temperature and the recruitment of Atlantic cod (*Gadus morhua*). Can J Fish Aquat Sci 56:2069–2077. doi:[10.1139/cjfas-56-11-2069](https://doi.org/10.1139/cjfas-56-11-2069)
- Planque B, Fromentin JM, Cury P, Drinkwater KF, Jennings S, Perry RI, Kifani S (2010) How does fishing alter marine populations and ecosystems sensitivity to climate? J Mar Syst 79:403–417
- Pörtner HO, Peck MA (2010) Climate change effects on fishes and fisheries: towards a cause-and-effect understanding. J Fish Biol 77:1745–1779. doi:[10.1111/j.1095-8649.2010.02783.x](https://doi.org/10.1111/j.1095-8649.2010.02783.x)
- Pörtner HO, Hardewig I, Sartoris FJ, Van Dijk PLM (1998) Energetic aspects of cold adaptation: critical temperatures in metabolic, ionic and acid-base regulation? Soc Exp Biol Seminar Ser 66:88–120
- Pörtner HO, Bernal B, Blust R, Brix O, Colosimo A, De Wachter B, Giuliani A, Johansen T, Fischer T, Knust R, Lannig G, Naevdal G, Nedenes A, Nyhammer G, Sartoris FJ, Serendero I, Sirabella P, Thorkildsen S, Zakhartsev M (2001) Climate induced temperature effects on growth performance, fecundity and recruitment in marine fish: developing a hypothesis for cause and effect relationships in Atlantic cod (*Gadus morhua*) and common eelpout (*Zoarces viviparus*). Cont Shelf Res 21:1975–1997. doi:[10.1016/S0278-4343\(01\)00038-3](https://doi.org/10.1016/S0278-4343(01)00038-3)
- Pörtner HO, Bock C, Knust R, Lannig G, Lucassen M, Mark FC, Sartoris FJ (2008) Cod and climate in a latitudinal cline: physiological analyses of climate effects in marine fishes. Clim Res 37:253–270. doi:[10.3354/cr00766](https://doi.org/10.3354/cr00766)
- Provencher L, Munro J, Dutil JD (1993) Osmotic performance and survival of Atlantic cod (*Gadus morhua*) at low salinities. Aquaculture 116:219–231. doi:[10.1016/0044-8486\(93\)90010-v](https://doi.org/10.1016/0044-8486(93)90010-v)
- Purchase CF, Brown JA (2000) Interpopulation differences in growth rates and food conversion efficiencies of young Grand Banks and Gulf of Maine Atlantic cod (*Gadus morhua*). Can J Fish Aquat Sci 57:2223–2229. doi:[10.1139/cjfas-57-11-2223](https://doi.org/10.1139/cjfas-57-11-2223)
- Rabalais NN, Diaz RJ, Levin LA, Turner RE, Gilbert D, Zhang J (2010) Dynamics and distribution of natural and human-caused hypoxia. Biogeosciences 7:585–619
- Rajasilta M, Eklund J, Hanninen J, Kurkilahti M, Kaaria J, Rannikko P, Soikkeli M (1993) Spawning of herring (*Clupea harengus membras* L.) in the Archipelago Sea. ICES J Mar Sci 50:233–246. doi:[10.1006/jmsc.1993.1026](https://doi.org/10.1006/jmsc.1993.1026)
- Ratz HJ, Lloret J (2003) Variation in fish condition between Atlantic cod (*Gadus morhua*) stocks, the effect on their productivity and management implications. Fish Res 60:369–380. doi:[10.1016/S0165-7836\(02\)00132-7](https://doi.org/10.1016/S0165-7836(02)00132-7)
- Restrepo VR, Powers JE (1999) Precautionary control rules in US fisheries management: specification and performance. ICES J Mar Sci 56:846–852. doi:[10.1006/jmsc.1999.0546](https://doi.org/10.1006/jmsc.1999.0546)
- Rijnsdorp AD, Peck MA, Engelhard GH, Mollmann C, Pinnegar JK (2009) Resolving the effect of climate change on fish populations. ICES J Mar Sci 66:1570–1583. doi:[10.1093/icesjms/fsp056](https://doi.org/10.1093/icesjms/fsp056)
- Robichaud D, Rose GA (2004) Migratory behaviour and range in Atlantic cod: inference from a century of tagging. Fish Fish 5:185–214
- Rogers CA (1976) Effects of temperature and salinity on survival of winter flounder embryos. Fish Bull 74:52–58
- Rose GA, Atkinson BA, Baird J, Bishop CA, Kulka DW (1994) Changes in distribution of Atlantic cod and thermal variations in Newfoundland waters, 1980–1992. In: Jakobsen J, Atthorsson OS, Beverton RJH et al (eds) Cod and climate change—proceedings of a symposium. ICES Mar Sci Symp 198:542–552
- Rosenberg AA, Bolster WJ, Alexander KE, Leavenworth WB, Cooper AB, McKenzie MG (2005) The history of ocean resources: modeling cod biomass using historical records. Front Ecol Environ 3:84–90
- Rosenberg AA, Swasey JH, Bowman M (2006) Rebuilding US fisheries: progress and problems. Front Ecol Environ 4:303–308
- Runge JA, Kovach AI, Churchill JH, Kerr LA, Morrison JR, Beardsley RC, Berlinsky DL, Chen CS, Cadrin SX, Davis CS, Ford KH, Grabowski JH, Howell WH, Ji RB, Jones RJ, Pershing AJ, Record NR, Thomas AC, Sherwood GD, Tallack SML, Townsend DW (2010) Understanding

- climate impacts on recruitment and spatial dynamics of Atlantic cod in the Gulf of Maine: integration of observations and modeling. *Prog Oceanogr* 87:251–263. doi:[10.1016/j.pocean.2010.09.016](https://doi.org/10.1016/j.pocean.2010.09.016)
- Safina C, Rosenberg AA, Myers RA, Quinn TJ II, Collie JS (2005) U.S. ocean fish recovery: staying the course. *Science* 309:707–708. doi:[10.1126/science.1113725](https://doi.org/10.1126/science.1113725)
- Sale PF, Hanks I, Kritzer JP (2006) The merging of metapopulation theory and marine ecology: establishing the historical context. In: Kritzer JP, Sale PF (eds) *Marine metapopulations*. Academic Press, Boston, pp 3–22
- Sartoris FJ, Bock C, Serendero I, Lannig G, Pörtner HO (2003) Temperature-dependent changes in energy metabolism, intracellular pH and blood oxygen tension in the Atlantic cod. *J Fish Biol* 62:1239–1253. doi:[10.1046/j.1095-8649.2003.00099.x](https://doi.org/10.1046/j.1095-8649.2003.00099.x)
- Sayer MDJ, Reader JP, Dalziel TRK (1993) Fresh-water acidification—effects on the early life stages of fish. *Rev Fish Biol Fish* 3:298
- Sherwood GD, Grabowski JH (2010) Exploring the life-history implications of colour variation in offshore Gulf of Maine cod (*Gadus morhua*). *ICES J Mar Sci* 67:1640–1649
- Sherwood GD, Grabowski JH (2015) A comparison of cod life-history parameters inside and outside of four year-round groundfish closed areas in New England, USA. *ICES J Mar Sci*. doi:[10.1093/icesjms/fsv215](https://doi.org/10.1093/icesjms/fsv215)
- Simpson MR, Walsh SJ (2004) Changes in the spatial structure of Grand Bank yellowtail flounder: testing MacCall's basin hypothesis. *J Sea Res* 51:199–210. doi:[10.1016/j.seares.2003.08.007](https://doi.org/10.1016/j.seares.2003.08.007)
- Simpson SD, Jennings S, Johnson MP, Blanchard JL, Schon PJ, Sims DW, Genner MJ (2011) Continental shelf-wide response of a fish assemblage to rapid warming of the sea. *Curr Biol* 21:1565–1570. doi:[10.1016/j.cub.2011.08.016](https://doi.org/10.1016/j.cub.2011.08.016)
- Sissenwine MP (1974) Variability in recruitment and equilibrium catch of southern New England yellowtail flounder fishery. *ICES J Mar Sci* 36:15–26
- Smedbol KR, Wroblewski JS (2002) Metapopulation theory and northern cod population structure: interdependency of subpopulations in recovery of a groundfish population. *Fish Res* 55:161–174. doi:[10.1016/S0165-7836\(01\)00289-2](https://doi.org/10.1016/S0165-7836(01)00289-2)
- Smith WG, Morse WW (1985) Retention of larval haddock *Melanogrammus aeglefinus* in the Georges Bank region, a gyre-influenced spawning area. *Mar Ecol Prog Ser* 24:1–13. doi:[10.3354/meps024001](https://doi.org/10.3354/meps024001)
- Smith WG, Berrien P, McMillan DG, Wells A (1981) The distribution, abundance and production of Atlantic cod and haddock larvae off northeastern United States in 1978–1979 and 1979–1980. *ICES Demersal Fish Committee C.M.* 1981/G:52
- Steinacher M, Joos F, Froelicher TL, Plattner GK, Doney SC (2009) Imminent ocean acidification in the Arctic projected with the NCAR global coupled carbon cycle-climate model. *Biogeosciences* 6:515–533
- Steinacher M, Joos F, Froelicher TL, Bopp L, Cadule P, Cocco V, Doney SC, Gehlen M, Lindsay K, Moore JK, Schneider B, Segschneider J (2010) Projected 21st century decrease in marine productivity: a multi-model analysis. *Biogeosciences* 7:979–1005
- Steneck RS, Wilson JA (2010) A fisheries play in an ecosystem theater: challenges of managing ecological and social drivers of marine fisheries at multiple spatial scales. *Bull Mar Sci* 86:387–411
- Steneck RS, Vavrinc J, Leland AV (2004) Accelerating trophic-level dysfunction in kelp forest ecosystems of the western North Atlantic. *Ecosystems* 7:323–332
- Steves BP, Cowen RK, Malchoff MH (2000) Settlement and nursery habitats for demersal fishes on the continental shelf of the New York Bight. *Fish Bull* 98:167–188
- Stierhoff KL, Targett TE, Miller K (2006) Ecophysiological responses of juvenile summer and winter flounder to hypoxia: experimental and modeling analyses of effects on estuarine nursery quality. *Mar Ecol Prog Ser* 325:255–266. doi:[10.3354/meps325255](https://doi.org/10.3354/meps325255)
- Stoner AW, Manderson JP, Pessutti JP (2001) Spatially explicit analysis of estuarine habitat for juvenile winter flounder: combining generalized additive models and geographic information systems. *Mar Ecol Prog Ser* 213:253–271
- Sullivan MC, Cowen RK, Steves BP (2005) Evidence for atmosphere-ocean forcing of yellowtail flounder (*Limanda ferruginea*) recruitment in the Middle Atlantic Bight. *Fish Oceanogr* 14:386–399. doi:[10.1111/j.1365-2419.2005.00343.x](https://doi.org/10.1111/j.1365-2419.2005.00343.x)
- Sumaila UR, Cheung WWL, Lam VWY, Pauly D, Herrick S (2011) Climate change impacts on the biophysics and economics of world fisheries. *Nat Clim Change* 1:449–456. doi:[10.1038/nclimate1301](https://doi.org/10.1038/nclimate1301)
- Sutcliffe WH, Drinkwater K, Muir BS (1977) Correlations of fish catch and environmental factors in Gulf of Maine. *J Fish Res Board Can* 34:19–30
- Sutcliffe WH, Loucks RH, Drinkwater KF, Coote AR (1983) Nutrient flux onto the Labrador Shelf from Hudson Strait and its biological consequences. *Can J Fish Aquat Sci* 40:1692–1701
- Taylor DL, Collie JS (2003) Effect of temperature on the functional response and foraging behavior of the sand shrimp *Crangon septemspinosa* preying on juvenile winter flounder *Pseudopleuronectes americanus*. *Mar Ecol Prog Ser* 263:217–234. doi:[10.3354/meps263217](https://doi.org/10.3354/meps263217)
- Taylor DL, Danila DJ (2005) Predation on winter flounder (*Pseudopleuronectes americanus*) eggs by the sand shrimp (*Crangon septemspinosa*). *Can J Fish Aquat Sci* 62:1611–1625. doi:[10.1139/f05-047](https://doi.org/10.1139/f05-047)
- Topp RW (1968) An estimate of fecundity of the winter flounder, *Pseudopleuronectes americanus*. *J Fish Res Board Can* 25:1299–1302
- Vert-pre KA, Amoroso RO, Jensen OP, Hilborn R (2013) Frequency and intensity of productivity regime shifts in marine fish stocks. *Proc Natl Acad Sci USA* 110:1779–1784. doi:[10.1073/pnas.1214879110](https://doi.org/10.1073/pnas.1214879110)
- Walsh SJ, Morgan MJ (2004) Observations of natural behaviour of yellowtail flounder derived from data storage tags. *ICES J Mar Sci* 61:1151–1156. doi:[10.1016/j.icesjms.2004.07.005](https://doi.org/10.1016/j.icesjms.2004.07.005)
- Walsh SJ, Simpson M, Morgan MJ (2004) Continental shelf nurseries and recruitment variability in American plaice and yellowtail flounder on the Grand Bank: insights into stock resiliency. *J Sea Res* 51:271–286. doi:[10.1016/j.seares.2003.10.003](https://doi.org/10.1016/j.seares.2003.10.003)
- Walsh J, Wuebbles D, Hayhoe K, Kossin J, Kunkel K, Stephens G, Thorne P, Vose R, Wehner M, Willis J, Anderson D,

- Doney S, Feely R, Hennon P, Kharin V, Knutson T, Landerer F, Lenton T, Kennedy J, Somerville R (2014) Our changing climate. In: Melillo JM, Richmond TC, Yohe GW (eds) Climate change impacts in the United States: the third national climate assessment. U.S. Global Change Research Program, Washington, DC, pp 19–67. doi:[10.7930/J0KW5CXT](https://doi.org/10.7930/J0KW5CXT)
- Westin L, Nissling A (1991) Effects of salinity on spermatozoa motility, percentage of fertilized eggs and egg development of Baltic cod (*Gadus morhua*), and implications for cod stock fluctuations in the Baltic. Mar Biol 108:5–9. doi:[10.1007/bf01313465](https://doi.org/10.1007/bf01313465)
- Wilber DH, Clarke DG, Gallo J, Alcoba CJ, Dileo AM, Zappala SE (2013) Identification of winter flounder (*Pseudopleuronectes americanus*) estuarine spawning habitat and factors influencing egg and larval distributions. Estuaries Coasts 36:1304–1318. doi:[10.1007/s12237-013-9642-z](https://doi.org/10.1007/s12237-013-9642-z)
- Williams GC (1975) Viable embryogenesis of winter flounder *Pseudopleuronectes americanus* from -1.8°C to 15°C . Mar Biol 33:71–74. doi:[10.1007/bf00395003](https://doi.org/10.1007/bf00395003)
- Wirgin I, Kovach AI, Maceda L, Roy NK, Waldman J, Berlin-sky DL (2007) Stock identification of Atlantic cod in US waters using microsatellite and single nucleotide polymorphism DNA analyses. Trans Am Fish Soc 136:375–391. doi:[10.1577/t06-068.1](https://doi.org/10.1577/t06-068.1)
- Witman JD, Etter RJ, Smith F (2004) The relationship between regional and local species diversity in marine benthic communities: a global perspective. Proc Nat Acad Sci 101:15664–15669
- Worm B, Hilborn R, Baum JK, Branch TA, Collie JS, Costello C, Fogarty MJ, Fulton EA, Hutchings JA, Jennings S, Jensen OP, Lotze HK, Mace PM, McClanahan TR, Minto C, Palumbi SR, Parma AM, Ricard D, Rosenberg AA, Watson R, Zeller D (2009) Rebuilding global fisheries. Science 325:578–585