

Mathematics/Statistics Bootcamp

Part I: Calculus

Fan Bu¹ Kyle Burris¹

¹Department of Statistical Science
Duke University

Graduate Orientation, August 2018

Overview

Limit and Continuity

Derivative

Definition and Differentiation Rules

Application of Derivatives

Integrals

Definite Integrals

Improper Integrals

Sequences and Series

Multivariate Calculus

Limit and Continuity

Limit

Suppose $-\infty < a, L < +\infty$ and $f(x) : X \rightarrow Y$ is a real-valued function, then

$$\lim_{x \rightarrow a} f(x) = L$$

if for any $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - L| < \epsilon \text{ whenever } |x - a| < \delta.$$

Left-hand limit: $\lim_{x \rightarrow a^-} f(x) = L$ if for any $\epsilon > 0$, there exists $\delta > 0$ such that $|f(x) - L| < \epsilon$ whenever $a - \delta < x < a$.

Right-hand limit: $\lim_{x \rightarrow a^+} f(x) = L$ if for any $\epsilon > 0$, there exists $\delta > 0$ such that $|f(x) - L| < \epsilon$ whenever $a < x < a + \delta$.

Limit: An Example




Figure: Plot of $y = f(x)$.

- ▶ What is $\lim_{x \rightarrow 5^-} f(x)$?
- ▶ What is $\lim_{x \rightarrow 5^+} f(x)$?
- ▶ What is $\lim_{x \rightarrow 5} f(x)$?

Infinite Limit/Limit at Infinity

► How to define $\lim_{x \rightarrow a} f(x) = \infty$ for $-\infty < a < +\infty$?

► How to define $\lim_{x \rightarrow \infty} f(x) = a$ for $-\infty < a < +\infty$?

Continuity

A function f is continuous at a number a if

$$\lim_{x \rightarrow a} f(x) = f(a).$$


It implies 3 things:

1. $f(a)$ is defined ($a \in X$);
2. $\lim_{x \rightarrow a} f(x)$ exists;
3. $\lim_{x \rightarrow a} f(x) = f(a)$.


Right continuous: $\lim_{x \rightarrow a^-} f(x) = f(a)$.

Left continuous: $\lim_{x \rightarrow a^+} f(x) = f(a)$.

Continuity: Examples



This function is discontinuous
at $x = 5$.



This function is discontinuous
(but right continuous) at any
integer x .

Other Useful Continuity Notions

- ▶ **Uniform continuity:**

For all $\epsilon > 0$ there exists a $\delta > 0$ such that for any $x, y \in X$ with $|x - y| < \delta$ we have $|f(x) - f(y)| < \epsilon$.

(A global property rather than a local one: as long as x, y are sufficiently close, we can guarantee that $f(x), f(y)$ are close.)

- ▶ **Lipschitz continuity:**

There exists a constant L such that for any $x, y \in X$, we have $|f(x) - f(y)| \leq L|x - y|$.

Continuity: Exercise

Which statement(s) of the following is incorrect?

- A $f(x) = x^2$, $x \in \mathbb{R}$ is not uniformly continuous, but $f(x) = x^2$, $x \in [0, 245]$ is Lipschitz continuous;
- B $f(x) = x^{1/3}$, $x \in [0, 1]$ is Lipschitz continuous;
- C $f(x) = x^{3/2} \sin(\frac{1}{x})$ ($x \neq 0$), $f(0) = 0$, $x \in [0, 1]$ is uniformly continuous as well as Lipschitz continuous;
- D Every continuous function defined on $[-1, 1]$ is uniformly continuous;
- E A continuous function f is defined on $[-5, 5]$ and satisfies $f(-2) = -3$, $f(1) = 4$, then there must exist a number $x_0 \in [-2, 1]$ such that $f(x_0) = 0$.


Derivative

Definition of Derivative

The derivative of function f at $a \in X$, denoted by $f'(a)$ is


$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

if this limit exists (“differentiable”).



$$(a) \quad f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

= slope of tangent at P
= slope of curve at P



$$(b) \quad f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

= slope of tangent at P
= slope of curve at P

Figure: Geometric interpretations of the derivative.

Differentiation Rules

Derivatives of some common functions:


- ▶ $f(x) = \text{const}$, then $f'(x) = 0$;
- ▶ $f(x) = x^\alpha$, $\alpha \neq 0$, then $f'(x) = \alpha x^{\alpha-1}$;
- ▶ $(e^x)' = e^x$, $(\ln x)' = 1/x$ ($x > 0$);
- ▶ $(\sin x)' = \cos x$, $(\cos x)' = -\sin x$, $(\tan x)' = 1/\cos^2 x$;
- ▶ $(\sin^{-1} x)' = 1/\sqrt{1-x^2}$, $(\cos^{-1} x)' = -1/\sqrt{1-x^2}$,
 $(\tan^{-1} x)' = 1/1+x^2$.

If both $f(x)$ and $g(x)$ are differentiable:

- ▶ $(cf(x))' = cf'(x)$, $(f(x) + g(x))' = f'(x) + g'(x)$;
- ▶ $(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$;
- ▶ $\left(\frac{f(x)}{g(x)}\right)' = \frac{f'(x)g(x) - f(x)g'(x)}{g^2(x)}$ (assume $g(x) > 0$);
- ▶ The **chain rule**: if $F = f \circ g$, then $F'(x) = f'(g(x))g'(x)$.

Derivative: Exercises

1. The following figure shows the graphs of f , f' , f'' , and f''' . Identify each curve.



2. $f(x) = \frac{1}{\sqrt{\gamma}} \exp\left(-\frac{(x-\mu)^2}{\gamma}\right)$ where constants $\gamma > 0$ and $\mu \in \mathbb{R}$, and $x \in \mathbb{R}$. Calculate $f'(x)$ and find $x_0 \in \mathbb{R}$ such that the tangent line of $f(x)$ at x_0 is horizontal.

3. Find $\lim_{x \rightarrow 0} (1+x)^{1/x}$.

Solution to Exercise 3

Let $f(x) = \ln x$, then

$$\begin{aligned} f'(1) &= \lim_{x \rightarrow 0} \frac{\ln(1+x) - \ln 1}{x} \\ &= \lim_{x \rightarrow 0} \frac{1}{x} \ln(1+x) \\ &= \lim_{x \rightarrow 0} \ln(1+x)^{1/x}. \end{aligned}$$

Since $f'(1) = 1$, $\lim_{x \rightarrow 0} (1+x)^{1/x} = e^1 = e$.

Minimum and Maximum

Theorem (Fermat's Theorem)

If f has a local minimum or maximum at c and $f'(c)$ exists, then $f'(c) = 0$.

Note: the converse is not true.

Theorem (The Second Derivative Test)

If f has second derivative on $(c - \epsilon_0, c + \epsilon_0)$ for a certain $\epsilon_0 > 0$, then


- ▶ *if $f'(c) = 0$ and $f''(c) > 0$, f has a local minimum at c ;*
- ▶ *if $f'(c) = 0$ and $f''(c) < 0$, f has a local maximum at c .*

Minimum and Maximum: Example

Let v_1 be the velocity of light in air and v_2 the velocity of light in water. According to Fermat's Principle, a ray of light will travel from a point A in the air to a point B in the water by a path ACB that **minimizes** the time taken. Then

$$\frac{\sin \theta_1}{\sin \theta_2} = \frac{v_1}{v_2},$$

where θ_1 is the angle of incidence and θ_2 is the angle of refraction. This equation is known as **Snell's Law**.




Convexity

A function defined on a convex set X , $f : X \rightarrow \mathbb{R}$ is convex if for any $x, y \in X$ and $t \in [0, 1]$,

$$f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y).$$

Visually, a convex function has a “curve up” shape:



Convexity and Derivatives

Suppose $f(x)$ is twice differentiable on interval I , then

- ▶ f is convex on I if and only if $f'(x)$ is monotonically non-decreasing on I ;
- ▶ f is convex on I if and only if $f''(x) \geq 0$ for $x \in I$ (often used to test for convexity).

A nice property of convexity:

Any local minimum of a convex function is also a global minimum;
a strictly convex function has at most one global minimum.
(Therefore convexity is much desired in optimization.)

Review Exercises: Morning Session

1. Are all Lipschitz continuous functions defined on $[-1, 1]$ also differentiable on $(-1, 1)$? If not, give a counterexample.

2. Calculate $\lim_{x \rightarrow 0} \frac{\sin x}{x}$ and $\lim_{x \rightarrow 0} \frac{\tan x}{x}$.

3. Let $f(x) = \frac{1}{x}, x > 0$. For every positive integer n , find $f^{(n)}(x)$.

4. Which following functions are convex?

- ▶ $f_1(x) = |x|, x \in [-1, 1];$
- ▶ $f_2(x) = \ln(x^2 + 1), x \in \mathbb{R};$
- ▶ $f_3(x) = e^{-x}, x \in \mathbb{R}.$

5. $f(x) = \frac{1}{\sqrt{\gamma}} \exp\left(-\frac{(x-\mu)^2}{\gamma^2}\right)$ where constants $\gamma > 0$ and $\mu \in \mathbb{R}$, and $x \in \mathbb{R}$. Find all the global maximums of $f(x)$.

Taylor Expansion

Talor Series, by *3Blue1Brown*

Challenge Exercises: Morning Session

1. Find the Taylor expansion of $f(x) = \tan x$ around 0 and use the result to show that $\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$.

2. Calculate $\sum_{x=0}^{\infty} \frac{e^{-\lambda} \lambda^x}{x!}$. Here $\lambda > 0$ is a constant and $x \in \mathbb{N}$. ($0! = 1$.)

3. Let $f(x) = x^{\alpha-1}(1-x)^{\beta-1}$, $x \in (0, 1)$, where $\alpha, \beta > 0$ are constants. Find all the minimums and maximums (if there are any) of f .

4. $f(x)$ is a convex function on \mathbb{R} , $P = (x_0, f(x_0))$ is a point on $f(x)$, and l is the tangent line of the curve of $f(x)$ at P . Choose the correct statement(s).

- A l must lie above or on the curve of $f(x)$;
- B l must lie below or on the curve of $f(x)$;
- C l can intersect with $f(x)$ at multiple points.

Integrals

Properties of Definite Integrals

Let $a \leq d \leq b \in \mathbb{R}$:

- ▶ If $c \in \mathbb{R}$ is a constant, then $\int_a^b c dx = c(b - a)$;
- ▶ $\int_a^b cf(x) dx = c \int_a^b f(x) dx$;
- ▶ $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$;
- ▶ $\int_a^d f(x) dx + \int_d^b f(x) dx = \int_a^b f(x) dx$;
- ▶ If $f(x) \geq g(x)$ for $a \leq x \leq b$, then $\int_a^b f(x) dx \geq \int_a^b g(x) dx$;
- ▶ If $m \leq f(x) \leq M$ for $a \leq x \leq b$, then $m(b - a) \leq \int_a^b f(x) dx \leq M(b - a)$.

The Fundamental Theorem of Calculus

If f is continuous on $[a, b]$, then:

- ▶ function $g(x) = \int_a^x f(x)dx$, $a \leq x \leq b$ is continuous on $[a, b]$ and differentiable on (a, b) , and $g'(x) = f(x)$;
- ▶ $\int_a^b f(x)dx = F(b) - F(a)$, where F is any anti-derivative of f ($F' = f$).

A mini-exercise: find $\frac{d}{dx} \int_1^x \sin x^4 dx$.

Useful Rules for Integration

- ▶ **Substitution rule:** If $u = g(x)$ is continuously differentiable on $[a, b]$ and f is continuous on the range of u , then

$$\int_a^b f(g(x))g'(x)dx = \int_{g(a)}^{g(b)} f(u)du.$$

- ▶ **Integration by parts:** If functions u and v are both continuously differentiable on $[a, b]$, then

$$\int_a^b u(x)v'(x)dx = [u(x)v(x)]|_a^b - \int_a^b v(x)u'(x)dx.$$

Integration: Exercises

1. Calculate $\int_1^e \frac{\ln x}{x} dx$.

2. Calculate $\int_0^1 x \cos x dx$.

Improper Integrals

1. **Infinite intervals:** if $\int_a^t f(x)dx$ exists for every $t \geq a$ then

$$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx$$

provided that this limit exists (convergent); similarly, one may define $\int_{-\infty}^a f(x)dx$, and if both $\int_a^\infty f(x)dx$ and $\int_{-\infty}^a f(x)dx$ are convergent, then $\int_{-\infty}^\infty f(x)dx = \int_{-\infty}^a f(x)dx + \int_a^\infty f(x)dx$.

2. **Discontinuous integrand:** if f is continuous on $[a, b)$ and is discontinuous at b , then

$$\int_a^b f(x)dx = \lim_{t \rightarrow b^-} \int_a^t f(x)dx$$

if this limit exists; similarly, if f is continuous on $(a, b]$ and is discontinuous at a , $\int_a^b f(x)dx = \lim_{t \rightarrow a^+} \int_t^b f(x)dx$.

Improper Integrals: Exercises

1. For what values of $p \in \mathbb{R}$ is the integral

$$\int_1^{\infty} \frac{1}{x^p} dx$$

convergent?

2. Evaluate $\int_2^5 \frac{1}{\sqrt{x-2}} dx$.

Sequences and Series

Basics of Sequences

A **sequence** is a list of numbers written in a definite order. We often denote a sequence $\{a_1, a_2, a_3, \dots\}$ by $\{a_n\}$ or $\{a_n\}_{n=1}^{\infty}$.

A sequence $\{a_n\}$ has **limit** L (written as $\lim_{n \rightarrow \infty} a_n = L$, or $a_n \rightarrow L$ as $n \rightarrow \infty$) if for every $\epsilon > 0$ there is a corresponding integer N such that $|a_n - L| < \epsilon$ whenever $n > N$.

If for every $n \in \mathbb{N}$, $a_n \leq a_{n+1}$ (increasing) or $a_n \geq a_{n+1}$ (decreasing), then the sequence $\{a_n\}$ is **monotonic**. If there exists a number $M > 0$ such that $|a_n| \leq M$ for every n then the sequence $\{a_n\}$ is **bounded**.

Monotonic Sequence Theorem: Every bounded, monotonic sequence is convergent (has a limit).

Sequences: Exercises

1. Find the limit of the sequence a_n where $a_n = \left(1 + \frac{1}{n}\right)^n$, $n \in \mathbb{N}$.
2. Let $a_n = \frac{2^n}{n!}$, $n \in \mathbb{N}$. Is the sequence a_n convergent? If so, what is its limit?

Basics of Series

A **series** can be thought of as the infinite sum of a sequence a_n , written as $\sum_{n=1}^{\infty} a_n$ or $\sum a_n$. More formally, it can be defined by taking the limit of partial sums $\{s_n\}$, where $s_n = \sum_{i=1}^n a_i$: if $\lim_{n \rightarrow \infty} s_n$ exists then the series $\sum a_n$ is convergent, otherwise it is divergent.

If $\sum_{n=1}^{\infty} a_n$ is convergent, then $\lim_{n \rightarrow \infty} a_n = 0$.

An important example - **the geometric series**:

$$\sum_{n=1}^{\infty} ar^{n-1} = a + ar + ar^2 + \dots$$

is convergent if $|r| < 1$ and its sum is $\sum_{n=1}^{\infty} ar^{n-1} = \frac{a}{1-r}$. If $|r| \geq 1$, the geometric series is divergent.

Series: Exercises

1. Show that $\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$ is convergent and find its sum.

2. Find $\sum_{n=0}^{\infty} x^n$ where $|x| < 1$.

Convergence of Series

Commonly used tests for convergence:

1. **The comparison test:** Suppose that $\sum a_n$ and $\sum b_n$ are series with positive terms.
 - (i) If $\sum b_n$ is convergent and $a_n \leq b_n$ for all n , then $\sum a_n$ is also convergent;
 - (ii) If $\sum b_n$ is divergent and $a_n \geq b_n$ for all n , then $\sum a_n$ is also divergent.
 - ▶ Video example: **famous proof that the harmonic series diverges**, by *Khan Academy*.
2. **The limit comparison test:** Suppose that $\sum a_n$ and $\sum b_n$ are series with positive terms. If $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$ where c is a finite positive number, then either both series converge or both diverge.
3. **The integral test** (by *Khan Academy*).
4. **The alternating series test:** If the alternating series $\sum_{n=1}^{\infty} (-1)^{n-1} b_n$ ($b_n > 0$) satisfies (i) $b_{n+1} \leq b_n$ for all n , and (ii) $\lim_{n \rightarrow \infty} b_n = 0$ then the series is convergent.

Review Exercises: Afternoon Session

1. Evaluate the following definite integrals:

- ▶ $\int_0^4 \frac{x}{\sqrt{x^2+9}} dx;$
- ▶ $\int_0^1 \frac{2x}{1+x^4} dx.$

2. Evaluate the following definite integrals:

- ▶ $\int_0^1 x \ln x dx;$
- ▶ $\int_0^\pi e^x \sin x dx.$

3. If f is continuous on \mathbb{R} , show that

$$\int_a^b f(x+c) dx = \int_{a+c}^{b+c} f(x) dx.$$

4. Do the following series converge? Calculate the value of the infinite sum for each convergent series.

- (a) $\sum_{n=2}^{\infty} 5^{n-1} \left(\frac{9}{10}\right)^n;$
- (b) $\sum_{n=1}^{\infty} \left(\frac{3}{2}\right)^n \frac{1}{9^{n+2}}.$

5. True or false?

- ▶ If $x_n \rightarrow 0$ as $n \rightarrow \infty$, then $\sum_{n=1}^{\infty} x_n$ is convergent;
- ▶ $\sum_{n=1}^{\infty} x^n e^{-nx}$ is convergent for any $x > 0$;
- ▶ $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$ is convergent.

Multivariate Calculus

Partial Derivatives

If u is a function of n variables, $u = f(x_1, x_2, \dots, x_n)$, its partial derivative with respect to the i th variable x_i is

$$\frac{\partial u}{\partial x_i} = \lim_{h \rightarrow 0} \frac{f(x_1, \dots, x_{i-1}, x_i + h, x_{i+1}, \dots, x_n) - f(x_1, \dots, x_i, \dots, x_n)}{h}.$$

(Strategy: treat all the other variables as constants and take the derivative with respect to the variable of interest.)

Suppose $u = f(x_1, x_2, \dots, x_n)$ is defined on \mathbb{R}^n . If $\frac{\partial^2 u}{\partial x_i \partial x_j}$ and $\frac{\partial^2 u}{\partial x_j \partial x_i}$ are both continuous on \mathbb{R}^n , then $\frac{\partial^2 u}{\partial x_i \partial x_j} = \frac{\partial^2 u}{\partial x_j \partial x_i}$.

The Gradient Vector and Hessian Matrix

Suppose $f(x_1, x_2, \dots, x_n)$ is a function of n variables such that all the partial derivatives exist, then the gradient vector of f is

$$\nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right).$$

If all the second-order partial derivatives of f also exist, the Hessian matrix of f is

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{bmatrix}.$$

Change of Variables

Take the two-variable case as an example:

Suppose $z = f(x, y)$ is function of x, y and $x = u(s, t)$, $y = v(s, t)$ with respect two other variables s, t , then $z = g(s, t)$ as a function of s, t , where

$$g(s, t) = f(u(s, t), v(s, t))|J|.$$

Here J is the **Jacobian** of the transformation $x = u(s, t)$, $y = v(s, t)$:

$$J = \begin{vmatrix} \frac{\partial x}{\partial s} & \frac{\partial x}{\partial t} \\ \frac{\partial y}{\partial s} & \frac{\partial y}{\partial t} \end{vmatrix} = \frac{\partial x}{\partial s} \frac{\partial y}{\partial t} - \frac{\partial x}{\partial t} \frac{\partial y}{\partial s}.$$

Suppose that we want to integrate $f(x, y)$ over a region R . Under the transformation $x = u(s, t)$, $y = v(s, t)$ the regions becomes S and the integral becomes:

$$\iint_R f(x, y) dx dy = \iint_S f(u(s, t), v(s, t)) |J| ds dt.$$

Challenge Exercises: Afternoon Session

1. Given that $\int_1^\infty \frac{1}{x} dx = \infty$ and $\sum_{n=1}^\infty \frac{1}{n} = \infty$, use two methods to show that $\sum_{n=1}^\infty \frac{1}{n^p} = \infty$ for any $p \in (0, 1)$.

2. Find $\lim_{x \rightarrow 3} \frac{x}{x-3} \int_3^x \frac{\sin t}{t} dt$.

3. The Gamma function $\Gamma(x)$ is defined for any real number $x > 0$ as
 $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$. Show that $\Gamma(x+1) = x\Gamma(x)$.

4. Evaluate $\int_0^\infty e^{x^2} dx$.
(Hint: start with $\int_0^\infty \int_0^\infty e^{x^2+y^2} dx dy$.)

5. $f(x, y) = \frac{1}{\sqrt{2\pi y^2}} e^{-\frac{(x-s)^2}{2y^2}}$,
where $s \in \mathbb{R}$ is a constant.
Obtain the Hessian matrix of f .

The End