

# Mathematics/Statistics Bootcamp

## Part I: Calculus

Fan Bu<sup>1</sup>   Kyle Burris<sup>1</sup>

<sup>1</sup>Department of Statistical Science  
Duke University

Graduate Orientation, August 2018

# Overview

Limit and Continuity

Derivative

Definition and Differentiation Rules

Application of Derivatives

Integrals

Definite Integrals

Improper Integrals

Series

Multivariate Calculus

# Limit and Continuity

# Limit

Suppose  $-\infty < a, L < +\infty$  and  $f(x) : X \rightarrow Y$  is a real-valued function, then

$$\lim_{x \rightarrow a} f(x) = L$$

if for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that

$$|f(x) - L| < \epsilon \text{ whenever } 0 < |x - a| < \delta.$$

(The value of  $f(x)$  approaches  $L$  when  $x$  approaches  $a$ .)

**Left-hand limit:**  $\lim_{x \rightarrow a^-} f(x) = L$  if for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $|f(x) - L| < \epsilon$  whenever  $a - \delta < x < a$ .

**Right-hand limit:**  $\lim_{x \rightarrow a^+} f(x) = L$  if for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $|f(x) - L| < \epsilon$  whenever  $a < x < a + \delta$ .

## Limit: An Example




Figure: Plot of  $y = f(x)$ .

- ▶ What is  $\lim_{x \rightarrow 5^-} f(x)$  ?
- ▶ What is  $\lim_{x \rightarrow 5^+} f(x)$  ?
- ▶ What is  $\lim_{x \rightarrow 5} f(x)$  ?

# Infinite Limit/Limit at Infinity

► How to define  $\lim_{x \rightarrow a} f(x) = \infty$  for  $-\infty < a < +\infty$  ?

► How to define  $\lim_{x \rightarrow \infty} f(x) = a$  for  $-\infty < a < +\infty$  ?

# Infinite Limit/Limit at Infinity

- ▶ How to define  $\lim_{x \rightarrow a} f(x) = \infty$  for  $-\infty < a < +\infty$  ?

For any  $M > 0$ , there exists  $\delta > 0$  such that

$$f(x) > M \text{ whenever } 0 < |x - a| < \delta.$$

(The value of  $f(x)$  approaches  $\infty$  when  $x$  approaches  $a$ .)

- ▶ How to define  $\lim_{x \rightarrow \infty} f(x) = a$  for  $-\infty < a < +\infty$  ?

For any  $\epsilon > 0$ , there exists a  $M > 0$  such that

$$|f(x) - a| < \epsilon \text{ whenever } x > M.$$

(The value of  $f(x)$  approaches  $a$  when  $x$  approaches  $\infty$ .)

# Continuity

A function  $f$  is continuous at a number  $a$  if

$$\lim_{x \rightarrow a} f(x) = f(a).$$

It implies 3 things:


1.  $f(a)$  is defined ( $a \in X$ );
2.  $\lim_{x \rightarrow a} f(x)$  exists;
3.  $\lim_{x \rightarrow a} f(x) = f(a)$ .

**Right continuous:**  $\lim_{x \rightarrow a^-} f(x) = f(a)$ .


**Left continuous:**  $\lim_{x \rightarrow a^+} f(x) = f(a)$ .



# Continuity: Examples



This function is discontinuous  
at  $x = 5$ .



This function is discontinuous  
(but right continuous) at any  
integer  $x$ .


# Derivative

# Definition of Derivative


The derivative of function  $f$  at  $a \in X$ , denoted by  $f'(a)$  is

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

if this limit exists (“differentiable”).



$$\begin{aligned} \text{(a) } f'(a) &= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \\ &= \text{slope of tangent at } P \\ &= \text{slope of curve at } P \end{aligned}$$



$$\begin{aligned} \text{(b) } f'(a) &= \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \\ &= \text{slope of tangent at } P \\ &= \text{slope of curve at } P \end{aligned}$$

Figure: Geometric interpretations of the derivative.

# Differentiation Rules

Derivatives of some common functions:

- ▶  $f(x) = \text{const}$ , then  $f'(x) = 0$ ;
- ▶  $f(x) = x^\alpha$ ,  $\alpha \neq 0$ , then  $f'(x) = \alpha x^{\alpha-1}$ ;
- ▶  $(e^x)' = e^x$ ,  $(\ln x)' = 1/x$  ( $x > 0$ );
- ▶  $(\sin x)' = \cos x$ ,  $(\cos x)' = -\sin x$ ,  $(\tan x)' = 1/\cos^2 x$ ;
- ▶  $(\sin^{-1} x)' = 1/\sqrt{1-x^2}$ ,  $(\cos^{-1} x)' = -1/\sqrt{1-x^2}$ ,  
 $(\tan^{-1} x)' = 1/1+x^2$ .

If both  $f(x)$  and  $g(x)$  are differentiable:

- ▶  $(cf(x))' = cf'(x)$ ,  $(f(x) + g(x))' = f'(x) + g'(x)$ ;
- ▶  $(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$ ;
- ▶  $\left(\frac{f(x)}{g(x)}\right)' = \frac{f'(x)g(x) - f(x)g'(x)}{g^2(x)}$  (assume  $g(x) > 0$ );
- ▶ The **chain rule**: if  $F = f \circ g$ , then  $F'(x) = f'(g(x))g'(x)$ .

# Derivative: Exercises

1. Find the derivatives of the following functions

►  $f(x) = xe^x;$

►  $f(x) = 1 - \cos^2 x;$

►  $f(x) = \frac{\ln x}{x}.$

2. Find  $\lim_{x \rightarrow 0}(1+x)^{1/x}.$

## Solution to Exercise 2

Let  $f(x) = \ln x$ , then

$$\begin{aligned} f'(1) &= \lim_{x \rightarrow 0} \frac{\ln(1+x) - \ln 1}{x} \\ &= \lim_{x \rightarrow 0} \frac{1}{x} \ln(1+x) \\ &= \lim_{x \rightarrow 0} \ln(1+x)^{1/x}. \end{aligned}$$

Since  $f'(1) = 1$ ,  $\lim_{x \rightarrow 0} (1+x)^{1/x} = e^1 = e$ .

# Minimum and Maximum

## Theorem (Fermat's Theorem)

*If  $f$  has a local minimum or maximum at  $c$  and  $f'(c)$  exists, then  $f'(c) = 0$ .*

**Note:** the converse is not true.

## Theorem (The Second Derivative Test)

*If  $f$  has second derivative on  $(c - \epsilon_0, c + \epsilon_0)$  for a certain  $\epsilon_0 > 0$ , then*


- ▶ *if  $f'(c) = 0$  and  $f''(c) > 0$ ,  $f$  has a local minimum at  $c$ ;*
- ▶ *if  $f'(c) = 0$  and  $f''(c) < 0$ ,  $f$  has a local maximum at  $c$ .*

# Convexity

A function defined on a convex set  $X$ ,  $f : X \rightarrow \mathbb{R}$  is convex if for any  $x, y \in X$  and  $t \in [0, 1]$ ,

$$f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y).$$

Visually, a convex function has a “curve up” shape:





# Convexity and Derivatives

Suppose  $f(x)$  is twice differentiable on interval  $I$ , then

- ▶  $f$  is convex on  $I$  if and only if  $f'(x)$  is monotonically non-decreasing on  $I$ ;
- ▶  $f$  is convex on  $I$  if and only if  $f''(x) \geq 0$  for  $x \in I$  (often used to test for convexity).

A nice property of convexity:

Any local minimum of a convex function is also a global minimum;  
a strictly convex function has at most one global minimum.  
(Therefore convexity is much desired in optimization.)

## Review Exercises: Morning Session

1. Calculate  $\lim_{x \rightarrow 0} \frac{\sin x}{x}$  and  $\lim_{x \rightarrow 0} \frac{\tan x}{x}$ .

2. Which following functions are convex?

- A  $f_1(x) = |x|, x \in [-1, 1]$ ;
- B  $f_2(x) = \ln(x^2 + 1), x \in \mathbb{R}$ ;
- C  $f_3(x) = e^{-x}, x \in \mathbb{R}$ .

3. Let  $f(x) = \frac{1}{x}, x > 0$ . For every positive integer  $n$ , find  $f^{(n)}(x)$ .

4.  $f(x) = \frac{1}{\sqrt{\gamma}} \exp\left(-\frac{(x-\mu)^2}{\gamma^2}\right)$  where constants  $\gamma > 0$  and  $\mu \in \mathbb{R}$ , and  $x \in \mathbb{R}$ . Find all the global maximums of  $f(x)$ .

# Taylor Expansion

**Taylor Series**, by *3Blue1Brown*

# Challenge Exercises: Morning Session

# Integrals

# Properties of Definite Integrals

Let  $a \leq d \leq b \in \mathbb{R}$ :

- ▶ If  $c \in \mathbb{R}$  is a constant, then  $\int_a^b c dx = c(b - a)$ ;
- ▶  $\int_a^b cf(x) dx = c \int_a^b f(x) dx$ ;
- ▶  $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ ;
- ▶  $\int_a^d f(x) dx + \int_d^b f(x) dx = \int_a^b f(x) dx$ ;
- ▶ If  $f(x) \geq g(x)$  for  $a \leq x \leq b$ , then  $\int_a^b f(x) dx \geq \int_a^b g(x) dx$ ;
- ▶ If  $m \leq f(x) \leq M$  for  $a \leq x \leq b$ , then  $m(b - a) \leq \int_a^b f(x) dx \leq M(b - a)$ .

# The Fundamental Theorem of Calculus

If  $f$  is continuous on  $[a, b]$ , then:

- ▶ function  $g(x) = \int_a^x f(s)ds$ ,  $a \leq x \leq b$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and  $g'(x) = f(x)$ ;
- ▶  $\int_a^b f(x)dx = F(b) - F(a)$ , where  $F$  is any anti-derivative of  $f$  ( $F' = f$ ).

A mini-exercise: find  $\frac{d}{dx} \int_1^x \sin t^4 dt$ .

# Useful Rules for Integration

- ▶ **Substitution rule:** If  $u = g(x)$  is continuously differentiable on  $[a, b]$  and  $f$  is continuous on the range of  $u$ , then

$$\int_a^b f(g(x))g'(x)dx = \int_{g(a)}^{g(b)} f(u)du.$$

- ▶ **Integration by parts:** If functions  $u$  and  $v$  are both continuously differentiable on  $[a, b]$ , then

$$\int_a^b u(x)v'(x)dx = [u(x)v(x)]|_a^b - \int_a^b v(x)u'(x)dx.$$



# Integration: Exercises

1. Calculate  $\int_1^e \frac{\ln x}{x} dx$ .

2. Calculate  $\int_0^\pi x \cos x dx$ .

# Improper Integrals

1. **Infinite intervals:** if  $\int_a^t f(x)dx$  exists for every  $t \geq a$  then

$$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx$$

provided that this limit exists (convergent); similarly, one may define  $\int_{-\infty}^a f(x)dx$ , and if both  $\int_a^\infty f(x)dx$  and  $\int_{-\infty}^a f(x)dx$  are convergent, then  $\int_{-\infty}^\infty f(x)dx = \int_{-\infty}^a f(x)dx + \int_a^\infty f(x)dx$ .

2. **Discontinuous integrand:** if  $f$  is continuous on  $[a, b)$  and is discontinuous at  $b$ , then

$$\int_a^b f(x)dx = \lim_{t \rightarrow b^-} \int_a^t f(x)dx$$

if this limit exists; similarly, if  $f$  is continuous on  $(a, b]$  and is discontinuous at  $a$ ,  $\int_a^b f(x)dx = \lim_{t \rightarrow a^+} \int_t^b f(x)dx$ .

# Improper Integrals: Mini-Exercise

For what values of  $p \in \mathbb{R}$  is the integral

$$\int_1^{\infty} \frac{1}{x^p} dx$$

convergent?

# Series

# Basics of Series

A **series** can be thought of as the infinite sum of a sequence  $\{a_n\}$ , written as  $\sum_{n=1}^{\infty} a_n$  or  $\sum a_n$ . More formally, it can be defined by taking the limit of partial sums  $\{s_n\}$ , where  $s_n = \sum_{i=1}^n a_i$ : if  $\lim_{n \rightarrow \infty} s_n$  exists then the series  $\sum a_n$  is convergent, otherwise it is divergent.

If  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .

An important example - **the geometric series**:

$$\sum_{n=1}^{\infty} ar^{n-1} = a + ar + ar^2 + \dots$$

is convergent if  $|r| < 1$  and its sum is  $\sum_{n=1}^{\infty} ar^{n-1} = \frac{a}{1-r}$ . If  $|r| \geq 1$ , the geometric series is divergent.

# Convergence of Series

Commonly used tests for convergence:

1. **The comparison test:** Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.
  - (i) If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent;
  - (ii) If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.
    - ▶ Video example: **famous proof that the harmonic series diverges**, by *Khan Academy*.
2. **The integral test** (by *Khan Academy*).

# Review Exercises: Afternoon Session

1. Evaluate the following definite integrals:

▶  $\int_0^4 \frac{x}{\sqrt{x^2+9}} dx;$

▶  $\int_0^1 x \ln x dx;$

2. If  $f$  is continuous on  $\mathbb{R}$ , show that

$$\int_a^b f(x+c) dx = \int_{a+c}^{b+c} f(x) dx.$$

3. Do the following series converge? Calculate the value of the infinite sum for each convergent series.

(a)  $\sum_{n=2}^{\infty} 5^{n-1} \left(\frac{9}{10}\right)^n;$

(b)  $\sum_{n=1}^{\infty} \left(\frac{3}{2}\right)^n \frac{1}{9^{n+2}}.$

4. True or false?

▶ If  $x_n \rightarrow 0$  as  $n \rightarrow \infty$ , then  $\sum_{n=1}^{\infty} x_n$  is convergent;

▶  $\sum_{n=1}^{\infty} x^n e^{-nx}$  is convergent for any  $x > 0$ ;

▶  $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$  is convergent.

# Multivariate Calculus



# Partial Derivatives

If  $u$  is a function of  $n$  variables,  $u = f(x_1, x_2, \dots, x_n)$ , its partial derivative with respect to the  $i$ th variable  $x_i$  is

$$\frac{\partial u}{\partial x_i} = \lim_{h \rightarrow 0} \frac{f(x_1, \dots, x_{i-1}, x_i + h, x_{i+1}, \dots, x_n) - f(x_1, \dots, x_i, \dots, x_n)}{h}.$$

(Strategy: treat all the other variables as constants and take the derivative with respect to the variable of interest.)

Suppose  $u = f(x_1, x_2, \dots, x_n)$  is defined on  $\mathbb{R}^n$ . If  $\frac{\partial^2 u}{\partial x_i \partial x_j}$  and  $\frac{\partial^2 u}{\partial x_j \partial x_i}$  are both continuous on  $\mathbb{R}^n$ , then  $\frac{\partial^2 u}{\partial x_i \partial x_j} = \frac{\partial^2 u}{\partial x_j \partial x_i}$ .

# The Gradient Vector and Hessian Matrix

Suppose  $f(x_1, x_2, \dots, x_n)$  is a function of  $n$  variables such that all the partial derivatives exist, then the gradient vector of  $f$  is

$$\nabla f = \left( \frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right).$$

If all the second-order partial derivatives of  $f$  also exist, the Hessian matrix of  $f$  is

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{bmatrix}.$$

## Change of Variables

Take the two-variable case as an example:

Suppose  $z = f(x, y)$  is function of  $x, y$  and  $x = u(s, t)$ ,  $y = v(s, t)$  with respect two other variables  $s, t$ , then  $z = g(s, t)$  as a function of  $s, t$ , where

$$g(s, t) = f(u(s, t), v(s, t))|J|.$$

Here  $J$  is the **Jacobian** of the transformation  $x = u(s, t)$ ,  $y = v(s, t)$ :

$$J = \begin{vmatrix} \frac{\partial x}{\partial s} & \frac{\partial x}{\partial t} \\ \frac{\partial y}{\partial s} & \frac{\partial y}{\partial t} \end{vmatrix} = \frac{\partial x}{\partial s} \frac{\partial y}{\partial t} - \frac{\partial x}{\partial t} \frac{\partial y}{\partial s}.$$

Suppose that we want to integrate  $f(x, y)$  over a region  $R$ . Under the transformation  $x = u(s, t)$ ,  $y = v(s, t)$  the regions becomes  $S$  and the integral becomes:

$$\iint_R f(x, y) dx dy = \iint_S f(u(s, t), v(s, t)) |J| ds dt.$$

# Multivariate Calculus Review Exercises

1. Let  $f(x, y, z) = ye^x \ln z + z \tan z$  ( $z \in (0, \frac{\pi}{2})$ ) and  $g(x, y, z) = x^3y + y^3z + z^5 + \sqrt{xyz}$ .  
Compute  $\frac{\partial f}{\partial y}$ ,  $\frac{\partial f}{\partial z}$ ,  $\frac{\partial g}{\partial x}$ , and  $\frac{\partial g}{\partial y}$ .

2.  $f(x, y) = \frac{1}{\sqrt{2\pi y^2}} e^{-\frac{(x-s)^2}{2y^2}}$ , where  $s \in \mathbb{R}$  is a constant, and  $y > 0$ . Obtain the gradient vector and Hessian matrix of  $g(x, y) = \ln f(x, y)$ .

3. Evaluate the double integral  $\int_1^3 \int_0^2 (xy + x^2y^3) dy dx$ .

# Challenge Exercises: Afternoon Session