

Risk Summary

Preeclampsia

Medium Score: 0.6395082154771051

Model Interpretation: Risk-based triage only (not diagnosis).

Macrosomia

Medium Score: 0.5943816940585895

Model Interpretation: Risk-based coaching support.

Summary: Patient TEST_MED_MED at 32 weeks gestation has a medium risk for preeclampsia due to elevated systolic blood pressure and positive proteinuria. A medium risk for macrosomia is noted, driven by a high pre-pregnancy BMI and borderline fasting glucose, despite positive lifestyle efforts. An immediate doctor alert is recommended.

Patient Snapshot (Recent Inputs)

Parameter	Value	Notes
Blood Pressure	142/79 mmHg	Hypertensive threshold ≥140/90, severe ≥160/110.
Proteinuria (dipstick)	1	≥1+ supports PE suspicion in context of HTN.
Symptoms	Headache=0, Vision=0, RUQ Pain=0, Swelling=0	Severe features increase escalation priority.
Glucose	Fasting=99 mg/dL 1hr PP=126 mg/dL	Targets typically fasting <95, 1hr <140 (pregnancy goals).
HbA1c	5.6	Higher HbA1c indicates sustained glycemic elevation.
Maternal Context	Age=38 BMI=42.4 GDM=0	Risk modifiers for fetal overgrowth + metabolic risk.
Activity / Sleep	Post-meal walk=29 min Sleep=6.1 hrs	Lifestyle factors influence glycemic response.
Food Log	paneer salad	Used for dietary correlation patterns.

Reason for Alert

Action Requested: ALERT_DOCTOR

Clinical Rationale: Patient TEST_MED_MED, 32 weeks gestation, presents with a medium preeclampsia risk tier. Key vitals include systolic BP 142 mmHg, diastolic BP 79 mmHg, and proteinuria dipstick 1. No severe symptoms (headache, vision changes, abdominal pain, swelling) reported. There is also a medium macrosomia risk tier, with pre-pregnancy BMI 42.4 kg/m², fasting glucose 99 mg/dL, and HbA1c 5.6%. Patient is not diagnosed with GDM. Review of preeclampsia indicators is recommended urgently.

Suggested Next Steps

- Review trend: BP, symptoms, proteinuria, glucose readings
- Consider confirmatory tests (protein/creatinine ratio, LFTs, platelets) if clinically indicated
- Consider GDM diet review or insulin adjustment if glucose targets exceeded
- Provide patient education + return precautions

