

Risk Summary

Preeclampsia

Medium Score: 0.6395082154771051

Model Interpretation: Risk-based triage only (not diagnosis).

Macrosomia

Medium Score: 0.5943816940585895

Model Interpretation: Risk-based coaching support.

Summary: The patient is at 32 weeks gestation with a 'medium' risk for preeclampsia and 'medium' risk for macrosomia. Key findings include an elevated systolic blood pressure (142 mmHg) and proteinuria (1 on dipstick), contributing to the preeclampsia risk. While severe symptoms are not reported, these findings warrant a doctor's review.

Patient Snapshot (Recent Inputs)

Parameter	Value	Notes
Blood Pressure	142/79 mmHg	Hypertensive threshold ≥140/90, severe ≥160/110.
Proteinuria (dipstick)	1	≥1+ supports PE suspicion in context of HTN.
Symptoms	Headache=0, Vision=0, RUQ Pain=0, Swelling=0	Severe features increase escalation priority.
Glucose	Fasting=99 mg/dL 1hr PP=126 mg/dL	Targets typically fasting <95, 1hr <140 (pregnancy goals).
HbA1c	5.6	Higher HbA1c indicates sustained glycemic elevation.
Maternal Context	Age=38 BMI=42.4 GDM=0	Risk modifiers for fetal overgrowth + metabolic risk.
Activity / Sleep	Post-meal walk=29 min Sleep=6.1 hrs	Lifestyle factors influence glycemic response.
Food Log	paneer salad	Used for dietary correlation patterns.

Reason for Alert

Action Requested: ALERT_DOCTOR

Clinical Rationale: Preeclampsia Risk: Medium (score: 0.6395). Macrosomia Risk: Medium (score: 0.5944). Patient is 32 weeks gestational age. Key vitals include systolic BP 142 mmHg, diastolic BP 79 mmHg, and proteinuria dipstick 1. Fasting glucose 99 mg/dL. No severe headache, vision changes, upper abdominal pain, or swelling of face/hands reported. Elevated systolic BP and proteinuria contribute to the preeclampsia risk. Requesting urgent review for further evaluation and management.

Suggested Next Steps

- Review trend: BP, symptoms, proteinuria, glucose readings
- Consider confirmatory tests (protein/creatinine ratio, LFTs, platelets) if clinically indicated
- Consider GDM diet review or insulin adjustment if glucose targets exceeded
- Provide patient education + return precautions

