

✓ Congratulations! You passed!

Grade received 100% To pass 80% or higher

Go to next item

Practice quiz: Gradient descent for logistic regression

Latest Submission Grade 100%

1.

1 / 1 point

Gradient descent for logistic regression

repeat {

$$w_j = w_j - \alpha \left[\frac{1}{m} \sum_{i=1}^m (f_{\vec{w},b}(\vec{x}^{(i)}) - y^{(i)}) x_j^{(i)} \right]$$
$$b = b - \alpha \left[\frac{1}{m} \sum_{i=1}^m (f_{\vec{w},b}(\vec{x}^{(i)}) - y^{(i)}) \right]$$

} simultaneous updates

$$f_{\vec{w},b}(\vec{x}) = \frac{1}{1 + e^{-(\vec{w} \cdot \vec{x} + b)}}$$

Which of the following two statements is a more accurate statement about gradient descent for logistic regression?

- ☐ The update steps are identical to the update steps for linear regression.
- ☒ The update steps look like the update steps for linear regression, but the definition of $f_{\vec{w},b}(\mathbf{x}^{(i)})$ is different.

✓ Correct

For logistic regression, $f_{\vec{w},b}(\mathbf{x}^{(i)})$ is the sigmoid function instead of a straight line.