

# Rubik's cubes and permutation group theory

---

**Lawrence Chen**

October 8, 2022

**Honours presentation**



# Contents

## Some basic group theory

- What is a group?

- Order and generators

- Permutations

- Group actions

## The Rubik's group

- Representing the cube and its moves

- Moves vs states for Rubik's cube

- The Rubik's group of permutations

- Orbits and stabilisers

- Orders of moves

- Jake's theorems

## Analysing the Rubik's group

- Bases and stabiliser chains

- How many valid states are there?

- Can this restickering be solved?

- Solving a Rubik's cube...

## Concluding remarks

- References

## Questions about Rubik's cube i

- How can we represent *move sequences* and *states* of a cube?

## Questions about Rubik's cube i

- How can we represent *move sequences* and *states* of a cube?
- How can we tell *how many* states a Rubik's cube can take?

## Questions about Rubik's cube i

- How can we represent *move sequences* and *states* of a cube?
- How can we tell *how many* states a Rubik's cube can take?
- If we repeat a move, do we eventually *get back to the start*?

## Questions about Rubik's cube i

- How can we represent *move sequences* and *states* of a cube?
- How can we tell *how many* states a Rubik's cube can take?
- If we repeat a move, do we eventually *get back to the start*?
- If a Rubik's cube is *restickered*, is it *solvable*?

## Questions about Rubik's cube i

- How can we represent *move sequences* and *states* of a cube?
- How can we tell *how many* states a Rubik's cube can take?
- If we repeat a move, do we eventually *get back to the start*?
- If a Rubik's cube is *restickered*, is it *solvable*?
- How can we use maths to *solve* a Rubik's cube?

## Questions about Rubik's cube i

- How can we represent *move sequences* and *states* of a cube?
- How can we tell *how many* states a Rubik's cube can take?
- If we repeat a move, do we eventually *get back to the start*?
- If a Rubik's cube is *restickered*, is it *solvable*?
- How can we use maths to *solve* a Rubik's cube?

*Answer:* using permutations and *permutation group theory*!



# Questions about Rubik's cube i

- How can we represent *move sequences* and *states* of a cube?
- How can we tell *how many* states a Rubik's cube can take?
- If we repeat a move, do we eventually *get back to the start*?
- If a Rubik's cube is *restickered*, is it *solvable*?
- How can we use maths to *solve* a Rubik's cube?

*Answer: using permutations and permutation group theory!*

## **(J. A. Paulos, Innumeracy)**

*Ideal Toy Company stated on the package of the original Rubik cube that there were more than three billion possible states the cube could attain. It's analogous to McDonald's proudly announcing that they've sold more than 120 hamburgers.*

## **Some basic group theory**

---

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

(i) (**identity**) there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

- (i) (**identity**) there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;
- (ii) (**inverses**) for all  $g \in G$ , there is  $g^{-1} \in G$  with  $g^{-1}g = gg^{-1} = 1$ ;

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

- (i) **(identity)** there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;
- (ii) **(inverses)** for all  $g \in G$ , there is  $g^{-1} \in G$  with  $g^{-1}g = gg^{-1} = 1$ ;
- (iii) **(associative)**  $(gh)k = g(hk)$  for all  $g, h, k \in G$ .

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

- (i) **(identity)** there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;
- (ii) **(inverses)** for all  $g \in G$ , there is  $g^{-1} \in G$  with  $g^{-1}g = gg^{-1} = 1$ ;
- (iii) **(associative)**  $(gh)k = g(hk)$  for all  $g, h, k \in G$ .

## Example (Integers under addition)

The integers  $(\mathbb{Z}, +)$  form an **abelian** group: identity

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

- (i) **(identity)** there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;
- (ii) **(inverses)** for all  $g \in G$ , there is  $g^{-1} \in G$  with  $g^{-1}g = gg^{-1} = 1$ ;
- (iii) **(associative)**  $(gh)k = g(hk)$  for all  $g, h, k \in G$ .

## Example (Integers under addition)

The integers  $(\mathbb{Z}, +)$  form an **abelian** group: identity 0, inverses



# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

- (i) **(identity)** there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;
- (ii) **(inverses)** for all  $g \in G$ , there is  $g^{-1} \in G$  with  $g^{-1}g = gg^{-1} = 1$ ;
- (iii) **(associative)**  $(gh)k = g(hk)$  for all  $g, h, k \in G$ .

## Example (Integers under addition)

The integers  $(\mathbb{Z}, +)$  form an **abelian** group: identity 0, inverses  $-k$  for  $k \in \mathbb{Z}$ , associative.

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

- (i) **(identity)** there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;
- (ii) **(inverses)** for all  $g \in G$ , there is  $g^{-1} \in G$  with  $g^{-1}g = gg^{-1} = 1$ ;
- (iii) **(associative)**  $(gh)k = g(hk)$  for all  $g, h, k \in G$ .

## Example (Integers under addition)

The integers  $(\mathbb{Z}, +)$  form an **abelian** group: identity 0, inverses  $-k$  for  $k \in \mathbb{Z}$ , associative.

## Example (Cyclic group)

The set  $C_n = \{a^0, a^1, a^2, \dots, a^{n-1}\}$  with rules  $a^k a^\ell = a^{k+\ell}$ ,  $a^n = a^0$  forms group: identity

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

- (i) **(identity)** there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;
- (ii) **(inverses)** for all  $g \in G$ , there is  $g^{-1} \in G$  with  $g^{-1}g = gg^{-1} = 1$ ;
- (iii) **(associative)**  $(gh)k = g(hk)$  for all  $g, h, k \in G$ .

## Example (Integers under addition)

The integers  $(\mathbb{Z}, +)$  form an **abelian** group: identity 0, inverses  $-k$  for  $k \in \mathbb{Z}$ , associative.

## Example (Cyclic group)

The set  $C_n = \{a^0, a^1, a^2, \dots, a^{n-1}\}$  with rules  $a^k a^\ell = a^{k+\ell}$ ,  $a^n = a^0$  forms group: identity  $1 = a^0$ , inverses

# What is a group? i

## Definition (group)

A **group** is a set  $G \neq \emptyset$  with operation  $G \times G \rightarrow G$ ,  $(g, h) \mapsto gh$ ,

- (i) **(identity)** there is  $1 \in G$  with  $1g = g1 = g$  for all  $g \in G$ ;
- (ii) **(inverses)** for all  $g \in G$ , there is  $g^{-1} \in G$  with  $g^{-1}g = gg^{-1} = 1$ ;
- (iii) **(associative)**  $(gh)k = g(hk)$  for all  $g, h, k \in G$ .

## Example (Integers under addition)

The integers  $(\mathbb{Z}, +)$  form an **abelian** group: identity 0, inverses  $-k$  for  $k \in \mathbb{Z}$ , associative.

## Example (Cyclic group)

The set  $C_n = \{a^0, a^1, a^2, \dots, a^{n-1}\}$  with rules  $a^k a^\ell = a^{k+\ell}$ ,  $a^n = a^0$  forms group: identity  $1 = a^0$ , inverses  $a^{-k}$  for  $a^k \in C_n$ , associative.

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is



# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is 2, order of  $a^3$  is

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is 2, order of  $a^3$  is 4.

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is 2, order of  $a^3$  is 4.

## Definition (generator)

Set  $X$  **generates**  $G$  if every  $g \in G$  is  $g = x_1^{\pm 1} \cdots x_r^{\pm 1}$  for some  $r \in \mathbb{N}$ ,  $x_i \in X$  **generators**; write  $G = \langle X \rangle$ .

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is 2, order of  $a^3$  is 4.

## Definition (generator)

Set  $X$  **generates**  $G$  if every  $g \in G$  is  $g = x_1^{\pm 1} \cdots x_r^{\pm 1}$  for some  $r \in \mathbb{N}$ ,  $x_i \in X$  **generators**; write  $G = \langle X \rangle$ . (If  $|X| = 1$ ,  $G$  is **cyclic**.)

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is 2, order of  $a^3$  is 4.

## Definition (generator)

Set  $X$  **generates**  $G$  if every  $g \in G$  is  $g = x_1^{\pm 1} \cdots x_r^{\pm 1}$  for some  $r \in \mathbb{N}$ ,  $x_i \in X$  **generators**; write  $G = \langle X \rangle$ . (If  $|X| = 1$ ,  $G$  is **cyclic**.)

## Example (Cyclic group)

Consider group  $C_6 = \{1, a, a^2, a^3, a^4, a^5\}$ :

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is 2, order of  $a^3$  is 4.

## Definition (generator)

Set  $X$  **generates**  $G$  if every  $g \in G$  is  $g = x_1^{\pm 1} \cdots x_r^{\pm 1}$  for some  $r \in \mathbb{N}$ ,  $x_i \in X$  **generators**; write  $G = \langle X \rangle$ . (If  $|X| = 1$ ,  $G$  is **cyclic**.)

## Example (Cyclic group)

Consider group  $C_6 = \{1, a, a^2, a^3, a^4, a^5\}$ :  $C_6 = \langle a \rangle$ .

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is 2, order of  $a^3$  is 4.

## Definition (generator)

Set  $X$  **generates**  $G$  if every  $g \in G$  is  $g = x_1^{\pm 1} \cdots x_r^{\pm 1}$  for some  $r \in \mathbb{N}$ ,  $x_i \in X$  **generators**; write  $G = \langle X \rangle$ . (If  $|X| = 1$ ,  $G$  is **cyclic**.)

## Example (Cyclic group)

Consider group  $C_6 = \{1, a, a^2, a^3, a^4, a^5\}$ :  $C_6 = \langle a \rangle$ . If  $b = a^2$ ,  $c = a^3$  then  $C_6 = \langle b, c \rangle$  since

# Order and generators i

## Definition (order)

**Order** of  $g \in G$  is least  $k \in \mathbb{Z}_+$  with  $g^k = g \cdots g = 1$  (otherwise  $\infty$ ).

## Example (Cyclic group)

Consider group  $C_4 = \{1, a, a^2, a^3\}$ : order of 1 is 1, order of  $a$  is 4, order of  $a^2$  is 2, order of  $a^3$  is 4.

## Definition (generator)

Set  $X$  **generates**  $G$  if every  $g \in G$  is  $g = x_1^{\pm 1} \cdots x_r^{\pm 1}$  for some  $r \in \mathbb{N}$ ,  $x_i \in X$  **generators**; write  $G = \langle X \rangle$ . (If  $|X| = 1$ ,  $G$  is **cyclic**.)

## Example (Cyclic group)

Consider group  $C_6 = \{1, a, a^2, a^3, a^4, a^5\}$ :  $C_6 = \langle a \rangle$ . If  $b = a^2$ ,  $c = a^3$  then  $C_6 = \langle b, c \rangle$  since  $a = cb^{-1}$  so  $a^k = cb^{-1} \cdots cb^{-1} = c^k b^{-k}$ .



## **Definition (permutation)**

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

# Permutations i

## Definition (permutation)

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

Write  $1 = ()$  for identity. Write  $i^\sigma$  not  $\sigma(i)$  for *image*.

# Permutations i

## Definition (permutation)

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

Write  $1 = ()$  for identity. Write  $i^\sigma$  not  $\sigma(i)$  for *image*.

*Cycle notation:*  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$  is:

$$\begin{array}{cccccc} & 1 & 2 & 3 & 4 & 5 & 6 \\ \sigma & & & & & & \\ & 1 & 2 & 3 & 4 & 5 & 6 \end{array}$$

It means


# Permutations i

## Definition (permutation)

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

Write  $1 = ()$  for identity. Write  $i^\sigma$  not  $\sigma(i)$  for *image*.

*Cycle notation:*  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$  is:



It means

$$1^\sigma = 4,$$


# Permutations i

## Definition (permutation)

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

Write  $1 = ()$  for identity. Write  $i^\sigma$  not  $\sigma(i)$  for *image*.

*Cycle notation:*  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$  is:



It means

$$1^\sigma = 4, 4^\sigma = 5,$$


# Permutations i

## Definition (permutation)

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

Write  $1 = ()$  for identity. Write  $i^\sigma$  not  $\sigma(i)$  for *image*.

*Cycle notation:*  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$  is:



It means

$$1^\sigma = 4, 4^\sigma = 5, 5^\sigma = 1,$$


# Permutations i

## Definition (permutation)

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

Write  $1 = ()$  for identity. Write  $i^\sigma$  not  $\sigma(i)$  for *image*.

*Cycle notation:*  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$  is:



It means

$$1^\sigma = 4, 4^\sigma = 5, 5^\sigma = 1, 2^\sigma = 6,$$


# Permutations i

## Definition (permutation)

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

Write  $1 = ()$  for identity. Write  $i^\sigma$  not  $\sigma(i)$  for *image*.

*Cycle notation:*  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$  is:



It means

$$1^\sigma = 4, 4^\sigma = 5, 5^\sigma = 1, 2^\sigma = 6, 6^\sigma = 2,$$




# Permutations i

## Definition (permutation)

**Permutation** of  $[n] := \{1, \dots, n\}$  is bijection  $\sigma : [n] \rightarrow [n]$ .

Write  $1 = ()$  for identity. Write  $i^\sigma$  not  $\sigma(i)$  for *image*.

*Cycle notation:*  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$  is:




It means

$$1^\sigma = 4, 4^\sigma = 5, 5^\sigma = 1, 2^\sigma = 6, 6^\sigma = 2, 3^\sigma = 3.$$


## Permutations ii

*Inverses:* For  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$ :



## Permutations ii

*Inverses:* For  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$ :




Inverse is  $\sigma^{-1} = (1, 5, 4)(2, 6) \in \text{Sym}(6)$ .

*Product/composition:* for  $\sigma, \tau \in \text{Sym}(n)$ ,  $\sigma\tau$  means “first  $\sigma$ , then  $\tau$ ”,  
so  $i^{\sigma\tau} = (i^\sigma)^\tau$ .


## Permutations iii

*Product/composition:* for  $\sigma, \tau \in \text{Sym}(n)$ ,  $\sigma\tau$  means “first  $\sigma$ , then  $\tau$ ”, so  $i^{\sigma\tau} = (i^\sigma)^\tau$ . E.g.  $\sigma = (1, 2, 3)$ ,





## Permutations iii

*Product/composition:* for  $\sigma, \tau \in \text{Sym}(n)$ ,  $\sigma\tau$  means “first  $\sigma$ , then  $\tau$ ”, so  $i^{\sigma\tau} = (i^\sigma)^\tau$ . E.g.  $\sigma = (1, 2, 3), \tau = (1, 3)(2, 4) \in \text{Sym}(4)$ ,



## Permutations iii



*Product/composition:* for  $\sigma, \tau \in \text{Sym}(n)$ ,  $\sigma\tau$  means “first  $\sigma$ , then  $\tau$ ”, so  $i^{\sigma\tau} = (i^\sigma)^\tau$ . E.g.  $\sigma = (1, 2, 3), \tau = (1, 3)(2, 4) \in \text{Sym}(4)$ ,



$$\sigma\tau = (1, 2, 3)(1, 3)(2, 4) = (1,$$

## Permutations iii

*Product/composition:* for  $\sigma, \tau \in \text{Sym}(n)$ ,  $\sigma\tau$  means “first  $\sigma$ , then  $\tau$ ”, so  $i^{\sigma\tau} = (i^\sigma)^\tau$ . E.g.  $\sigma = (1, 2, 3), \tau = (1, 3)(2, 4) \in \text{Sym}(4)$ ,





$$\sigma\tau = (1, 2, 3)(1, 3)(2, 4) = (1, 4,$$



## Permutations iii



*Product/composition:* for  $\sigma, \tau \in \text{Sym}(n)$ ,  $\sigma\tau$  means “first  $\sigma$ , then  $\tau$ ”, so  $i^{\sigma\tau} = (i^\sigma)^\tau$ . E.g.  $\sigma = (1, 2, 3), \tau = (1, 3)(2, 4) \in \text{Sym}(4)$ ,



$$\sigma\tau = (1, 2, 3)(1, 3)(2, 4) = (1, 4, 2)$$

## Permutations iii



*Product/composition:* for  $\sigma, \tau \in \text{Sym}(n)$ ,  $\sigma\tau$  means “first  $\sigma$ , then  $\tau$ ”, so  $i^{\sigma\tau} = (i^\sigma)^\tau$ . E.g.  $\sigma = (1, 2, 3), \tau = (1, 3)(2, 4) \in \text{Sym}(4)$ ,



$$\sigma\tau = (1, 2, 3)(1, 3)(2, 4) = (1, 4, 2)$$

## Permutations iii

*Product/composition:* for  $\sigma, \tau \in \text{Sym}(n)$ ,  $\sigma\tau$  means “first  $\sigma$ , then  $\tau$ ”, so  $i^{\sigma\tau} = (i^\sigma)^\tau$ . E.g.  $\sigma = (1, 2, 3), \tau = (1, 3)(2, 4) \in \text{Sym}(4)$ ,




$$\sigma\tau = (1, 2, 3)(1, 3)(2, 4) = (1, 4, 2) \in \text{Sym}(4).$$

*Inverse of product:* Is  $(\sigma\tau)^{-1} = \sigma^{-1}\tau^{-1}$ ?

## Permutations iv

*Inverse of product:* Is  $(\sigma\tau)^{-1} = \sigma^{-1}\tau^{-1}$ ?


$$\sigma^{-1} = (1, 3, 2),$$



## Permutations iv

*Inverse of product:* Is  $(\sigma\tau)^{-1} = \sigma^{-1}\tau^{-1}$ ?



$$\sigma^{-1} = (1, 3, 2), \tau^{-1} = (1, 3)(2, 4),$$



## Permutations iv

*Inverse of product:* Is  $(\sigma\tau)^{-1} = \sigma^{-1}\tau^{-1}$ ?



$\sigma^{-1} = (1, 3, 2)$ ,  $\tau^{-1} = (1, 3)(2, 4)$ ,  $(\sigma\tau)^{-1} = (1, 2, 4)$ .



## Permutations iv

*Inverse of product:* Is  $(\sigma\tau)^{-1} = \sigma^{-1}\tau^{-1}$ ?

$\sigma^{-1} = (1, 3, 2)$ ,  $\tau^{-1} = (1, 3)(2, 4)$ ,  $(\sigma\tau)^{-1} = (1, 2, 4)$ .





$$\sigma^{-1}\tau^{-1} = (1, 3, 2)(1, 3)(2, 4) = (2, 3, 4) \neq (\sigma\tau)^{-1},$$



## Permutations iv

*Inverse of product:* Is  $(\sigma\tau)^{-1} = \sigma^{-1}\tau^{-1}$ ?

$\sigma^{-1} = (1, 3, 2)$ ,  $\tau^{-1} = (1, 3)(2, 4)$ ,  $(\sigma\tau)^{-1} = (1, 2, 4)$ .



$$\sigma^{-1}\tau^{-1} = (1, 3, 2)(1, 3)(2, 4) = (2, 3, 4) \neq (\sigma\tau)^{-1},$$

$$\tau^{-1}\sigma^{-1} = (1, 3)(2, 4)(1, 3, 2) = (1, 2, 4) = (\sigma\tau)^{-1}.$$

Set of permutations under *product* is **symmetric group**  $\text{Sym}(n)$ :  
identity  $1 = ()$ , inverses (since bijection), associative.

What is size of  $\text{Sym}(n)$ ?

Set of permutations under *product* is **symmetric group**  $\text{Sym}(n)$ :  
identity  $1 = ()$ , inverses (since bijection), associative.

What is size of  $\text{Sym}(n)$ ? *Answer:*  $n!$

## Example (Order of permutation)

Consider  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$ . Then  $1^{\sigma^3} = 4^{\sigma^2} = 5^{\sigma} = 1$ ,

## Permutations v

Set of permutations under *product* is **symmetric group**  $\text{Sym}(n)$ :  
identity  $1 = ()$ , inverses (since bijection), associative.

What is size of  $\text{Sym}(n)$ ? *Answer:*  $n!$

### Example (Order of permutation)

Consider  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$ . Then  $1^{\sigma^3} = 4^{\sigma^2} = 5^{\sigma} = 1$ ,  
 $4^{\sigma^3} = 4$ ,  $5^{\sigma^3} = 5$ ,  $2^{\sigma^2} = 2$ ,  $6^{\sigma^2} = 6$  so

## Permutations v

Set of permutations under *product* is **symmetric group**  $\text{Sym}(n)$ :  
identity  $1 = ()$ , inverses (since bijection), associative.

What is size of  $\text{Sym}(n)$ ? *Answer:*  $n!$

### Example (Order of permutation)

Consider  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$ . Then  $1^{\sigma^3} = 4^{\sigma^2} = 5^{\sigma} = 1$ ,  
 $4^{\sigma^3} = 4$ ,  $5^{\sigma^3} = 5$ ,  $2^{\sigma^2} = 2$ ,  $6^{\sigma^2} = 6$  so  $\sigma^6 = () = 1$ ; order of  $\sigma$  is 6.

# Permutations v

Set of permutations under *product* is **symmetric group**  $\text{Sym}(n)$ :  
identity  $1 = ()$ , inverses (since bijection), associative.

What is size of  $\text{Sym}(n)$ ? *Answer:*  $n!$

## Example (Order of permutation)

Consider  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$ . Then  $1^{\sigma^3} = 4^{\sigma^2} = 5^{\sigma} = 1$ ,  
 $4^{\sigma^3} = 4$ ,  $5^{\sigma^3} = 5$ ,  $2^{\sigma^2} = 2$ ,  $6^{\sigma^2} = 6$  so  $\sigma^6 = () = 1$ ; order of  $\sigma$  is 6.

*Fact:* order of  $\sigma \in \text{Sym}(n)$  is lcm of cycle lengths.

# Permutations v

Set of permutations under *product* is **symmetric group**  $\text{Sym}(n)$ :  
identity  $1 = ()$ , inverses (since bijection), associative.

What is size of  $\text{Sym}(n)$ ? *Answer:*  $n!$

## Example (Order of permutation)

Consider  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$ . Then  $1^{\sigma^3} = 4^{\sigma^2} = 5^{\sigma} = 1$ ,  
 $4^{\sigma^3} = 4$ ,  $5^{\sigma^3} = 5$ ,  $2^{\sigma^2} = 2$ ,  $6^{\sigma^2} = 6$  so  $\sigma^6 = () = 1$ ; order of  $\sigma$  is 6.

*Fact:* order of  $\sigma \in \text{Sym}(n)$  is lcm of cycle lengths.

## Definition (subgroup)

Subset  $H$  of group  $G$  is **subgroup** if it is group under same operation;  
write  $H \leq G$ . (Need to check: nonempty, closure, inverses.)

# Permutations v

Set of permutations under *product* is **symmetric group**  $\text{Sym}(n)$ :  
identity  $1 = ()$ , inverses (since bijection), associative.

What is size of  $\text{Sym}(n)$ ? *Answer:*  $n!$

## Example (Order of permutation)

Consider  $\sigma = (1, 4, 5)(2, 6) \in \text{Sym}(6)$ . Then  $1^{\sigma^3} = 4^{\sigma^2} = 5^{\sigma} = 1$ ,  
 $4^{\sigma^3} = 4$ ,  $5^{\sigma^3} = 5$ ,  $2^{\sigma^2} = 2$ ,  $6^{\sigma^2} = 6$  so  $\sigma^6 = () = 1$ ; order of  $\sigma$  is 6.

*Fact:* order of  $\sigma \in \text{Sym}(n)$  is lcm of cycle lengths.

## Definition (subgroup)

Subset  $H$  of group  $G$  is **subgroup** if it is group under same operation;  
write  $H \leq G$ . (Need to check: nonempty, closure, inverses.)

## Definition (permutation group)

A **permutation group** of *degree*  $n$  is a subgroup of  $\text{Sym}(n)$ .



### Definition (group action)

If  $G$  is group and  $\Omega \neq \emptyset$  is set, a  **$G$ -action** is a map  $\Omega \times G \rightarrow \Omega$ ,  $(\alpha, g) \mapsto \alpha^g$  s.t.  $\alpha^1 = \alpha$  and  $\alpha^{gh} = (\alpha^g)^h$  for  $\alpha \in \Omega$  and  $g, h \in G$ .

*Idea:*  $\alpha \in \Omega$  is *state*, apply *move*  $g \in G$  to get state  $\alpha^g \in \Omega$ , in way that respects group operation.

# Group actions i

## Definition (group action)

If  $G$  is group and  $\Omega \neq \emptyset$  is set, a  **$G$ -action** is a map  $\Omega \times G \rightarrow \Omega$ ,  $(\alpha, g) \mapsto \alpha^g$  s.t.  $\alpha^1 = \alpha$  and  $\alpha^{gh} = (\alpha^g)^h$  for  $\alpha \in \Omega$  and  $g, h \in G$ .

*Idea:*  $\alpha \in \Omega$  is *state*, apply *move*  $g \in G$  to get state  $\alpha^g \in \Omega$ , in way that respects group operation.

## Example (adding time)

$\mathbb{Z}$  acts on  $\Omega = \{12:00, 1:00, \dots, 11:00\}$  by  $(\alpha:00)^k = [\alpha + k]_{12:00}$  for  $\alpha:00 \in \Omega$  and  $k \in \mathbb{Z}$ .

# Group actions i

## Definition (group action)

If  $G$  is group and  $\Omega \neq \emptyset$  is set, a  **$G$ -action** is a map  $\Omega \times G \rightarrow \Omega$ ,  $(\alpha, g) \mapsto \alpha^g$  s.t.  $\alpha^1 = \alpha$  and  $\alpha^{gh} = (\alpha^g)^h$  for  $\alpha \in \Omega$  and  $g, h \in G$ .

*Idea:*  $\alpha \in \Omega$  is *state*, apply *move*  $g \in G$  to get state  $\alpha^g \in \Omega$ , in way that respects group operation.

## Example (adding time)

$\mathbb{Z}$  acts on  $\Omega = \{12:00, 1:00, \dots, 11:00\}$  by  $(\alpha:00)^k = [\alpha + k]_{12:00}$  for  $\alpha:00 \in \Omega$  and  $k \in \mathbb{Z}$ .

E.g. 5:00 plus 9 hrs is  $(5:00)^9 = [5 + 9]_{12:00} = 2:00$ .

# Group actions i

## Definition (group action)

If  $G$  is group and  $\Omega \neq \emptyset$  is set, a  **$G$ -action** is a map  $\Omega \times G \rightarrow \Omega$ ,  $(\alpha, g) \mapsto \alpha^g$  s.t.  $\alpha^1 = \alpha$  and  $\alpha^{gh} = (\alpha^g)^h$  for  $\alpha \in \Omega$  and  $g, h \in G$ .

*Idea:*  $\alpha \in \Omega$  is *state*, apply *move*  $g \in G$  to get state  $\alpha^g \in \Omega$ , in way that respects group operation.

## Example (adding time)

$\mathbb{Z}$  acts on  $\Omega = \{12:00, 1:00, \dots, 11:00\}$  by  $(\alpha:00)^k = [\alpha + k]_{12:00}$  for  $\alpha:00 \in \Omega$  and  $k \in \mathbb{Z}$ .

E.g. 5:00 plus 9 hrs is  $(5:00)^9 = [5 + 9]_{12:00} = 2:00$ .

## Example (natural action)

$G \leq \text{Sym}(n)$  acts on  $\Omega = [n]$  by  $\alpha^g = \alpha^g$  (image) for  $\alpha \in [n]$ ,  $g \in G$ .

# Group actions i

## Definition (group action)

If  $G$  is group and  $\Omega \neq \emptyset$  is set, a  **$G$ -action** is a map  $\Omega \times G \rightarrow \Omega$ ,  $(\alpha, g) \mapsto \alpha^g$  s.t.  $\alpha^1 = \alpha$  and  $\alpha^{gh} = (\alpha^g)^h$  for  $\alpha \in \Omega$  and  $g, h \in G$ .

*Idea:*  $\alpha \in \Omega$  is *state*, apply *move*  $g \in G$  to get state  $\alpha^g \in \Omega$ , in way that respects group operation.

## Example (adding time)

$\mathbb{Z}$  acts on  $\Omega = \{12:00, 1:00, \dots, 11:00\}$  by  $(\alpha:00)^k = [\alpha + k]_{12:00}$  for  $\alpha:00 \in \Omega$  and  $k \in \mathbb{Z}$ .

E.g. 5:00 plus 9 hrs is  $(5:00)^9 = [5 + 9]_{12:00} = 2:00$ .

## Example (natural action)

$G \leq \text{Sym}(n)$  acts on  $\Omega = [n]$  by  $\alpha^g = \alpha^g$  (image) for  $\alpha \in [n]$ ,  $g \in G$ .

## Example (right regular action)

Group  $G$  acts on  $\Omega = G$  (itself) via  $\alpha^g = \alpha g$  for  $\alpha, g \in G$ .

### Definition (orbit)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$ .

*Idea:* states  $\alpha^g \in \Omega$  reachable from fixed  $\alpha \in \Omega$  by moves  $g \in G$ .

### Definition (orbit)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$ .

*Idea:* states  $\alpha^g \in \Omega$  reachable from fixed  $\alpha \in \Omega$  by moves  $g \in G$ .

### Definition (stabiliser)

If  $G$  acts on  $\Omega$ , then **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Idea:* moves  $g \in G$  that fix given  $\alpha \in \Omega$ .

### Definition (orbit)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$ .

*Idea:* states  $\alpha^g \in \Omega$  reachable from fixed  $\alpha \in \Omega$  by moves  $g \in G$ .

### Definition (stabiliser)

If  $G$  acts on  $\Omega$ , then **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Idea:* moves  $g \in G$  that fix given  $\alpha \in \Omega$ .

### Example (Adding time)

$\mathbb{Z}$ -orbit of 11:00 is



### Definition (orbit)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$ .

*Idea:* states  $\alpha^g \in \Omega$  reachable from fixed  $\alpha \in \Omega$  by moves  $g \in G$ .

### Definition (stabiliser)

If  $G$  acts on  $\Omega$ , then **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Idea:* moves  $g \in G$  that fix given  $\alpha \in \Omega$ .

### Example (Adding time)

$\mathbb{Z}$ -orbit of 11:00 is  $\Omega = \{12:00, \dots, 11:00\}$  (e.g.  $(11:00)^{-2} = 9:00$ ).

$\mathbb{Z}$ -stabiliser of 11:00 is

## Group actions ii

### Definition (orbit)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$ .

*Idea:* states  $\alpha^g \in \Omega$  reachable from fixed  $\alpha \in \Omega$  by moves  $g \in G$ .

### Definition (stabiliser)

If  $G$  acts on  $\Omega$ , then **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Idea:* moves  $g \in G$  that fix given  $\alpha \in \Omega$ .

### Example (Adding time)

$\mathbb{Z}$ -orbit of 11:00 is  $\Omega = \{12:00, \dots, 11:00\}$  (e.g.  $(11:00)^{-2} = 9:00$ ).

$\mathbb{Z}$ -stabiliser of 11:00 is  $12\mathbb{Z} = \{12k : k \in \mathbb{Z}\}$  (add multiples of 12 hrs).

## Group actions ii

### Definition (orbit)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$ .

*Idea:* states  $\alpha^g \in \Omega$  reachable from fixed  $\alpha \in \Omega$  by moves  $g \in G$ .

### Definition (stabiliser)

If  $G$  acts on  $\Omega$ , then **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Idea:* moves  $g \in G$  that fix given  $\alpha \in \Omega$ .

### Example (Adding time)

$\mathbb{Z}$ -orbit of 11:00 is  $\Omega = \{12:00, \dots, 11:00\}$  (e.g.  $(11:00)^{-2} = 9:00$ ).

$\mathbb{Z}$ -stabiliser of 11:00 is  $12\mathbb{Z} = \{12k : k \in \mathbb{Z}\}$  (add multiples of 12 hrs).

### Example (right regular action)

$G$  acts on  $\Omega = G$  via  $\alpha^g = \alpha g$  for  $\alpha, g \in G$ . Orbit of  $\alpha \in G$  is

## Group actions ii

### Definition (orbit)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$ .

*Idea:* states  $\alpha^g \in \Omega$  reachable from fixed  $\alpha \in \Omega$  by moves  $g \in G$ .

### Definition (stabiliser)

If  $G$  acts on  $\Omega$ , then **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Idea:* moves  $g \in G$  that fix given  $\alpha \in \Omega$ .

### Example (Adding time)

$\mathbb{Z}$ -orbit of 11:00 is  $\Omega = \{12:00, \dots, 11:00\}$  (e.g.  $(11:00)^{-2} = 9:00$ ).

$\mathbb{Z}$ -stabiliser of 11:00 is  $12\mathbb{Z} = \{12k : k \in \mathbb{Z}\}$  (add multiples of 12 hrs).

### Example (right regular action)

$G$  acts on  $\Omega = G$  via  $\alpha^g = \alpha g$  for  $\alpha, g \in G$ . Orbit of  $\alpha \in G$  is  $\Omega = G$  ( $\alpha^{\alpha^{-1}\beta} = \beta \in G$ ); stabiliser of  $\alpha$  is

## Group actions ii

### Definition (orbit)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$ .

*Idea:* states  $\alpha^g \in \Omega$  reachable from fixed  $\alpha \in \Omega$  by moves  $g \in G$ .

### Definition (stabiliser)

If  $G$  acts on  $\Omega$ , then **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Idea:* moves  $g \in G$  that fix given  $\alpha \in \Omega$ .

### Example (Adding time)

$\mathbb{Z}$ -orbit of 11:00 is  $\Omega = \{12:00, \dots, 11:00\}$  (e.g.  $(11:00)^{-2} = 9:00$ ).

$\mathbb{Z}$ -stabiliser of 11:00 is  $12\mathbb{Z} = \{12k : k \in \mathbb{Z}\}$  (add multiples of 12 hrs).

### Example (right regular action)

$G$  acts on  $\Omega = G$  via  $\alpha^g = \alpha g$  for  $\alpha, g \in G$ . Orbit of  $\alpha \in G$  is  $\Omega = G$  ( $\alpha^{\alpha^{-1}\beta} = \beta \in G$ ); stabiliser of  $\alpha$  is  $\{1\} = 1$  ( $\alpha g = \alpha \implies g = 1$ ).

### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is

### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is  $1^G = \{1, 2, 4\}$ , stabiliser of 1 is

### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is  $1^G = \{1, 2, 4\}$ , stabiliser of 1 is  $G_1 = \{()\} = 1$ . Orbit of 3 is



### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is  $1^G = \{1, 2, 4\}$ , stabiliser of 1 is  $G_1 = \{()\} = 1$ . Orbit of 3 is  $3^G = \{3\}$ , stabiliser of 3 is

### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is  $1^G = \{1, 2, 4\}$ , stabiliser of 1 is  $G_1 = \{()\} = 1$ . Orbit of 3 is  $3^G = \{3\}$ , stabiliser of 3 is  $G_3 = G$ .

### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is  $1^G = \{1, 2, 4\}$ , stabiliser of 1 is  $G_1 = \{()\} = 1$ . Orbit of 3 is  $3^G = \{3\}$ , stabiliser of 3 is  $G_3 = G$ .

*Note:*  $|1^G||G_1| = 3 \cdot 1 = 3$

### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is  $1^G = \{1, 2, 4\}$ , stabiliser of 1 is  $G_1 = \{()\} = 1$ . Orbit of 3 is  $3^G = \{3\}$ , stabiliser of 3 is  $G_3 = G$ .

*Note:*  $|1^G||G_1| = 3 \cdot 1 = 3 = |G|$ ,

### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is  $1^G = \{1, 2, 4\}$ , stabiliser of 1 is  $G_1 = \{()\} = 1$ . Orbit of 3 is  $3^G = \{3\}$ , stabiliser of 3 is  $G_3 = G$ .

*Note:*  $|1^G||G_1| = 3 \cdot 1 = 3 = |G|$ ,  $|3^G||G_3| = 1 \cdot 3 = 3 = |G|$ .

### Definition (orbit, stabiliser)

If  $G$  acts on  $\Omega$ , then **orbit** of  $\alpha \in \Omega$  is  $\alpha^G := \{\alpha^g : g \in G\}$  and **stabiliser** of  $\alpha \in \Omega$  is  $G_\alpha := \{g \in G : \alpha^g = \alpha\}$ .

*Note:* stabiliser  $G_\alpha$  is subgroup of  $G$ . (So  $G_\alpha$  acts on  $\Omega$ .)

### Example (Natural action)

$G = \{(), (1, 2, 4), (1, 4, 2)\} \leq \text{Sym}(4)$  acts on  $\Omega = [4]$  naturally.

Orbit of 1 is  $1^G = \{1, 2, 4\}$ , stabiliser of 1 is  $G_1 = \{()\} = 1$ . Orbit of 3 is  $3^G = \{3\}$ , stabiliser of 3 is  $G_3 = G$ .

*Note:*  $|1^G||G_1| = 3 \cdot 1 = 3 = |G|$ ,  $|3^G||G_3| = 1 \cdot 3 = 3 = |G|$ .

### Theorem (orbit-stabiliser)


*If  $G$  acts on  $\Omega$ , then for  $\alpha \in \Omega$ ,  $|\alpha^G||G_\alpha| = |G|$ .*

# The Rubik's group

---

## Representing the cube and its moves i

A Rubik's cube has 6 large faces (each with  $3 \times 3$  smaller faces).





## Representing the cube and its moves i

A Rubik's cube has 6 large faces (each with  $3 \times 3$  smaller faces).


In **solved state 1**, label smaller faces (except each centre) using [48]:

			1	2	3							
			4	<i>U</i>	5							
			6	7	8							
9	10	11	17	18	19	25	26	27	33	34	35	
12	<i>L</i>	13	20	<i>F</i>	21	28	<i>R</i>	29	36	<i>B</i>	37	
14	15	16	22	23	24	30	31	32	38	39	40	
			41	42	43							
			44	<i>D</i>	45							
			46	47	48							

## Representing the cube and its moves i

A Rubik's cube has 6 large faces (each with  $3 \times 3$  smaller faces).

In **solved state 1**, label smaller faces (except each centre) using [48]:



**6 generators** (moves in CC): *U, L, F, R, B, D* (rot. *clockwise*).


## Representing the cube and its moves ii

From *solved state 1*, consider  $F$  which rotates front face clockwise:

			1	2	3							
			4	$U$	5							
			6	7	8							
9	10	11	17	18	19	25	26	27	33	34	35	
12	$L$	13	20	$F$	21	28	$R$	29	36	$B$	37	
14	15	16	22	23	24	30	31	32	38	39	40	
			41	42	43							
			44	$D$	45							
			46	47	48							

## Representing the cube and its moves ii


From *solved state 1*, consider  $F$  which rotates front face clockwise:



Under  $F$ :  $17 \mapsto 19 \mapsto 24 \mapsto 22 \mapsto 17$ ,  $18 \mapsto 21 \mapsto 23 \mapsto 20 \mapsto 18$ ,  $6 \mapsto 25 \mapsto 43 \mapsto 16 \mapsto 6$ ,  $7 \mapsto 28 \mapsto 42 \mapsto 13 \mapsto 7$ ,  $8 \mapsto 30 \mapsto 41 \mapsto 11 \mapsto 8$ , else fixed. So

## Representing the cube and its moves ii

From *solved state 1*, consider  $F$  which rotates front face clockwise:



Under  $F$ :  $17 \mapsto 19 \mapsto 24 \mapsto 22 \mapsto 17$ ,  $18 \mapsto 21 \mapsto 23 \mapsto 20 \mapsto 18$ ,  $6 \mapsto 25 \mapsto 43 \mapsto 16 \mapsto 6$ ,  $7 \mapsto 28 \mapsto 42 \mapsto 13 \mapsto 7$ ,  $8 \mapsto 30 \mapsto 41 \mapsto 11 \mapsto 8$ , else fixed. So

$$F = (17, 19, 24, 22)(18, 21, 23, 20)(6, 25, 43, 16)(7, 28, 42, 13)(8, 30, 41, 11) \in \text{Sym}(48).$$

## Representing the cube and its moves iii

Generators as permutations of labels [48]:

- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$
- $L = (9, 11, 16, 14)(10, 13, 15, 12)(1, 17, 41, 40)(4, 20, 44, 37)(6, 22, 46, 35)$
- $F = (17, 19, 24, 22)(18, 21, 23, 20)(6, 25, 43, 16)(7, 28, 42, 13)(8, 30, 41, 11)$
- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $B = (33, 35, 40, 38)(34, 37, 39, 36)(3, 9, 46, 32)(2, 12, 47, 29)(1, 14, 48, 27)$
- $D = (41, 43, 48, 46)(42, 45, 47, 44)(14, 22, 30, 38)(15, 23, 31, 39)(16, 24, 32, 40)$

## Representing the cube and its moves iii

Generators as permutations of labels [48]:

- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$
- $L = (9, 11, 16, 14)(10, 13, 15, 12)(1, 17, 41, 40)(4, 20, 44, 37)(6, 22, 46, 35)$
- $F = (17, 19, 24, 22)(18, 21, 23, 20)(6, 25, 43, 16)(7, 28, 42, 13)(8, 30, 41, 11)$
- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $B = (33, 35, 40, 38)(34, 37, 39, 36)(3, 9, 46, 32)(2, 12, 47, 29)(1, 14, 48, 27)$
- $D = (41, 43, 48, 46)(42, 45, 47, 44)(14, 22, 30, 38)(15, 23, 31, 39)(16, 24, 32, 40)$

**(Valid) move** is sequence of generators and inverses. E.g.

$$RUR^{-1}U^{-1},$$

## Representing the cube and its moves iii

Generators as permutations of labels [48]:

- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$
- $L = (9, 11, 16, 14)(10, 13, 15, 12)(1, 17, 41, 40)(4, 20, 44, 37)(6, 22, 46, 35)$
- $F = (17, 19, 24, 22)(18, 21, 23, 20)(6, 25, 43, 16)(7, 28, 42, 13)(8, 30, 41, 11)$
- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $B = (33, 35, 40, 38)(34, 37, 39, 36)(3, 9, 46, 32)(2, 12, 47, 29)(1, 14, 48, 27)$
- $D = (41, 43, 48, 46)(42, 45, 47, 44)(14, 22, 30, 38)(15, 23, 31, 39)(16, 24, 32, 40)$

**(Valid) move** is sequence of generators and inverses. E.g.

$$RUR^{-1}U^{-1}, URU^{-1}L^{-1}UR^{-1}U^{-1}L,$$



## Representing the cube and its moves iii

Generators as permutations of labels [48]:

- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$
- $L = (9, 11, 16, 14)(10, 13, 15, 12)(1, 17, 41, 40)(4, 20, 44, 37)(6, 22, 46, 35)$
- $F = (17, 19, 24, 22)(18, 21, 23, 20)(6, 25, 43, 16)(7, 28, 42, 13)(8, 30, 41, 11)$
- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $B = (33, 35, 40, 38)(34, 37, 39, 36)(3, 9, 46, 32)(2, 12, 47, 29)(1, 14, 48, 27)$
- $D = (41, 43, 48, 46)(42, 45, 47, 44)(14, 22, 30, 38)(15, 23, 31, 39)(16, 24, 32, 40)$

**(Valid) move** is sequence of generators and inverses. E.g.

$RUR^{-1}U^{-1}, URU^{-1}L^{-1}UR^{-1}U^{-1}L, RUR^{-1}URU^2R^{-1}U^2.$

**Empty move** is  $1 = ()$  (valid:  $1 = RR^{-1}$ ).

## Representing the cube and its moves iii

Generators as permutations of labels [48]:

- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$
- $L = (9, 11, 16, 14)(10, 13, 15, 12)(1, 17, 41, 40)(4, 20, 44, 37)(6, 22, 46, 35)$
- $F = (17, 19, 24, 22)(18, 21, 23, 20)(6, 25, 43, 16)(7, 28, 42, 13)(8, 30, 41, 11)$
- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $B = (33, 35, 40, 38)(34, 37, 39, 36)(3, 9, 46, 32)(2, 12, 47, 29)(1, 14, 48, 27)$
- $D = (41, 43, 48, 46)(42, 45, 47, 44)(14, 22, 30, 38)(15, 23, 31, 39)(16, 24, 32, 40)$

**(Valid) move** is sequence of generators and inverses. E.g.

$RUR^{-1}U^{-1}, URU^{-1}L^{-1}UR^{-1}U^{-1}L, RUR^{-1}URU^2R^{-1}U^2.$

**Empty move** is  $1 = ()$  (valid:  $1 = RR^{-1}$ ).

**Solving** is applying valid move to get to solved state 1.

## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

*Recall:*  $\sigma = \tau$  in  $\text{Sym}(n)$  iff  $i^\sigma = i^\tau$  for all  $i \in [n]$ .

## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

*Recall:*  $\sigma = \tau$  in  $\text{Sym}(n)$  iff  $i^\sigma = i^\tau$  for all  $i \in [n]$ .

Moves *don't generally commute*:  $RU \neq UR$  since

- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$

$$19^{RU} =$$

## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

*Recall:*  $\sigma = \tau$  in  $\text{Sym}(n)$  iff  $i^\sigma = i^\tau$  for all  $i \in [n]$ .

*Moves don't generally commute:*  $RU \neq UR$  since

- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$

$$19^{RU} = (19^R)^U =$$

## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

*Recall:*  $\sigma = \tau$  in  $\text{Sym}(n)$  iff  $i^\sigma = i^\tau$  for all  $i \in [n]$ .

*Moves don't generally commute:*  $RU \neq UR$  since

- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$

$$19^{RU} = (19^R)^U = 3^U =$$

## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

*Recall:*  $\sigma = \tau$  in  $\text{Sym}(n)$  iff  $i^\sigma = i^\tau$  for all  $i \in [n]$ .

*Moves don't generally commute:*  $RU \neq UR$  since

- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$

$$19^{RU} = (19^R)^U = 3^U = 8 \quad \text{but} \quad 19^{UR} =$$



## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

*Recall:*  $\sigma = \tau$  in  $\text{Sym}(n)$  iff  $i^\sigma = i^\tau$  for all  $i \in [n]$ .

Moves *don't generally commute*:  $RU \neq UR$  since

- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$

$$19^{RU} = (19^R)^U = 3^U = 8 \quad \text{but} \quad 19^{UR} = (19^U)^R =$$

## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

*Recall:*  $\sigma = \tau$  in  $\text{Sym}(n)$  iff  $i^\sigma = i^\tau$  for all  $i \in [n]$ .

Moves *don't generally commute*:  $RU \neq UR$  since

- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$

$$19^{RU} = (19^R)^U = 3^U = 8 \quad \text{but} \quad 19^{UR} = (19^U)^R = 11^R =$$

## Representing the cube and its moves iv

*In cubing community:* moves called *move sequences*. Generators called *moves*. *Inverse elementary moves* written  $U', L', F', R', B', D'$  (instead of  $U^{-1}$  etc.); powers written  $U2, R2$  etc. (instead of  $U^2, R^2$ ).

*Recall:*  $\sigma = \tau$  in  $\text{Sym}(n)$  iff  $i^\sigma = i^\tau$  for all  $i \in [n]$ .

*Moves don't generally commute:*  $RU \neq UR$  since

- $R = (25, 27, 32, 30)(26, 29, 31, 28)(3, 38, 43, 19)(5, 36, 45, 21)(8, 33, 48, 24)$
- $U = (1, 3, 8, 6)(2, 5, 7, 4)(9, 33, 25, 17)(10, 34, 26, 18)(11, 35, 27, 19)$

$$19^{RU} = (19^R)^U = 3^U = 8 \quad \text{but} \quad 19^{UR} = (19^U)^R = 11^R = 11.$$

## Moves vs states for Rubik's cube i

(Valid) state is result of applying *valid move* to *solved state* 1.

			1	2	3							
			4	<i>U</i>	5							
			6	7	8							
9	10	11	17	18	19	25	26	27	33	34	35	
12	<i>L</i>	13	20	<i>F</i>	21	28	<i>R</i>	29	36	<i>B</i>	37	
14	15	16	22	23	24	30	31	32	38	39	40	
			41	42	43							
			44	<i>D</i>	45							
			46	47	48							

## Moves vs states for Rubik's cube i

(Valid) state is result of applying *valid move* to *solved state* 1.

			1	2	3								
			4	<i>U</i>	5								
			16	13	11								
9	10	41	22	20	17	6	26	27	33	34	35		
12	<i>L</i>	42	23	<i>F</i>	18	7	<i>R</i>	29	36	<i>B</i>	37		
14	15	43	24	21	19	8	31	32	38	39	40		
			30	28	25								
			44	<i>D</i>	45								
			46	47	48								

This new state is valid, as result of applying *F* to solved state.

## Moves vs states for Rubik's cube ii

*Restickering* is valid state iff it can be *solved*. How to check?

Let  $\mathcal{S}$  be valid **states**; let state  $x \in \mathcal{S}$  be element of  $\text{Sym}(48)$  giving permutation of labels to solved state  $1 \in \mathcal{S}$ .

(I.e.  $i^x$  is label at  $x$ -position of  $i$  in solved state 1.)

## Moves vs states for Rubik's cube ii

*Restickering* is valid state iff it can be *solved*. How to check?

Let  $\mathcal{S}$  be valid **states**; let state  $x \in \mathcal{S}$  be element of  $\text{Sym}(48)$  giving permutation of labels to solved state  $1 \in \mathcal{S}$ .

(I.e.  $i^x$  is label at  $x$ -position of  $i$  in solved state 1.)

Let  $\mathcal{G}$  be valid **moves**; let move  $\sigma \in \mathcal{G}$  be element of  $\text{Sym}(48)$  giving corresponding permutation of labels.

(I.e.  $i^\sigma$  is label at position  $\sigma$  maps  $i$  into.)

## Moves vs states for Rubik's cube ii

*Restickering* is valid state iff it can be *solved*. How to check?

Let  $\mathcal{S}$  be valid **states**; let state  $x \in \mathcal{S}$  be element of  $\text{Sym}(48)$  giving permutation of labels to solved state  $1 \in \mathcal{S}$ .

(I.e.  $i^x$  is label at  $x$ -position of  $i$  in solved state 1.)

Let  $\mathcal{G}$  be valid **moves**; let move  $\sigma \in \mathcal{G}$  be element of  $\text{Sym}(48)$  giving corresponding permutation of labels.

(I.e.  $i^\sigma$  is label at position  $\sigma$  maps  $i$  into.)

- State  $x \in \mathcal{S}$  corresponds to move  $x \in \mathcal{G}$  required to get solved state 1 into state  $x$ .



## Moves vs states for Rubik's cube ii

*Restickering* is valid state iff it can be *solved*. How to check?

Let  $\mathcal{S}$  be valid **states**; let state  $x \in \mathcal{S}$  be element of  $\text{Sym}(48)$  giving permutation of labels to solved state  $1 \in \mathcal{S}$ .

(I.e.  $i^x$  is label at  $x$ -position of  $i$  in solved state 1.)

Let  $\mathcal{G}$  be valid **moves**; let move  $\sigma \in \mathcal{G}$  be element of  $\text{Sym}(48)$  giving corresponding permutation of labels.

(I.e.  $i^\sigma$  is label at position  $\sigma$  maps  $i$  into.)

- State  $x \in \mathcal{S}$  corresponds to move  $x \in \mathcal{G}$  required to get solved state 1 into state  $x$ .
- Move  $\sigma \in \mathcal{G}$  corresponds to state  $\sigma \in \mathcal{S}$  reached by applying move  $\sigma$  to solved state 1.

## Moves vs states for Rubik's cube ii

*Restickering* is valid state iff it can be *solved*. How to check?

Let  $\mathcal{S}$  be valid **states**; let state  $x \in \mathcal{S}$  be element of  $\text{Sym}(48)$  giving permutation of labels to solved state  $1 \in \mathcal{S}$ .

(I.e.  $i^x$  is label at  $x$ -position of  $i$  in solved state 1.)

Let  $\mathcal{G}$  be valid **moves**; let move  $\sigma \in \mathcal{G}$  be element of  $\text{Sym}(48)$  giving corresponding permutation of labels.

(I.e.  $i^\sigma$  is label at position  $\sigma$  maps  $i$  into.)

- State  $x \in \mathcal{S}$  corresponds to move  $x \in \mathcal{G}$  required to get solved state 1 into state  $x$ .
- Move  $\sigma \in \mathcal{G}$  corresponds to state  $\sigma \in \mathcal{S}$  reached by applying move  $\sigma$  to solved state 1.

So moves  $\leftrightarrow$  states; as sets,  $\mathcal{S} = \mathcal{G}$ . *Solved state* is  $1 = () \in \text{Sym}(48)$ .

# The Rubik's group of permutations i

Set of moves  $\mathcal{G}$  forms group: composition of valid moves

# The Rubik's group of permutations i

Set of moves  $\mathcal{G}$  forms group: composition of valid moves is valid move; identity move is

# The Rubik's group of permutations i

Set of moves  $\mathcal{G}$  forms group: composition of valid moves is valid move; identity move is  $1 = () \in \mathcal{G}$ , inverse moves

# The Rubik's group of permutations i

Set of moves  $\mathcal{G}$  forms group: composition of valid moves is valid move; identity move is  $1 = () \in \mathcal{G}$ , inverse moves exist (undo elementary moves/inverses); associative.

Set of moves  $\mathcal{G}$  forms group: composition of valid moves is valid move; identity move is  $1 = () \in \mathcal{G}$ , inverse moves exist (undo elementary moves/inverses); associative.

**Definition (Rubik's group)**

$\mathcal{G} \leq \text{Sym}(48)$  is permutation group of degree 48, called the **Rubik's group**; it acts naturally on  $[48]$ .

Set of moves  $\mathcal{G}$  forms group: composition of valid moves is valid move; identity move is  $1 = () \in \mathcal{G}$ , inverse moves exist (undo elementary moves/inverses); associative.

**Definition (Rubik's group)**

$\mathcal{G} \leq \text{Sym}(48)$  is permutation group of degree 48, called the **Rubik's group**; it acts naturally on  $[48]$ . *Note:*  $G = \langle U, L, F, R, B, D \rangle$ .



# The Rubik's group of permutations i

Set of moves  $\mathcal{G}$  forms group: composition of valid moves is valid move; identity move is  $1 = () \in \mathcal{G}$ , inverse moves exist (undo elementary moves/inverses); associative.

## Definition (Rubik's group)

$\mathcal{G} \leq \text{Sym}(48)$  is permutation group of degree 48, called the **Rubik's group**; it acts naturally on  $[48]$ . *Note:*  $G = \langle U, L, F, R, B, D \rangle$ .

For move  $\sigma \in \mathcal{G}$  and state  $x \in \mathcal{S}$ , applying  $\sigma$  to  $x$  gives state  $x^\sigma = x\sigma \in \mathcal{S}$ . This is *regular action* of  $\mathcal{G}$ . (Consider states  $x \in \mathcal{G}$ .)

# The Rubik's group of permutations i

Set of moves  $\mathcal{G}$  forms group: composition of valid moves is valid move; identity move is  $1 = () \in \mathcal{G}$ , inverse moves exist (undo elementary moves/inverses); associative.

## Definition (Rubik's group)

$\mathcal{G} \leq \text{Sym}(48)$  is permutation group of degree 48, called the **Rubik's group**; it acts naturally on  $[48]$ . *Note:*  $G = \langle U, L, F, R, B, D \rangle$ .

For move  $\sigma \in \mathcal{G}$  and state  $x \in \mathcal{S}$ , applying  $\sigma$  to  $x$  gives state  $x^\sigma = x\sigma \in \mathcal{S}$ . This is *regular action* of  $\mathcal{G}$ . (Consider states  $x \in \mathcal{G}$ .)

Clearly  $\mathcal{G}$  finite (states  $\leftrightarrow$  moves; also  $|\mathcal{G}| \leq 48!$ ). But what is  $|\mathcal{G}|$ ?

## The Rubik's group of permutations ii

GAP code to define generators and  $\mathcal{G} = \langle U, L, F, R, B, D \rangle$  (as G):

```
1 U := ( 1, 3, 8, 6)( 2, 5, 7, 4)( 9,33,25,17)(10,34,26,18)
      (11,35,27,19);
2 L := ( 9,11,16,14)(10,13,15,12)( 1,17,41,40)( 4,20,44,37)(
      6,22,46,35);
3 F := (17,19,24,22)(18,21,23,20)( 6,25,43,16)( 7,28,42,13)(
      8,30,41,11);
4 R := (25,27,32,30)(26,29,31,28)( 3,38,43,19)( 5,36,45,21)(
      8,33,48,24);
5 B := (33,35,40,38)(34,37,39,36)( 3, 9,46,32)( 2,12,47,29)(
      1,14,48,27);
6 D := (41,43,48,46)(42,45,47,44)(14,22,30,38)(15,23,31,39)
      (16,24,32,40);
7 G := Group( U, L, F, R, B, D );
```

## The Rubik's group of permutations ii

GAP code to define generators and  $\mathcal{G} = \langle U, L, F, R, B, D \rangle$  (as G):

```
1 U := ( 1, 3, 8, 6)( 2, 5, 7, 4)( 9,33,25,17)(10,34,26,18)
      (11,35,27,19);
2 L := ( 9,11,16,14)(10,13,15,12)( 1,17,41,40)( 4,20,44,37)(
      6,22,46,35);
3 F := (17,19,24,22)(18,21,23,20)( 6,25,43,16)( 7,28,42,13)(
      8,30,41,11);
4 R := (25,27,32,30)(26,29,31,28)( 3,38,43,19)( 5,36,45,21)(
      8,33,48,24);
5 B := (33,35,40,38)(34,37,39,36)( 3, 9,46,32)( 2,12,47,29)(
      1,14,48,27);
6 D := (41,43,48,46)(42,45,47,44)(14,22,30,38)(15,23,31,39)
      (16,24,32,40);
7 G := Group( U, L, F, R, B, D );
```

Order cmd:  $|\mathcal{G}| = 43\,252\,003\,274\,489\,856\,000 \approx 4.3 \cdot 10^{19}$ . *How?*

# Orbits and stabilisers i

			1	2	3						
			4	<i>U</i>	5						
			6	7	8						
9	10	11	17	18	19	25	26	27	33	34	35
12	<i>L</i>	13	20	<i>F</i>	21	28	<i>R</i>	29	36	<i>B</i>	37
14	15	16	22	23	24	30	31	32	38	39	40
			41	42	43						
			44	<i>D</i>	45						
			46	47	48						

```
1 gap> Orbit( G, 1 );
2 [ 1, 6, 40, 27, 8, 35, 16, 41, 32, 25, 48, 3, 11, 24, 46, 33, 43, 17,
   30, 14, 19, 9, 22, 38 ]
3 gap> Orbit( G, 2 );
4 [ 2, 5, 12, 7, 36, 10, 47, 4, 28, 45, 34, 13, 29, 44, 20, 42, 26, 21,
   37, 15, 31, 18, 23, 39 ]
```


Two  $\mathcal{G}$ -orbits: corner pieces  $1^{\mathcal{G}}$ , edge pieces  $2^{\mathcal{G}}$ .

## Orbits and stabilisers i

1	2	3	4	5	6	7	8	9	10
4	U	5	16	17	18	19	20	21	22
6	7	8	25	26	27	28	R	29	30
9	10	11	33	34	35	36	B	37	38
12	L	13	40	41	42	43	44	45	46
14	15	16	49	50	51	52	53	54	55
17	18	19	58	59	60	61	62	63	64
20	21	22	67	68	69	70	71	72	73
23	24	25	76	77	78	79	80	81	82
26	27	28	85	86	87	88	89	90	91
29	30	31	94	95	96	97	98	99	100

Moves in  $\mathcal{H} = \mathcal{G}_{1,3,6,8} = (((\mathcal{G}_1)_3)_6)_8$  fix white corners 1, 3, 6, 8.

# Orbits and stabilisers i



Moves in  $\mathcal{H} = \mathcal{G}_{1,3,6,8} = (((\mathcal{G}_1)_3)_6)_8$  fix white corners 1, 3, 6, 8.

```

1 gap> G_1368 := Stabilizer( G, [ 1, 3, 6, 8 ], OnTuples );
2 <permutation group of size 3178424696832000 with 12 generators>
3 gap> Orbit( G_1368, 17 );
4 [ 17 ]
5 gap> Orbit( G_1368, 24 );
6 [ 24, 30, 43, 32, 38, 46, 48, 40, 14, 41, 16, 22 ]
7 gap> Set( Orbit( G_1368, 2 ) ) = Set( Orbit( G, 2 ) );
8 true

```

Some  $\mathcal{H}$ -orbits:  $17^{\mathcal{H}} = \{17\}$ , bottom corner pieces  $24^{\mathcal{H}}$ , edge pieces  $2^{\mathcal{H}} = 2^{\mathcal{G}}$ .

## Orders of moves i

Use GAP to compute products, order (using Order cmd).

```
1 gap> R*U*R^(-1)*U^(-1);  
2 (1,27,35,33,9,3)(2,21,5)(8,30,25,43,19,24)(26,34,28)  
3 gap> Order( last );  
4 6
```

How many times must we repeat move  $\sigma \in \mathcal{G}$  to have no effect?



## Orders of moves i

Use GAP to compute products, order (using Order cmd).

```
1 gap> R*U*R^(-1)*U^(-1);  
2 (1,27,35,33,9,3)(2,21,5)(8,30,25,43,19,24)(26,34,28)  
3 gap> Order( last );  
4 6
```

How many times must we repeat move  $\sigma \in \mathcal{G}$  to have no effect? I.e. for state  $x \in \mathcal{S}$ , smallest  $k \in \mathbb{Z}_+$  with  $x\sigma^k = x\sigma^k = x \iff \sigma^k = 1$ .

## Orders of moves i

Use GAP to compute products, order (using Order cmd).

```
1 gap> R*U*R^(-1)*U^(-1);  
2 (1,27,35,33,9,3)(2,21,5)(8,30,25,43,19,24)(26,34,28)  
3 gap> Order( last );  
4 6
```

How many times must we repeat move  $\sigma \in \mathcal{G}$  to have no effect? I.e. for state  $x \in \mathcal{S}$ , smallest  $k \in \mathbb{Z}_+$  with  $x\sigma^k = x\sigma^k = x \iff \sigma^k = 1$ .

*Recall:* order of  $\sigma$  is lcm of cycle lengths.

- Any generator ( $U, L, F, R, B, D$ ) has cycles of length 4, 4, 4, 4, 4:  
order is  $\text{lcm}(4, 4, 4, 4, 4) = 4$ .

## Orders of moves i

Use GAP to compute products, order (using Order cmd).

```
1 gap> R*U*R^(-1)*U^(-1);  
2 (1,27,35,33,9,3)(2,21,5)(8,30,25,43,19,24)(26,34,28)  
3 gap> Order( last );  
4 6
```

How many times must we repeat move  $\sigma \in \mathcal{G}$  to have no effect? I.e. for state  $x \in \mathcal{S}$ , smallest  $k \in \mathbb{Z}_+$  with  $x\sigma^k = x\sigma^k = x \iff \sigma^k = 1$ .

*Recall:* order of  $\sigma$  is lcm of cycle lengths.

- Any generator ( $U, L, F, R, B, D$ ) has cycles of length 4, 4, 4, 4, 4: order is  $\text{lcm}(4, 4, 4, 4, 4) = 4$ .
- Commutator  $RUR^{-1}U^{-1}$   
 $= (1, 27, 35, 33, 9, 3)(2, 21, 5)(8, 30, 25, 43, 19, 24)(26, 34, 28):$   
order is  $\text{lcm}(6, 3, 6, 3) = 6$ .

- *Sune*  $RUR^{-1}URU^2R^{-1}U^2$

$$= (1, 9, 35)(2, 5, 7)(3, 33, 27)(8, 25, 19)(18, 34, 26):$$

order is  $\text{lcm}(3, 3, 3, 3, 3) = 3$ .

- *Sune*  $RUR^{-1}URU^2R^{-1}U^2$

$$= (1, 9, 35)(2, 5, 7)(3, 33, 27)(8, 25, 19)(18, 34, 26):$$

order is  $\text{lcm}(3, 3, 3, 3, 3) = 3$ .

- *Lawrence's move*  $RU$  has cycles of length 15, 7, 3, 7: order is  $\text{lcm}(15, 7, 3, 7) = 105$ .

- *Sune*  $RUR^{-1}URU^2R^{-1}U^2$

$$= (1, 9, 35)(2, 5, 7)(3, 33, 27)(8, 25, 19)(18, 34, 26):$$

order is  $\text{lcm}(3, 3, 3, 3, 3) = 3$ .

- *Lawrence's move*  $RU$  has cycles of length 15, 7, 3, 7: order is  $\text{lcm}(15, 7, 3, 7) = 105$ .
- *Clayton's move*  $UL'$  has cycles of length 9, 7, 9, 7: order is  $\text{lcm}(9, 7, 9, 7) = 63$ .

Watch video demonstration by my friend Wes :D

- *Sune*  $RUR^{-1}URU^2R^{-1}U^2$

$$= (1, 9, 35)(2, 5, 7)(3, 33, 27)(8, 25, 19)(18, 34, 26):$$

order is  $\text{lcm}(3, 3, 3, 3, 3) = 3$ .

- *Lawrence's move*  $RU$  has cycles of length 15, 7, 3, 7: order is  $\text{lcm}(15, 7, 3, 7) = 105$ .
- *Clayton's move*  $UL'$  has cycles of length 9, 7, 9, 7: order is  $\text{lcm}(9, 7, 9, 7) = 63$ .

Watch video demonstration by my friend Wes :D

Element of order 5?

## Orders of moves ii

- *Sune*  $RUR^{-1}URU^2R^{-1}U^2$

$$= (1, 9, 35)(2, 5, 7)(3, 33, 27)(8, 25, 19)(18, 34, 26):$$

order is  $\text{lcm}(3, 3, 3, 3, 3) = 3$ .

- *Lawrence's move*  $RU$  has cycles of length 15, 7, 3, 7: order is  $\text{lcm}(15, 7, 3, 7) = 105$ .
- *Clayton's move*  $UL'$  has cycles of length 9, 7, 9, 7: order is  $\text{lcm}(9, 7, 9, 7) = 63$ .

Watch video demonstration by my friend Wes :D

Element of order 5? *Answer:*  $(RU)^{21}$  since  $((RU)^{21})^5 = (RU)^{105} = 1$ .



### **Theorem (Jake Vandenberg's conjecture)**

*There is no Rubik's cube move that cycles through all states.*

### **Theorem (Jake Vandenberg's conjecture)**

*There is no Rubik's cube move that cycles through all states.*

*Recall:* states  $\leftrightarrow$  moves. Rubik's group  $\mathcal{G}$  acts on states by applying move  $\sigma \in \mathcal{G}$  to state  $x \in \mathcal{G}$  to get state  $x^\sigma = x\sigma \in \mathcal{G}$ .

### Theorem (Jake Vandenberg's conjecture)

*There is no Rubik's cube move that cycles through all states.*

*Recall:* states  $\leftrightarrow$  moves. Rubik's group  $\mathcal{G}$  acts on states by applying move  $\sigma \in \mathcal{G}$  to state  $x \in \mathcal{G}$  to get state  $x^\sigma = x\sigma \in \mathcal{G}$ .

*Equivalent question:* for starting state, WLOG  $1 = ()$ , is there  $\sigma \in \mathcal{G}$  with  $\{1^{\sigma^k} : k \in \mathbb{Z}\} = \{1\sigma^k : k \in \mathbb{Z}\} = \{\sigma^k : k \in \mathbb{Z}\} = \mathcal{G}$ ?

### **Theorem (Jake Vandenberg's conjecture)**

*There is no Rubik's cube move that cycles through all states.*

*Recall:* states  $\leftrightarrow$  moves. Rubik's group  $\mathcal{G}$  acts on states by applying move  $\sigma \in \mathcal{G}$  to state  $x \in \mathcal{G}$  to get state  $x^\sigma = x\sigma \in \mathcal{G}$ .

*Equivalent question:* for starting state, WLOG  $1 = ()$ , is there  $\sigma \in \mathcal{G}$  with  $\{1^{\sigma^k} : k \in \mathbb{Z}\} = \{1\sigma^k : k \in \mathbb{Z}\} = \{\sigma^k : k \in \mathbb{Z}\} = \mathcal{G}$ ? In group theory language:

### **Theorem (Jake Vandenberg's conjecture)**

*The Rubik's group  $\mathcal{G}$  is not cyclic. (I.e. no  $\sigma \in \mathcal{G}$  with  $\mathcal{G} = \langle \sigma \rangle$ .)*

### **Theorem (Jake Vandenberg's conjecture)**

*There is no Rubik's cube move that cycles through all states.*

*Recall:* states  $\leftrightarrow$  moves. Rubik's group  $\mathcal{G}$  acts on states by applying move  $\sigma \in \mathcal{G}$  to state  $x \in \mathcal{G}$  to get state  $x^\sigma = x\sigma \in \mathcal{G}$ .

*Equivalent question:* for starting state, WLOG  $1 = ()$ , is there  $\sigma \in \mathcal{G}$  with  $\{1^{\sigma^k} : k \in \mathbb{Z}\} = \{1\sigma^k : k \in \mathbb{Z}\} = \{\sigma^k : k \in \mathbb{Z}\} = \mathcal{G}$ ? In group theory language:

### **Theorem (Jake Vandenberg's conjecture)**

*The Rubik's group  $\mathcal{G}$  is not cyclic. (I.e. no  $\sigma \in \mathcal{G}$  with  $\mathcal{G} = \langle \sigma \rangle$ .)*

### **Proof.**

If  $\mathcal{G}$  is cyclic, then  $\mathcal{G}$  is abelian.

### **Theorem (Jake Vandenberg's conjecture)**

*There is no Rubik's cube move that cycles through all states.*

*Recall:* states  $\leftrightarrow$  moves. Rubik's group  $\mathcal{G}$  acts on states by applying move  $\sigma \in \mathcal{G}$  to state  $x \in \mathcal{G}$  to get state  $x^\sigma = x\sigma \in \mathcal{G}$ .

*Equivalent question:* for starting state, WLOG  $1 = ()$ , is there  $\sigma \in \mathcal{G}$  with  $\{1^{\sigma^k} : k \in \mathbb{Z}\} = \{1\sigma^k : k \in \mathbb{Z}\} = \{\sigma^k : k \in \mathbb{Z}\} = \mathcal{G}$ ? In group theory language:

### **Theorem (Jake Vandenberg's conjecture)**

*The Rubik's group  $\mathcal{G}$  is not cyclic. (I.e. no  $\sigma \in \mathcal{G}$  with  $\mathcal{G} = \langle \sigma \rangle$ .)*

### **Proof.**

If  $\mathcal{G}$  is cyclic, then  $\mathcal{G}$  is abelian. But  $\mathcal{G}$  is not abelian:  $RU \neq UR$ .  $\square$

### **Theorem (Jake Vandenberg's theorem)**

*There is no Rubik's cube move that when repeated, if starting from the solved state, never returns to the solved state.*

### **Theorem (Jake Vandenberg's theorem)**

*There is no Rubik's cube move that when repeated, if starting from the solved state, never returns to the solved state.*

$k$ -fold repetition of move  $\sigma \in G$ , applied to solved state  $1 = ()$ , gives  $1^{\sigma^k} = 1\sigma^k = \sigma^k$ . Returning to solved state:  $\sigma^k = 1$  (for  $k \in \mathbb{Z}_+$ ).



### Theorem (Jake Vandenberg's theorem)

*There is no Rubik's cube move that when repeated, if starting from the solved state, never returns to the solved state.*

$k$ -fold repetition of move  $\sigma \in G$ , applied to solved state  $1 = ()$ , gives  $1^{\sigma^k} = 1\sigma^k = \sigma^k$ . Returning to solved state:  $\sigma^k = 1$  (for  $k \in \mathbb{Z}_+$ ).

*Equivalent question:* does any  $\sigma \in G$  have infinite order?

### **Theorem (Jake Vandenberg's theorem)**

*There is no Rubik's cube move that when repeated, if starting from the solved state, never returns to the solved state.*

$k$ -fold repetition of move  $\sigma \in G$ , applied to solved state  $1 = ()$ , gives  $1^{\sigma^k} = 1\sigma^k = \sigma^k$ . Returning to solved state:  $\sigma^k = 1$  (for  $k \in \mathbb{Z}_+$ ).

*Equivalent question:* does any  $\sigma \in G$  have infinite order?

### **Proposition (corollary of Lagrange's theorem)**

*If  $G$  is finite group and  $g \in G$ , then  $g^{|G|} = 1$ .*

### **Theorem (Jake Vandenberg's theorem)**

*There is no Rubik's cube move that when repeated, if starting from the solved state, never returns to the solved state.*

$k$ -fold repetition of move  $\sigma \in G$ , applied to solved state  $1 = ()$ , gives  $1^{\sigma^k} = 1\sigma^k = \sigma^k$ . Returning to solved state:  $\sigma^k = 1$  (for  $k \in \mathbb{Z}_+$ ).

*Equivalent question: does any  $\sigma \in G$  have infinite order?*

### **Proposition (corollary of Lagrange's theorem)**

*If  $G$  is finite group and  $g \in G$ , then  $g^{|G|} = 1$ .*

### **Corollary (Jake Vandenberg's theorem)**

*There is no  $\sigma \in \mathcal{G}$  with infinite order (since  $\mathcal{G}$  is finite).*

# **Analysing the Rubik's group**

---

### Definition (Base, stabiliser chain)

If  $G \leq \text{Sym}(n)$ , distinct elts  $B = [\beta_1, \dots, \beta_r] \subseteq [n]$  is **base** for  $G$  if  $G_{\beta_1, \dots, \beta_r} = 1$ . (Recall:  $G_{\beta_1, \dots, \beta_r} = \{g \in G : \beta_1^g = \beta_1, \dots, \beta_r^g = \beta_r\}$ .)

### Definition (Base, stabiliser chain)

If  $G \leq \text{Sym}(n)$ , distinct elts  $B = [\beta_1, \dots, \beta_r] \subseteq [n]$  is **base** for  $G$  if  $G_{\beta_1, \dots, \beta_r} = 1$ . (Recall:  $G_{\beta_1, \dots, \beta_r} = \{g \in G : \beta_1^g = \beta_1, \dots, \beta_r^g = \beta_r\}$ .)

Corresponding **stabiliser chain** is

$$G = G^0 \geq G^1 \geq \dots \geq G^r = 1$$

where  $G^i = G_{\beta_i}^{i-1} = G_{\beta_1, \dots, \beta_i}$ .

## Definition (Base, stabiliser chain)

If  $G \leq \text{Sym}(n)$ , distinct elts  $B = [\beta_1, \dots, \beta_r] \subseteq [n]$  is **base** for  $G$  if  $G_{\beta_1, \dots, \beta_r} = 1$ . (Recall:  $G_{\beta_1, \dots, \beta_r} = \{g \in G : \beta_1^g = \beta_1, \dots, \beta_r^g = \beta_r\}$ .)

Corresponding **stabiliser chain** is

$$G = G^0 \geq G^1 \geq \dots \geq G^r = 1$$

where  $G^i = G_{\beta_i}^{i-1} = G_{\beta_1, \dots, \beta_i}$ .

Base  $B$  contains elts of  $[n]$  such that only  $1 \in G$  fixes every  $\beta_i \in B$ .

(Short base desirable: how to compute minimum base?)

## Definition (Base, stabiliser chain)

If  $G \leq \text{Sym}(n)$ , distinct elts  $B = [\beta_1, \dots, \beta_r] \subseteq [n]$  is **base** for  $G$  if  $G_{\beta_1, \dots, \beta_r} = 1$ . (Recall:  $G_{\beta_1, \dots, \beta_r} = \{g \in G : \beta_1^g = \beta_1, \dots, \beta_r^g = \beta_r\}$ .)

Corresponding **stabiliser chain** is

$$G = G^0 \geq G^1 \geq \dots \geq G^r = 1$$

where  $G^i = G_{\beta_i}^{i-1} = G_{\beta_1, \dots, \beta_i}$ .

Base  $B$  contains elts of  $[n]$  such that only  $1 \in G$  fixes every  $\beta_i \in B$ .

(Short base desirable: how to compute minimum base?)

Stabiliser chain can be implemented computationally; useful in algorithms (membership testing, random element generation, factorisation into generators).



### Example (Rubik's group)

Using GAP:

```
1 gap> BaseOfGroup( G );
2 [ 1, 3, 6, 8, 2, 4, 5, 7, 12, 13, 14, 15, 16, 21,
   23, 24, 29, 31 ]
3 gap> Size( last );
4 18
```

### Example (Rubik's group)

Using GAP:

```
1 gap> BaseOfGroup( G );
2 [ 1, 3, 6, 8, 2, 4, 5, 7, 12, 13, 14, 15, 16, 21,
   23, 24, 29, 31 ]
3 gap> Size( last );
4 18
```

Base of  $\mathcal{G}$  of size 18 is

$$B = [1, 3, 6, 8, 2, 4, 5, 7, 12, 13, 14, 15, 16, 21, 23, 24, 29, 31].$$

### Example (Rubik's group)

Using GAP:

```
1 gap> BaseOfGroup( G );  
2 [ 1, 3, 6, 8, 2, 4, 5, 7, 12, 13, 14, 15, 16, 21,  
   23, 24, 29, 31 ]  
3 gap> Size( last );  
4 18
```

Base of  $\mathcal{G}$  of size 18 is

$$B = [1, 3, 6, 8, 2, 4, 5, 7, 12, 13, 14, 15, 16, 21, 23, 24, 29, 31].$$

If move  $\sigma \in \mathcal{G}$  fixes every  $\beta_i \in B$  then  $\sigma = 1$  is empty move.

## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

### Example (rotation group of cube)

Compute order of rotation group  $G \leq \text{Sym}(8)$  for cube:

## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

### Example (rotation group of cube)

Compute order of rotation group  $G \leq \text{Sym}(8)$  for cube: base of adjacent vertices say 1, 2 (once fixed, can't rotate, so  $G_{1,2} = 1$ ).

## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

### Example (rotation group of cube)

Compute order of rotation group  $G \leq \text{Sym}(8)$  for cube: base of adjacent vertices say 1, 2 (once fixed, can't rotate, so  $G_{1,2} = 1$ ).

$|1^G| = 8$  (all vertices);

## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

### Example (rotation group of cube)

Compute order of rotation group  $G \leq \text{Sym}(8)$  for cube: base of adjacent vertices say 1, 2 (once fixed, can't rotate, so  $G_{1,2} = 1$ ).

$|1^G| = 8$  (all vertices); in  $G_1$ ,  $|2^{G_1}| = 3$  (vertices adjacent to 1); so



## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

### Example (rotation group of cube)

Compute order of rotation group  $G \leq \text{Sym}(8)$  for cube: base of adjacent vertices say 1, 2 (once fixed, can't rotate, so  $G_{1,2} = 1$ ).

$|1^G| = 8$  (all vertices); in  $G_1$ ,  $|2^{G_1}| = 3$  (vertices adjacent to 1); so

$$|G| = |1^G| |2^{G_1}| = 8 \cdot 3 = 24.$$

## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

Orbits and stabilisers can be easily computed (e.g. using GAP).

## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

Orbits and stabilisers can be easily computed (e.g. using GAP).

Implementing base and stabiliser chain for Rubik's group  $\mathcal{G}$  (using `BaseOfGroup` and `StabChain` cmds), GAP computes:

## How many valid states are there? i

### Theorem (size of perm group)

If  $B = [\beta_1, \dots, \beta_r]$  is base for  $G \leq \text{Sym}(n)$  with stabiliser chain  $G = G^0 \geq G^1 \geq \dots \geq G^r = 1$ , then

$$|G| = |\beta_1^{G^0}| |\beta_2^{G^1}| \cdots |\beta_r^{G^{r-1}}|.$$

Orbits and stabilisers can be easily computed (e.g. using GAP).

Implementing base and stabiliser chain for Rubik's group  $\mathcal{G}$  (using `BaseOfGroup` and `StabChain` cmds), GAP computes:


### Corollary

$$|\mathcal{G}| = 43\,252\,003\,274\,489\,856\,000 \approx 4.3 \cdot 10^{19}.$$

## Can this restickering be solved? i

### Theorem (Wes's conjecture)


*“I’m 99% sure you can’t swap two [adjacent] edge pieces without affecting another piece?!”*



## Can this restickering be solved? i

### Theorem (Wes's conjecture)

*“I’m 99% sure you can’t swap two [adjacent] edge pieces without affecting another piece?!”*




WLOG consider solved state. *Equivalent question*: does only restickering two adjacent edge pieces give solvable state?

# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*




WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?

By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28).

# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*



WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?


By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28). Four ways: given by where red pieces go (2 options each).



# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*




WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?

By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28). Four ways: given by where red pieces go (2 options each).

# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*




WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?

By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28). Four ways: given by where red pieces go (2 options each).

# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*




WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?

By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28). Four ways: given by where red pieces go (2 options each).

# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*




WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?

By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28). Four ways: given by where red pieces go (2 options each).

# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*




WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?

By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28). Four ways: given by where red pieces go (2 options each).

# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*




WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?

By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28). Four ways: given by where red pieces go (2 options each).

# Can this restickering be solved? i

## Theorem (Wes's conjecture)

*"I'm 99% sure you can't swap two [adjacent] edge pieces without affecting another piece?!"*



WLOG consider solved state. *Equivalent question:* does only restickering two adjacent edge pieces give solvable state?

By symmetry, just check one pair, say red/white (18/7) and red/blue (21/28). Four ways: given by where red pieces go (2 options each).

## Can this restickering be solved? ii

These restickerings should be invalid states.



## Can this restickering be solved? ii

These restickerings should be invalid states. In group theory language:

**Theorem (Wes's conjecture)**

$(18, 21)(7, 28) \notin \mathcal{G}$ ,  $(18, 28)(7, 21) \notin \mathcal{G}$ ,  $(18, 21, 7, 28) \notin \mathcal{G}$ , and  $(18, 28, 7, 21) \notin \mathcal{G}$ .

## Can this restickering be solved? ii

These restickerings should be invalid states. In group theory language:

### Theorem (Wes's conjecture)

$(18, 21)(7, 28) \notin \mathcal{G}$ ,  $(18, 28)(7, 21) \notin \mathcal{G}$ ,  $(18, 21, 7, 28) \notin \mathcal{G}$ , and  $(18, 28, 7, 21) \notin \mathcal{G}$ .

### Proof.

By GAP:

```
1 gap> (18,21)(7,28) in G or (18,28)(7,21) in G or  
      (18,21,7,28) in G or (18,28,7,21) in G;  
2 false
```

(GAP uses stabiliser chains to verify membership!)

□

## Can this restickering be solved? ii

These restickerings should be invalid states. In group theory language:

### Theorem (Wes's conjecture)

$(18, 21)(7, 28) \notin \mathcal{G}$ ,  $(18, 28)(7, 21) \notin \mathcal{G}$ ,  $(18, 21, 7, 28) \notin \mathcal{G}$ , and  $(18, 28, 7, 21) \notin \mathcal{G}$ .

### Proof.

By GAP:

```
1 gap> (18,21)(7,28) in G or (18,28)(7,21) in G or  
      (18,21,7,28) in G or (18,28,7,21) in G;  
2 false
```

(GAP uses stabiliser chains to verify membership!)

□

Can generalise to any two edge pieces (more cases)!

## Solving a Rubik's cube... i

We can use GAP to solve Rubik's cube state:

# Solving a Rubik's cube... i

We can use GAP to solve Rubik's cube state:

```
1 gap> H := FreeGroup("u","l","f","r","b","d");
2 <free group on the generators [ u, l, f, r, b, d ]>
3 gap> h := GroupHomomorphismByImages( H, G, GeneratorsOfGroup( H ),
    GeneratorsOfGroup( G ) );
4 [ u, l, f, r, b, d ] -> [ (1,3,8,6)(2,5,7,4)(9,33,25,17)(10,34,26,18)
    (11,35,27,19),
5 (1,17,41,40)(4,20,44,37)(6,22,46,35)(9,11,16,14)(10,13,15,12),
    (6,25,43,16)(7,28,42,13)(8,30,41,11)(17,19,24,
6 22)(18,21,23,20), (3,38,43,19)(5,36,45,21)(8,33,48,24)
    (25,27,32,30)(26,29,31,28),
7 (1,14,48,27)(2,12,47,29)(3,9,46,32)(33,35,40,38)(34,37,39,36),
    (14,22,30,38)(15,23,31,39)(16,24,32,40)(41,43,48,
8 46)(42,45,47,44) ]
```

$(F = \langle u, \ell, f, r, b, d \rangle$  is free group on 6 generators. Then  $f : F \rightarrow \mathcal{G}$  is hom given by  $u \mapsto U, l \mapsto L, f \mapsto F, r \mapsto R, b \mapsto B, d \mapsto D.$ )

Use GAP to generate random state  $x \in \mathcal{G}$  (uses stabiliser chain):

Use GAP to generate random state  $x \in \mathcal{G}$  (uses stabiliser chain):

```
1 gap> x := Random( G );  
2 (1,27,32,6,43,14,22)(2,28,13,37,18,15,47,42,31)(3,38,17,24,46,41,9)  
   (5,26)(7,44,39,23,45,34,21,20,12)(11,30,40,16,35,33,48)(29,36)
```

Randomly generated state (uniform distribution on  $\mathcal{G}$ ):


$$x = (1, 27, 32, 6, 43, 14, 22)(2, 28, 13, 37, 18, 15, 47, 42, 31) \\ (3, 38, 17, 24, 46, 41, 9)(5, 26)(7, 44, 39, 23, 45, 34, 21, 20, 12) \\ (11, 30, 40, 16, 35, 33, 48)(29, 36)$$

## Solving a Rubik's cube... iii

$$x = (1, 27, 32, 6, 43, 14, 22)(2, 28, 13, 37, 18, 15, 47, 42, 31)$$

$$(3, 38, 17, 24, 46, 41, 9)(5, 26)(7, 44, 39, 23, 45, 34, 21, 20, 12)$$

$$(11, 30, 40, 16, 35, 33, 48)(29, 36)$$





Factorisation into 78 generators and inverses:

## Solving a Rubik's cube... iv

Factorisation into 78 generators and inverses:

```
1 gap> PreImagesRepresentative( h, x );
2 1*f^-1*1^-1*f*u*f*u^-1*f^2*1*f*1^-1*u^-1*1^-1*u*1*u^-1*1*u*f*u^-1*f
   ^-1*1^-2*u*1*f^-1*1*f*(1^-1*u)^2*b^-1*u*b*1*u*1^-1*f^-1*1^-1*f*1
   ^2*u*1^-1*u*1*b^-1*u^-1*b*1*d*f^2\
3 *d^-1*1*f^-1*u*1^-1*f*u^-1*1*d^-1*1*b*d*u^-2*b^-1*r^-1*b*u^-1*r*f^-1*
   u*d^-2
4 gap> Length( last );
5 78
```

$$\begin{aligned} x = & LF^{-1}L^{-1}FUFU^{-1}F^2LFL^{-1}U^{-1}L^{-1}ULU^{-1}LUFU^{-1}F^{-1}L^{-2}U \\ & LF^{-1}LF(L^{-1}U)^2B^{-1}UBLUL^{-1}F^{-1}L^{-1}FL^2UL^{-1}ULB^{-1}U^{-1}BL \\ & DF^2D^{-1}LF^{-1}UL^{-1}FU^{-1}LD^{-1}LBDU^{-2}B^{-1}R^{-1}BU^{-1}RF^{-1}UD^{-2} \end{aligned}$$

## Solving a Rubik's cube... iv

Factorisation into 78 generators and inverses:

```
1 gap> PreImagesRepresentative( h, x );
2 1*f^-1*1^-1*f*u*f*u^-1*f^2*1*f*1^-1*u^-1*1^-1*u*1*u^-1*1*u*f*u^-1*f
   ^-1*1^-2*u*1*f^-1*1*f*(1^-1*u)^2*b^-1*u*b*1*u*1^-1*f^-1*1^-1*f*1
   ^2*u*1^-1*u*1*b^-1*u^-1*b*1*d*f^2\
3 *d^-1*1*f^-1*u*1^-1*f*u^-1*1*d^-1*1*b*d*u^-2*b^-1*r^-1*b*u^-1*r*f^-1*
   u*d^-2
4 gap> Length( last );
5 78
```

$$\begin{aligned}x = & LF^{-1}L^{-1}FUFU^{-1}F^2LFL^{-1}U^{-1}L^{-1}ULU^{-1}LUFU^{-1}F^{-1}L^{-2}U \\ & LF^{-1}LF(L^{-1}U)^2B^{-1}UBLUL^{-1}F^{-1}L^{-1}FL^2UL^{-1}ULB^{-1}U^{-1}BL \\ & DF^2D^{-1}LF^{-1}UL^{-1}FU^{-1}LD^{-1}LBDU^{-2}B^{-1}R^{-1}BU^{-1}RF^{-1}UD^{-2}\end{aligned}$$

(GAP uses stabiliser chains to factorise!)

## Solving a Rubik's cube... v

Check this is correct:

```
1 gap> x = L*F^(-1)*L^(-1)*F*U*F*U^(-1)*F^2*L*F*L^(-1)*U^(-1)*L^(-1)*U*
  L*U^(-1)*L*U*F*U^(-1)*F^(-1)*L^(-2)*U*L*F^(-1)*L*F*(L^(-1)*U)^2*
  B^(-1)*U*B*L*U*L^(-1)*F^(-1)*L^(-1)*F*L^2*U*L^(-1)*U*L*B^(-1)*U
  ^(-1)*B*L*D*F^2*D^(-1)*L*F^(-1)*U*L^(-1)*F*U^(-1)*L*D^(-1)*L*B*D
  *U^(-2)*B^(-1)*R^(-1)*B*U^(-1)*R*F^(-1)*U*D^(-2);
2 true
```

## Solving a Rubik's cube... v

Check this is correct:

```
1 gap> x = L*F^(-1)*L^(-1)*F*U*F*U^(-1)*F^2*L*F*L^(-1)*U^(-1)*L^(-1)*U*
  L*U^(-1)*L*U*F*U^(-1)*F^(-1)*L^(-2)*U*L*F^(-1)*L*F*(L^(-1)*U)^2*
  B^(-1)*U*B*L*U*L^(-1)*F^(-1)*L^(-1)*F*L^2*U*L^(-1)*U*L*B^(-1)*U
  ^(-1)*B*L*D*F^2*D^(-1)*L*F^(-1)*U*L^(-1)*F*U^(-1)*L*D^(-1)*L*B*D
  *U^(-2)*B^(-1)*R^(-1)*B*U^(-1)*R*F^(-1)*U*D^(-2);
2 true
```

To solve state  $x$ , apply move  $x^{-1} \in \mathcal{G}$  since

## Solving a Rubik's cube... v

Check this is correct:

```
1 gap> x = L*F^(-1)*L^(-1)*F*U*F*U^(-1)*F^2*L*F*L^(-1)*U^(-1)*L^(-1)*U*  
L*U^(-1)*L*U*F*U^(-1)*F^(-1)*L^(-2)*U*L*F^(-1)*L*F*(L^(-1)*U)^2*  
B^(-1)*U*B*L*U*L^(-1)*F^(-1)*L^(-1)*F*L^2*U*L^(-1)*U*L*B^(-1)*U  
^(-1)*B*L*D*F^2*D^(-1)*L*F^(-1)*U*L^(-1)*F*U^(-1)*L*D^(-1)*L*B*D  
*U^(-2)*B^(-1)*R^(-1)*B*U^(-1)*R*F^(-1)*U*D^(-2);  
2 true
```

To solve state  $x$ , apply move  $x^{-1} \in \mathcal{G}$  since  $x^{x^{-1}} = xx^{-1} = 1$ :

$$\begin{aligned}x^{-1} = & D^2U^{-1}FR^{-1}UB^{-1}RBU^2D^{-1}B^{-1}L^{-1}DL^{-1}UF^{-1}LU^{-1}FL^{-1}DF^{-2}D^{-1} \\ & L^{-1}B^{-1}UBL^{-1}U^{-1}LU^{-1}L^{-2}F^{-1}LFLU^{-1}L^{-1}B^{-1}U^{-1}B(U^{-1}L)^2F^{-1}L^{-1}F \\ & L^{-1}U^{-1}L^2FUF^{-1}U^{-1}L^{-1}UL^{-1}U^{-1}LULF^{-1}L^{-1}F^{-2}UF^{-1}U^{-1}F^{-1}LFL^{-1}\end{aligned}$$

(Just invert each term in factorisation above and reverse, thus 78 steps.)

## Solving a Rubik's cube... v

Check this is correct:

```
1 gap> x = L*F^(-1)*L^(-1)*F*U*F*U^(-1)*F^2*L*F*L^(-1)*U^(-1)*L^(-1)*U*  
L*U^(-1)*L*U*F*U^(-1)*F^(-1)*L^(-2)*U*L*F^(-1)*L*F*(L^(-1)*U)^2*  
B^(-1)*U*B*L*U*L^(-1)*F^(-1)*L^(-1)*F*L^2*U*L^(-1)*U*L*B^(-1)*U  
^(-1)*B*L*D*F^2*D^(-1)*L*F^(-1)*U*L^(-1)*F*U^(-1)*L*D^(-1)*L*B*D  
*U^(-2)*B^(-1)*R^(-1)*B*U^(-1)*R*F^(-1)*U*D^(-2);  
2 true
```

To solve state  $x$ , apply move  $x^{-1} \in \mathcal{G}$  since  $x^{x^{-1}} = xx^{-1} = 1$ :

$$\begin{aligned}x^{-1} = & D^2U^{-1}FR^{-1}UB^{-1}RBU^2D^{-1}B^{-1}L^{-1}DL^{-1}UF^{-1}LU^{-1}FL^{-1}DF^{-2}D^{-1} \\ & L^{-1}B^{-1}UBL^{-1}U^{-1}LU^{-1}L^{-2}F^{-1}LFLU^{-1}L^{-1}B^{-1}U^{-1}B(U^{-1}L)^2F^{-1}L^{-1}F \\ & L^{-1}U^{-1}L^2FUF^{-1}U^{-1}L^{-1}UL^{-1}U^{-1}LULF^{-1}L^{-1}F^{-2}UF^{-1}U^{-1}F^{-1}LFL^{-1}\end{aligned}$$

(Just invert each term in factorisation above and reverse, thus 78 steps.)

Not very efficient, since it solves one piece in base  $B$  at a time (proceeding up stabiliser chain)... but it works!

## **Concluding remarks**

---



- Analyzing Rubik's cube with GAP: <https://www.gap-system.org/Doc/Examples/rubik.html>
- J.A. Paulos – Innumeracy (book)