

Support Vector Machine

Prof. Dr. Christina Bauer

christina.bauer@th-deg.de


Faculty of Computer Science

ALTERNATIVE VIEW OF LOGISTIC REGRESSION

$$h_{\theta}(x) = \frac{1}{1 + e^{-\theta^T x}}$$

If $y = 1$, we want $h_{\theta}(x) \sim 1 \rightarrow \theta^T x \gg 0$

If $y = 0$, we want $h_{\theta}(x) \sim 0 \rightarrow \theta^T x \ll 0$




ALTERNATIVE VIEW OF LOGISTIC REGRESSION


Cost of one example:

$$\begin{aligned} & -(y \log(h_\theta(x)) + (1-y) \log(1-h_\theta(x))) \\ &= -y \log\left(\frac{1}{1+e^{-\theta^T x}}\right) - (1-y) \log\left(1 - \frac{1}{1+e^{-\theta^T x}}\right) \end{aligned}$$

If $y = 1 \rightarrow [z = \theta^T x] \gg 0$





If $y = 0 \rightarrow [z = \theta^T x] \ll 0$



SUPPORT VECTOR MACHINE

Cost function Logistic Regression

$$\min_{\theta} \frac{1}{m} \left[\sum_{i=1}^m y^{(i)} (-\log(h_{\theta}(x^{(i)}))) + (1-y^{(i)}) (-\log(1-h_{\theta}(x^{(i)}))) \right] + \frac{\lambda}{2m} \sum_{j=1}^n \theta_j^2$$

 $\text{cost}_1(\theta^T x)$  $\text{cost}_0(\theta^T x)$


Support Vector Machine

$$\min_{\theta} \frac{1}{m} \left[\sum_{i=1}^m y^{(i)} \text{cost}_1(\theta^T x) + (1-y^{(i)}) \text{cost}_0(\theta^T x) \right] + \frac{\lambda}{2m} \sum_{j=1}^n \theta_j^2 \rightarrow \cancel{\frac{1}{m}}$$

C instead of λ

$$\min_{\theta} C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T x) + (1-y^{(i)}) \text{cost}_0(\theta^T x)] + \frac{1}{2} \sum_{j=1}^n \theta_j^2$$

QUESTION




Consider the following minimization problems:

1. $\min_{\theta} \frac{1}{m} \left[\sum_{i=1}^m y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)}) \text{cost}_0(\theta^T x) \right] + \frac{\lambda}{2m} \sum_{j=1}^n \theta_j^2$
2. $\min_{\theta} C \sum_{i=1}^m \left[y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)}) \text{cost}_0(\theta^T x) \right] + \frac{1}{2} \sum_{j=1}^n \theta_j^2$

These two optimization problems will give the same value of θ (i.e., the same value of θ gives the optimal solution to both problems) if:

- A: $C=\lambda$
- B: $C=-\lambda$
- C: $C=1/\lambda$
- D: $C=2/\lambda$

QUESTION



Consider the following minimization problems:

1. $\min_{\theta} \frac{1}{m} \left[\sum_{i=1}^m y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)}) \text{cost}_0(\theta^T x) \right] + \frac{\lambda}{2m} \sum_{j=1}^n \theta_j^2$
2. $\min_{\theta} C \sum_{i=1}^m \left[y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)}) \text{cost}_0(\theta^T x) \right] + \frac{1}{2} \sum_{j=1}^n \theta_j^2$

These two optimization problems will give the same value of θ (i.e., the same value of θ gives the optimal solution to both problems) if:

A: $C=\lambda$

B: $C=-\lambda$

~~C: $C=1/\lambda$~~

D: $C=2/\lambda$


SVM HYPOTHESIS

$$\min_{\theta} C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)})\text{cost}_0(\theta^T x)] + \frac{1}{2} \sum_{j=1}^n \theta_j^2$$


$$h_{\theta}(x) \begin{cases} 1 & \text{if } \theta^T x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

SVM – LARGE MARGIN CLASSIFIERS

$$\min_{\theta} C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)})\text{cost}_1(\theta^T x)] + \frac{1}{2} \sum_{j=1}^n \theta_j^2$$



If $y = 1$, we want $\theta^T x \geq 1$ (not just ≥ 0)



If $y = 0$, we want $\theta^T x \leq -1$ (not just < 0)


SVM – DECISION BOUNDARY

$$\min_{\theta} C \sum_{i=1}^m \underline{[y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)})\text{cost}_1(\theta^T x)]} + \frac{1}{2} \sum_{j=1}^n \theta_j^2$$

(With C being a large number \rightarrow first term of function should be near 0)


Whenever $y^{(i)} = 1$:

$$\theta^T x^{(i)} \geq 1 \text{ (not just } \geq 0 \text{)}$$



Whenever $y^{(i)} = 0$:

$$\theta^T x^{(i)} \leq -1$$



Optimization problem:


$$\min_{\theta} \epsilon * \theta + \frac{1}{2} \sum_{j=1}^n \theta_j^2$$

s.t.

$$\theta^T x^{(i)} \geq 1 \text{ if } y^{(i)} = 1$$


$$\theta^T x^{(i)} \leq -1 \text{ if } y^{(i)} = 0$$

SVM – DECISION BOUNDARY: LINEARLY SEPARABLE





→ Large margin classifier

SVM – LARGE MARGIN CLASSIFIER IN THE PRESENCE OF OUTLIERS



- C very large → decision boundary will change to red one (good idea?)
- C not too large → decision boundary will stay at the purple line
- If C is not too large SVM is more robust to outliers (red point) and if your data is not linearly separable (orange points)

VECTOR INNER PRODUCTS



$$\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$$

$\|\mathbf{u}\|$ = length of vector $\mathbf{u} = \sqrt{u_1^2 + u_2^2} \in \mathbb{R}$

p = length of projection of \mathbf{v} onto $\mathbf{u} \in \mathbb{R}$

inner product: $\mathbf{u}^T \mathbf{v} = \mathbf{v}^T \mathbf{u}$

$$\mathbf{u}^T \mathbf{v} = p * \|\mathbf{u}\|$$

$$\mathbf{u}^T \mathbf{v} = u_1 * v_1 + u_2 * v_2$$

SVM DECISION BOUNDARY

Optimization problem:

$$\min_{\theta} \frac{1}{2} \sum_{j=1}^n \theta_j^2$$

→ Simplification: $\theta_0 = 0$; $n = 2$

$$\rightarrow \frac{1}{2}(\theta_1^2 + \theta_2^2) \rightarrow \frac{1}{2}(\sqrt{\theta_1^2 + \theta_2^2})^2$$

→ $\sqrt{\theta_1^2 + \theta_2^2} = \|\theta\|$ (length of the vector theta)

$$\rightarrow \frac{1}{2} \|\theta\|^2$$


s.t.

$$\theta^T x^{(i)} \geq 1 \text{ if } y^{(i)} = 1$$

$$\theta^T x^{(i)} \leq -1 \text{ if } y^{(i)} = 0$$

→ $\theta^T x^{(i)}$ → inner product

$$\rightarrow p^{(i)} * \|\theta\| = \theta_1 x_1^{(i)} + \theta_2 x_2^{(i)}$$



SVM DECISION BOUNDARY

Optimization problem (C is very large):

$$\min_{\theta} \frac{1}{2} \sum_{j=1}^n \theta_j^2 = \frac{1}{2} \|\theta\|^2$$

→ Minimize the length of the vector θ

s.t.



$$p^{(i)} * \|\theta\| \geq 1 \text{ if } y^{(i)} = 1$$

$$p^{(i)} * \|\theta\| \leq -1 \text{ if } y^{(i)} = 0$$

$$\rightarrow \theta^T x^{(i)} = p^{(i)} * \|\theta\|$$

→ $p^{(i)}$ is the projection of $x^{(i)}$ onto the vector θ

→ Simplification: $\theta_0 = 0$ (Decision Boundary passes through (0;0))



→ The upper decision boundary will not be chosen

→ $p^{(1)} * \|\theta\| \geq 1$; if $p^{(1)}$ is small $\|\theta\|$ has to be large

→ $p^{(2)} * \|\theta\| \leq -1$; if $p^{(2)}$ is small negative number $\|\theta\|$ has to be large

→ $\min_{\theta} \frac{1}{2} \|\theta\|^2 \rightarrow$ the norm of θ should be small


→ SVM hypothesis → large margin

→ $p^{(1)} * \|\theta\| \geq 1 \rightarrow p^{(1)}$ is bigger now $\|\theta\|$ can be smaller (equivalent for $p^{(2)}$)

NON-LINEAR DECISION BOUNDARY

- Predict e.g. $y = 1$ if $\theta_0 + \theta_1 \underline{x_1} + \theta_2 \underline{x_2} + \theta_3 \underline{x_1 x_2} + \theta_4 \underline{x_1^2} + \theta_4 \underline{x_2^2} + \theta_5 \underline{x_2^3} + \dots \geq 0$

$$h_{\theta}(x) = \begin{cases} 1 & \text{if } \theta^T x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$



→ Predict e.g. $y = 1$ if $\theta_0 + \theta_1 f_1 + \theta_2 f_2 + \theta_3 f_3 + \theta_4 f_4 + \theta_4 f_5 + \theta_5 f_6 + \dots \geq 0$

- Is there a different/better choice of the features f_1, f_2, f_3, \dots ?


KERNEL

Given x , compute new features depending on proximity to landmarks $l^{(1)}, l^{(2)}, l^{(3)}$

$$f_1 = \text{similarity}(x, l^{(1)}) = e^{-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}}$$

$$f_2 = \text{similarity}(x, l^{(2)}) = e^{-\frac{\|x - l^{(2)}\|^2}{2\sigma^2}}$$

$$f_3 = \text{similarity}(x, l^{(3)}) = e^{-\frac{\|x - l^{(3)}\|^2}{2\sigma^2}}$$



Similarity = kernel (here: Gaussian kernel) $\rightarrow k(x, l)$

KERNELS AND SIMILARITY

$$f_1 = \text{similarity}(x, l^{(1)}) = e^{-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}} \quad (\text{where } \|x - l\|^2 = \sum_{j=1}^n (x_j - l_j)^2)$$

If $x \sim l^{(1)}$:

$$f_1 \sim e^{-\frac{0^2}{2\sigma^2}} \sim 1$$

If x is far from $l^{(1)}$:

$$f_1 \sim e^{-\frac{\text{large number}^2}{2\sigma^2}} \sim 0$$




$$l^{(1)} \rightarrow f_1$$

$$l^{(2)} \rightarrow f_2$$

$$l^{(3)} \rightarrow f_3$$

KERNELS

- σ^2 defines how narrow the gaussian kernels are





Example with $l^{(1)} = ((3);(5))$ and $f_1 = e^{-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}}$

QUESTION


Consider a 1-D example with one feature x_1 . Suppose $l^{(1)}=5$.

This is a plot of $f_1 = e^{-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}}$ when $\sigma^2 = 1$.


Suppose we now change $\sigma^2 = 4$. Which of the following is a plot of f_1 with the new value of σ^2 ?




A:



B:



C:





QUESTION


Consider a 1-D example with one feature x_1 . Suppose $l^{(1)}=5$.

This is a plot of $f_1 = e^{-\frac{\|x-l^{(1)}\|^2}{2\sigma^2}}$ when $\sigma^2 = 1$.


Suppose we now change $\sigma^2 = 4$. Which of the following is a plot of f_1 with the new value of σ^2 ?




A:




B:



~~C:~~



KERNELS



Predict "1" when $\theta_0 + \theta_1 f_1 + \theta_2 f_2 + \theta_3 f_3 \geq 0$


- Example:
- $\theta_0 = -0.5$; $\theta_1 = 1$; $\theta_2 = 1$; $\theta_3 = 0$

$$\begin{aligned} x^{(1)} &\rightarrow f_1 \sim 1; f_2 \sim 0; f_3 \sim 0 \\ &\rightarrow \theta_0 + \theta_1 * 1 + \theta_2 * 0 + \theta_3 * 0 \\ &\rightarrow -0.5 + 1 * 1 + 1 * 0 + 0 * 0 \\ &\rightarrow 0.5 \geq 0 \rightarrow x^{(1)} \text{ predict 1} \end{aligned}$$

$$x^{(2)} \rightarrow f_1 \sim 0; f_2 \sim 0; f_3 \sim 0 \rightarrow -0.5 + 1 * 0 + 1 * 0 + 0 * 0 \rightarrow -0.5 < 0$$

CHOOSING THE LANDMARKS

- Given x :
- $f_1 = \text{similarity}(x, l^{(1)}) = e^{-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}}$
- Predict “1” when $\theta_0 + \theta_1 f_1 + \theta_2 f_2 + \theta_3 f_3 \geq 0$
- Where to get $l^{(1)}, l^{(2)}, l^{(3)}$?
- Idea: set $l^{(1)}, \dots, l^{(m)}$ to $x^{(1)}, \dots, x^{(m)}$



SVM WITH KERNELS

- Given $(x^{(1)}, y^{(1)}), (x^{(2)}, y^{(2)}), \dots, (x^{(m)}, y^{(m)})$
- Choose $l^{(1)} = x^{(1)}, l^{(2)} = x^{(2)}, \dots, l^{(m)} = x^{(m)}$

- Given example x :
 - $f_1 = \text{similarity}(x, l^{(1)})$ (where $l^{(1)} = x^{(1)}$)
 - $f_2 = \text{similarity}(x, l^{(2)})$

- For training example $(x^{(i)}, y^{(i)})$

$x^{(i)} \rightarrow$

$$f_1^{(i)} = \text{similarity}(x^{(i)}, l^{(1)})$$

$$f_2^{(i)} = \text{similarity}(x^{(i)}, l^{(2)})$$

...

$$f_m^{(i)} = \text{similarity}(x^{(i)}, l^{(m)})$$

\rightarrow Somewhere in this list: $f_i^{(i)} = \text{similarity}(x^{(i)}, l^{(i)})$ (where $l^{(i)} = x^{(i)}$)

\rightarrow With Gaussian kernel: $e^{-\frac{0}{2\sigma^2}} = 1$

$$f = \begin{bmatrix} f_0 \\ f_1 \\ \vdots \\ f_m \end{bmatrix} \quad f_0 = 1$$

$$x^{(i)} = f^{(i)} = \begin{bmatrix} f_0^{(i)} \\ f_1^{(i)} \\ \vdots \\ f_m^{(i)} \end{bmatrix} \quad f_0^{(i)} = 1$$

SVM WITH KERNELS

- Hypothesis: Given x , compute features $f \in \mathbb{R}^{m+1}$ (m = training set size and number of landmarks)
 - Predict „1“ if $\theta^T f \geq 0$ (with $\theta \in \mathbb{R}^{m+1}$)

- Training:

- Before:

$$\min_{\theta} C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)})\text{cost}_0(\theta^T x)] + \frac{1}{2} \sum_{j=1}^n \theta_j^2$$


- Now:

$$\min_{\theta} C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T f^{(i)}) (1-y^{(i)})\text{cost}_1(\theta^T f^{(i)})] + \frac{1}{2} \sum_{j=1}^m \theta_j^2$$


Note: $\sum_{j=1}^m \theta_j^2 = \theta^T \theta$; often implemented as $\theta^T M \theta$ (M depends on the kernel)

SVM PARAMETERS

- $C = \left(\frac{1}{\lambda}\right)$:
 - Large C: lower bias, high variance (think of small λ)
 - Small C: higher bias, low variance (think of large λ)
- σ^2
 - Large σ^2 : Features f_i vary more smoothly
 - \rightarrow higher bias, lower variance
 - Small σ^2 : Features f_i vary less smoothly
 - \rightarrow lower bias, higher variance




QUESTION



Suppose you train an SVM and find it overfits your training data. Which of these would be a reasonable next step? Check all that apply.

- A: Increase C
- B: Decrease C
- C: Increase σ^2
- D: Decrease σ^2

QUESTION



Suppose you train an SVM and find it overfits your training data. Which of these would be a reasonable next step? Check all that apply.

A: Increase C

~~B: Decrease C~~

~~C: Increase σ^2~~

D: Decrease σ^2

USING A SVM

- Use a SVM software package to solve for the parameters θ
- Specify:
 - Parameter C
 - Kernel
- E. g. no kernel (“linear kernel”)
 - Predict “ $y = 1$ ” if $\theta^T x \geq 0$
 - Linear Decision Boundary
 - Applied if n is large and m is small (risk of overfitting)
- Gaussian kernel
 - $f_i = e^{-\frac{\|x - l^{(i)}\|^2}{2\sigma^2}}$, where $l^{(i)} = x^{(i)}$
 - Choose σ^2
 - Applied if n is small and m is large

KERNEL FUNCTION EXAMPLE

Function $f = \text{kernel}(x_1, x_2)$

$$f_i = \exp\left(-\frac{\|x_1 - x_2\|^2}{2\sigma^2}\right)$$

return

Note: Do perform feature scaling before using the Gaussian kernel


$$\|x - l\|^2 \rightarrow (x_1 - l_1)^2 + (x_2 - l_2)^2 + \dots + (x_n - l_n)^2$$

→ If one x is very large it will “dominate” the value

OTHER KERNEL

- Note: Not all similarity functions make valid kernels.
→ Need to satisfy technical condition called “Mercer’s Theorem” to make sure SVM packages’ optimizations run correctly and do not diverge
- Other valid kernels:
 - Polynomial kernel: $(x^T l + \text{constant})^{\text{degree}}$ e.g. $(x^T l)^3, (x^T l + 1)^3, (x^T l + 5)^4$
 - String kernel, chi-square kernel, histogram intersection kernel,...

QUESTION



Suppose you are trying to decide among a few different choices of kernels and are also choosing parameters such as C , σ^2 , etc. How should you make the choice?


A: Choose whatever performs best on the training data.

B: Choose whatever performs best on the cross-validation data.

C: Choose whatever performs best on the test data.

D: Choose whatever gives the largest SVM margin.

QUESTION



Suppose you are trying to decide among a few different choices of kernels and are also choosing parameters such as C , σ^2 , etc. How should you make the choice?

A: Choose whatever performs best on the training data.

~~B~~: Choose whatever performs best on the cross-validation data.


C: Choose whatever performs best on the test data.

D: Choose whatever gives the largest SVM margin.

MULTICLASS CLASSIFICATION

- Many SVM packages already have built-in multi-class classification functionality
- Otherwise, use one-vs-all method (Train K SVMs, one to distinguish $y=i$ from the rest, for $i = 1, 2, \dots, K$), get $\theta^{(1)}, \theta^{(2)}, \dots, \theta^{(K)}$, pick class i with largest $(\theta^{(i)})^T x$

- $Y \in \{1, 2, \dots, K\}$



LOGISTIC REGRESSION VS. SVM - RECOMMENDATION

- If n is large (relative to m) (e. g. text classification)
 - Use logistic regression, or SVM without a kernel (the “linear kernel”)
- If n is small and m is intermediate
 - Use SVM with a Gaussian Kernel
- If n is small and m is large
 - Create/add more features, then use logistic regression or SVM without a kernel.
- Note: a neural network is likely to work well for any of these situations, but may be slower to train.

QUIZ – QUESTION 1


- Suppose you have trained an SVM classifier with a Gaussian kernel, and it learned the following decision boundary on the training set:
- You suspect that the SVM is underfitting your dataset. Should you try increasing or decreasing C ? Increasing or decreasing σ^2 ?

A: It would be reasonable to try increasing C . It would also be reasonable to try increasing σ^2 .

B: It would be reasonable to try decreasing C . It would also be reasonable to try increasing σ^2 .

C: It would be reasonable to try decreasing C . It would also be reasonable to try decreasing σ^2 .

D: It would be reasonable to try increasing C . It would also be reasonable to try decreasing σ^2 .



QUIZ – QUESTION 1


- Suppose you have trained an SVM classifier with a Gaussian kernel, and it learned the following decision boundary on the training set:
- You suspect that the SVM is underfitting your dataset. Should you try increasing or decreasing C ? Increasing or decreasing σ^2 ?

A: It would be reasonable to try increasing C . It would also be reasonable to try increasing σ^2 .

B: It would be reasonable to try decreasing C . It would also be reasonable to try increasing σ^2 .


C: It would be reasonable to try decreasing C . It would also be reasonable to try decreasing σ^2 .

~~D: It would be reasonable to try increasing C . It would also be reasonable to try decreasing σ^2 .~~




QUIZ – QUESTION 2


- The formula for the Gaussian kernel is given by $f_1 = \text{similarity}(x, l^{(1)}) = e^{-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}}$
- The figure shows a plot of $f_1 = \text{similarity}(x, l^{(1)})$ with $\sigma^2 = 1$
- Which of the following is a plot of f_1 when $\sigma^2 = 0.25$?




A:




B:



C:




D:




QUIZ – QUESTION 2


- The formula for the Gaussian kernel is given by $f_1 = \text{similarity}(x, l^{(1)}) = e^{-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}}$
- The figure shows a plot of $f_1 = \text{similarity}(x, l^{(1)})$ with $\sigma^2 = 1$
- Which of the following is a plot of f_1 when $\sigma^2 = 0.25$?




~~A:~~




B:





C:



D:



QUIZ – QUESTION 3



- The SVM solves $\min_{\theta} C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)})\text{cost}_0(\theta^T x)] + \frac{1}{2} \sum_{j=1}^n \theta_j^2$
- The first term in the objective is: $C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)})\text{cost}_0(\theta^T x)]$
- This first term will be zero if two of the following four conditions hold true.
Which are the two conditions that would guarantee that this term equals zero?



A: For every example with $y^{(i)}=1$, we have that $\theta^T x^{(i)} \geq 1$.

B: For every example with $y^{(i)}=0$, we have that $\theta^T x^{(i)} \leq -1$.

C: For every example with $y^{(i)}=1$, we have that $\theta^T x^{(i)} \geq 0$.

D: For every example with $y^{(i)}=0$, we have that $\theta^T x^{(i)} \leq 0$.

QUIZ – QUESTION 3



- The SVM solves $\min_{\theta} C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)})\text{cost}_0(\theta^T x)] + \frac{1}{2} \sum_{j=1}^n \theta_j^2$
- The first term in the objective is: $C \sum_{i=1}^m [y^{(i)} \text{cost}_1(\theta^T x) (1-y^{(i)})\text{cost}_0(\theta^T x)]$
- This first term will be zero if two of the following four conditions hold true.
Which are the two conditions that would guarantee that this term equals zero?

~~A~~: For every example with $y^{(i)}=1$, we have that $\theta^T x^{(i)} \geq 1$.

~~B~~: For every example with $y^{(i)}=0$, we have that $\theta^T x^{(i)} \leq -1$.

C: For every example with $y^{(i)}=1$, we have that $\theta^T x^{(i)} \geq 0$.

D: For every example with $y^{(i)}=0$, we have that $\theta^T x^{(i)} \leq 0$.

QUIZ – QUESTION 4

Suppose you have a dataset with $n = 10$ features and $m = 5000$ examples. After training your logistic regression classifier with gradient descent, you find that it has underfit the training set and does not achieve the desired performance on the training or cross validation sets.

Which of the following might be promising steps to take? Check all that apply.

- A: Use an SVM with a linear kernel, without introducing new features.
- B: Increase the regularization parameter λ .
- C: Use an SVM with a Gaussian Kernel.
- D: Create/add new polynomial features.

QUIZ – QUESTION 4

Suppose you have a dataset with $n = 10$ features and $m = 5000$ examples. After training your logistic regression classifier with gradient descent, you find that it has underfit the training set and does not achieve the desired performance on the training or cross validation sets.

Which of the following might be promising steps to take? Check all that apply.

A: Use an SVM with a linear kernel, without introducing new features.

B: Increase the regularization parameter λ .

☒ C: Use an SVM with a Gaussian Kernel.

☒ D: Create/add new polynomial features.

QUIZ – QUESTION 5

Which of the following statements are true? Check all that apply.

A: It is important to perform feature normalization before using the Gaussian kernel.

B: If the data are linearly separable, an SVM using a linear kernel will return the same parameters θ regardless of the chosen value of C (i.e., the resulting value of θ does not depend on C).

C: The maximum value of the Gaussian kernel (i.e., $\text{sim}(x, l^{(1)})$) is 1.

D: Suppose you are using SVMs to do multi-class classification and would like to use the one-vs-all approach. If you have K different classes, you will train $K - 1$ different SVMs.

QUIZ – QUESTION 5

Which of the following statements are true? Check all that apply.

☒ A: It is important to perform feature normalization before using the Gaussian kernel.

☐ B: If the data are linearly separable, an SVM using a linear kernel will return the same parameters θ regardless of the chosen value of C (i.e., the resulting value of θ does not depend on C).

☒ C: The maximum value of the Gaussian kernel (i.e., $\text{sim}(x, l^{(1)})$) is 1.

☐ D: Suppose you are using SVMs to do multi-class classification and would like to use the one-vs-all approach. If you have K different classes, you will train $K - 1$ different SVMs.