

Machine Learning - Assignment 3

Sudip Sinha

May 4, 2014

Exercise 2

Given

- $VV^T = V^T V = I$
- D is an invertible diagonal matrix

Proof

Let $B = VD^{-1}V^T$

- $AB = VDV^TVD^{-1}V^T = VD(V^TV)D^{-1}V^T = V DID^{-1}V^T = V(DD^{-1})V^T = VIV^T = I$
- $BA = VD^{-1}V^TVDV^T = VD^{-1}(V^TV)DV^T = VD^{-1}IDV^T = V(D^{-1}D)V^T = VIV^T = I$

Thus, $A^{-1} = B = VD^{-1}V^T$.

Exercise 3

We just have to show that the square of the norm is a convex function.

By the definition of norm,

$$\begin{aligned} \|tx + (1-t)y\| &\leq \|tx\| + \|(1-t)y\| = t\|x\| + (1-t)\|y\| \\ \implies \|tx + (1-t)y\|^2 &\leq (t\|x\| + (1-t)\|y\|)^2 \leq t^2\|x\|^2 + (1-t)^2\|y\|^2 \end{aligned}$$

Thus, the least squares loss function $\|y - Xw\|^2$ is a convex function of w .

Exercise 4

Let $W = \text{diag}(r_i)$. Therefore, $W^T = W$

Thus, the problem can be reformulated as

$$w = \arg \min_w (e), \text{ where } e = (y - Xw)^T W (y - Xw)$$

Using the rules of Matrix Calculus given [here](#):

$$\begin{aligned} \frac{\partial e}{\partial w} &= (y - Xw)^T W \frac{\partial}{\partial w} (y - Xw) + (y - Xw)^T W^T \frac{\partial}{\partial w} (y - Xw) \\ &= 2(y - Xw)^T W^T (-X) \quad (\text{Since } W^T = W) \\ &= 2(X^T W (Xw - y))^T \end{aligned}$$

For minimum error:

$$\begin{aligned} \frac{\partial e}{\partial w} &= 0^T \\ \implies 2(X^T W (Xw - y))^T &= 0^T \\ \implies X^T W (Xw - y) &= 0 \\ \implies X^T W Xw &= X^T y \\ \implies w &= (X^T W X)^{-1} X^T y \end{aligned}$$

Exercise 6

Notations

Symbol	Dimension	Description
X	$m \times n$	The design matrix
y	$m \times 1$	The actual values of the predicted variable
$O(\cdot)$	-	The Big O notation

Computational Complexity¹

	Output	Complexity	Operation
LLS	$X^T X$	$O(mn^2)$	Matrix multiplication
	$X^T y$	$O(mn)$	Matrix multiplication
	$X^T X w = X^T y$	$O(n^3)$	Solving linear system of equations

Now, we shall assume that $n \ll m$. In this case, $O(mn^2)$ dominates over $O(n^3)$. Thus the computational complexity of LLS is $O(mn^2)$.

	Output	Complexity	Operation
kNN	$D_i = \Sigma_i (X_{test} - X_{train,i})^2$	$O(mn)$	Subtraction, squaring and summing (each $O(n)$) / row of train
	Sorted index vector δ	$O(m \ln m)$	Sorting
	Predicted class	$O(k)$	Mode

Now, we shall assume that $k \ll m$ and $\ln m \ll n$. In this case, $O(mn)$ dominates over the other complexities.

Space Complexity

	Object	Space
LLS	X	mn
	y	m
	X^T	mn
	$X^T X$	n^2
	$X^T y$	n
	w	n

Thus the space complexity of LLS is $O(mn)$.

	Object	Space
kNN	X	mn
	y	m
	$X^T y$	n
	δ	n

Thus the space complexity of LLS is $O(mn)$.

Final results

	Computation	Time
LLS	$O(mn^2)$	$O(mn)$
kNN	$O(mn)$	$O(mn)$

¹Taken from [Computational complexity of mathematical operations in Wikipedia](#)