
Body Cream (O/W)

Vitamin F Ethylester





Phase	Trade Name	INCI Name	w/w %	Supplier
А	Montanov 202	Arachidyl Alcohol, Behenyl Alcohol, Arachidyl Glucoside	3,00	Seppic
	Lanol P	Glycol Palmitate	5,00	Seppic
	Lipopeg 100-S	PEG-100 Stearate	2,10	Lipo Chemicals
	Ceraphyl 375	Isostearyl Neopentanoate	5,00	Ashland
	Lanol 99	Isononyl Isononanoate	9,00	Seppic
	Vitamin F Ethylester CLR	Ethyl Linoleate, Ethyl Linolenate, Ethyl Oleate	1,00	CLR
	Copherol F 1300 C	Tocopherol	0,10	BASF
В	Water	Water	ad 100	
С	LexFeel D4	Neopentyl Glycol Diheptanoate, Isododecane	8,00	Inolex
	Simulgel EG	Sodium Acrylate/Sodium Acryloyldimethyltaurate Copolymer, Isohexadecane, Polysorbate 80	2,50	Seppic
D	Ethanol (96%)	Alcohol Denat.	4,00	
	Euxyl PE 9010*	Phenoxyethanol, Ethylhexylglycerin	0,80	Schülke

Manufacture:

Heat up A and B to 75°C separately. Add B to A while stirring. Homogenize for 3 minutes by Ultra Turrax. Cool down while stirring. Add C and D below 30°C and stir until homogenous.



The recommendations and formulations given are based on our knowledge and experience in the field of technical application. The are, to the best of our belief, correct, but are offered without obligation. Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.