

## Some Useful Constants

Speed of light constant	: $c$
Boltzmann constant	: $k_B$
Electron charge	: $e$
Planck's constant	: $h$
Rest mass of electron	: $m_e$
Rest mass of proton	: $m_p$
Permeability of free space	: $\mu_0$
Permittivity of free space	: $\epsilon_0$

All other symbols have their usual meanings unless otherwise specified.

### Q. 1 – Q. 20 carry one mark each

**Q. 1** The value of the contour integral  $\left| \oint_C \mathbf{r} \times d\theta \right|$ , for a circle  $C$  of radius  $r$  with center at the origin is:

- (a)  $2\pi r$                       (b)  $\frac{r^2}{2}$                       (c)  $\pi r^2$                       (d)  $r$

**Q. 2** An electrostatic field  $\mathbf{E}$  exists in a given region  $R$ . Choose the **wrong** statement.

- (a) Circulation of  $\mathbf{E}$  is zero.  
 (b)  $\mathbf{E}$  can always be expressed as the gradient of a scalar field.  
 (c) The potential difference between any two arbitrary points in the region  $R$  is zero.  
 (d) The work done in a closed path lying entirely in  $R$  is zero.

**Q. 3** The Lagrangian of a free particle in spherical polar co-ordinates is given by

$$L = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\theta}^2 + r^2 \sin^2 \theta \dot{\phi}^2)$$

The quantity that is conserved is:

- (a)  $\frac{\partial L}{\partial \dot{r}}$     (b)  $\frac{\partial L}{\partial \dot{\theta}}$

(c)  $\frac{\partial L}{\partial \dot{\phi}}$

(d)  $\frac{\partial L}{\partial \dot{\phi}} + r\dot{\theta}$

**Q. 4** A conducting loop L of surface area S is moving with a velocity  $\mathbf{v}$  in a magnetic field  $\mathbf{B}(\mathbf{r}, t) = B_o t^2 \hat{z}$ .  $B_o$  is a positive constant of suitable dimensions. The emf induced,  $V_{\text{emf}}$ , in the loop is given by

(a)  $-\int_S \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{S}$

(c)  $-\int_S \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{S} - \oint_L (\mathbf{v} \times \mathbf{B}) \cdot d\mathbf{L}$

(b)  $\oint_L (\mathbf{v} \times \mathbf{B}) \cdot d\mathbf{L}$

(d)  $-\int_S \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{S} + \oint_L (\mathbf{v} \times \mathbf{B}) \cdot d\mathbf{L}$

**Q. 5** The eigenvalues of the matrix  $A = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$  are

(a) real and distinct

(c) complex and coinciding

(b) complex and distinct

(d) real and coinciding

**Q. 6**  $\sigma_i (i = 1, 2, 3)$  represent the Pauli spin matrices. Which one of the following is **NOT** true?

(a)  $\sigma_i \sigma_j + \sigma_j \sigma_i = 2\delta_{ij}$

(c) The eigenvalues of  $\sigma_i$  are  $\pm 1$

(b)  $\text{Tr}(\sigma_i) = 0$

(d)  $\det(\sigma_i) = 1$

**Q. 7** Which one of the functions given below represents the bound state eigenfunction of the operator  $-\frac{d^2}{dx^2}$  in the region,  $0 \leq x < \infty$ , with the eigenvalue  $-4$ ?

(a)  $A_o e^{2x}$

(c)  $A_o e^{-2x}$

(b)  $A_o \cosh 2x$

(d)  $A_o \sinh 2x$

**Q. 8** Pick the **WRONG** statement.

(a) The nuclear force is independent of electric charge

(b) The Yukawa potential is proportional to  $r^{-1} \exp\left(-\frac{mc}{\hbar} r\right)$ , where r is the separation between two nucleons

(c) The range of nuclear force is of the order of  $10^{-15} \text{ m} - 10^{-14} \text{ m}$

(d) The nucleons interact among each other by the exchange of mesons

**Q. 9** If p and q are the position and momentum variables, which one of the following is **NOT** a canonical transformation?

- (a)  $Q = \alpha q$  and  $P = \frac{1}{\alpha}p$ , for  $\alpha \neq 0$
- (b)  $Q = \alpha q + \beta p$  and  $P = \beta q + \alpha p$  for  $\alpha, \beta$  real and  $\alpha^2 - \beta^2 = 1$
- (c)  $Q = p$  and  $P = q$
- (d)  $Q = p$  and  $P = -q$


**Q. 10** The Common Mode Rejection Ratio (CMRR) of a differential amplifier using an operational amplifier is 100 dB. The output voltage for a differential input of  $200 \mu\text{V}$  is 2 V. The common mode gain is

- (a) 10 (c) 30 dB
- (b) 0.1 (d) 10 dB

**Q. 11** In an insulating solid which one of the following physical phenomena is a consequence of Pauli's exclusion principle?

- (a) Ionic conductivity
- (b) Ferromagnetism
- (c) Paramagnetism
- (d) Ferroelectricity

**Q. 12** Which one of the following curves gives the solution of the differential equation  $k_1 \frac{dx}{dt} + k_2 x = k_3$ , where  $k_1, k_2$  and  $k_3$  are positive constants with initial conditions  $x = 0$  at  $t = 0$ ?



**Q. 13** Identify which one is a first order phase transition?

- (a) A liquid to gas transition at its critical temperature.

- (b) A liquid to gas transition close to its triple point.
- (c) A paramagnetic to ferromagnetic transition in the absence of a magnetic field.
- (d) A metal to superconductor transition in the absence of a magnetic field.

**Q. 14** Group I lists some physical phenomena while Group II gives some physical parameters. Match the phenomena with the corresponding parameter.

**Group I**

- P. Doppler Broadening
- Q. Natural Broadening
- R. Rotational spectrum
- S. Total internal reflection

**Group II**

- 1. Moment of inertia
- 2. Refractive index
- 3. Lifetime of the energy level
- 4. Pressure

- (a) P-4, Q-3, R-1, S-2
- (b) P-3, Q-2, R-1, S-4
- (c) P-2, Q-3, R-4, S-1
- (d) P-1, Q-4, R-2, S-3

**Q. 15** The separation between the first Stokes and corresponding anti-Stokes lines of the rotational Raman spectrum in terms of the rotational constant,  $B$  is

- (a)  $2B$
- (b)  $4B$
- (c)  $6B$
- (d)  $12B$

**Q. 16** A superconducting ring is cooled in the presence of a magnetic field below its critical temperature ( $T_C$ ). The total magnetic flux that passes through the ring is

- (a) zero
- (b)  $n \frac{h}{2e}$
- (c)  $\frac{nh}{4\pi e}$
- (d)  $\frac{ne^2}{hc}$

**Q. 17** In a cubic crystal, atoms of mass  $M_1$  lie on one set of planes and atoms of mass  $M_2$  lie on planes interleaved between those of the first set. If  $C$  is the force constant between nearest neighbour planes, the frequency of lattice vibrations for the optical phonon branch with wavevector  $k = 0$  is

(a)  $\sqrt{2C\left(\frac{1}{M_1} + \frac{1}{M_2}\right)}$

(b)  $\sqrt{C\left(\frac{1}{2M_1} + \frac{1}{M_2}\right)}$

(c)  $\sqrt{C\left(\frac{1}{M_1} + \frac{1}{2M_2}\right)}$

(d) 0

**Q. 18** In the quark model which one of the following represents a proton?

(a)  $udd$

(b)  $uud$

(c)  $u\bar{b}$

(d)  $c\bar{c}$




Figure 1: Circuit diagram

**Q. 19** The circuit shown above

(a) is a common-emitter amplifier (c) is an oscillator

(b) uses a pnp transistor

(d) has a voltage gain less than one

**Q. 20** Consider a nucleus with  $N$  neutrons and  $Z$  protons. If  $m_p$ ,  $m_n$  and  $BE$  represent the mass of the proton, the mass of the neutron and the binding energy of the nucleus respectively and  $c$  is the velocity of light in free space, the mass of the nucleus is given by

(a)  $Nm_n + Zm_p$

(c)  $Nm_n + Zm_p + \frac{BE}{c^2}$

(b)  $Nm_p + Zm_n$

$$(d) Nm_p + Zm_n + \frac{BE}{c^2}$$

**Q. 21 to Q. 60 carry two marks each.**

**Q. 21** The magnetic field (in  $A\ m^{-1}$ ) inside a long solid cylindrical conductor of radius  $a = 0.1$  m is,  $\mathbf{H} = \frac{10^4}{r} \left[ \frac{1}{\alpha^2} \sin(\alpha r) - \frac{r}{\alpha} \cos(\alpha r) \right] \hat{\phi}$ , where  $\alpha = \frac{\pi}{2a}$ . What is the total current (in A) in the conductor?

- (a)  $\frac{\pi}{2a}$                       (b)  $\frac{800}{\pi}$                       (c)  $\frac{400}{\pi}$                       (d)  $\frac{300}{\pi}$

**Q. 22** Which one of the following current densities,  $\mathbf{J}$ , can generate the magnetic vector potential  $\mathbf{A} = (y^2\hat{i} + x^2\hat{j})$ ?

- (a)  $\frac{2}{\mu_0}(x\hat{i} + y\hat{j})$                       (c)  $\frac{2}{\mu_0}(\hat{i} - \hat{j})$   
 (b)  $-\frac{2}{\mu_0}(\hat{i} + \hat{j})$                       (d)  $\frac{2}{\mu_0}(x\hat{i} - y\hat{j})$

**Q. 23** The value of the integral  $\oint_C \frac{e^z}{z^2 - 3z + 2} dz$ , where the contour  $C$  is the circle  $|z| = 3/2$  is

- (a)  $2\pi ie$                       (b)  $\pi ie$                       (c)  $-2\pi ie$                       (d)  $-\pi ie$

**Q. 24** In a non-conducting medium characterized by  $\epsilon = \epsilon_0$ ,  $\mu = \mu_0$  and conductivity  $\sigma = 0$ , the electric field (in  $V\ m^{-1}$ ) is given by  $\mathbf{E} = 20 \sin[10^8 t - kz]\hat{j}$ . The magnetic field,  $\mathbf{H}$  (in  $A\ m^{-1}$ ), is given by

- (a)  $20k \cos[10^8 t - kz]\hat{i}$   
 (b)  $\frac{20k}{10^8 \mu_0} \sin[10^8 t - kz]\hat{j}$   
 (c)  $-\frac{20k}{10^8 \mu_0} \sin[10^8 t - kz]\hat{i}$   
 (d)  $-20k \cos[10^8 t - kz]\hat{j}$

**Q. 25** A cylindrical rod of length  $L$  and radius  $r$ , made of an inhomogeneous dielectric, is placed with its axis along the  $z$  direction with one end at the origin as shown below.




Figure 2: Cylindrical rod of length  $L$  and radius  $r$

If the rod carries a polarization,  $\mathbf{P} = (5z^2 + 7)\hat{k}$ , the volume bound charge inside the dielectric is

- (a) Zero      (b)  $10\pi r^2 L$       (c)  $-5\pi r^2 L$       (d)  $-5\pi r^2 L^2$

**Q. 26** Let  $T_{ij} = \sum_k \epsilon_{ijk} a_k$  and  $\beta_k = \sum_{i,j} \epsilon_{ijk} T_{ij}$ , where  $\epsilon_{ijk}$  is the Levi-Civita density, defined to be zero if two of the indices coincide and +1 and -1 depending on whether  $ijk$  is even or odd permutation of 1, 2, 3. Then  $\beta_3$  is equal to

- (a)  $2a_3$       (b)  $-2a_3$       (c)  $a_3$       (d)  $-a_3$

**Q. 27** The dependence of the magnetic susceptibility ( $\chi$ ) of a material with temperature ( $T$ ) can be represented by  $\chi \propto \frac{1}{T-\theta}$ , where  $\theta$  is the Curie-Weiss temperature. The plot of magnetic susceptibility versus temperature is sketched in the figure, as curves P, Q and R with curve Q having  $\theta = 0$ . Which one of the following statements is correct?




Figure 3: Magnetic susceptibility ( $\chi$ ) of a material vs. Temperature

- (a) Curve R represents a paramagnet and Q a ferromagnet  
 (b) Curve Q represents a ferromagnet and P an antiferromagnet

- (c) Curve R represents an antiferromagnet and Q a paramagnet
- (d) Curve R represents an antiferromagnet and Q a ferromagnet

**Q. 28** The dielectric constant of a material at optical frequencies is mainly due to

- (a) ionic polarizability
- (b) electronic polarizability
- (c) dipolar polarizability
- (d) ionic and dipolar polarizability

**Q. 29** An electron of wavevector  $\mathbf{k}_e$ , velocity  $\mathbf{v}_e$  and effective mass  $m_e$  is removed from a filled energy band. The resulting hole has wavevector  $\mathbf{k}_h$ , velocity  $\mathbf{v}_h$  and effective mass  $m_h$ . Which one of the following statements is correct?

- (a)  $\mathbf{k}_h = -\mathbf{k}_e$ ;  $\mathbf{v}_h = -\mathbf{v}_e$ ;  $m_h = -m_e$
- (b)  $\mathbf{k}_h = \mathbf{k}_e$ ;  $\mathbf{v}_h = \mathbf{v}_e$ ;  $m_h = m_e$
- (c)  $\mathbf{k}_h = \mathbf{k}_e$ ;  $\mathbf{v}_h = -\mathbf{v}_e$ ;  $m_h = -m_e$
- (d)  $\mathbf{k}_h = -\mathbf{k}_e$ ;  $\mathbf{v}_h = \mathbf{v}_e$ ;  $m_h = -m_e$



- Q. 30** In a diatomic molecule, the internuclear separation of the ground and first excited electronic state are the same as shown in the figure. If the molecule is initially in the lowest vibrational state of the ground state, then the absorption spectrum will appear as





Figure 4: Potential energy curves for a diatomic molecule



- Q. 31** Five energy levels of a system including the ground state are shown below. Their lifetimes and the allowed electric dipole transitions are also marked.




Figure 5: Energy levels of a system

Which one of the following transitions is the most suitable for a continuous wave (CW) laser?

- (a)  $1 \rightarrow 0$       (b)  $2 \rightarrow 0$       (c)  $4 \rightarrow 2$       (d)  $4 \rightarrow 3$
- Q. 32** Assuming the mean life time of a muon (in its rest frame) to be  $2 \times 10^{-6} s$ , its life time in the laboratory frame, when it is moving with a velocity  $0.95c$  is

- (a)  $6.4 \times 10^{-6} s$       (c)  $2.16 \times 10^{-6} s$   
 (b)  $0.62 \times 10^{-6} s$       (d)  $0.19 \times 10^{-6} s$

- Q. 33** Cesium has a nuclear spin of  $7/2$ . The hyperfine spectrum of the D lines of the cesium atom will consist of

- (a) 10 lines      (b) 4 lines      (c) 6 lines      (d) 14 lines

- Q. 34** The probability that an energy level  $\varepsilon$  at a temperature  $T$  is unoccupied by a fermion of chemical potential  $\mu$  is given by

- (a)  $\frac{1}{e^{(\varepsilon-\mu)/k_B T} + 1}$       (c)  $\frac{1}{e^{(\mu-\varepsilon)/k_B T} + 1}$   
 (b)  $\frac{1}{e^{(\varepsilon-\mu)/k_B T} - 1}$       (d)  $\frac{1}{e^{(\mu-\varepsilon)/k_B T} - 1}$

- Q. 35** Consider the following expression for the mass of a nucleus with  $Z$  protons and  $A$  nucleons:  $M(A, Z) = \frac{1}{c^2}(f(A) + yZ + zZ^2)$  Here  $f(A)$  is a

function of  $A$ ,

$$y = -4a_A,$$

$$z = a_c A^{-1/3} + 4a_A A^{-1},$$

$a_A$  and  $a_c$  are constants of suitable dimensions. For a fixed  $A$ , the expression of  $Z$  for the most stable nucleus is

<p>(a) <math>Z = \frac{A/2}{1 + \left(\frac{a_c}{a_A}\right) A^{2/3}}</math></p> <p>(b) <math>Z = \frac{A/2}{1 + \left(\frac{a_c}{4a_A}\right) A^{2/3}}</math></p>	<p>(c) <math>Z = \frac{A}{1 + \left(\frac{a_c}{4a_A}\right) A^{2/3}}</math></p> <p>(d) <math>Z = \frac{A}{1 + A^{2/3}}</math></p>
--	---

**Q. 36** The de Broglie wavelength of particles of mass  $m$  with average momentum  $p$  at a temperature  $T$  in three dimensions is given by

<p>(a) <math>\lambda = \frac{h}{\sqrt{2mk_B T}}</math></p> <p>(b) <math>\lambda = \frac{h}{\sqrt{3mk_B T}}</math></p>	<p>(c) <math>\lambda = \frac{h}{\sqrt{2k_B T}}</math></p> <p>(d) <math>\lambda = \frac{h}{\sqrt{3m}}</math></p>
---	---

**Q. 37**




Figure 6: Circuit for Question 37.

Assuming an ideal voltage source, Thevenin's resistance and Thevenin's voltage respectively for the above circuit are

- (a)  $15\ \Omega$  and  $7.5\ \text{V}$  (c)  $10\ \Omega$  and  $10\ \text{V}$   
 (b)  $20\ \Omega$  and  $5\ \text{V}$  (d)  $30\ \Omega$  and  $15\ \text{V}$

**Q. 38** Let  $|n\rangle$  and  $|p\rangle$  denote the isospin states with  $I = 1/2$ ,  $I_3 = 1/2$  and  $I = 1/2$ ,  $I_3 = -1/2$  of a nucleon respectively. Which one of the following two-nucleon states has  $I = 0$ ,  $I_3 = 0$ ?

- (a)  $\frac{1}{\sqrt{2}}(|nn\rangle - |pp\rangle)$  (c)  $\frac{1}{\sqrt{2}}(|np\rangle - |pn\rangle)$   
 (b)  $\frac{1}{\sqrt{2}}(|nn\rangle + |pp\rangle)$  (d)  $\frac{1}{\sqrt{2}}(|np\rangle + |pn\rangle)$

**Q. 39** An amplifier of gain 1000 is made into a feedback amplifier by feeding 9.9 % of its output voltage in series with the input opposing. If  $f_L = 20\ \text{Hz}$  and  $f_H = 200\ \text{kHz}$  for the amplifier without feedback, then due to the feedback [cite: 8]

- (a) the gain decreases by 10 times (as) the  $f_H$  increases by 100 times [cite: 8]  
 (b) the output resistance increases (as) the input resistance decreases by 100 times [cite: 8]  
 by 10 times [cite: 8]

**Q. 40**




Figure 7: Circuit for Question 40.

Pick the correct statement based on the above circuit.

- (a) The maximum Zener current,  $I_{z(\max)}$ , when  $R_L = 10\ \text{k}\Omega$  is  $15\ \text{mA}$   
 (b) The minimum Zener current,  $I_{z(\min)}$ , when  $R_L = 10\ \text{k}\Omega$  is  $5\ \text{mA}$



**Q. 45** Consider the set of vectors in three-dimensional real vector space  $\mathbf{R}^3$ ,  $S = \{(1, 1, 1), (1, -1, 1), (1, 1, -1)\}$ . Which one of the following statements is true?

- (a)  $S$  is not a linearly independent set.
- (b)  $S$  is a basis for  $\mathbf{R}^3$ .
- (c) The vectors in  $S$  are orthogonal.
- (d) An orthogonal set of vectors cannot be generated from  $S$ .

**Q. 46** For a Fermi gas of  $N$  particles in three dimensions at  $T = 0$  K, the Fermi energy,  $E_F$ , is proportional to

- (a)  $N^{2/3}$                       (b)  $N^{3/2}$                       (c)  $N^3$                       (d)  $N^2$

**Q. 47** The Lagrangian of a diatomic molecule is given by  $L = \frac{m}{2}(\dot{x}_1^2 + \dot{x}_2^2) - \frac{k}{2}x_1x_2$ , where  $m$  is the mass of each of the atoms and  $x_1$  and  $x_2$  are the displacements of atoms measured from the equilibrium position and  $k > 0$ . The normal frequencies are

- (a)  $\pm \left(\frac{k}{m}\right)^{1/2}$                       (b)  $\pm \left(\frac{k}{m}\right)^{1/4}$                       (c)  $\pm \left(\frac{k}{2m}\right)^{1/4}$                       (d)  $\pm \left(\frac{k}{2m}\right)^{1/2}$

**Q. 48** A particle is in the normalized state  $|\psi\rangle$  which is a superposition of the energy eigenstates  $|E_o = 10 \text{ eV}\rangle$  and  $|E_1 = 30 \text{ eV}\rangle$ . The average value of energy of the particle in the state  $|\psi\rangle$  is 20 eV. The state  $|\psi\rangle$  is given by

- (a)  $\frac{1}{2}|E_o = 10 \text{ eV}\rangle + \frac{\sqrt{3}}{4}|E_1 = 30 \text{ eV}\rangle$
- (b)  $\frac{1}{\sqrt{3}}|E_o = 10 \text{ eV}\rangle + \sqrt{\frac{2}{3}}|E_1 = 30 \text{ eV}\rangle$
- (c)  $\frac{1}{2}|E_o = 10 \text{ eV}\rangle - \frac{\sqrt{3}}{4}|E_1 = 30 \text{ eV}\rangle$
- (d)  $\frac{1}{\sqrt{2}}|E_o = 10 \text{ eV}\rangle - \frac{1}{\sqrt{2}}|E_1 = 30 \text{ eV}\rangle$

**Q. 49** The Lagrangian of a particle of mass  $m$  moving in one dimension is  $L = \exp(\alpha t) \left[ \frac{m\dot{x}^2}{2} - \frac{kx^2}{2} \right]$ , where  $\alpha$  and  $k$  are positive constants. The equation of motion of the particle is

(a)  $\ddot{x} + \alpha\dot{x} = 0$

(c)  $\ddot{x} - \alpha\dot{x} + \frac{k}{m}x = 0$

(b)  $\ddot{x} + \frac{k}{m}x = 0$

(d)  $\ddot{x} + \alpha\dot{x} + \frac{k}{m}x = 0$

**Q. 50** Two monochromatic waves having frequencies  $\omega$  and  $\omega + \Delta\omega$  ( $\Delta\omega \ll \omega$ ) and corresponding wavelengths  $\lambda$  and  $\lambda - \Delta\lambda$  ( $\Delta\lambda \ll \lambda$ ) of same polarization, traveling along x-axis are superimposed on each other. The phase velocity and group velocity of the resultant wave are respectively given by

(a)  $\frac{\omega\lambda}{2\pi}, \frac{\Delta\omega\lambda^2}{2\pi\Delta\lambda}$

(c)  $\frac{\omega\Delta\lambda}{2\pi}, \frac{\Delta\omega\Delta\lambda}{2\pi}$

(b)  $\omega\lambda, \frac{\Delta\omega\lambda^2}{\Delta\lambda}$

(d)  $\omega\lambda, \omega\Delta\lambda$

### Common Data Questions

#### Common Data for Questions 51 and 52:

Consider a two level quantum system with energies  $\varepsilon_1 = 0$  and  $\varepsilon_2 = \varepsilon$ .

**Q. 51** The Helmholtz free energy of the system is given by

(a)  $-k_B T \ln(1 + e^{-\varepsilon/k_B T})$

(c)  $\frac{3}{2}k_B T$

(b)  $k_B T \ln(1 + e^{-\varepsilon/k_B T})$

(d)  $\varepsilon - k_B T$

**Q. 52** The specific heat of the system is given by

(a)  $\frac{\varepsilon}{k_B T} \frac{e^{-\varepsilon/k_B T}}{(1 + e^{-\varepsilon/k_B T})^2}$

(c)  $-\frac{\varepsilon^2 e^{-\varepsilon/k_B T}}{(1 + e^{-\varepsilon/k_B T})^2}$

(b)  $\frac{\varepsilon^2}{k_B T^2} \frac{e^{-\varepsilon/k_B T}}{1 + e^{-\varepsilon/k_B T}}$

(d)  $\frac{\varepsilon^2}{k_B T^2} \frac{e^{-\varepsilon/k_B T}}{(1 + e^{-\varepsilon/k_B T})^2}$

#### Common Data for Questions 53 and 54:

A free particle of mass  $m$  moves along the x direction. At  $t = 0$ , the normalized wave function of the particle is given by  $\psi(x, 0) = \frac{1}{(2\pi\alpha)^{1/4}} \exp\left[-\frac{x^2}{4\alpha^2} + ix\right]$ , where  $\alpha$  is a real constant.

**Q. 53** The expectation value of the momentum, in this state is

- (a)  $\hbar\alpha$                       (b)  $\hbar\sqrt{\alpha}$                       (c)  $\alpha$                       (d)  $\frac{\hbar}{\sqrt{\alpha}}$

**Q. 54** The expectation value of the particle energy is

- (a)  $\frac{\hbar^2}{2m} \frac{1}{2\alpha^{3/2}}$                       (c)  $\frac{\hbar^2}{2m} \frac{4\alpha^2 + 1}{4\alpha^{3/2}}$   
 (b)  $\frac{\hbar^2}{2m} \alpha^2$                       (d)  $\frac{\hbar^2}{8m\alpha^{3/2}}$

**Common Data for Questions 55 and 56:**

Consider the Zeeman splitting of a single electron system for the  $3d \rightarrow 3p$  electric dipole transition.

**Q. 55** The Zeeman spectrum is

- (a) randomly polarized                      (c) only  $\sigma$  polarized  
 (b) only  $\pi$  polarized                      (d) both  $\pi$  and  $\sigma$  polarized

**Q. 56** The fine structure line having the longest wavelength will split into

- (a) 17 components                      (c) 8 components  
 (b) 10 components                      (d) 4 components

**Linked Answer Questions**

**Statement for Linked Answer Questions 57 and 58:**

The primitive translation vectors of the face centered cubic (fcc) lattice are  $\mathbf{a}_1 = \frac{a}{2}(\hat{j} + \hat{k})$ ;  $\mathbf{a}_2 = \frac{a}{2}(\hat{i} + \hat{k})$ ;  $\mathbf{a}_3 = \frac{a}{2}(\hat{i} + \hat{j})$ .

**Q. 57** The primitive translation vectors of the fcc reciprocal lattice are

- (a)  $\mathbf{b}_1 = \frac{2\pi}{a}(-\hat{i} + \hat{j} + \hat{k})$ ;  $\mathbf{b}_2 = \frac{2\pi}{a}(\hat{i} - \hat{j} + \hat{k})$ ;  $\mathbf{b}_3 = \frac{2\pi}{a}(\hat{i} + \hat{j} - \hat{k})$   
 (b)  $\mathbf{b}_1 = \frac{\pi}{a}(-\hat{i} + \hat{j} + \hat{k})$ ;  $\mathbf{b}_2 = \frac{\pi}{a}(\hat{i} - \hat{j} + \hat{k})$ ;  $\mathbf{b}_3 = \frac{\pi}{a}(\hat{i} + \hat{j} - \hat{k})$   
 (c)  $\mathbf{b}_1 = \frac{\pi}{2a}(-\hat{i} + \hat{j} + \hat{k})$ ;  $\mathbf{b}_2 = \frac{\pi}{2a}(\hat{i} - \hat{j} + \hat{k})$ ;  $\mathbf{b}_3 = \frac{\pi}{2a}(\hat{i} + \hat{j} - \hat{k})$   
 (d)  $\mathbf{b}_1 = \frac{3\pi}{a}(-\hat{i} + \hat{j} + \hat{k})$ ;  $\mathbf{b}_2 = \frac{3\pi}{a}(\hat{i} - \hat{j} + \hat{k})$ ;  $\mathbf{b}_3 = \frac{3\pi}{a}(\hat{i} + \hat{j} - \hat{k})$

**Q. 58** The volume of the primitive cell of the fcc reciprocal lattice is



(a)  $4\left(\frac{2\pi}{a}\right)^3$       (b)  $4\left(\frac{\pi}{a}\right)^3$       (c)  $4\left(\frac{\pi}{2a}\right)^3$       (d)  $4\left(\frac{3\pi}{a}\right)^3$

**Statement for Linked Answer Questions 59 and 60:**

The Karnaugh map of a logic circuit is shown below:


	$\bar{R}$	$R$
$\bar{P}\bar{Q}$	1	1
$\bar{P}Q$	1	
$PQ$		
$P\bar{Q}$	1	1

Figure 8: Karnaugh map of a logic circuit.

**Q. 59** The minimized logic expression for the above map is

- (a)  $Y = \bar{P}\bar{R} + \bar{Q}$       (c)  $Y = \bar{Q} + PR$   
 (b)  $Y = \bar{Q} \cdot PR$       (d)  $Y = Q \cdot \bar{P}\bar{R}$

**Q. 60** The corresponding logic implementation using gates is given as:



**END OF THE QUESTION PAPER**