

<!DOCTYPE DOCUMENT SYSTEM "icon.dtd" [ ]>

<DOCUMENT author="perl program" datec="020199" title="leaflet" book="ABPI"  
edition="1996-97" section="company">

<PB n=1>

<CB n=1>

<FIGURE></FIGURE>

<CB n=2>

<DIV type=0>

<HEAD weight="bold" >

Patient Information Leaflet on Amoxil syrupon Amoxil

</HEAD>

<BLOCK>

Amoxil syrup is prescribed mainly for children, but may also be prescribed for adults. This leaflet is written with child patients and their parents in mind; but the information enclosed is just as important if you are taking the medicine yourself, as an adult, or are helping another adult to take it.

</BLOCK>

<BLOCK>

Please read this leaflet before this medicine is started. It is essential to follow your doctor's advice. If there is anything you do not understand, please ask your doctor or pharmacist (chemist).

</BLOCK>

<BLOCK>

<EMPH slant = "italic">

Keep this leaflet until all the prescribed course of Amoxil has been finished. You may want to read it again.

</EMPH>

</BLOCK>

<DIV type=1>

<HEAD weight="bold" justified="center" >

The name of this medicine is AMOXIL

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

What is Amoxil?

</HEAD>

<BLOCK>

It is one of a group of antibiotic medicines called penicillins. The active ingredient in Amoxil is amoxycillin.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

What does Amoxil do?

</HEAD>

<BLOCK>

It is used to treat a range of bacterial infections.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

IMPORTANT THINGS TO KNOW ABOUT AMOXIL

</HEAD>

<LIST type="ordered">

<ITEM>

<BLOCK>

This medicine is unsuitable for people who are allergic to penicillin. (You will find more information about this later in the leaflet.)

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Give Amoxil to your child as directed by your doctor. Read the dispensing label .

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Keep giving your child Amoxil until the prescribed course is finished. Do not stop just because your child seems better.

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Like all medicines, Amoxil can sometimes cause unwanted effects. (You will find more information about this later in the leaflet.)

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

ALWAYS KEEP MEDICINES OUT OF THE REACH OF CHILDREN.

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

WOMEN: Before taking ANY medicines, you should tell our doctor if you are pregnant (or think you may be) or if you are breast feeding.

</BLOCK>

</ITEM>

</LIST>

<BLOCK>

<EMPH weight="bold">

Now turn over for details on how to give this medicine to your child.

</EMPH>

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

BEFORE GIVING YOUR CHILD THIS MEDICINE

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

Is my child allergic to penicillin?

</HEAD>

<BLOCK>

If you think your child may be allergic to penicillin, check with your doctor before giving this medicine.

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >  
GIVING YOUR CHILD THIS MEDICINE  
</HEAD>

<DIV type=2>

<HEAD weight="bold" >  
How much should I give my child and how often?  
</HEAD>

<BLOCK>  
Follow your doctor's directions about how and when to give your child Amoxil.  
Please read the dispensing label carefully. Your pharmacist or doctor can help if  
you are not sure.  
</BLOCK>

<BLOCK>  
Amoxil syrup should be given using the 5 ml spoon provided by the pharmacist. It  
is best for your child to take the medicine direct from the spoon. However, if  
necessary, the dose may be stirred into a few spoonfuls of cold water or infant  
drink (not milk). Make sure the whole dose is swallowed each time.  
</BLOCK>

<BLOCK>  
Space the doses as evenly as possible through the day.  
<EMPH slant = "italic">  
For example  
</EMPH>  
, if you have been told to give your child Amoxil 3 times a day:  
</BLOCK>

<PB n=2>  
<CB n=1>

<BLOCK justified=center>  
Dose 1 <CB n=2> Dose 2 <CB n=1> Dose 3 <PB n=3> <CB n=2> <FIGURE></FIGURE> <CB  
n=1> <FIGURE></FIGURE> <CB n=2> <FIGURE></FIGURE> <PB n=4> <CB n=1> about 7am <CB  
n=2> about 3pm <CB n=1> about 11pm  
</BLOCK>

<BLOCK>  
You may give Amoxil to your child before, with or after food, unless the label  
advises a specific time. It is not normally necessary to arrange to give your  
child a dose in the middle of the night unless your doctor has told you to do so.  
</BLOCK>

</DIV>  
<DIV type=2>

<HEAD weight="bold" >  
How long should I carry on giving it to my child?  
</HEAD>

<BLOCK>

Keep giving your child Amoxil until the prescribed course is finished. Do not stop just because your child seems better. If you stop too soon, the infection may start up again.

</BLOCK>

<BLOCK>

If your child still seems unwell at the end of the prescribed course of treatment, tell your doctor.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

What should I do if I forget to give my child a dose?

</HEAD>

<BLOCK>

If you forget to give your child a dose at the right time, take it as soon as you remember. Then carry on as before, but do not take two doses within an hour or so of each other.

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

AFTER GIVING YOUR CHILD THIS MEDICINE

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

What unwanted effects might it have?

</HEAD>

<BLOCK>

A few people may find that Amoxil gives them diarrhoea or indigestion. If this happens, these symptoms are usually mild. If your child suffers badly, tell your doctor.

</BLOCK>

<BLOCK>

Tell your doctor if your child develops rashes, or itching, or any other unusual problems, while they are taking this medicine.

</BLOCK>

<BLOCK>

If your doctor decides to stop the treatment, pour any left-over **medicine** down the sink.

</DIV>  
<DIV type=2>

<HEAD weight="bold" >  
What if I have given my child too much?  
</HEAD>

<BLOCK>  
If a large amount has been taken, give your child plenty of water to drink and consult your doctor or local hospital casualty department at once.  
</BLOCK>

</DIV>

</DIV>  
<DIV type=1>

<HEAD weight="bold" justified="center" >  
STORING **THIS MEDICINE**  
</HEAD>

<DIV type=2>

<HEAD weight="bold" >  
How should I store **Amoxil** **syrup**?  
</HEAD>

<BLOCK>  
Keep **it** in a dry place. Store all **medicines** out of the reach of children ☹ preferably in a locked cupboard or medicine cabinet.  
</BLOCK>

<BLOCK>

<EMPH slant = "italic">  
Remember  
</EMPH>

**this medicine** has been prescribed just for your child. Never offer your **medicine** to other people. **It** may not be suitable for them even if their symptoms seem the same as your child's.  
</BLOCK>

</DIV>

</DIV>  
<DIV type=1>

<HEAD weight="bold" justified="center" >  
MORE INFORMATION

</HEAD>

<BLOCK>

Amoxil syrup is available in two strengths containing either 125 mg or 250 mg of amoxycillin as amoxycillin trihydrate per 5 ml.

</BLOCK>

<BLOCK>

Non-proprietary name: Amoxycillin Syrup.

</BLOCK>

<BLOCK>

Amoxil syrup also contain inactive ingredients to make the medicine acceptable. Amongst these are sodium benzoate and sorbitol,

</BLOCK>

<BLOCK>

Manufacturer and Product Licence Holder and Manufacturer:

</BLOCK>

<BLOCK>

Beecham Group plc trading as Bencard, Great West Road,

</BLOCK>

<BLOCK>

Brentford, TW8 9BD, England.

</BLOCK>

<DIV type=2>

<HEAD weight="bold" >

How can I obtain more information about Amoxil?

</HEAD>

<BLOCK>

In this short leaflet, we can only give you the most important patient information about Amoxil. If you would like to know more, ask your doctor or pharmacist.

</BLOCK>

<FIGURE></FIGURE>

<BLOCK>

<EMPH weight="bold">

Amoxil

</EMPH>

and the Bencard logo are trade marks.

</BLOCK>

<BLOCK>

Leaflet prepared in October 1989.

</BLOCK>

</DIV>

</DIV>

</DIV>

</DOCUMENT>