

<!DOCTYPE DOCUMENT SYSTEM "icon.dtd" []>

<DOCUMENT author="perl program" datec="020199" title="leaflet" book="ABPI" edition="1996-97" section="company">

<PB n=1>

<CB n=1>

<FIGURE></FIGURE>

<CB n=2>

<DIV type=0>

<HEAD weight="bold" >

Patient Information Leaflet on Amoxil Paediatric Suspension

</HEAD>

<BLOCK>

Please read this leaflet before you start to take your medicine.

</BLOCK>

<BLOCK>

It is essential to follow your doctor's advice. If there is anything you do not understand, please ask your doctor or pharmacist (chemist).

</BLOCK>

<BLOCK>

<EMPH slant = "italic">

Keep this leaflet until your baby has finished all the prescribed course of Amoxil. You may want to read it again.

</EMPH>

</BLOCK>

<DIV type=1>

<HEAD weight="bold" justified="center" >

The name of your baby's medicine is AMOXIL

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

What is Amoxil?

</HEAD>

<BLOCK>

It is one of a group of antibiotic medicines called penicillins. The active ingredient in Amoxil is amoxycillin.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

What does Amoxil do?

</HEAD>

<BLOCK>

It is used to treat a range of bacterial infections.

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

IMPORTANT THINGS TO KNOW ABOUT AMOXIL

</HEAD>

<LIST type="ordered">

<ITEM>

<BLOCK>

This medicine is unsuitable for babies who are allergic to penicillin. (You will find more information about this later in the leaflet.)

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Give Amoxil to your baby as directed by your doctor. Read the dispensing label .

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Keep giving your baby Amoxil until the prescribed course is finished. Do not stop just because your baby seems better.

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Like all medicines, Amoxil can sometimes cause unwanted effects. (You will find more information about this later in the leaflet.)

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

ALWAYS KEEP MEDICINES OUT OF THE REACH OF CHILDREN.

</BLOCK>

</ITEM>

</LIST>

<BLOCK>

<EMPH weight="bold">

Now turn over for details on how to give this medicine to your baby.

</EMPH>

</BLOCK>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

BEFORE GIVING YOUR BABY THIS MEDICINE

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

Is my baby allergic to penicillin?

</HEAD>

<BLOCK>

If you think your baby may be allergic to penicillin, check with your doctor before giving this medicine.

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

GIVING YOUR BABY THIS MEDICINE

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

How much should I give my baby and how often?

</HEAD>

<BLOCK>

Follow your doctor's directions about how and when to give your baby Amoxil.

Please read the dispensing label carefully. Your pharmacist or doctor can help if you are not sure.

<BLOCK>

Space the doses as evenly as possible through the day.

<EMPH slant = "italic">

For example

</EMPH>

, if you have been told to give your baby Amoxil 3 times a day:

</BLOCK>

<PB n=2>

<CB n=1>

<BLOCK justified=center>

Dose 1 <CB n=2> Dose 2 <CB n=1> Dose 3 <PB n=3> <CB n=2> <FIGURE></FIGURE> <CB n=1> <FIGURE></FIGURE> <CB n=2> <FIGURE></FIGURE> <PB n=4> <CB n=1> about 7am <CB n=2> about 3pm <CB n=1> about 11pm

</BLOCK>

<BLOCK>

You may give Amoxil to your baby before, with or after food, unless the label advises a specific time. It is not normally necessary to arrange to give your baby a dose in the middle of the night unless your doctor has told you to do so.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

How long should I carry on giving it to my baby?

</HEAD>

<BLOCK>

Keep giving your baby Amoxil until the prescribed course is finished. Do not stop just because your baby seems better. If you stop too soon, the infection may start up again. If your baby still seems unwell at the end of the prescribed course of treatment, tell your doctor.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

What should I do if I forget to give my baby a dose?

</HEAD>

<BLOCK>

If you forget to give your baby a dose at the right time, take it as soon as you remember. Then carry on as before, but do not take two doses within an hour or so of each other.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

How should I give this medicine to my baby?

</HEAD>

<BLOCK>

<EMPH weight="bold">

NB

</EMPH>

</BLOCK>

<PB n=5>

<CB n=2>

<LIST type="ordered">

<ITEM>

<BLOCK>

Shake the bottle

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

<EMPH slant = "italic">

(a) If the dropper is not in the bottle :

</EMPH>

</BLOCK>

</ITEM>

</LIST>

<BLOCK>

First remove cap from bottle and discard cap. Then remove dropper from wrapping.

</BLOCK>

<BLOCK>

Follow steps 3, 4, 5, 6, 7 and 8.

</BLOCK>

<CB n=1>

<DIV type=3>

<HEAD slant="italic" >
(b) If the dropper is in the bottle:0
</HEAD>

<BLOCK>
Unscrew cap to remove dropper from bottle.
</BLOCK>

<BLOCK>
Follow steps 3, 4, 5, 6, 7 and 8.
</BLOCK>

<PB n=6>
<CB n=2>
<FIGURE></FIGURE>
<PB n=7>
<CB n=1>

<LIST type="ordered">

<ITEM>

<BLOCK>
Squeeze the rubber bulb tightly
</BLOCK>

</ITEM>

<ITEM>

<BLOCK>
Still squeezing the bulb, push the dropper into the medicine in the bottle.
</BLOCK>

</ITEM>

<ITEM>

<BLOCK>
Stop squeezing the bulb, the dropper will fill above the red line marked on it.
</BLOCK>

</ITEM>

<ITEM>

<BLOCK>
Take out the dropper. Squeeze the bulb gently until the medicine is exactly in line with red dose mark.
</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Put the medicine onto the baby's tongue by squeezing the bulb until all the medicine in the dropper has been pushed out. Do not put it on the front of the tongue or it might dribble out of the mouth.

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Wash out the dropper with water. Shake to remove any drops of water. Replace the dropper in the bottle and screw it on tightly.

</BLOCK>

</ITEM>

</LIST>

</DIV>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

AFTER GIVING YOUR BABY THIS MEDICINE

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

What unwanted effects might it have?

</HEAD>

<BLOCK>

A few people may find that Amoxil gives their baby diarrhoea or indigestion (colic). If this happens, the symptoms are usually mild. If this persists or if the baby is suffering badly, tell your doctor.

</BLOCK>

<BLOCK>

Tell your doctor if your baby develops a rash, or itching, or any other unusual problems, while you are giving this medicine.

</BLOCK>

<BLOCK>

If your doctor decides to stop the treatment, return any left-over medicine to the pharmacist.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

What if I have given my baby too much?

</HEAD>

<BLOCK>

If a large amount has been taken, give your baby plenty of water to drink and consult your doctor or local hospital casualty department at once.

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

STORING THIS MEDICINE

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

How should I store Amoxil paediatric suspension?

</HEAD>

<BLOCK>

Keep your baby's Amoxil paediatric suspension in a cool place. Store all medicines out of the reach of children - preferably in a locked cupboard or medicine cabinet.

</BLOCK>

<BLOCK>

<EMPH slant = "italic">

Remember

</EMPH>

this medicine has been prescribed just for your baby. Never offer this medicine to other people. It may not be suitable for them even if their symptoms seem the same as your baby's.

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

MORE INFORMATION

</HEAD>

<BLOCK>

Each 1.25 ml of Amoxil paediatric suspension contains amoxycillin trihydrate, equivalent to 125 mg of amoxycillin.

</BLOCK>

<BLOCK>

Non-proprietary name: Amoxycillin Paediatric Oral Suspension.

</BLOCK>

<BLOCK>

Amoxil paediatric suspension also contains inactive ingredients to make the medicine acceptable. Amongst these are sodium benzoate and sucrose.

</BLOCK>

<BLOCK>

Amoxil paediatric suspension does not contain tartrazine or any other azo dyes.

</BLOCK>

<BLOCK>

Manufacturer and Product Licence Holder:

</BLOCK>

<BLOCK>

Beecham Group plc trading as Bencard, Great West Road, Brentford, TW8 9BD, England.

</BLOCK>

<DIV type=2>

<HEAD weight="bold" >

How can I obtain more information about Amoxil?

</HEAD>

<BLOCK>

In this short leaflet, we can only give you the most important patient information about Amoxil. If you would like to know more, ask your doctor or pharmacist.

</BLOCK>

<FIGURE></FIGURE>

<BLOCK>

<EMPH weight="bold">

Amoxil

</EMPH>

and the Bencard logo are trade marks.

</BLOCK>

<BLOCK>

Leaflet prepared in August 1989.

</BLOCK>

</DIV>

</DIV>

</DIV>

</DOCUMENT>