

<!DOCTYPE DOCUMENT SYSTEM "icon.dtd" []>

<DOCUMENT author="perl program" datec="020199" title="leaflet" book="ABPI"
edition="1996-97" section="company">

<PB n=1>

<CB n=1>

<FIGURE></FIGURE>

<CB n=2>

<DIV type=0>

<HEAD weight="bold" >

Patient Information Leaflet on Amoxil capsuleson Amoxil

</HEAD>

<BLOCK>

Please read this leaflet before you start to take your medicine.

</BLOCK>

<BLOCK>

It is essential to follow the advice of your doctor (or dentist). If there is anything you do not understand, please ask your doctor or pharmacist (chemist).

<EMPH slant = "italic">

Keep this leaflet until you have finished all your prescribed course of Amoxil. You may want to read it again.

</EMPH>

</BLOCK>

<BLOCK>

If you are helping someone else to take this medicine, please read this leaflet before you give the first dose.

</BLOCK>

<DIV type=1>

<HEAD weight="bold" justified="center" >

The name of your medicine is AMOXIL

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

What is Amoxil?

</HEAD>

<BLOCK>

It is one of a group of antibiotic medicines called penicillins. The active ingredient in Amoxil is amoxycillin.

</BLOCK>

</DIV>

<DIV type=2>

<HEAD weight="bold" >

What does Amoxil do?

</HEAD>

<BLOCK>

It is used to treat a range of bacterial infections.

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

IMPORTANT THINGS TO KNOW ABOUT AMOXIL

</HEAD>

<LIST type="ordered">

<ITEM>

<BLOCK>

This medicine is unsuitable for people who are allergic to penicillin. (You will find more information about this later in the leaflet.)

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

<EMPH weight="bold">

Before taking ANY medicines, you should tell your doctor if you are pregnant (or think you may be), or if you are breast feeding.

</EMPH>

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Take Amoxil as directed by your doctor. Read the dispensing label .

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Keep taking your Amoxil until your prescribed course is finished. Do not stop just because you feel better.

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

Like all medicines, Amoxil can sometimes cause unwanted effects. (You will find more information about this later in the leaflet.)

</BLOCK>

</ITEM>

<ITEM>

<BLOCK>

ALWAYS KEEP MEDICINES OUT OF THE REACH OF CHILDREN.

</BLOCK>

</ITEM>

</LIST>

<BLOCK>

For more information about Amoxil, please read the rest of this leaflet.

</BLOCK>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

BEFORE TAKING YOUR MEDICINE

</HEAD>

<DIV type=2>

<HEAD weight="bold" >

Am I allergic to penicillin?

</HEAD>

<BLOCK>

Do you think or have you been told that you are allergic to penicillin? If so, do not take Amoxil; tell your doctor. You may be given another medicine instead.

</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >

TAKING YOUR MEDICINE

</HEAD>

<DIV type=2>

<HEAD weight="bold" >
How much should I take and how often?
</HEAD>

<BLOCK>
Follow your doctor's directions about how and when to take your medicine. Please read the dispensing label carefully. Your pharmacist or doctor can help if you are not sure. Amoxil capsules should be swallowed with water.
</BLOCK>

<BLOCK>
Space the doses as evenly as possible through the day.
<EMPH slant = "italic">
For example
</EMPH>
, if you have been told to take your Amoxil 3 times a day:
</BLOCK>

<PB n=2>
<CB n=1>

<BLOCK justified=center>
Dose 1 <CB n=2> Dose 2 <CB n=1> Dose 3 <PB n=3> <CB n=2> <FIGURE></FIGURE> <CB n=1> <FIGURE></FIGURE> <CB n=2> <FIGURE></FIGURE> <PB n=4> <CB n=1> about 7am <CB n=2> about 3pm <CB n=1> about 11pm
</BLOCK>

<BLOCK>
You may take Amoxil before, with or after food, unless the label advises a specific time. It is not normally necessary to arrange to take a dose in the middle of the night unless your doctor has told you to do so.
</BLOCK>

</DIV>
<DIV type=2>

<HEAD weight="bold" >
How long should I carry on taking it?
</HEAD>

<BLOCK>
Keep taking your Amoxil until your prescribed course is finished. Do not stop just because you feel better. If you stop too soon, the infection may start up again.
</BLOCK>

<BLOCK>
If you still feel unwell at the end of your prescribed course of treatment, tell your doctor.
</BLOCK>

</DIV>
<DIV type=2>

<HEAD weight="bold" >
What should I do if I forget to take a dose?
</HEAD>

<BLOCK>
If you forget to take a dose at the right time, take it as soon as you remember.
Then go on as before, but do not take two doses within an hour or so of each other.
</BLOCK>

</DIV>

</DIV>
<DIV type=1>

<HEAD weight="bold" justified="center" >
AFTER TAKING YOUR MEDICINE
</HEAD>

<DIV type=2>

<HEAD weight="bold" >
What unwanted effects might it have?
</HEAD>

<BLOCK>
A few people may find that Amoxil gives them diarrhoea or indigestion. If this happens, these symptoms are usually mild. If you suffer badly, tell your doctor.
</BLOCK>

<BLOCK>
Tell your doctor if you develop rashes, or itching, or any other unusual problems, while you are taking this medicine.
</BLOCK>

<BLOCK>
If your doctor decides to stop the treatment, return any left-over medicine to the pharmacist.
</BLOCK>

</DIV>
<DIV type=2>

<HEAD weight="bold" >
What if I have taken too much?
</HEAD>

<BLOCK>

If a large amount has been taken, drink plenty of water and consult your doctor or local hospital casualty department at once.

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >
STORING YOUR MEDICINE
</HEAD>

<DIV type=2>

<HEAD weight="bold" >
How should I store Amoxil capsules?
</HEAD>

<BLOCK>
Keep them in a dry place. Store all medicines out of the reach of children preferably in a locked cupboard or medicine cabinet.
</BLOCK>

<BLOCK>

<EMPH slant = "italic">
Remember
</EMPH>
this medicine has been prescribed just for you. Never offer your medicine to other people. It may not be suitable for them even if their symptoms seem the same as yours.
</BLOCK>

</DIV>

</DIV>

<DIV type=1>

<HEAD weight="bold" justified="center" >
MORE INFORMATION
</HEAD>

<BLOCK>
Amoxil capsules are available in two strengths containing either 250 mg or 500 mg of amoxycillin as amoxycillin trihydrate.
</BLOCK>

<BLOCK>
Non-proprietary name: Amoxycillin Capsules.
</BLOCK>

<BLOCK>

Amoxil capsules do not contain tartrazine or any other azo dyes.

</BLOCK>

<BLOCK>

Manufacturer and Product Licence Holder: Beecham Group plc trading as Bencard,
Great West Road, Brentford, TW8 9BD, England.

</BLOCK>

<DIV type=2>

<HEAD weight="bold" >

How can I obtain more information about Amoxil?

</HEAD>

<BLOCK>

In this short leaflet, we can only give you the most important patient information
about Amoxil. If you would like to know more, ask your doctor or pharmacist.

</BLOCK>

<FIGURE></FIGURE>

<BLOCK>

<EMPH weight="bold">

Amoxil

</EMPH>

and the Bencard logo are trade marks.

</BLOCK>

<BLOCK>

Leaflet prepared in August 1988.

</BLOCK>

</DIV>

</DIV>

</DIV>

</DOCUMENT>