Main Category	Category	Terms
Disease	OMIM_DISEASE	Breast cancer, lobular, somatic
Disease	OMIM_DISEASE	Blepharocheilodontic syndrome 1
Disease	OMIM_DISEASE	Diffuse gastric and lobular breast cancer syndrome with or without cleft lip and/or palate
Disease	OMIM_DISEASE	Ovarian cancer, somatic
Disease	OMIM_DISEASE	Endometrial carcinoma, somatic
Disease	UP_KW_DISEASE	Ectodermal dysplasia
Disease	UP_KW_DISEASE	Disease variant
Functional Annotations	UP_KW_BIOLOGICAL_PROCESS	Cell adhesion
Functional Annotations	UP_KW_CELLULAR_COMPONENT	Golgi apparatus
Functional Annotations	UP_KW_CELLULAR_COMPONENT	Membrane
Functional Annotations	UP_KW_CELLULAR_COMPONENT	Cytoplasm
Functional Annotations	UP_KW_CELLULAR_COMPONENT	Cell junction
Functional Annotations	UP_KW_CELLULAR_COMPONENT	Endosome
Functional Annotations	UP_KW_CELLULAR_COMPONENT	Cell membrane
Functional Annotations	UP_KW_PTM	Cleavage on pair of basic residues
Functional Annotations	UP_KW_PTM	Glycoprotein
Functional Annotations	UP_KW_PTM	Phosphoprotein
Functional Annotations	UP_KW_PTM	Ubl conjugation
Functional Annotations	UP_KW_PTM	Disulfide bond
Functional Annotations	UP_SEQ_FEATURE	CARBOHYD:N-linked (GlcNAc...) asparagine
Functional Annotations	UP_SEQ_FEATURE	CARBOHYD:O-linked (Man...) serine
Functional Annotations	UP_SEQ_FEATURE	CARBOHYD:O-linked (Man...) threonine
Functional Annotations	UP_SEQ_FEATURE	COMPBIAS:Acidic residues

Main Category	Category	Terms
Functional Annotations	UP_SEQ_FEATURE	COMPBIAS:Basic and acidic residues
Functional Annotations	UP_SEQ_FEATURE	DISULFID:Interchain
Functional Annotations	UP_SEQ_FEATURE	DOMAIN:Cadherin
Functional Annotations	UP_SEQ_FEATURE	DOMAIN:Cadherin 1
Functional Annotations	UP_SEQ_FEATURE	DOMAIN:Cadherin 2
Functional Annotations	UP_SEQ_FEATURE	DOMAIN:Cadherin 3
Functional Annotations	UP_SEQ_FEATURE	DOMAIN:Cadherin 4
Functional Annotations	UP_SEQ_FEATURE	DOMAIN:Cadherin 5
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:GGG->AAA: Binds to CTNNB1 but abolishes interaction of CTNNB1 with PSEN1. Abolishes gamma-secretase cleavage.
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:L->D: Reduces localization to desmosome cell-cell junctions. Also reduces the localization of DSG2 and DSP to desmosome cell-cell junctions.
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:N->Q: CDH1 becomes a substrate for ERAD and is retro-translocated from ER to cytoplasm.
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:P->A,E,V: No longer binds L.monocytogenes InlA.
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:P->E: No longer adheres to L.monocytogenes InlA coated beads, nor do cells take up InlA coated beads.
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:W->A,S: No longer binds L.monocytogenes InlA.
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:W->D: Protein is partially unfolded, no longer binds L.monocytogenes InlA.
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:Y->F: Abolishes binding to CBLL1.
Functional Annotations	UP_SEQ_FEATURE	REGION:Disordered
Functional Annotations	UP_SEQ_FEATURE	REGION:Required for binding CTNND1 and PSEN1
Functional Annotations	UP_SEQ_FEATURE	REGION:Required for binding alpha, beta and gamma catenins
Functional Annotations	UP_SEQ_FEATURE	SITE:Cleavage; by S.pyogenes SpeB

Main Category	Category	Terms
Functional Annotations	UP_SEQ_FEATURE	SITE:Cleavage; by a metalloproteinase
Functional Annotations	UP_SEQ_FEATURE	SITE:Cleavage; by caspase-3
Functional Annotations	UP_SEQ_FEATURE	SITE:Cleavage; by gamma-secretase/PS1
Functional Annotations	UP_SEQ_FEATURE	TOPO_DOM:Cytoplasmic
Functional Annotations	UP_SEQ_FEATURE	TOPO_DOM:Extracellular
Functional Annotations	UP_SEQ_FEATURE	TRANSMEM:Helical
Gene Ontology	GOTERM_BP_DIRECT	cell morphogenesis
Gene Ontology	GOTERM_BP_DIRECT	desmosome assembly
Gene Ontology	GOTERM_BP_DIRECT	cell-cell junction assembly
Gene Ontology	GOTERM_BP_DIRECT	cell adhesion
Gene Ontology	GOTERM_BP_DIRECT	homophilic cell-cell adhesion
Gene Ontology	GOTERM_BP_DIRECT	synapse assembly
Gene Ontology	GOTERM_BP_DIRECT	response to xenobiotic stimulus
Gene Ontology	GOTERM_BP_DIRECT	response to toxic substance
Gene Ontology	GOTERM_BP_DIRECT	regulation of gene expression
Gene Ontology	GOTERM_BP_DIRECT	calcium-dependent cell-cell adhesion
Gene Ontology	GOTERM_BP_DIRECT	cell migration
Gene Ontology	GOTERM_BP_DIRECT	pituitary gland development
Gene Ontology	GOTERM_BP_DIRECT	negative regulation of cell-cell adhesion
Gene Ontology	GOTERM_BP_DIRECT	negative regulation of cell migration
Gene Ontology	GOTERM_BP_DIRECT	negative regulation of axon extension

Main Category	Category	Terms
Gene Ontology	GOTERM_BP_DIRECT	neuron projection development
Gene Ontology	GOTERM_BP_DIRECT	cell junction assembly
Gene Ontology	GOTERM_BP_DIRECT	adherens junction organization
Gene Ontology	GOTERM_BP_DIRECT	positive regulation of protein import into nucleus
Gene Ontology	GOTERM_BP_DIRECT	cell-cell adhesion mediated by cadherin
Gene Ontology	GOTERM_BP_DIRECT	positive regulation of DNA-templated transcription
Gene Ontology	GOTERM_BP_DIRECT	regulation of transport
Gene Ontology	GOTERM_BP_DIRECT	cellular response to lithium ion
Gene Ontology	GOTERM_BP_DIRECT	response to heparin
Gene Ontology	GOTERM_BP_DIRECT	cellular response to indole-3-methanol
Gene Ontology	GOTERM_BP_DIRECT	protein localization to plasma membrane
Gene Ontology	GOTERM_BP_DIRECT	cell-cell adhesion
Gene Ontology	GOTERM_BP_DIRECT	regulation of protein catabolic process at postsynapse, modulating synaptic transmission
Gene Ontology	GOTERM_BP_DIRECT	response to Gram-positive bacterium
Gene Ontology	GOTERM_BP_DIRECT	positive regulation of protein localization
Gene Ontology	GOTERM_CC_DIRECT	extracellular region
Gene Ontology	GOTERM_CC_DIRECT	cytoplasm
Gene Ontology	GOTERM_CC_DIRECT	endosome
Gene Ontology	GOTERM_CC_DIRECT	Golgi apparatus
Gene Ontology	GOTERM_CC_DIRECT	trans-Golgi network
Gene Ontology	GOTERM_CC_DIRECT	plasma membrane

Main Category	Category	Terms
Gene Ontology	GOTERM_CC_DIRECT	adherens junction
Gene Ontology	GOTERM_CC_DIRECT	cytoplasmic side of plasma membrane
Gene Ontology	GOTERM_CC_DIRECT	actin cytoskeleton
Gene Ontology	GOTERM_CC_DIRECT	membrane
Gene Ontology	GOTERM_CC_DIRECT	lateral plasma membrane
Gene Ontology	GOTERM_CC_DIRECT	catenin complex
Gene Ontology	GOTERM_CC_DIRECT	flotillin complex
Gene Ontology	GOTERM_CC_DIRECT	lamellipodium
Gene Ontology	GOTERM_CC_DIRECT	cell junction
Gene Ontology	GOTERM_CC_DIRECT	desmosome
Gene Ontology	GOTERM_CC_DIRECT	cortical actin cytoskeleton
Gene Ontology	GOTERM_CC_DIRECT	apical junction complex
Gene Ontology	GOTERM_CC_DIRECT	perinuclear region of cytoplasm
Gene Ontology	GOTERM_CC_DIRECT	extracellular exosome
Gene Ontology	GOTERM_CC_DIRECT	anchoring junction
Gene Ontology	GOTERM_CC_DIRECT	postsynapse
Gene Ontology	GOTERM_CC_DIRECT	glutamatergic synapse
Gene Ontology	GOTERM_MF_DIRECT	calcium ion binding
Gene Ontology	GOTERM_MF_DIRECT	protein binding
Gene Ontology	GOTERM_MF_DIRECT	beta-catenin binding
Gene Ontology	GOTERM_MF_DIRECT	ankyrin binding

Main Category	Category	Terms
Gene Ontology	GOTERM_MF_DIRECT	GTPase activating protein binding
Gene Ontology	GOTERM_MF_DIRECT	identical protein binding
Gene Ontology	GOTERM_MF_DIRECT	gamma-catenin binding
Gene Ontology	GOTERM_MF_DIRECT	cadherin binding
Gene Ontology	GOTERM_MF_DIRECT	metal ion binding
Gene Ontology	GOTERM_MF_DIRECT	cell adhesion molecule binding
Gene Ontology	GOTERM_MF_DIRECT	cell-cell adhesion mediator activity
Gene Ontology	GOTERM_MF_DIRECT	protein tyrosine kinase binding
General Annotations	CHROMOSOME	16
General Annotations	CYTOGENETIC_LOCATION	16q22.1
General Annotations	ENTREZ_GENE_SUMMARY	This gene encodes a classical cadherin of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion protein is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function of this gene is thought to contribute to cancer progression by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. This gene is present in a gene cluster with other members of the cadherin family on chromosome 16. [provided by RefSeq, Nov 2015]
General Annotations	UP_COMMENT_ALTERNATIVE_PRODUCTS	Event=Alternative splicing; Named isoforms=2; Name=1; IsoId=P12830-1; Sequence=Displayed; Name=2; IsoId=P12830-2; Sequence=VSP_055586;
General Annotations	UP_COMMENT_DISEASE	Blepharocheilodontic syndrome 1 (BCDS1) [MIM:119580]: A form of blepharocheilodontic syndrome, a rare autosomal dominant disorder. It is characterized by lower eyelid ectropion, upper eyelid distichiasis, euryblepharon, bilateral cleft lip and palate, and features of ectodermal dysplasia, including hair anomalies, conical teeth and tooth agenesis. An additional rare manifestation is imperforate anus. There is considerable phenotypic variability among affected

Main Category	Category	Terms
		individuals. {ECO:0000269|PubMed:28301459}. Note=The disease is caused by variants affecting the gene represented in this entry.
General Annotations	UP_COMMENT_DISEASE	Breast cancer, lobular (LBC) [MIM:137215]: A type of breast cancer that begins in the milk-producing glands (lobules) of the breast. {ECO:0000269|PubMed:17660459}. Note=The gene represented in this entry may be involved in disease pathogenesis.
General Annotations	UP_COMMENT_DISEASE	Diffuse gastric and lobular breast cancer syndrome (DGLBC) [MIM:137215]: A cancer predisposition syndrome with increased susceptibility to diffuse gastric cancer. Diffuse gastric cancer is a malignant disease characterized by poorly differentiated infiltrating lesions resulting in thickening of the stomach. Malignant tumors start in the stomach, can spread to the esophagus or the small intestine, and can extend through the stomach wall to nearby lymph nodes and organs. It also can metastasize to other parts of the body. In addition to gastric cancer, most female mutation carriers develop lobular carcinoma of the breast. {ECO:0000269|PubMed:10319582, ECO:0000269|PubMed:12216071, ECO:0000269|PubMed:9537325}. Note=Disease susceptibility is associated with variants affecting the gene represented in this entry. Heterozygous CDH1 germline mutations are responsible for familial cases of diffuse gastric cancer. Somatic mutations has also been found in patients with sporadic diffuse gastric cancer and lobular breast cancer.
General Annotations	UP_COMMENT_DISEASE	Endometrial cancer (ENDMC) [MIM:608089]: A malignancy of endometrium, the mucous lining of the uterus. Most endometrial cancers are adenocarcinomas, cancers that begin in cells that make and release mucus and other fluids. Note=Disease susceptibility is associated with variants affecting the gene represented in this entry.
General Annotations	UP_COMMENT_DISEASE	Ovarian cancer (OC) [MIM:167000]: The term ovarian cancer defines malignancies originating from ovarian tissue. Although many histologic types of ovarian tumors have been described, epithelial ovarian carcinoma is the most common form. Ovarian cancers are often asymptomatic and the recognized signs and symptoms, even of late-stage disease, are vague. Consequently, most patients are diagnosed with advanced disease. {ECO:0000269|PubMed:8075649}. Note=Disease susceptibility is associated with variants affecting the gene represented in this entry.
General Annotations	UP_COMMENT_DOMAIN	Three calcium ions are usually bound at the interface of each cadherin domain and strengthen the connections, imparting a strong curvature to the full-length ectodomain. {ECO:0000269|PubMed:2349235}.
General Annotations	UP_COMMENT_FUNCTION	(Microbial infection) Serves as a receptor for Listeria monocytogenes; internalin A (InlA) binds to this protein and promotes uptake of the bacteria. {ECO:0000269|PubMed:10406800, ECO:0000269|PubMed:17540170,

Main Category	Category	Terms
		ECO:0000269|PubMed:8601315}.
General Annotations	UP_COMMENT_FUNCTION	Cadherins are calcium-dependent cell adhesion proteins (PubMed:11976333). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells (PubMed:11976333). Promotes organization of radial actin fiber structure and cellular response to contractile forces, via its interaction with AMOTL2 which facilitates anchoring of radial actin fibers to CDH1 junction complexes at the cell membrane (By similarity). Plays a role in the early stages of desmosome cell-cell junction formation via facilitating the recruitment of DSG2 and DSP to desmosome plaques (PubMed:29999492). Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7. {ECO:0000250|UniProtKB:F1PAA9, ECO:0000269|PubMed:11976333, ECO:0000269|PubMed:16417575, ECO:0000269|PubMed:29999492}.
General Annotations	UP_COMMENT_FUNCTION	Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells. Promotes organization of radial actin fiber structure and cellular response to contractile forces, via its interaction with AMOTL2 which facilitates anchoring of radial actin fibers to CDH1 junction complexes at the cell membrane. Plays a role in the early stages of desmosome cell-cell junction formation via facilitating the recruitment of DSG2 and DSP to desmosome plaques. Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7. {ECO:0000256|ARBA:ARBA00059145}.
General Annotations	UP_COMMENT_FUNCTION	E-Cad/CTF2 promotes non-amyloidogenic degradation of Abeta precursors. Has a strong inhibitory effect on APP C99 and C83 production. {ECO:0000256|ARBA:ARBA00025086}.
General Annotations	UP_COMMENT_FUNCTION	E-Cad/CTF2 promotes non-amyloidogenic degradation of Abeta precursors. Has a strong inhibitory effect on APP C99 and C83 production. {ECO:0000269|PubMed:16417575}.
General Annotations	UP_COMMENT_INDUCTION	Induced in the hours following cyclic mechanical strain in keratinocytes (PubMed:31835537). Expression is repressed by MACROD1 (PubMed:17893710). {ECO:0000269|PubMed:17893710, ECO:0000269|PubMed:31835537}.
General Annotations	UP_COMMENT_INTERACTION	A5D8W4; Q9NP61: ARFGAP3; NbExp=3; IntAct=EBI-7793316, EBI-2875816; A5D8W4; P01106: MYC; NbExp=3; IntAct=EBI-7793316, EBI-447544; A5D8W4; Q13309: SKP2; NbExp=2; IntAct=EBI-7793316, EBI-456291;

Main Category	Category	Terms
General Annotations	UP_COMMENT_INTERACTION	P12830; O94983-5: CAMTA2; NbExp=3; IntAct=EBI-727477, EBI-10176008; P12830; P30260: CDC27; NbExp=2; IntAct=EBI-727477, EBI-994813; P12830; P12830: CDH1; NbExp=7; IntAct=EBI-727477, EBI-727477; P12830; Q6UXH1-2: CRELD2; NbExp=3; IntAct=EBI-727477, EBI-21670927; P12830; P35221: CTNNA1; NbExp=8; IntAct=EBI-727477, EBI-701918; P12830; P35222: CTNNB1; NbExp=22; IntAct=EBI-727477, EBI-491549; P12830; O60716: CTNND1; NbExp=9; IntAct=EBI-727477, EBI-701927; P12830; P49184: DNASE1L1; NbExp=3; IntAct=EBI-727477, EBI-20894690; P12830; Q9H8V3: ECT2; NbExp=3; IntAct=EBI-727477, EBI-1054039; P12830; P00533: EGFR; NbExp=4; IntAct=EBI-727477, EBI-297353; P12830; P16422: EPCAM; NbExp=3; IntAct=EBI-727477, EBI-1171184; P12830; P11362: FGFR1; NbExp=3; IntAct=EBI-727477, EBI-1028277; P12830; Q7Z6Z7-2: HUWE1; NbExp=3; IntAct=EBI-727477, EBI-10975491; P12830; Q9UHB6: LIMA1; NbExp=2; IntAct=EBI-727477, EBI-351479; P12830; P43356: MAGEA2B; NbExp=3; IntAct=EBI-727477, EBI-5650739; P12830; Q7Z434: MAVS; NbExp=3; IntAct=EBI-727477, EBI-995373; P12830; P08581: MET; NbExp=2; IntAct=EBI-727477, EBI-1039152; P12830; Q8NCR3: MFI; NbExp=3; IntAct=EBI-727477, EBI-744790; P12830; Q8TB02: MGC39372; NbExp=3; IntAct=EBI-727477, EBI-25834188; P12830; Q9NPC7: MYNN; NbExp=3; IntAct=EBI-727477, EBI-3446748; P12830; Q9HC98-4: NEK6; NbExp=3; IntAct=EBI-727477, EBI-11750983; P12830; Q8N323: NXPE1; NbExp=3; IntAct=EBI-727477, EBI-25834085; P12830; Q8N2W9: PIAS4; NbExp=3; IntAct=EBI-727477, EBI-473160; P12830; Q8NBT0: POC1A; NbExp=3; IntAct=EBI-727477, EBI-2557132; P12830; P62136: PPP1CA; NbExp=2; IntAct=EBI-727477, EBI-357253; P12830; Q96QH2: PRAM1; NbExp=3; IntAct=EBI-727477, EBI-2860740; P12830; P17612: PRKACA; NbExp=3; IntAct=EBI-727477, EBI-476586; P12830; Q15139: PRKD1; NbExp=7; IntAct=EBI-727477, EBI-1181072; P12830; Q9Y617: PSAT1; NbExp=3; IntAct=EBI-727477, EBI-709652; P12830; P50454: SERPINH1; NbExp=3; IntAct=EBI-727477, EBI-350723; P12830; Q02978: SLC25A11; NbExp=3; IntAct=EBI-727477, EBI-359174; P12830; P12931: SRC; NbExp=2; IntAct=EBI-727477, EBI-621482; P12830; P48775: TDO2; NbExp=3; IntAct=EBI-727477, EBI-743494; P12830; P37173: TGFBR2; NbExp=3; IntAct=EBI-727477, EBI-296151; P12830; Q9BT49: THAP7; NbExp=3; IntAct=EBI-727477, EBI-741350; P12830; Q68CZ2: TNS3; NbExp=2; IntAct=EBI-727477, EBI-1220488; P12830; Q9ULW0: TPX2; NbExp=3; IntAct=EBI-727477, EBI-1037322; P12830; Q96EF9: ZHX1-C8orf76; NbExp=3; IntAct=EBI-727477, EBI-25830993; P12830; Q9BQ29; NbExp=3; IntAct=EBI-727477, EBI-22013570; P12830; A0A0S2C3P2: cagA; Xeno; NbExp=3; IntAct=EBI-727477, EBI-55025805; P12830; Q9JIY2: Cbll1; Xeno; NbExp=21; IntAct=EBI-727477, EBI-7644904; P12830; P0DJM0: inlA; Xeno; NbExp=3; IntAct=EBI-727477, EBI-1035388;

Main Category	Category	Terms
General Annotations	UP_COMMENT_MISCELLANEOUS	May play a role in blister formation in Pemphigus vulgaris patients, expression is lost at sites of cell-cell contact between the basal and suprabasal layers in blisters and immediately adjacent to blisters. {ECO:0000269|PubMed:22294297}.
General Annotations	UP_COMMENT_PTM	(Microbial infection) Cleaved by S.pyogenes SpeB protease; leading to its degradation (PubMed:23532847). Degradation by SpeB promotes bacterial translocation across the host epithelial barrier (PubMed:23532847). {ECO:0000269|PubMed:23532847}.
General Annotations	UP_COMMENT_PTM	During apoptosis or with calcium influx, cleaved by a membrane- bound metalloproteinase (ADAM10), PS1/gamma-secretase and caspase-3 (PubMed:10597309, PubMed:11076937, PubMed:11953314). Processing by the metalloproteinase, induced by calcium influx, causes disruption of cell-cell adhesion and the subsequent release of beta-catenin into the cytoplasm (PubMed:10597309). The residual membrane-tethered cleavage product is rapidly degraded via an intracellular proteolytic pathway (PubMed:10597309). Cleavage by caspase-3 releases the cytoplasmic tail resulting in disintegration of the actin microfilament system (PubMed:11076937). The gamma-secretase-mediated cleavage promotes disassembly of adherens junctions (PubMed:11953314). During development of the cochlear organ of Corti, cleavage by ADAM10 at adherens junctions promotes pillar cell separation (By similarity). {ECO:0000250|UniProtKB:P09803, ECO:0000269|PubMed:10597309, ECO:0000269|PubMed:11076937, ECO:0000269|PubMed:11953314}.
General Annotations	UP_COMMENT_PTM	N-glycosylation at Asn-637 is essential for expression, folding and trafficking. Addition of bisecting N-acetylglucosamine by MGAT3 modulates its cell membrane location (PubMed:19403558). {ECO:0000269|PubMed:18491227, ECO:0000269|PubMed:19403558}.
General Annotations	UP_COMMENT_PTM	O-glycosylated. O-manosylated by TMTC1, TMTC2, TMTC3 or TMTC4. Thr-285 and Thr-509 are O-mannosylated by TMTC2 or TMTC4 but not TMTC1 or TMTC3. {ECO:0000250|UniProtKB:P09803}.
General Annotations	UP_COMMENT_PTM	Ubiquitinated by a SCF complex containing SKP2, which requires prior phosphorylation by CK1/CSNK1A1. Ubiquitinated by CBLL1/HAKAI, requires prior phosphorylation at Tyr-754. {ECO:0000269|PubMed:21283129, ECO:0000269|PubMed:22252131, ECO:0000269|PubMed:22770219}.
General Annotations	UP_COMMENT_SEQUENCE_CAUTION	Sequence=AAA61259.1; Type=Frameshift; Evidence={ECO:0000305};
General Annotations	UP_COMMENT_SUBCELLULAR_LOCATION	Cell junction, adherens junction {ECO:0000256|ARBA:ARBA00004536}. Cell junction, desmosome {ECO:0000256|ARBA:ARBA00004568}. Cell membrane {ECO:0000256|ARBA:ARBA00004236}.

Main Category	Category	Terms
General Annotations	UP_COMMENT_SUBCELLULAR_LOCATION	Cell junction, adherens junction {ECO:0000256|ARBA:ARBA00004536}. Cell junction, desmosome {ECO:0000256|ARBA:ARBA00004568}. Cell membrane {ECO:0000256|ARBA:ARBA00004251, ECO:0000256|RuleBase:RU003318}; Single- pass type I membrane protein {ECO:0000256|ARBA:ARBA00004251, ECO:0000256|RuleBase:RU003318}. Cytoplasm {ECO:0000256|ARBA:ARBA00004496}. Endosome {ECO:0000256|ARBA:ARBA00004177}. Golgi apparatus, trans-Golgi network {ECO:0000256|ARBA:ARBA00004601}.
General Annotations	UP_COMMENT_SUBCELLULAR_LOCATION	Cell junction, adherens junction {ECO:0000256|ARBA:ARBA00004536}. Cell junction, desmosome {ECO:0000256|ARBA:ARBA00004568}. Cell membrane {ECO:0000256|ARBA:ARBA00004251}; Single-pass type I membrane protein {ECO:0000256|ARBA:ARBA00004251}.
General Annotations	UP_COMMENT_SUBCELLULAR_LOCATION	Cell junction, adherens junction {ECO:0000269|PubMed:18343367, ECO:0000269|PubMed:22294297, ECO:0000269|PubMed:27760340, ECO:0000269|PubMed:28169360}. Cell membrane {ECO:0000269|PubMed:19403558, ECO:0000269|PubMed:20859650, ECO:0000269|PubMed:28301459}; Single-pass type I membrane protein. Endosome {ECO:0000269|PubMed:15689490}. Golgi apparatus, trans-Golgi network {ECO:0000269|PubMed:15689490}. Cytoplasm {ECO:0000269|PubMed:22294297}. Cell junction, desmosome {ECO:0000269|PubMed:25208567, ECO:0000269|PubMed:29999492, ECO:0000269|PubMed:33596089}. Note=Colocalizes with DLGAP5 at sites of cell-cell contact in intestinal epithelial cells. Anchored to actin microfilaments through association with alpha-, beta- and gamma- catenin. Sequential proteolysis induced by apoptosis or calcium influx, results in translocation from sites of cell-cell contact to the cytoplasm. Colocalizes with RAB11A endosomes during its transport from the Golgi apparatus to the plasma membrane. Recruited to desmosomes at the initial assembly phase and also accumulates progressively at mature desmosome cell-cell junctions (PubMed:25208567, PubMed:29999492). Localizes to cell-cell contacts as keratinocyte differentiation progresses (By similarity). {ECO:0000250|UniProtKB:P09803, ECO:0000269|PubMed:25208567, ECO:0000269|PubMed:29999492}.
General Annotations	UP_COMMENT_SUBCELLULAR_LOCATION	Cell junction, desmosome {ECO:0000256|ARBA:ARBA00004568}. Cell membrane {ECO:0000256|ARBA:ARBA00004236}.
General Annotations	UP_COMMENT_SUBUNIT	(Microbial infection) Interacts with L.monocytogenes InlA (PubMed:12526809, PubMed:17540170, PubMed:17715295). The formation of the complex between InlA and cadherin-1 is calcium-dependent (PubMed:12526809). {ECO:0000269|PubMed:12526809, ECO:0000269|PubMed:17540170, ECO:0000269|PubMed:17715295}.
General	UP_COMMENT_SUBUNIT	Homodimer; disulfide-linked (PubMed:11856755).

Main Category	Category	Terms
Annotations		Component of an E-cadherin/ catenin adhesion complex composed of at least E- cadherin/CDH1, beta-catenin/CTNNB1 or gamma-catenin/JUP, and potentially alpha-catenin/CTNNA1; the complex is located to adherens junctions (PubMed:16126725, PubMed:7982500). Found in a complex composed of CDH1, RAP1A and PKP3; PKP3 acts as a scaffold protein within the complex, the complex is required for CDH1 localization to mature desmosome cell junctions (PubMed:25208567). Interacts with the TRPV4 and CTNNB1 complex (By similarity). Interacts with CTNND1 (PubMed:15240885). The stable association of CTNNA1 is controversial as CTNNA1 was shown not to bind to F-actin when assembled in the complex (By similarity). Alternatively, the CTNNA1-containing complex may be linked to F-actin by other proteins such as LIMA1 (By similarity). Interaction with PSEN1, cleaves CDH1 resulting in the disassociation of cadherin-based adherens junctions (CAJs) (PubMed:11226248, PubMed:16126725). Interacts with AJAP1 and DLGAP5 (PubMed:14595118, PubMed:14699157). Interacts with TBC1D2 (PubMed:20116244). Interacts with LIMA1 (PubMed:18093941). Interacts with CAV1. Interacts with PIP5K1C (PubMed:17261850). Interacts with RAB8B (By similarity). Interacts with RAPGEF2 (By similarity). Interacts with DDR1; this stabilizes CDH1 at the cell surface and inhibits its internalization (PubMed:20432435). Interacts with KLRG1 (PubMed:19604491). Forms a ternary complex composed of ADAM10, CADH1 and EPHA4; within the complex, CADH1 is cleaved by ADAM10 which disrupts adherens junctions (By similarity). Interacts with SPEF1 (PubMed:31473225). Interacts with CTNNB1 and PKP2 (PubMed:11790773). Interacts with AMOTL2; the interaction may facilitate binding of radial actin fibers to cell junction complexes (By similarity). Interacts with DSG3; the interaction is required for CDH1 localization to developing adherens junctions (PubMed:22294297). {ECO:0000250, ECO:0000250|UniProtKB:F1PAA9, ECO:0000250|UniProtKB:P09803, ECO:0000250|UniProtKB:Q9R0T4, ECO:0000269|PubMed:11226248, ECO:0000269|PubMed:11790773, ECO:0000269|PubMed:11856755, ECO:0000269|PubMed:12526809, ECO:0000269|PubMed:14595118, ECO:0000269|PubMed:14699157, ECO:0000269|PubMed:15240885, ECO:0000269|PubMed:16126725, ECO:0000269|PubMed:17261850, ECO:0000269|PubMed:18093941, ECO:0000269|PubMed:19604491, ECO:0000269|PubMed:20116244, ECO:0000269|PubMed:20432435, ECO:0000269|PubMed:22294297, ECO:0000269|PubMed:25208567, ECO:0000269|PubMed:31473225, ECO:0000269|PubMed:7982500}.
General Annotations	UP_COMMENT_TISSUE_SPECIFICITY	Expressed in granuloma macrophages (at protein level) (PubMed:27760340). Expressed in the skin (at protein level) (PubMed:22294297). Expressed in the liver (PubMed:3263290). {ECO:0000269|PubMed:22294297, ECO:0000269|PubMed:27760340, ECO:0000269|PubMed:3263290}.

Main Category	Category	Terms
Identifiers	ENTREZ_GENE_ID	999
Identifiers	ENSEMBL_GENE_ID	ENSG00000039068
Identifiers	UNIPROT_ID	Q9UII7_HUMAN
Identifiers	UNIPROT_ID	J3QKP1_HUMAN
Identifiers	UNIPROT_ID	H3BVI7_HUMAN
Identifiers	UNIPROT_ID	H3BNC6_HUMAN
Identifiers	UNIPROT_ID	D3XNU5_HUMAN
Identifiers	UNIPROT_ID	CADH1_HUMAN
Identifiers	UNIPROT_ID	B3GN61_HUMAN
Identifiers	UNIPROT_ID	A5D8W4_HUMAN
Identifiers	UNIPROT_ID	A0A6C0N5X1_HUMAN
Identifiers	UNIPROT_ID	A0A3G2JNR5_HUMAN
Identifiers	UNIPROT_ID	A0A140HMB1_HUMAN
Identifiers	UNIPROT_ID	A0A0U2ZQU7_HUMAN
Identifiers	UNIPROT_ID	A0A0E3XJU3_HUMAN
Interactions	PUBCHEM	US20240208940, Example 11
Interactions	PUBCHEM	US20240208940, Example 12
Interactions	PUBCHEM	US20240208940, Example 13
Interactions	PUBCHEM	US20240208940, Example 14
Interactions	PUBCHEM	US20240208940, Example 17
Interactions	PUBCHEM	US20240208940, Example 19
Interactions	PUBCHEM	US20240208940, Example 30
Interactions	PUBCHEM	US20240208940, Example 31
Interactions	PUBCHEM	US20240208940, Example 32
Interactions	PUBCHEM	US20240208940, Example 33
Interactions	PUBCHEM	US20240208940, Example 34
Interactions	PUBCHEM	US20240208940, Example 35
Interactions	PUBCHEM	US20240208940, Example 36
Interactions	PUBCHEM	US20240208940, Example 37
Interactions	PUBCHEM	N-[3-[(3,4-dihydroxyphenyl)methylamino]propyl]-5-phenyl-1,2-oxazole-3-carboxamide;2,2,2-trifluoroacetic acid
Interactions	PUBCHEM	N-(3-(2-hydroxynicotinamido)propyl)-5-phenylisoxazole-3-carboxamide

Main Category	Category	Terms
Interactions	PUBCHEM	N-(3-((2-azidopyrimidin-4-yl)amino)propyl)-5-phenylisoxazole-3-carboxamide
Interactions	PUBCHEM	5-phenyl-N-(2-(thiazole-2-carboxamido)ethyl)isoxazole-3-carboxamide
Interactions	PUBCHEM	N-[3-[(5-cyano-2-pyridinyl)amino]propyl]-5-phenyl-1,2-oxazole-3-carboxamide;2,2,2-trifluoroacetic acid
Interactions	UP_KW_LIGAND	Calcium
Interactions	UP_KW_LIGAND	Metal-binding
Literature	GENERIF_SUMMARY	Products of the c-myc gene activate or repress the activity of the E-cadherin promoter
Literature	GENERIF_SUMMARY	Both expression of E-cadherin and its membranous localization are required for well-differentiated-type morphogenesis in gallbladder cancer cells.
Literature	GENERIF_SUMMARY	Mutations and polymorphisms in E-cadherin gene CDH1, such as S270A, may contribute to the onset of prostate cancer (PCA) and warrant further investigations in other populations.
Literature	GENERIF_SUMMARY	The E-cadherin gene was potentially inactivated in a significant number of synovial sarcomas.
Literature	GENERIF_SUMMARY	Loss of E-cadherin may result in the disruption of the function of the cell-cell adhesion complex, which may cause weak cell-cell adhesion and confer invasive properties on a tumor.
Literature	GENERIF_SUMMARY	Cadherin-mediated cell sorting not determined by binding or adhesion specificity
Literature	GENERIF_SUMMARY	restoration of E-cadherin dependent adhsion in human protate carcinoma cells by expression of receptor protein-tyrosine phosphatase, PTPmu
Literature	GENERIF_SUMMARY	E-cadherin might be used as additional cell markers of Schwann cell-derived tumors.
Literature	GENERIF_SUMMARY	distinct methylation pattern in bladder cancer with frequent methylation of RARbeta, DAPK, E-cadherin, and p16.
Literature	GENERIF_SUMMARY	D257A & D370A mutations result in abnormal protein localization, changes in the actin cytoskeleton, reduced homophilic cell adhesion, and altered cell morphology; tumor-associated D370A mutation, but not the D257A mutation, induced increased cell motility
Literature	GENERIF_SUMMARY	biological function of Cys(9) within the first repeat (the E-cadherin-derived N-terminal repeat)
Literature	GENERIF_SUMMARY	These results demonstrate that integrin-dependent cell to extracellular matrix adhesion reinforces E-cadherin-dependent cell-cell adhesion in Caco-2 cells.
Literature	GENERIF_SUMMARY	E-cadherin play important roles in esophageal carcinogenesis.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	The SLUG zinc-finger protein represses E-cadherin in breast cancer.
Literature	GENERIF_SUMMARY	Lateral dimerization of the E-cadherin extracellular domain is necessary but not sufficient for adhesive activity
Literature	GENERIF_SUMMARY	results suggest that the inadequate trophoblastic invasion, induced by antiphospholipid antibodies, can be the result of decreased alpha1 integrin and VE-cadherin and increased alpha5 integrin and E-cadherin expression in the trophoblast
Literature	GENERIF_SUMMARY	characterization of DNA polymorphism in promoter regions
Literature	GENERIF_SUMMARY	mucoepidermoid carcinoma of the thyroid (MECT)displays consistent neoexpression of P-cadherin and major alterations in the expression of E-cadherin and the three catenins
Literature	GENERIF_SUMMARY	Tumor-associated mutations of E-cadherin enhanced random cell movement of transfected MDA-MB-435S mammary carcinoma cells as compared to wild-type (wt) E-cadherin-expressing cells.
Literature	GENERIF_SUMMARY	restoration of cell adhesion by overexpression of nectin in HSC-39 cells
Literature	GENERIF_SUMMARY	Differential expression in human brain tumors
Literature	GENERIF_SUMMARY	E-cadherin expression in dental epithelium followed an apical-coronal gradient that was opposite to that observed for N-cadherin.
Literature	GENERIF_SUMMARY	CAS/CSE 1 stimulates E-cadhrin-dependent cell polarity in HT-29 human colon epithelial cells.
Literature	GENERIF_SUMMARY	promoter methylation studied in 80 patients with head and neck squamous cell carcinoma (HNSCC)
Literature	GENERIF_SUMMARY	role in regulating expression of the leucocyte common antigen-related tyrosine phosphatase
Literature	GENERIF_SUMMARY	Recovery of cellular E-cadherin precedes replenishment of estrogen receptor and estrogen-dependent proliferation of breast cancer cells rescued from a death stimulus.
Literature	GENERIF_SUMMARY	aberrant methylation preferentially occurs in invasive ductal breast cancer associated with poor prognosis and is one of the mechanisms of expression silence in breast cancers
Literature	GENERIF_SUMMARY	Src-induced disruption of E-cadherin localization requires specific integrin signalling.
Literature	GENERIF_SUMMARY	Data suggest that E-cadherin-mediated signaling through PI3-kinase can regulate the invasive behavior of cells by modulating proteinase secretion.
Literature	GENERIF_SUMMARY	Defected E-cadherin expression might play a role in the

Main Category	Category	Terms
		development of malignant phenotype in NSCLC
Literature	GENERIF_SUMMARY	ZEB1 plays a role in repressing E-cadherin and MUC1 in epithelial cells [ZEB-1]
Literature	GENERIF_SUMMARY	oxidative stress induces tyrosine phosphorylation and cellular redistribution of occludin-ZO-1 and E-cadherin-beta-catenin complexes by a tyrosine-kinase-dependent mechanism
Literature	GENERIF_SUMMARY	Downregulation of E-cadherin in melanocytes and melanoma cells by endothelin-1
Literature	GENERIF_SUMMARY	Altered expression of E-cadherin in hepatocellular carcinoma
Literature	GENERIF_SUMMARY	CpG hypermethylation was an important mechanism of E-cadherin gene inactivation in bladder cancer and also that specific CpG sites consistently presented higher methylation levels than others.
Literature	GENERIF_SUMMARY	E-cadherin promoter is subject to epigenetic control in colorectal ulceration and plays a role in the progression from chronic inflammation to colorectal cancer.
Literature	GENERIF_SUMMARY	E-cadherin polymorphism in prostate neoplasms
Literature	GENERIF_SUMMARY	CDH1 c-160a promotor polymorphism is not associated with risk of stomach cancer.
Literature	GENERIF_SUMMARY	germline E-cadherin mutations responsible for the predisposition to diffuse gastric cancer (DGC) among the Japanese
Literature	GENERIF_SUMMARY	Restoration of E-cadherin/beta-catenin expression in pancreatic cancer cells inhibits growth by induction of apoptosis.
Literature	GENERIF_SUMMARY	dimerization of E-cadherin is important for its cell adhesive properties
Literature	GENERIF_SUMMARY	In vivo recruitment of RB and AP-2 proteins on the E-cadherin promoter in MCF7 cells
Literature	GENERIF_SUMMARY	investigates whether E-cadherin is a substrate for calpain and whether calpain-dependent proteolysis was associated with prostate cancer progression
Literature	GENERIF_SUMMARY	E-cadherin is hypermethylated and shows low expression in colon and rectal neoplasms
Literature	GENERIF_SUMMARY	results indicate that expression of E-cadherin in IST-1 trophoblast cells results in a contact-mediated inhibition of motility and invasion and suggest an important role for E-cadherin down-regulation in the intermediate trophoblast during implantation
Literature	GENERIF_SUMMARY	in vivo evidence in cancer cells of the differential association between CpG methylation, MBPs, and histone modification in the entire CpG island of the human E-cadherin (CDH1) gene

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Involvement of the adhesive ligand pair CD103-E-cadherin in human thymocyte cell proliferation
Literature	GENERIF_SUMMARY	These data reveal a cooperative interaction between p120 catenin and E-cadherin and a novel role for p120 that is likely indispensable in normal cells.
Literature	GENERIF_SUMMARY	association of CDH1 haplotypes with susceptibility to sporadic diffuse gastric cancer
Literature	GENERIF_SUMMARY	clinical significance of E-cadherin and proliferating cell nuclear antigen expression in adenoid cystic carcinoma
Literature	GENERIF_SUMMARY	The reduction of E-cadherin and beta-catenin expression was related to invasiveness and proliferative status of prolactinomas and correlated with the more aggressive behavior of prolactinomas in men compared with in women.
Literature	GENERIF_SUMMARY	role in negative regulation of CD44-hyaluronan interaction
Literature	GENERIF_SUMMARY	promotion of E-cadherin and suppression of beta-catenin/T cell factor may be an important mechanism underlying the chemopreventive action of Ca(2+) in colon cancer
Literature	GENERIF_SUMMARY	Loss of E-cadherin expression followed by expression of the mts1 gene may be an important event for increasing cell proliferation, motility and invasion activity in the progression of gallbladder cancer.
Literature	GENERIF_SUMMARY	The presence of E-cadherin mutations can significantly alter the accumulation of the apoptosis-regulating p53 protein, whereas no correlation with the p53 mutation status or with Ki-67 staining was observed.
Literature	GENERIF_SUMMARY	Expression of e-cadherin and beta-catenin in human esophageal squamous cell carcinoma
Literature	GENERIF_SUMMARY	Abnormal E-cadherin and alpha-catenin and beta-catenin in pancreatic carcinoma tissues. Abnormal E-cadherin and alpha-catenin with differentiation, lymph node and liver metastases. Aberrant beta-catenin with lymph node and liver metastases.
Literature	GENERIF_SUMMARY	phosphorylation-dependent localization of CDH1 in vertebrate cells.
Literature	GENERIF_SUMMARY	CpG methylation and thus loss of E-cadherin is associated with metastasising laryngeal cancer.
Literature	GENERIF_SUMMARY	matrix metalloproteinase-2 and 9 and membrane-type 1 matrix metalloproteinase mRNA expression in endometriosis was higher than in normal endometrium whereas E-cadherin, alpha- and beta-catenin mRNA expression was not suppressed in endometriosis
Literature	GENERIF_SUMMARY	Pulse-labeled beta-catenin replaces the beta-catenin bound to the cell surface prebiotinylated E-cadherin immediately after synthesis or arrives at the plasma membrane in a complex with the E-cadherin precursor.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Hypermethylation of an E-cadherin (CDH1) promoter region in high grade transitional cell carcinoma of the bladder comprising carcinoma in situ.
Literature	GENERIF_SUMMARY	study verified reduced expressions of adenomatous polyposis coli and E-cadherin proteins in colorectal cancer cells and suggests that normal protein expressions in benign epithelium are progressively and independently lost in sporadic colorectal cancers
Literature	GENERIF_SUMMARY	possibility that multiple pathways exist for E-cadherin entry into cells that are likely to reflect cell context and regulation.
Literature	GENERIF_SUMMARY	decreased expression of the E-cadherin catenin complex and methylation of the E-cadherin gene promoter region was found only in growth hormone cell adenomas with prominent fibrous bodies
Literature	GENERIF_SUMMARY	E-cadherin and claudins/occludin have roles in the regulation of tight junctions during the epithelium-mesenchyme transition, but are repressed by snail
Literature	GENERIF_SUMMARY	The molecular requirements for E-cadherin to activate Rac signaling thus appear distinct from those that stimulate PI3-kinase, and we postulate that p120-ctn may play a central role in the E-cadherin-Rac signaling pathway.
Literature	GENERIF_SUMMARY	Abnormal expression of E-cadherin was seen in mucoepidermoid carcinoma.
Literature	GENERIF_SUMMARY	High expression of E-cadherin is associated with the development of lymphatic tumor emboli
Literature	GENERIF_SUMMARY	dedifferentiation and a decreased expression of E-cadherin and ZO-1 are closely related to liver metastasis
Literature	GENERIF_SUMMARY	polymorphism of CDH-1 3'-UTR is a valid genetic marker for calcium stone disease.
Literature	GENERIF_SUMMARY	Findings identify E-cadherin as a novel substrate for matrilysin and indicate that shedding of E-cadherin ectodomain is required for epithelial repair.
Literature	GENERIF_SUMMARY	E-cadherin A/A genotype may be associated with susceptibility to urothelial cancer, but not with the progression of disease.
Literature	GENERIF_SUMMARY	The loss of E-cadherin rather than beta-catenin mutation is the crucial event in determining the differentiation 'level' of thyroid carcinomas
Literature	GENERIF_SUMMARY	CDH1 inactivation due to mutations, LOH and methylation is associated with invasive and in situ lobular breast cancer.
Literature	GENERIF_SUMMARY	E-cadherin inhibits human mammary and prostate tumor cell invasion. The invasion suppressor signal is mediated through the beta-catenin-binding domain of the E-cadherin cytoplasmic tail.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Methyl-CpG-binding protein 2 and promoter methylation cooperatively regulate E-cadherin gene expression in colorectal carcinoma
Literature	GENERIF_SUMMARY	In nasopharyngeal cancer, methylation of promoter is a major cause of down-regulation of E-cadherin which may finally lead to detachment and metastasis of neoplastic cells.
Literature	GENERIF_SUMMARY	Single nucleotide polymorphism in the E-cadherin gene promoter is associated with tumorigenesis and progression of gastric carcinoma
Literature	GENERIF_SUMMARY	Expression of HPV16 E6 in keratinocyes decreases expression of cell surface E-cadherin thereby depleting langerhans cells at the site of infection
Literature	GENERIF_SUMMARY	Data demonstrate a direct role of E-cadherin/catenin complex organization in the regulation of matrix metalloproteinases and suggest an implication of this regulation in the expression of an invasive phenotype by bronchial tumor cells.
Literature	GENERIF_SUMMARY	Bcl-2 expression decreases the level of functional E-cadherin thereby interfering with junction formation
Literature	GENERIF_SUMMARY	Results suggested that human embryos and blastocysts could express E-cadherin and the expression increased during their development.
Literature	GENERIF_SUMMARY	the -160 C/A polymorphism of E-cadherin is not directly involved in Korean gastric cancer development
Literature	GENERIF_SUMMARY	provides support that E-cadherin induction by WNT/beta-catenin signaling is an evolutionarily conserved pathway operative in lung cancer cells and that loss of expression may be important in lung cancer development or progression
Literature	GENERIF_SUMMARY	E-cadherin has a role in tumorigenicity and metastasis
Literature	GENERIF_SUMMARY	Hypoxia induces down-regulation of E-cadherin in ovarian carcinoma cells, via up-regulation of the transcriptional repressor SNAIL.
Literature	GENERIF_SUMMARY	E-cadherin-mediated cell adhesion and apoptosis of prostatic cancer cells are regulated by ligand activation of the androgen receptor
Literature	GENERIF_SUMMARY	E-cadherin affected the regulation of cell proliferation, differentiation and adhesion, and decreased chemosensitivity
Literature	GENERIF_SUMMARY	the comparable region of N-cadherin can substitute for this required region in E-cadherin and is required for suppression by the mutant form of N-cadherin that is capable of suppressing
Literature	GENERIF_SUMMARY	The frequencies of E-cadherin promoter hypermethylation appear to be correlated with more advanced stage of squamous carcinogenesis in skin.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	A recombinant E-cadherin lacking free sulfhydryl groups and its mutants with novel cysteines were expressed. The coexistence of the structurally identical adhesive and lateral dimers suggests some flexibility of the extracellular cadherin region.
Literature	GENERIF_SUMMARY	Pancreatic cancer likely occurs in case of the inactivation of E-cadherin and alpha-catenin genes and abnormal expression of proteins
Literature	GENERIF_SUMMARY	We conclude, therefore, that two major components of cell-cell interaction synergistically regulate cell cycle progression in HEK293 cells by regulating p21 expression in a beta-catenin/TCF-dependent manner.
Literature	GENERIF_SUMMARY	Intragenic deletion of CDH1 is the activating mechanism of the wild-type allele in hereditary diffuse gastric carcinoma.
Literature	GENERIF_SUMMARY	In human melanocytic tumors, a causative role of (loss of) E-cadherin or (gain of) N-cadherin for melanocytic tumor progression still remains to be proven.
Literature	GENERIF_SUMMARY	E-cadherin and epidermal growth factor receptor (EGFR) are associated in mammary epithelial cells and that E-cadherin engagement in these cells induces transient activation of EGFR, as previously seen in keratinocytes.
Literature	GENERIF_SUMMARY	an 80 kDa fragment of E-cadherin has a role in the metastatic progression of prostate cancer
Literature	GENERIF_SUMMARY	E-cadherin mediated adhesion and signaling in mammalian epithelial cells requires homolog of disc-large
Literature	GENERIF_SUMMARY	investigation of whether the -347G-->GA single nucleotide polymorphism affects the transcriptional activity of the E-cadherin gene
Literature	GENERIF_SUMMARY	Low expression of E-cadherin is associated with late cervical metastasis in stage I and II invasive squamous cell carcinoma of the oral tongue
Literature	GENERIF_SUMMARY	FAM associates with E-cadherin and beta-catenin during trafficking to the plasma membrane
Literature	GENERIF_SUMMARY	Aberrant methylation of CDH1 is associated with recurrent cervical cancer
Literature	GENERIF_SUMMARY	Promoter hypermethylation of E-cadherin and its abnormal expression in Epstein-Barr virus-associated gastric carcinoma
Literature	GENERIF_SUMMARY	endogenous PGE2 is involved in the hBD-2 and E-cadherin responses of human gingival epithelial cells to A. actinomycetemcomitans. HGEC exposed to A. actinomycetemcomitans showed a decrease in E-cadherin levels.
Literature	GENERIF_SUMMARY	EphA2 and E-cadherin may play an important role in tumor metastasis in colorectal cancer

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	single nucleotide polymorphism in CDH1 is a low-penetrant prostate cancer susceptibility gene
Literature	GENERIF_SUMMARY	Cdx1 or Cdx2 expression is sufficient to induce E-cadherin-dependent adhesion of COLO 205 cells associated with polarization and cell-cell membrane compaction, as well as induction of differentiated gene-expression pattern.
Literature	GENERIF_SUMMARY	Combined evaluation of VEGF and E-cadherin levels may become a useful indicator of NSCLC biological behavior and provide clinically important evidence on which to base treatment in NSCLC.
Literature	GENERIF_SUMMARY	E-cadherin plays an important role in regulating the invasive potential of prostate cancer cells through an unique paracrine mechanism.
Literature	GENERIF_SUMMARY	promoter hypermethylation and consequent downregulated of E-cadherin is significantly associated with gastric carcinomas of the diffuse histotype
Literature	GENERIF_SUMMARY	Results identify a necessary role for cortactin in the cadherin-actin cooperation that supports productive contact formation.
Literature	GENERIF_SUMMARY	E-cadherin-mediated adhesion mechanisms have a role in sperm-oolemma interactions
Literature	GENERIF_SUMMARY	Snail is involved in both the direct transcriptional repression of genes, such as E-cadherin and occludin, and post-transcriptional events, including downregulation of claudin-1
Literature	GENERIF_SUMMARY	SH2 and SH3 domains of Src mediate peripheral accumulation of phospho-myosin, leading to integrin adhesion complex assembly, whereas loss of SH2 or SH3 function restores normal regulation of E-cadherin and inhibits vimentin expression
Literature	GENERIF_SUMMARY	CDH1 methylation and E-cadherin expression are related and have a role in progression of diffuse gastric cancer
Literature	GENERIF_SUMMARY	E-cadherin has a role in preventing peritoneal dissemination in gastric cancer
Literature	GENERIF_SUMMARY	The absence of dermal invasion of nevus cells could be due to the expression of E-cadherin in these cells in reconstructed epidermis.
Literature	GENERIF_SUMMARY	E- to N-cadherin switching in epithelial carcinomas has a potential impact on metastatic progression.
Literature	GENERIF_SUMMARY	Cytoplasmic p120ctn may contribute to the invasive phenotype of E-cadherin-deficient breast cancer cells.
Literature	GENERIF_SUMMARY	determination of CDH1 promoter methylation in breast tumors with decreased or absent E-cadherin protein expression
Literature	GENERIF_SUMMARY	E-cadherin-mediated cell adhesion is required for

Main Category	Category	Terms
		keratinocyte-mediated control of melanocytic cells, which can override proliferative activity of beta-catenin.
Literature	GENERIF_SUMMARY	E-cadherin -347 G-->GA polymorphism may be associated with colorectal cancer.
Literature	GENERIF_SUMMARY	Thirteen novel CDH1 mutations found in diffuse gastric cancer families.
Literature	GENERIF_SUMMARY	Hypermethylation of CDH1 is statistically significantly associated with poor outcome in cervical cancer.
Literature	GENERIF_SUMMARY	the establishment of E-cadherin-based cell-cell adhesion requires Rap1
Literature	GENERIF_SUMMARY	E-cadherin and p16INK4a are commonly methylated in non-small cell lung cancer
Literature	GENERIF_SUMMARY	mutant E-cadherin significantly alters the dynamics of cell adhesion and motility in living cells
Literature	GENERIF_SUMMARY	Data showed that intestinal metaplasia is associated with E-cadherin down-regulation in gastric mucosa, whereas H. pylori infection does not seem to play a direct role in this process.
Literature	GENERIF_SUMMARY	E-cadherin and calretinin are sensitive and specific in differential diagnosis of benign and malignant serous effusion specimens.
Literature	GENERIF_SUMMARY	abnormal immunhistochemical E-cadherin and beta-catenin expression is associated with changes in pit pattern in invasive colorectal neoplasms
Literature	GENERIF_SUMMARY	Review summarizes divergent studies that provide evidence of the function of E-cadherin trafficking and the puzzle of how this adhesion molecule is regulated and managed throughout the life of the epithelial cell.
Literature	GENERIF_SUMMARY	E-cadherin mutations have no influence on expression of genes involved in Wnt-signaling, but they may promote their own expression by blocking upregulation of E-cadherin repressors.
Literature	GENERIF_SUMMARY	Most invasive and metastatic areas of oral squamous cell carcinoma showed reduced expression and methylation of E-cadherin.
Literature	GENERIF_SUMMARY	Decreased E-cadherin expression is an independent prognostic factor for disease progression and mortality in pathological stage I-III endometrial cancer.
Literature	GENERIF_SUMMARY	T lymphocyte development can proceed independently of alphaEbeta7/E-cadherin interactions
Literature	GENERIF_SUMMARY	Laryngeal cancer presented inactivation of Cx43 gene and E-cad gene and down regulation of Cx43 and E-cad proteins.
Literature	GENERIF_SUMMARY	Results point to a functional link between matrix metalloproteinase-3 and E-cadherin

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	role in tumor genesis and progress of hereditary gastric carcinoma
Literature	GENERIF_SUMMARY	E-cadherin downregulation is associated with the loss or absence of glandular epithelial differentiation in certain synovial sarcoma
Literature	GENERIF_SUMMARY	The E-Cadherin, the main system of adherens junction, present in the tight junctions in HepG2 cells.
Literature	GENERIF_SUMMARY	conjoint endocytosis and trafficking is a novel mechanism for the coregulation of E-cadherin and FGFR1 during cell signaling and morphogenesis
Literature	GENERIF_SUMMARY	the interaction of Pnn with the corepressor CtBP1 may modulate repression of E-cadherin transcription by CtBP1
Literature	GENERIF_SUMMARY	CDHE expression is repressed in the lower crypt of small intestine and is a biological marker of functional relevance.
Literature	GENERIF_SUMMARY	E-cadherin-deficiency in metastatic cancer may in some cases be due to p120 downregulation [review]
Literature	GENERIF_SUMMARY	E-cadherin may have a role in progression of squamous cell carcinoma
Literature	GENERIF_SUMMARY	heregulin stimulates aggregation and inhibits invasion of MCF-7/6 cells via activation of the E-cadherin/ catenin complex
Literature	GENERIF_SUMMARY	E-cadherin gene C-160A promoter polymorphism may not play a major role in the etiology of non-cardia gastric cancer in Chinese population.
Literature	GENERIF_SUMMARY	Change in the location of E-cadherin expression (from membrane cytoplasm) is strongly associated with an increased aggressiveness of colrectal cancer.
Literature	GENERIF_SUMMARY	Wnt signaling stabilizes Snail and beta-catenin proteins in tandem fashion so as to cooperatively engage transcriptional programs that control an epithelial-mesenchymal transition.
Literature	GENERIF_SUMMARY	Loss of expression of E-cadherin and beta-catenin may play an important role in the progression of pulmonary adenocarcinoma
Literature	GENERIF_SUMMARY	Aberrant E-cadherin is result of point mutation in exon 9 donor splice site causing skipping of exon 9 with consequent deletion of corresponding amino acids. Leads to disturbed cell adhesion. Mediates formation of cadherin/ catenin complex.
Literature	GENERIF_SUMMARY	These results provide a new example of Rho GTPase regulation of basolateral trafficking and demonstrate novel roles for Cdc42 and Rac1 in the post-Golgi transport of E-cadherin.
Literature	GENERIF_SUMMARY	a novel pathway for Rab11 dependent, dileucine-mediated, mu1B-independent sorting and basolateral

Main Category	Category	Terms
		trafficking, exemplified by E-cadherin
Literature	GENERIF_SUMMARY	Data suggest when carcinoma in a gastric biopsy is negative for E-cadherin staining, metastatic breast carcinoma should be considered.
Literature	GENERIF_SUMMARY	Altered E-cadherin adhesion complex is an early event affecting atypical lobular hyperplasia as well as lobular carcinoma in situ and occurs prior to progression to invasive disease.
Literature	GENERIF_SUMMARY	All 61 meningothelial meningiomas, 10 of 12 invasive meningiomas, and 3 of 5 anaplastic meningiomas were positive for both ECAD and beta-catenin, while these were both negative in all of the fibrous meningiomas.
Literature	GENERIF_SUMMARY	Tumors with reduced E-cadherin expression invaded deeper, had more lymph node metastasis, and had more lymphatic invasion than the tumors with preserved E-cadherin expression.
Literature	GENERIF_SUMMARY	cadherin switching (downregulation of E-cadherin and upregulation of N-cadherin) is necessary for increased motility but is not required for the morphological changes that accompany the epithelium-to-mesenchyme transition
Literature	GENERIF_SUMMARY	Disruption of E-cadherin-mediated cell-cell contacts at high cell density inhibits the induction of p27(kip1) and restores proliferation in contact-inhibited cells.
Literature	GENERIF_SUMMARY	association between decreased expression and metastatic Wilms' tumor
Literature	GENERIF_SUMMARY	dysfunction of E-cadherin and nm23 has a role in progression of NSCLC
Literature	GENERIF_SUMMARY	mutations cause inactivation of the cell adhesion protein E-cadherin; carriers of the CDH1 germline gene mutation develop an aggressive, diffuse, submucosal gastric cancer at an early age [review]
Literature	GENERIF_SUMMARY	E-cadherin expression was significantly associated with higher tumour grade in colorectal carcinoma.
Literature	GENERIF_SUMMARY	These findings suggest that alteration of the E-cadherin pathway can contribute to human clefting.
Literature	GENERIF_SUMMARY	examined here the effect of trans-interacting nectin on non-trans-interacting E-cadherin endocytosis
Literature	GENERIF_SUMMARY	E-cadherin may play a distinct role in the development of ovarian epithelial cancers.
Literature	GENERIF_SUMMARY	Strong E-cadherin expression in lymph node metastases was highly predictive of improved survival.
Literature	GENERIF_SUMMARY	The entire cadherin-catenin complex is involved when assessing its prognostic value in colonic adenocarcinom.
Literature	GENERIF_SUMMARY	disturbances in the distribution of beta-catenin and E-

Main Category	Category	Terms
		cadherin might affect the morphology of adenoid cystic carcinoma
Literature	GENERIF_SUMMARY	ADAM10 has a role in E-cadherin shedding and epithelial cell-cell adhesion, migration, and beta-catenin translocation
Literature	GENERIF_SUMMARY	Domain 5 of E-cadherin may play a critical role in the regulation of heterophilic adhesion to integrin alphaEbeta7; domains 1, 2, 3, and 4 are involved in homophilic adhesion.
Literature	GENERIF_SUMMARY	expression at transcriptional level is repressed by HBV X protein
Literature	GENERIF_SUMMARY	SIP1 sumoylation by Pc2 attenuates transcriptional repression of E-cadherin
Literature	GENERIF_SUMMARY	an E-cadherin germline mutation may have a role in development of hereditary diffuse gastric cancer
Literature	GENERIF_SUMMARY	E-cadherin distribution in human embryos is stage-dependent
Literature	GENERIF_SUMMARY	Patients with the -160C/C genotype might require H. pylori infection to promote inactivation of CDH1, suggesting that H. pylori infection might affect gastric carcinomas in initial stage because polymorphism is germ line.
Literature	GENERIF_SUMMARY	Reduced E-cadherin is a common genetic phenotype of gastric cancers and plays beneficial roles in tumor metastasis.
Literature	GENERIF_SUMMARY	E-cadherin may have a role in progression and recurrence of transitional cell carcinomas of the urinary bladder
Literature	GENERIF_SUMMARY	The E-cadherin is a cell-cell adhesion protein expressed in cytotrophoblasts, which is lost as they differentiate and syncytialise.
Literature	GENERIF_SUMMARY	loss of E-cadherin-mediated adhesion plays a causal role in the transition from low- to high-grade squamous cell carcinoma
Literature	GENERIF_SUMMARY	E-cadherin protein was lost and mRNA levels were significantly decreased in prostate cancer cells expressing kallikrein 4 and prostate specific antigen.
Literature	GENERIF_SUMMARY	report strong confirmation of the association between prostate cancer risk in FH+ cases and a functional CDH1 promoter SNP in an independent population
Literature	GENERIF_SUMMARY	inhibition of E-cadherin-dependent cell-cell adhesion led to the genetic reprogramming of tumor cells
Literature	GENERIF_SUMMARY	-160(C-->A) polymorphism in CDH1 gene promoter region may not be in association with genetic susceptibility to gastric cancer in Chinese population from Fujian.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	The TGFbeta(1)-induced destabilisation of E-cadherin-mediated cell-cell adhesion involves phosphorylation of beta-catenin, which is regulated by E-cadherin adhesion complex-associated PI3-kinase and PTEN.
Literature	GENERIF_SUMMARY	Increased expression of delta-catenin os associated with the down-regulation and redistibution of ECAD and p120ctn in prostatic cancer.
Literature	GENERIF_SUMMARY	Cell-to-cell communication mediated by E-cadherin contributes to the acquirement of a cardiomyogenic phenotype of human endothelial progenitor cells.
Literature	GENERIF_SUMMARY	With tumor progression and development of heterogeneity, the abnormal expressions of MMP-9, TIMP-2, and E-cadherin or DNA aneuploid rate or high SPF gradually increases, suggesting that they play a crucial role in gastric carcinoma progression
Literature	GENERIF_SUMMARY	Depressed E-cadherin correlates with HBxAg trans-activation function.
Literature	GENERIF_SUMMARY	CDH-1 is closely related to metalloproteinase and plays important but not well-understood role in onset and progression of primary open angle glaucoma. Role could be resolved by posttranslated products of gene and protein-protein interaction.
Literature	GENERIF_SUMMARY	expression of E-cadherin and beta-catenin is significantly down regulated in prostate cancer compared to surrounding benign appearing prostate, which correlates with increasing Gleason grade
Literature	GENERIF_SUMMARY	The CDH-1 gene 3'-UTR C/T polymorphism is associated with prostate cancer. The 'CC' homozygote indicates a relatively higher risk for developing prostate cancer than other genotypes.
Literature	GENERIF_SUMMARY	In testicular tumours, as in other neoplasms, dysadherin downregulates E-cadherin expression, at least in part
Literature	GENERIF_SUMMARY	E-cadherin controls enterocyte-specific expression of genes, such as the apoA-IV gene, through the control of hepatic nuclear factor 4alpha nuclear abundance
Literature	GENERIF_SUMMARY	These results suggest that Scrib stabilizes the coupling between E-cadherin and the catenins and are consistent with the idea that mammalian Scrib could behave as a tumor suppressor by regulating epithelial cell adhesion and migration.
Literature	GENERIF_SUMMARY	transforming growth factor-beta1 overexpression and alteration in E-cadherin/beta-catenin complexes in bladder urothelium might play a crucial role in urinary bladder carcinogenesis in humans exposed to long-term low-dose ionizing radiation
Literature	GENERIF_SUMMARY	Data indicate that E-cadherin may modulate Wnt-dependent gene expression in DLD-1 colorectal cancer cells by regulating the availability of beta-catenin.
Literature	GENERIF_SUMMARY	CDH1 may be a low-penetrant prostate cancer

Main Category	Category	Terms
		susceptibility gene.
Literature	GENERIF_SUMMARY	Increase in HOXA1 expression in mammary carcinoma cells at full confluence is E-cadherin-dependent.
Literature	GENERIF_SUMMARY	Results suggest that loss of E-cadherin function is linked to regulation of cell-cell and cell-matrix adhesion, based in part on cell surface expression of alpha2, alpha3 and beta1 integrins.
Literature	GENERIF_SUMMARY	decreased immunoreactivity of E-cadherin in esophageal squamous cell carcinoma was statistically correlated with presence of lymphatic metastases
Literature	GENERIF_SUMMARY	Our present results show that gastric differentiated-type carcinomas have different characteristics according to the phenotypic marker expression of the tumour in terms of histological findings, E-cadherin expression and pattern of chromosomal changes.
Literature	GENERIF_SUMMARY	Data suggest that tissue expression of E-cadherin could be a useful marker to predict the progression and metastasis of hepatocellular carcinoma.
Literature	GENERIF_SUMMARY	CDH1 promoter methylation, but not mutational inactivation, is part of an entire programme, resulting in epithelial-mesenchymal transition and increased invasiveness in breast cancer
Literature	GENERIF_SUMMARY	PCP-2 may play an important role in the maintenance of epithelial integrity, and a loss of its regulatory function may be an alternative mechanism for activating beta-catenin signaling.
Literature	GENERIF_SUMMARY	Loss of E-cadherin and Death Associated Protein kinase methylation and expression is more frequent in lymph node metastases of pancreatic adenocarcinoma
Literature	GENERIF_SUMMARY	Data show that dynamic microtubules regulate the local concentration of E-cadherin at cell-cell contacts.
Literature	GENERIF_SUMMARY	Serum levels may be a prognostic biological marker innonsmall cell lung cancer.
Literature	GENERIF_SUMMARY	Findings demonstrate a stimulatory role for E-cadherin in proliferative regulation, and identify a simple mechanism by which cell-cell contact may trigger or inhibit epithelial cell proliferation in different settings.
Literature	GENERIF_SUMMARY	Reduced E-cadherin expression is related to a poor prognosis through hematogenous metastasis in pulmonary adenocarcinoma.
Literature	GENERIF_SUMMARY	The high frequency of methylation of the CDH1 gene promoter suggests that the inactivation of tumor suppressor genes and of the genes related to the control of cellular proliferation through this mechanism is involved in gallbladder carcinogenesis.
Literature	GENERIF_SUMMARY	E-cadherin pathway is a novel and functionally important mediator by which changes in Kruppel-like KLF6 tumor suppressor can directly alter ovarian tumor

Main Category	Category	Terms
		invasion and metastasis.
Literature	GENERIF_SUMMARY	Activation of protease-activated receptor-2 (PAR2) interrupted adhesion of E-cadherin-expressing L cells and of primary airway epithelial cells to immobilized E-cadherin extracellular domain. Activation of PAR2 interrupts E-cadherin adhesion.
Literature	GENERIF_SUMMARY	E-cadherin methylation was associated with the inhibition of E-cadherin gene and protein expression in acute myeloid leukemia.
Literature	GENERIF_SUMMARY	These data suggest that endocytosis is the main pathway for the dissociation of E-cadherin adhesive dimers.
Literature	GENERIF_SUMMARY	E-cadherin has a role in head and neck squamous cell carcinoma [review]
Literature	GENERIF_SUMMARY	the degradation of E-cadherin in response to expression of R-cadherin is due to competition for p120(ctn)
Literature	GENERIF_SUMMARY	No significant association between E-cadherin expression and tumor grade, stage, age or gender of the patients, the number of recurrences, or overall survival could be seen.
Literature	GENERIF_SUMMARY	CDH1 mutations are associated with a risk of early-onset gastric cancer.
Literature	GENERIF_SUMMARY	Levels of expression of CtBP and p300 are critical for the action of SNAIL and ZEB1, which have a pivotal role in levels of epithelial-mesenchymsal transitionin human colon carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin immunoreactivity was detected in all the epithelial tissues examined, except adrenal cortical cells and granulosa cells.
Literature	GENERIF_SUMMARY	The mechanism underlying 5-Azacytidine's anti-metastasis activity is associated with restored functional expression of E-cadherin in ovarian cancer.
Literature	GENERIF_SUMMARY	E-cadherin/CDH1 gene methylation is an important cause for its gene silence in diffuse variant gastric carcinoma.
Literature	GENERIF_SUMMARY	Examination of E-cadherin expression and distribution in colorectal tumors can be extremely valuable in predicting disease recurrence.
Literature	GENERIF_SUMMARY	A novel mutation in the E-cadherin gene in a family with hereditary diffuse gastric cancer. The presence of a 1610delC germline mutation was confirmed.
Literature	GENERIF_SUMMARY	CDH1 serum levels may be a serological marker for staging of patients with testicular germ cell neoplasia.
Literature	GENERIF_SUMMARY	Our results indicate a functional role of Met-E-cadherin interaction in MCF-7 cells through the amplification of the signaling downstream of HGF-Met triggering that involved c-Src and phosphoinositide-3-kinase activities.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	A model is proposed to infer the pathogenic significance of CDH1 germline missense variants.
Literature	GENERIF_SUMMARY	The expression of E-cad was significantly lower in endocrine inactive group than that in endocrine active group of pituitary adenoma.
Literature	GENERIF_SUMMARY	CD24 regulates E-cadherin and TGF-beta3 expression in cultured oral epithelial cells
Literature	GENERIF_SUMMARY	dysfunction of E-cadherin due to its endocytosis may occur in some proportion of human breast carcinomas in which the PP2A-A protein is lost or significantly reduced
Literature	GENERIF_SUMMARY	the gain and not the loss of the E-cadherin axis contributes to the invasiveness of inflammatory breast cancer cells unique phenotype
Literature	GENERIF_SUMMARY	The expression of TGF-beta1 and E-cadherin are correlated with lymph node metastasis in supraglottic larynx squamous cell carcinoma.
Literature	GENERIF_SUMMARY	Roles of E-cadherin and beta-catenin complexes in signet ring cell carcinoma of the lung differ from their roles in gastric or colorectal cancers.
Literature	GENERIF_SUMMARY	loss of extracellular E-cadherin observed in normal duodenal & colonic mucosa in patients familial adenomatous polyposis might play a role in high susceptibility of these tissues for (pre-) malignant transformation
Literature	GENERIF_SUMMARY	E-cadherin may sensitize human melanoma cells towards apoptosis
Literature	GENERIF_SUMMARY	E-cadherin expression responds only to the combination treatment and not to single PPARgamma agonists, defining a new class of PPARgamma target genes
Literature	GENERIF_SUMMARY	SCCA2 regulates cell migration and invasion via E-cadherin expression, suggesting that SCCA2 may be involved in cancer behavior such as invasion or metastasis.
Literature	GENERIF_SUMMARY	Increased N-cadherin expression was found in pseudomyxoma peritonei.
Literature	GENERIF_SUMMARY	SMAD family member 1 is involved in the progression of pancreatic cancer and plays a role in mediating signal transduction from collagen type I to downregulate E-cadherin expression.
Literature	GENERIF_SUMMARY	In conclusion, ROS play a central role in mediating TPA-triggered sustained PKC and ERK signaling for regulation of gene expression of integrins and E-cahedrin that are responsible for EMT and migration of HepG2.
Literature	GENERIF_SUMMARY	VHL promotes E2 box-dependent E-cadherin transcription by HIF-mediated regulation of SIP1 and

Main Category	Category	Terms
		snail
Literature	GENERIF_SUMMARY	Real-time reverse transcription-PCR and Western blotting were used to assess levels of E-cadherin expression in esophageal cancer.
Literature	GENERIF_SUMMARY	first CDH1 germline mutation of an Italian family with hereditary diffuse gastric cancer is reported
Literature	GENERIF_SUMMARY	E-cadherin repression is necessary for c-Myc-induced cell transformation.
Literature	GENERIF_SUMMARY	alpha2-antiplasmin induction inhibits E-cadherin processing mediated by the plasminogen activator/ plasmin system, leading to suppression of progression of oral squamous cell carcinoma via upregulation of cell-cell adhesion
Literature	GENERIF_SUMMARY	Found in in situ lesions. May play some role in the prognosis/invasion of extramammary Paget's disease.
Literature	GENERIF_SUMMARY	Seven novel mutations within the CDH1 gene are associated with Hereditary diffuse gastric cancer.
Literature	GENERIF_SUMMARY	targeting constitutive expression of AP-2alpha in AP-2-positive KM12C colon cancer cells with small interfering RNA resulted in an increase in their invasive potential, downregulation of E-cadherin and increased expression of MMP-9
Literature	GENERIF_SUMMARY	These results indicate that perturbation of the E-cadherin/beta-catenin complex by H. pylori CagA plays an important role in the development of intestinal metaplasia.
Literature	GENERIF_SUMMARY	the recruitment of PI3K to the E-cadherin/beta-catenin/ p120-catenin complex via beta-catenin at the plasma membrane is required for calcium-induced phospholipase C-gamma1 activation and, ultimately, keratinocyte differentiation
Literature	GENERIF_SUMMARY	PIPKIgamma links E-cadherin to adaptor protein complexes
Literature	GENERIF_SUMMARY	E-cad expression to be correlated with E-cad methylation at highly statistically significant levels. Above a threshold of approximately 20% to 30% mean methylation, the expression of E-cad was effectively silenced.
Literature	GENERIF_SUMMARY	High Nonmembranous type E-cadherin expression correlated significantly with lymph node metastasis, distant metastasis, and recurrence regulated by hepatocyte growth factor
Literature	GENERIF_SUMMARY	Cadherin-E interaction with integrin alphaEbeta7 causes antitumor cytotoxic response by CD8+/CD103+ tumor-reactive T lymphocytes.
Literature	GENERIF_SUMMARY	C. albicans invades mucosal tissues by promoting the proteolytic degradation of E-cadherin in epithelial adherens junctions, mediated by transcription factor

Main Category	Category	Terms
		Rim101 and Sap5 proteins.
Literature	GENERIF_SUMMARY	Decreased E-cadherin expression is associated with advanced gallbladder cancers
Literature	GENERIF_SUMMARY	CDHE was expressed in both tumor types, especially in the cytoplasm and cell membrane of the cuboidal cells and to a lesser extent in the polygonal cell membrane, where it was expressed mostly in the cytoplasm.
Literature	GENERIF_SUMMARY	E-cadherin cleavage is enhanced by contact with C. albicans
Literature	GENERIF_SUMMARY	Data show tha E-cadherin homophilic binding independent of other cell contacts directly transduces growth inhibition by a beta-catenin-dependent mechanism that inhibits selective signaling functions of growth factor receptors.
Literature	GENERIF_SUMMARY	incubation with HGF mediated the release of MMP-7, resulting in extracellular cleavage of E-cadherin from stomach cancer cells
Literature	GENERIF_SUMMARY	Data show that E-cadherin/beta-catenin-based adherens junctions are dispensable for tight junction formation and apical lumen biogenesis but not for apical lumen remodeling.
Literature	GENERIF_SUMMARY	Decreased E-cadherin is asociated with thyroid cancer
Literature	GENERIF_SUMMARY	E-cadherin promoter hypermethylation was observed in the tumour of the P373L mutation carrier.
Literature	GENERIF_SUMMARY	None of the 30 infiltrating lobular carcinomas cells showed preserved E-cadherin expression.
Literature	GENERIF_SUMMARY	Together, these findings demonstrate a role for SNX1 in retrieval of E-cadherin from a degradative endosomal pathway and in membrane trafficking pathways that regulate E-cadherin recycling.
Literature	GENERIF_SUMMARY	Differentially regulated beta-catenin pools associate with the E-cadherin-gamma-secretase adherens junction complex; one pool regulated by gamma-secretase is important to intestinal epithelial cell homeostasis.
Literature	GENERIF_SUMMARY	Characterization of endogenous CDHE-CTNNB complexes with ELISA represents a dramatic improvement over other assays.
Literature	GENERIF_SUMMARY	While CDH1 methylation seems to be an early event in Hp gastritis, MLH1 methylation occurs late along with IM.
Literature	GENERIF_SUMMARY	E-cad promoter methylation may play a role in tumor cell differentiation and perineural invasion.
Literature	GENERIF_SUMMARY	Recurrent CDH1 mutations in families with hereditary diffuse gastric cancer are due to both independent mutational events and common ancestry. The presence of a founder mutation from Newfoundland is strongly

Main Category	Category	Terms
		supported.
Literature	GENERIF_SUMMARY	results of this study show no association exists between the E-cadherin genotype and the risk of tumor recurrence in Chinese patients with hepatocellular carcinoma
Literature	GENERIF_SUMMARY	Downregulation of E-cadhedrin expression is associataed with increased EGFR downstream signalling and a subsequent increase in expression of Th2-attracting chemokine TARC.
Literature	GENERIF_SUMMARY	The C/A polymorphism of CDHE has a direct effect on the risk of diffuse gastric cancer at young age in Mexican population.
Literature	GENERIF_SUMMARY	E-cadherin signaling is an important activator of c-Src at cell-cell contacts, providing a key input into a signaling pathway where quantitative changes in signal strength may result in qualitative differences in functional outcome.
Literature	GENERIF_SUMMARY	we explored the implication of three proteins (E-cadherin, a- and b-catenins) that form the cadherin-catenin complex, a receptorial structure strictly involved in tumoral vascular invasion and embolization in this biologic event
Literature	GENERIF_SUMMARY	Aberrant E-cadherin expressions were described in several tumors such as in bladder cancer. This was also found to be correlated with tumor invasion and survival.
Literature	GENERIF_SUMMARY	A tight control of E-cadherin expression depending on the differentiation stage of the progenitor cells, is suggested.
Literature	GENERIF_SUMMARY	In addition to suppressing late-stage tumor progression, E-cadherin-mediated adherens junctions may also contribute to the initial emergence of a cohesive morphogenic phenotype that is a hallmark of differentiated epithelial ovarian carcinoma.
Literature	GENERIF_SUMMARY	NFkappaB activation mediated by loss of E-cadherin may represent an essential, even initial event in colorectal neoplasm metastasis
Literature	GENERIF_SUMMARY	study demonstrates that E-cadherin represents a ligand for KLRG1 and that ligation of KLRG1 by E-cadherin inhibited effector cell functions of polyclonal NK cells
Literature	GENERIF_SUMMARY	E-cadherin thus requires both ankyrin-G and beta-2-spectrin for its cellular localization in early embryos as well as cultured epithelial cells.
Literature	GENERIF_SUMMARY	Junctional adhesion molecule-A is critical for the formation of pseudocanaliculi and modulates E-cadherin expression in hepatic cells
Literature	GENERIF_SUMMARY	The expression of aPKC-iota and E-cadherin may reflect the differentiation and invasive potential of cholangiocarcinoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Results show E-cadherin-mediated Listeria internalization requires Arp2/3 complex acts as an actin nucleator.
Literature	GENERIF_SUMMARY	The major subtypes of renal epithelial neoplasms display differential aberrant CDH1, PTGS2, and RASSF1A promoter methylation levels. This gene panel might contribute to a more accurate discrimination among common renal tumors.
Literature	GENERIF_SUMMARY	Ep-CAM cross signaling with N-cadherin involves Pi3K, resulting in the abrogation of the cadherin adhesion complexes in epithelial cells.
Literature	GENERIF_SUMMARY	High level E-cadherin expression and p120ctn localization were associated with esophageal squamous cell carcinoma differentiation and lymph node metastasis
Literature	GENERIF_SUMMARY	there are significant differences of expression of E-cadherin between primary breast cancer cells and their metastases
Literature	GENERIF_SUMMARY	Aberrant methylation of multiple genes (E-cadherin, estrogen receptor, RB1 , p16, p15, p14, and MGMT) is involved in gastric carcinogenesis.
Literature	GENERIF_SUMMARY	presence of the A allele of the -160C/A polymorphism in the E-cadherin gene may be a risk factor for prostate cancer in the Japanese population
Literature	GENERIF_SUMMARY	The expression of E-cadherin was negatively correlated with lymphatic metastasis, clinical stage, and cancer differentiation in gastric cardia carcinoma.
Literature	GENERIF_SUMMARY	a relation between the E-cadherin 3'-UTR C --> T polymorphism, the -160 A/-347 GA haplotype of two promoter polymorphisms and risk of endometriosis
Literature	GENERIF_SUMMARY	A novel germline CDH1 truncating mutation in the extracellular portion of the protein (517insA) was found in 2 relatives with lobular breast cancer. Germline CDH1 mutations can be associated with invasive LBC in the absence of diffuse gastric cancer.
Literature	GENERIF_SUMMARY	E-cadherin and beta-catenin have roles in progression of Epstein-Barr virus-associated gastric carcinoma
Literature	GENERIF_SUMMARY	The previously reported characteristics of this mutation, E-cadherin (V832M) do not apply to human epithelial cells expressing this mutant protein.
Literature	GENERIF_SUMMARY	Expression may be reduced in some early gastric carcinomas.
Literature	GENERIF_SUMMARY	Loss of E-cadherin expression is a risk factor for lymph node metastasis in early gastric cancer.
Literature	GENERIF_SUMMARY	The E-cadherin gene methylation presence in tumors with lowest invasive and metastatic potential suggests the early involvement of this epigenetic event in the multistep progression of the oral carcinogenesis

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	E-cadherin immunostaining was less intense in metaplastic cervical epithelial cells of women on tibolone, whereas hormone therapy and raloxifene were not associated with altered E-cadherin expression.
Literature	GENERIF_SUMMARY	These results demonstrate that lysophosphatidic acid regulates c-Met function through PKC delta and E-cadherin.
Literature	GENERIF_SUMMARY	E-cad expression in gastric carcinoma is relatively low in the internal obstruction of stagnant toxin type and the in-coordination between liver and stomach type.
Literature	GENERIF_SUMMARY	Analysis of interface of In1a protein with E-cadherin to identify single amino acid substitutions in InlA that would potentially improve the overall quality of interaction and hence increase the weak binding affinity of the complex
Literature	GENERIF_SUMMARY	These data suggest that serine/threonine phosphorylation of p120 influences the dynamics of E-cadherin in junctions.
Literature	GENERIF_SUMMARY	Our findings suggest that the hypermethylation of E-cadherin promoter might be involved in the process of gastric carcinoma through the specialized factors in H. pylori-induced enlarged fold gastritis.
Literature	GENERIF_SUMMARY	convincing evidence that cadherin adhesion is based on high-affinity cadherin-cadherin interactions
Literature	GENERIF_SUMMARY	MUC1 may affect cancer cell migration by increasing E-cadherin/beta-catenin complex formation and restoring E-cadherin membrane localization
Literature	GENERIF_SUMMARY	Re-expression of E-cadherin in HT29(US) cells restored the ability of caveolin-1 to down-regulate beta-catenin-Tcf/Lef-dependent transcription and survivin expression, as seen in HT29(ATCC) cells.
Literature	GENERIF_SUMMARY	Our results suggest that methylation of E-cadherin is a frequent, early event in gastric carcinoma progression. Two factors were ...associated with lymph node metastasis
Literature	GENERIF_SUMMARY	Kallikrein 6 induces E-cadherin shedding and promotes cell proliferation, migration, and invasion.
Literature	GENERIF_SUMMARY	Reduced expression of E-cadherin was associated with short overall survival incolorectal neoplasms
Literature	GENERIF_SUMMARY	Altogether, our results suggested that E-cadherin mediated cell-cell adhesion was essential for preventing the proteasome degradation of PTEN/
Literature	GENERIF_SUMMARY	Comparison of the E-cadherin structure with other type I cadherin structures reveals features that are likely to be critical to facilitate dimerization by strand exchange as well as dimer flexibility.
Literature	GENERIF_SUMMARY	ATCTG and CTTTG CDH1 haplotypes may be associated with an increased risk and decreased risk,

Main Category	Category	Terms
		respectively, of gastric cancer in Japan.
Literature	GENERIF_SUMMARY	tumor-specific downregulation of expression and methylation of CDH13 and CDH1, alone or in combination, may be involved in the development and invasive growth of pituitary adenomas.
Literature	GENERIF_SUMMARY	This paper focuses on changes in E-cadherin (CDH1), adenomatous polyposis coli (APC), and beta-catenin (CTNNB1) in 50 tumors of the central nervous system
Literature	GENERIF_SUMMARY	there is an aberrant nuclear localization of E-cadherin in CC-RCC harboring VHL mutations, suggesting potential prognostic value of VHL and E-cadherin in CC-RCC.
Literature	GENERIF_SUMMARY	findings suggest E-cadherin-dependent cell-cell contact, directly or indirectly, mediates signal to undergo apoptosis of CaSki cells during multicellular tumor spheroid formation & provides new information on role of E-cadherin in cervix cancer apoptosis
Literature	GENERIF_SUMMARY	E-cadherin expression is down-regulated in adenoid cystic carcinoma of the salivary glands, which is correlated to nerve invasion, regional recurrence and distant metastasis.
Literature	GENERIF_SUMMARY	Data suggest that Dia1 localizes to and controls E-cadherin-mediated junctions in a RhoA-dependent manner.
Literature	GENERIF_SUMMARY	Somatic E-cadherin mutations affect apoptosis regulation in that way that they can facilitate the disruption of adherens junctions thereby possibly influencing the response to cisplatin-based chemotherapy.
Literature	GENERIF_SUMMARY	Report a potential association of variants in the CCND1 and CDH1 genes with familial colorectal cancer using a unique study design with phenotypic extremes.
Literature	GENERIF_SUMMARY	The current findings suggested that simultaneous methylation of the E-cadherin and H-cadherin genes occurs in a subset of NSCLCs
Literature	GENERIF_SUMMARY	-160A of the E-cadherin gene is emerging as a low-penetrance tumor susceptibility allele for the development of gastric, lung, prostate, and urothelial cancers.
Literature	GENERIF_SUMMARY	functional significance of combined dysregulation of PKD1 and E-cadherin in prostate cancer; their effect on cell growth is mediated by beta-catenin.
Literature	GENERIF_SUMMARY	removal of N-glycans on E-cadherin resulted in elevated tyrosine phosphorylation level of beta-catenin and reduced beta- and alpha-catenins at adherens junctions
Literature	GENERIF_SUMMARY	genetic predisposition plays role in hereditary diffuse gastric cancer
Literature	GENERIF_SUMMARY	analysis of Epstein-Barr virus, beta-catenin, and E-cadherin in gastric carcinomas

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	CpG island hypermethylation of E-cad gene exists in gastric cancer, which down-regulates E-cad expression and might be involved in tumorigenesis and development of gastric cancer.
Literature	GENERIF_SUMMARY	In colorectal adenocarcinoma, MUC1 expression was negatively correlated to E-cad expression, and MUC2 and MUC5AC expression were positively correlated to E-cad expression.
Literature	GENERIF_SUMMARY	Therefore, a loss of cell-cell junctions, a key process that occurs during the epithelial-mesenchymal transition, should have a broad impact on ER alpha transcriptional functions.
Literature	GENERIF_SUMMARY	KCC3 down-regulates E-cadherin/beta-catenin complex formation by inhibiting transcription of E-cadherin gene and accelerating proteosome-dependent degradation of beta-catenin protein
Literature	GENERIF_SUMMARY	membranous overexpression of E-cadherin and beta-catenin are associated with the metastatic prostate cancer cells in bone and the high frequency of expression suggests their involvement in the intercellular adhesion of the metastatic cells in bone
Literature	GENERIF_SUMMARY	Galectin-3 and E-cadherin expressions are associated with lymph node metastasis of colon cancer.
Literature	GENERIF_SUMMARY	The C/C genotype of 3'-UTR C/T SNP and -160C/-374GA haplotype in E-cadherin gene may be a factor for risk of epithelial ovarian cancer in Chinese; the lower expression of E-cadherin might play a role in the pathogenesis of epithelial ovarian cancer.
Literature	GENERIF_SUMMARY	A high frequency of hypermethylation was detected in CDH1 and SFN genes in tumoral and normal cortex tissues.
Literature	GENERIF_SUMMARY	In most hereditary cancer syndromes, like hereditory diffuse gastric cancer and lobular carcinoma of the breast, multiple organ sites are affected by cancer and have been shown to be associated with germline mutations in CDH1 at 16q22.1.
Literature	GENERIF_SUMMARY	Silencing Cdc42 blocks activation of EGFR and Src induced by Ca2+ depletion, resulting in a reduction in E-cadherin degradation
Literature	GENERIF_SUMMARY	The expression of E-cadherin was significantly lower in supraglottic carcinoma than that in normal laryngeal mucosa.
Literature	GENERIF_SUMMARY	Using transient promoter assays, we show that Twist can down-regulate E-cadherin promoter activity by up to two folds.
Literature	GENERIF_SUMMARY	the CaR regulates cell survival and Ca(2+)(o)-induced differentiation in keratinocytes at least in part by activating the E-cadherin/PI3K pathway through a Src family tyrosine kinase-mediated signaling

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	in addition to the previously implicated tumor suppressor activity of E-cadherin, modified forms of this glycoprotein might also play a role in growth promotion
Literature	GENERIF_SUMMARY	Micrometastasis was significantly correlated with the depth of invasion, tumor size, operation method, Lauren classification, lymphovascular invasion and loss of E-cadherin expression in primary tumor.
Literature	GENERIF_SUMMARY	that the heterogeneous pattern of SCCA and E-cadherin in primary lesions is strongly associated with the high incidence of lymph node metastasis in cervical squamous cell carcinoma
Literature	GENERIF_SUMMARY	changes of Ecadherin (CDH1) and beta-catenin (CTNNB1) genes in two central nervous system germ cell tumors are reported; both germ cell tumors analyzed demonstrated LOH of the CDH1 gene
Literature	GENERIF_SUMMARY	the APC has a novel role in UV-induced cell death; APC/CDH1 is regulated through proteolysis
Literature	GENERIF_SUMMARY	E-cadherin expression is frequently reduced in laryngeal squamous epithelium of patients with laryngopharyneal reflux.
Literature	GENERIF_SUMMARY	Rac signaling to cyclin D1 is a crucial pro-proliferative effect of E-cadherin-mediated cell-cell adhesion
Literature	GENERIF_SUMMARY	in solid pseudopapillary neoplasm of the pancreas E-cadherin expression moved from membranous to intracytoplasmic localization
Literature	GENERIF_SUMMARY	The CDH1 C-160A polymorphism is not associated with the risk of CRC in the German population
Literature	GENERIF_SUMMARY	The present study indicates that CDH1 single nucleotide polymorphisms might modify susceptibility to esophageal squamous cell carcinomas and gastric cardia adenocarcinomas
Literature	GENERIF_SUMMARY	ADAM10-mediated E-cadherin proteolysis leads to the impaired cohesion of keratinocytes observed in eczematous dermatitis.
Literature	GENERIF_SUMMARY	Noncohesive pancreatic cancers were characterized by the loss of E-cadherin protein expression. Promoter hypermethylation is a possible mechanism of E-cadherin gene silencing in a subset of these cancers.
Literature	GENERIF_SUMMARY	The E-cadherin-repressed hNanos1 gene induces tumor cell invasion by upregulating MT1-MMP expression.
Literature	GENERIF_SUMMARY	The expressions of aPKC-iota and E-cadherin may reflect the differentiation and invasive potential of cholangiocarcinoma
Literature	GENERIF_SUMMARY	E-cadherin mutant missing N-glycans at Asn554, Asn566 and Asn618 failed to induce cell cycle arrest in G1 phase and to suppress cell proliferation in comparison with wild-type E-cadherin.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Increased histone acetylation might be responsible for PGC-1alpha-mediated transactivation of a minimal E-cadherin promoter.
Literature	GENERIF_SUMMARY	study describes activation of EGFR by mutant E-cadherin as a novel mechanism in tumor cells that explains the enhanced motility of tumor cells in the presence of an extracellular mutation of E-cadherin
Literature	GENERIF_SUMMARY	Expression of S100A4 protein and loss of E-cadherin protein expression are significantly associated with tumor progression in non-small cell lung cancer.
Literature	GENERIF_SUMMARY	The abnormal expression of CDX(2) and E-cadherin plays an important role in the development of gastric carcinoma.
Literature	GENERIF_SUMMARY	Vpu leads to the depression of both total and beta-catenin-associated E-cadherin levels through beta-TrCP-dependent stabilization of the transcriptional repressor Snail.
Literature	GENERIF_SUMMARY	PTPRK influences transactivating activity of beta-catenin in non-tumoral and neoplastic cells by regulating the balance between signaling and adhesive beta-catenin, thus providing biochemical basis for the hypothesis of PTPRK as a tumor suppressor gene.
Literature	GENERIF_SUMMARY	E-cad, CD44v6 and PCNA play important roles in invasion and metastasis of non-small cell lung cancer (NSCLC).
Literature	GENERIF_SUMMARY	Decreased E-cadherin may be important in the development of endometrial endometrioid carcinoma.
Literature	GENERIF_SUMMARY	Combined analysis of Cripto-1 and E-cadherin has significant value in evaluating the metastatic potential of gastric cancer and predicting patient prognosis.
Literature	GENERIF_SUMMARY	levels in colon carcinomas were not statistically different from levels in adjacent normal mucosa and were not correlated with tumor, nodes, and metastases stage
Literature	GENERIF_SUMMARY	The disassembly of E-cadherin complexes from junctions in human keratinocytes was induced by Rac1 and Rac3 via activation of the Rac target PAK1.
Literature	GENERIF_SUMMARY	These results lead to a new hypothesis that Snail and ZEB1 are downstream of CCN6 and play a critical role in CCN6-mediated regulation of E-cadherin in breast cancer.
Literature	GENERIF_SUMMARY	p35 co-expression targets E-cadherin to lysosomes and p35-triggered disappearance of E-cadherin precursor can be blocked specifically by lysosomal protease inhibitors, indicating p35 induces endocytosis and subsequent degradation of precursor E-cadherin
Literature	GENERIF_SUMMARY	results indicate that Src and E-cadherin may play an important role in epithelial-mesenchymal transformations, invasion, and aggressive

Main Category	Category	Terms
		clinicopathologic features of head and neck squamous carcinoma
Literature	GENERIF_SUMMARY	E-cadherin expressioni in adenomas suggest that this molecule might have role in adenoma formation though not necessarily be involved in neoplastic progression.
Literature	GENERIF_SUMMARY	LIV-1 may be a regulator of E-cadherin
Literature	GENERIF_SUMMARY	ASH1 inactivates DKK1 and DKK3, negative regulators of Wnt/beta-catenin signaling, E-cadherin, and integrin beta1 through ASH1-mediated deacetylation and repressive trimethylation of lysine 27 of histone H3 in the promoter regions of DKK1 and E-cadherin.
Literature	GENERIF_SUMMARY	No CDH1 polymorphisms or haplotypes were associated with GC risk, no differences of effect were seen by H pylori infection. Three CDH1 polymorphisms in the same haplotype interacted with smoking to increase GC risk in smokers but not in never smokers.
Literature	GENERIF_SUMMARY	The heterotypic trans-interaction of LI- and E-cadherin might play a role during development of the intestinal epithelium when the cells do not yet have elaborate membrane specializations.
Literature	GENERIF_SUMMARY	Methylation of CDH1 was present in 80% of NSCLC tissues but only in 14% of noncancerous tissues.
Literature	GENERIF_SUMMARY	Activin A downregulates E-cadherin expression in endometrial cells.
Literature	GENERIF_SUMMARY	This study provides into the mechanisms underlying the functional role of EZH2 overexpression in gastric cancer cells and a new modality of regulation of E-cadherin expression in silencing mechanisms of tumor suppressor genes.
Literature	GENERIF_SUMMARY	Positive staining for E-cadherin should not preclude a diagnosis of lobular in favor of ductal carcinoma.
Literature	GENERIF_SUMMARY	The -160A, 234 2I allele and haplotype A-A-T-2I were risk factors of Transitional Cell Carcinoma of the Bladder. Haplotype C-A-T-I might act as a protective factor for TCCB.
Literature	GENERIF_SUMMARY	transforming growth factor-beta1 indirectly reduces antigen-presenting cell density in EpM by affecting E-cadherin expression, which might explain the increased susceptibility of abnormal tissue differentiation.
Literature	GENERIF_SUMMARY	superficial intramucosal signet ring carcinoma, although widespread, is predominantly located in the proximal one third of the stomach in patients with E-cadherin gene mutations.
Literature	GENERIF_SUMMARY	Data show that the haplotype the E-cadherin gene may have impact on the risk of epithelial ovarian carcinoma. The CC genotype of 3'-UTR + 54CT may be a potential risk factor for the epithelial ovarian carcinoma.
Literature	GENERIF_SUMMARY	study of correlation between mutations & expression of

Main Category	Category	Terms
		E-cadherin, beta-catenin, occludin & claudin & complexity of colon carcinoma growth; perturbed expression & distribution of these proteins was found, but could not be linked to complexity of growth
Literature	GENERIF_SUMMARY	loss of expression of the miR-200 family members may play a critical role in the repression of E-cadherin by ZEB1 and ZEB2 during EMT, thereby enhancing migration and invasion during cancer progression.
Literature	GENERIF_SUMMARY	Rsults suggest that the lower epithelial alpha-catenin, E-cadherin and (or) ZO-1 expression in patients with atopic asthma contributes to a defective airway epithelial barrier and a higher influx of eosinophils in the epithelium.
Literature	GENERIF_SUMMARY	The nonsense-mediated-decay mRNA surveillance pathway downregulates aberrant E-cadherin transcripts in gastric cancer cells and in CDH1 mutation carriers.
Literature	GENERIF_SUMMARY	ADAM15 catalyzes the cleavage of E-cadherin to generate a soluble fragment that in turn binds to and stimulates ErbB receptor signaling
Literature	GENERIF_SUMMARY	Data show that loss of E-cadherin expression was independent prognostic factor for progression-free survival.
Literature	GENERIF_SUMMARY	aberrant or decreased expression of E-cadherin seems to be one of most promising markers of poor prognosis in localized prostate cancer
Literature	GENERIF_SUMMARY	promoter-hypermethylation of E-cad is significantly associated with its defective expression and tumor differentiation in non-small cell lung cancer
Literature	GENERIF_SUMMARY	Reduced expression of E-cadherin may have a role to play in the pathogenesis of invasive Paget's disease of the vulva.
Literature	GENERIF_SUMMARY	The CDH1 163+37235G>A polymorphism may represent a novel susceptibility variant for sporadic diffuse gastric cancer
Literature	GENERIF_SUMMARY	E-cadherin loss in tumors contributes to metastatic dissemination by inducing wide-ranging transcriptional and functional changes
Literature	GENERIF_SUMMARY	N-glycosylation at Asn-633 is essential for E-cadherin expression, folding and trafficking.
Literature	GENERIF_SUMMARY	results indicate that E-cadherin in the nevus cells of the giant congenital nevocellular nevi may affect nevus cell motility rather than intercellular attachment
Literature	GENERIF_SUMMARY	data did not support a crucial role of promoter methylation of the E-cadherin gene in the remarkable downregulation of E-cadherin expression in colorectal cancers
Literature	GENERIF_SUMMARY	E-cadherine and CD44 immunoexpression in oral squamous cell carcinoma as prognosis factors.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Loss of E-cadherin-mediated adhesion led to acquisition of phenotypic properties that augmented cell motility and directed transition from precancer to cancer in skin-like tissues.
Literature	GENERIF_SUMMARY	loss of E-cadherin is probably consequent on p120 loss or decrease. Aberrations and other alterations of the E-cadherin gene are unlikely to be responsible for the loss of E-cadherin in solid pseudopapillary tumors.
Literature	GENERIF_SUMMARY	Results suggest that a decrease in E-cadherin gene expression level in high-grade neuroepithelial tumors may be a hallmark of malignancy in dedifferentiated tumors and that it may be possibly correlated with their progression and dissemination.
Literature	GENERIF_SUMMARY	Core-fucosylated E-cadherin regulated nuclear beta-catenin accumulation in lung cancer cells.
Literature	GENERIF_SUMMARY	Loss of e-cadherin is associated with ovarian carcinoma
Literature	GENERIF_SUMMARY	E-cadherin may have a role in inhibition melanoma invasiveness by epigallocatechin-3-gallate
Literature	GENERIF_SUMMARY	Time-lapse microscopy in live cells showed that the recycling endosome is always requisite for E-cadherin sorting and trafficking.
Literature	GENERIF_SUMMARY	Results suggest that histone H3 deacetylation plays a crucial role in transcriptional repression of E-cadherin in colorectal cancers.
Literature	GENERIF_SUMMARY	rather than being translocated to the nucleus for regulating the target gene transcription, Smad7-stabilized-beta-catenin is shunted to the E-cadherin complex to modulate cell-cell adhesion.
Literature	GENERIF_SUMMARY	Rb depletion results in deregulation of E-cadherin.
Literature	GENERIF_SUMMARY	The Matrix Metalloproteinase (MMP-2, MMP-9) to E-cadherin (M/E ratio) characterizes an important aspect of the molecular phenotype associated with the histologic progression of prostate cancer among African American prostate cancer patients.
Literature	GENERIF_SUMMARY	Results show that InlA triggers two successive E-cadherin post-translational modifications, the Src-mediated tyrosine phosphorylation of E-cadherin followed by its ubiquitination by the ubiquitin-ligase Hakai.
Literature	GENERIF_SUMMARY	Hypermethylation of E-cadherin was found in 28 of 38 nasopharyngeal carcinoma
Literature	GENERIF_SUMMARY	Sequence analysis of the high-frequency region along E-cadherin exons 7-9 revealed a number of sequence alterations in the patient group as a whole. A C insertion at nt 76,598 was found in 2 African American patients.
Literature	GENERIF_SUMMARY	Decreased immunoexpression of beta-catenin and E-cadherin in serous ovarian tumors may be helpful in

Main Category	Category	Terms
		identifying the cases of higher metastatic potential and infiltration ability.
Literature	GENERIF_SUMMARY	Results show a correlation of CDH1 and APC promoter methylation with loss of E-cadherin and APC proteins and with activation of Wnt/beta-catenin signaling pathway.
Literature	GENERIF_SUMMARY	beta-catenin expression was significantly related to E-cadherin and MMP-7 expression
Literature	GENERIF_SUMMARY	Increased dysadherin expression is possibly one of the post-transcriptional mechanisms responsible for E-cadherin downregulation in thyroid papillary neoplasia.
Literature	GENERIF_SUMMARY	Only borderline results for the association of AXIN2 and CDH1 with oral clefts with and without tooth agenesis.
Literature	GENERIF_SUMMARY	loss of expression is related to bone metastasis in prostate cancer
Literature	GENERIF_SUMMARY	The significant difference in methylation of CDHE observed in liver neoplasms and liver cirrhosis was not observed in plasma samples.
Literature	GENERIF_SUMMARY	hypermethylation of CDH1 and integrin alpha4 genes may be used as recurrence-associated prognostic indicators in stage I and stage II esophageal squamous cell carcinoma, respectively.
Literature	GENERIF_SUMMARY	important role of the E-cadherin/beta-catenin/Tcf signaling pathway in atherosclerosis.
Literature	GENERIF_SUMMARY	In the gastric cancerous tissues, the expression percentage of Wnt-1, beta-catenin and E-cadherin is 54.4%, 45.6%, 47.2%, respectively, which is significantly higher than the percentage expression of these genes in normal tissues (p
Literature	GENERIF_SUMMARY	By inhibiting E-cadherin expression and enhancing ICAM-1 expression, LMP1 may reduce the cell-cell adhesive ability and improve the cell-matrix adhesive ability and migratory ability of nasopharyngeal carcinoma cells.
Literature	GENERIF_SUMMARY	Patients who had tumors with CDH1 hypermethylation had significantly better overall survival compared with patients who had tumors without hypermethylation (P
Literature	GENERIF_SUMMARY	significant association of concurrent expression of unmethylated E-cadherin gene and E-cadherin protein with metastatic prostate cancer cells in bone. May have role in intercellular adhesion in formation of metastatic lesions in bone.
Literature	GENERIF_SUMMARY	The abnormal expression of E-cadherin and beta-catenin were closely related with differentiation, clinical stage, metastasis in laryngeal carcinoma.
Literature	GENERIF_SUMMARY	cadherin-mediated cell-cell contact regulates early keratinocyte differentiation independently from changes in cell shape

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	The -347del allele of E-cadherin strongly links with the +178T and +234 13N ins alleles. The -347del/del genotype may increase susceptibility of sporadic gastric carcinoma among males in high-risk areas of China.
Literature	GENERIF_SUMMARY	study demonstrated that E-cadherin missense mutants are subjected to Endoplasmic Reticulum Quality Control (ERQC) and that their loss is due to endoplasmic reticulum-associated degradation
Literature	GENERIF_SUMMARY	ILK silencing inhibited binding of PARP-1 to SIRE
Literature	GENERIF_SUMMARY	Overall meta-analysis indicated that the -160A allele carriers (CA+AA) had a 21% elevated risk of prostate cancer, when compared with the homozygotes (CC) (OR=1.21; 95% CI
Literature	GENERIF_SUMMARY	the activated AR can downregulate E-cadherin expression to promote the activation of epithelial-mesenchymal transition and tumor metastasis.
Literature	GENERIF_SUMMARY	Discontinuous staining of N-cadherin and loss of E-cadherin expression in hepatocellular carcinoma predicts a high risk of recurrence after surgical treatment.
Literature	GENERIF_SUMMARY	In human HCC tissues, we observed a correlation among ROS induction, E-cadherin down-regulation, Snail up-regulation, and E-cadherin promoter methylation.
Literature	GENERIF_SUMMARY	The presence of E-cadherin in tumors correlated with the absence of metastases in regional lymph nodes and was observed, as a rule, in the patients at the early stages of the disease.
Literature	GENERIF_SUMMARY	Activated Akt seems to characterize well-differentiated invasive squamous laryngeal carcinomas, loss of E-cadherin and activation of beta-catenin correlated with high grade carcinomas.
Literature	GENERIF_SUMMARY	The results presented here describe the expression of E-cadherin in the male reproductive tract and gametes and strongly suggest its involvement in adhesion events during human fertilization.
Literature	GENERIF_SUMMARY	Tbx2 and Tbx3 may play a dual role during the radial to vertical growth phase transition by both inhibiting senescence via repression of p21(CIP1) expression, and enhancing melanoma invasiveness by decreasing E-cadherin levels.
Literature	GENERIF_SUMMARY	Reduced expression of E-cadherin/catenin complex in hepatocellular carcinomas.
Literature	GENERIF_SUMMARY	In hepatocellular carcinoma (HCC), there is a negative correlation between the positive expression of p130Cas and the normal expression of E-cadherin/beta-catenin. p130Cas plays important roles in the invasion, metastasis and prognosis of HCC.
Literature	GENERIF_SUMMARY	Tissue and urine expression of E-cadherin in bladder

Main Category	Category	Terms
		transitional cell carcinoma are very closely associated with the biological behavior of the neoplasm.
Literature	GENERIF_SUMMARY	E-cadherin confer significantly different characteristics on cells.
Literature	GENERIF_SUMMARY	E-cadherin plays an important role in uterine receptivity.
Literature	GENERIF_SUMMARY	We suggest that CDH1 cytoplasmic immunolocalization as a result of increased IGF-II levels identifies those nonmuscle invasive presentations most likely to recur
Literature	GENERIF_SUMMARY	The resistance of E-cadherin over-expressing cells to staurosporine may due to the up-regulation of Bcl-2/Bax ratio. When E-cadherin interference plasmids were transfected into MCF-7 cells, Bcl-2 expression was down-regulated.
Literature	GENERIF_SUMMARY	E-cadherin was strongly expressed in urothelial cell carcinoma in situ (CIS) of the bladder.
Literature	GENERIF_SUMMARY	Ras transformation has at least two independent actions to disrupt E-cadherin junctions, with effects to cause both mislocalization of E-cadherin away from the cell surface and profound decrease in the expression of E-cadherin
Literature	GENERIF_SUMMARY	the over-expression of EphA2 or the down-expression of E-cadherin is correlated with cancer progression and lymphogenous metastasis in gastric cancer.
Literature	GENERIF_SUMMARY	Expression of p53, BCL-2 and Ki-67 was upregulated in clinically localized prostate cancer compared with benign prostate tissue, with no alteration in E-cadherin expression.
Literature	GENERIF_SUMMARY	recruitment of E-cadherin expression by Connexin 43 and inhibits the malignant behaviour of lung cancer cells
Literature	GENERIF_SUMMARY	N/E- cadherin expression profile has a significant prognostic value in invasive bladder cancer.
Literature	GENERIF_SUMMARY	The C-160A polymorphism is not associated with gastric cancer risk in the Italian population.
Literature	GENERIF_SUMMARY	overexpression of SRF in colorectal carcinoma cells is associated with modulation of E-cadherin/beta-catenin expression and may play an important role in colorectal cancer metastasis.
Literature	GENERIF_SUMMARY	alterations in beta-catenin/E-cadherin complex play a critical role in spindle and/or corded (SPICO) cells' features
Literature	GENERIF_SUMMARY	In polarized epithelial cells, E-cadherin regulates AMPK phosphorylation by controlling the localization of the LKB1 complex.
Literature	GENERIF_SUMMARY	The reduced immunoexpression of E-cadhedrin in the membrane may be related to the high degree of cell indifferentiation in cases of oral squamous cell

Main Category	Category	Terms
		carcinoma with high scores.
Literature	GENERIF_SUMMARY	metastasis and survival of patients with bladder cancer are mediated, at least in part, by E-cadherin
Literature	GENERIF_SUMMARY	augmented production of HA and the decreased E-cadherin expression by keratinocytes stimulated with IL-4/IL-13 or IFN-gamma cause spongiosis in acute eczema
Literature	GENERIF_SUMMARY	There was increased expression of E-cadherin and beta-catenin in the eutopic and ectopic endometrium in adenomyosis.
Literature	GENERIF_SUMMARY	of promoter hypermethylation of TIMP3, CDH1, DAPK, RASSF1A, p16INK4A and MGMT, only the epigenetic silencing of TIMP3 and CDH1 predicted a better outcome in head and neck squamous cell carcinoma
Literature	GENERIF_SUMMARY	E-cadherin-mediated adherens junctions formation contributes to PI3K/AKT activation in epithelial ovarian cancer (EOC) by a mechanism that provides basis for therapeutic strategies that exploit PI3K inhibition to halt EOCs
Literature	GENERIF_SUMMARY	E-cadherin truncation and sLeX overexpression are early events which may facilitate the tumor cells in oral cancer to metastasize
Literature	GENERIF_SUMMARY	E-cadherin, beta-catenin, and ZEB1 have roles in malignant progression of cancer [review]
Literature	GENERIF_SUMMARY	the overexpression of epithelial cell adhesion molecule and reduced expression of E-cadherin in combination is more significant than a single marker for predicting poor survival.
Literature	GENERIF_SUMMARY	Cleavage of the extracellular domain and nuclear translocation of E-cadherin is a common event that may determine local invasion in pituitary adenomas.
Literature	GENERIF_SUMMARY	Abnormal p120-catenin expression correlates with abnormal E-cadherin expression and specific small GTPase overexpression, which contribute to the malignancy-related to NSCLC.
Literature	GENERIF_SUMMARY	Germline deletions of CDH1 are one of the mechanisms underlying hereditary diffuse gastric cancer.
Literature	GENERIF_SUMMARY	Cell surface E-cadherin expression level was significantly lower in leukemia cells than in normal hematopoietic progenitors.
Literature	GENERIF_SUMMARY	Polyamines regulate E-cadherin transcription through c-Myc modulating intestinal epithelial barrier function.
Literature	GENERIF_SUMMARY	Urinary soluble 80kDa fragment of E-cadherin has a potential clinical diagnostic value for diabetic nephropathy and E-cadherin may participate in the pathogenesis of DN.
Literature	GENERIF_SUMMARY	Hypermethylation of E-cadherin, p14 or RASSF1A in

Main Category	Category	Terms
		urine sediment DNA is a potential biomarker for detecting superficial, low grade cancer
Literature	GENERIF_SUMMARY	ZO-1, occludin, and E-cadherin are downregulated in carcinomas arising from various compartments of the biliary tract
Literature	GENERIF_SUMMARY	Cell separation or abrogation of E-cadherin-mediated cell-cell contacts both cause a dramatic increase in accumulation of the c-Jun protein.
Literature	GENERIF_SUMMARY	uterine non-endometrioid (papillary serous and clear cell) carcinomas were less likely to express E-cadherin compared with endometrioid carcinomas and normal endometrial tissue
Literature	GENERIF_SUMMARY	E-cadherin deficiency initiates gastric signet-ring cell carcinoma in mice and man.
Literature	GENERIF_SUMMARY	Data show that Wnt-5a stimulation of breast epithelial cells leads to increased cell-cell adhesion, and that Wnt-5a/casein kinase Ialpha (CKIalpha)-specific phosphorylation of beta-catenin leads to increased complex formation of beta-catenin/E-cadherin.
Literature	GENERIF_SUMMARY	Results suggest that Ecad mutant forms lack some specific interaction shown by the wild type hEcad, and that these missing (or altered) interactions would play a role in tumour formation.
Literature	GENERIF_SUMMARY	the different 2nd hits in CDH1 occurring in neoplastic lesions from hereditary diffuse gastric cancer patients was quantified
Literature	GENERIF_SUMMARY	data suggest that Dkk-1 stimulates the release of beta-catenin from cell membrane and facilitates cell migration which accompanies degradation of beta-catenin/E-cadherin
Literature	GENERIF_SUMMARY	Collagen type I may influence the expression of E-cadherin and beta-catenin in carcinoma ex-pleomorphic adenoma.
Literature	GENERIF_SUMMARY	E-cadherin expression was reduced in hepatocellular carcinoma.
Literature	GENERIF_SUMMARY	p190RhoGAP and p120ctn associated predominantly on the plasma membrane of cells overexpressing E-cadherin, and that E-cadherin-bound p120ctn contributed to RhoA inactivation by favoring p190RhoGAP-RhoA association.
Literature	GENERIF_SUMMARY	core fucosylation regulates the processing of oligosaccharides and turnover of E-cadherin
Literature	GENERIF_SUMMARY	NGAL overexpression altered subcellular localization of E-cadherin and catenins and decreased E-cadherin-mediated cell-cell adhesion.
Literature	GENERIF_SUMMARY	up-regulation of E-cadherin is an early defining event in ovarian cancer and may play a significant role in the initial development of the primary ovarian tumor

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	N- and E-cadherin expression in the human fetal ovary indicates likely roles in gonadal development from germ cell proliferation to primordial follicle formation, as well as in the development of the urogenital ducts of both genders.
Literature	GENERIF_SUMMARY	Loss plays an important role in progression of oral squamous cell carcinoma, that is, down regulation of its expression is associated with de-differentiation and metastasis
Literature	GENERIF_SUMMARY	E-cadherins engaged at junctions do not directly revert to free membrane diffusion
Literature	GENERIF_SUMMARY	This review highlights recent advances in defining the mechanisms that regulate E-cadherin expression in breast cancer.
Literature	GENERIF_SUMMARY	these results suggest a novel role for the ERalpha that plays the dual role of ligand-independent activator and ligand-dependent repressor of E-cadherin in breast cancer cells.
Literature	GENERIF_SUMMARY	changes in the Akt/beta-catenin pathway play key roles in the regulation of E-cadherin through the transactivation of the Slug gene in uterine carcinosarcomas.
Literature	GENERIF_SUMMARY	aberrant expression of E-cadherin may contribute to cell motility and invasion in breast tumors
Literature	GENERIF_SUMMARY	Downregulation of E-cadherin and diffuse membranous beta-catenin expression suggest a dysregulation of the E-cadherin/beta-catenin complex in Merkel cell carcinoma.
Literature	GENERIF_SUMMARY	results demonstrate an association between polymorphisms within the CDH1 (E-cadherin) gene and Crohn's disease, which leads to cytoplasmic accumulation of E-cadherin instead of its normal localisation at the plasma membrane/adherens junction.
Literature	GENERIF_SUMMARY	MGAT3 and MGAT5 competitively modified E-cadherin N-glycans.
Literature	GENERIF_SUMMARY	alpha(2)-, alpha(3)-, and beta(1)-integrins and E-cadherin expression in normal human lung and bronchopulmonary sequestration and congenital cystic adenomatoid malformation were evaluated using Western blot and immunohistochemistry.
Literature	GENERIF_SUMMARY	E-cadherin regulates both the PTEN protein levels and its recruitment to cell-cell junctions, which integrates proliferative and morphogenetic signaling in breast epithelial cells.
Literature	GENERIF_SUMMARY	reduced syndecan-1/E-cadherin expression may be good indicators of recurrence and prognosis in extrahepatic bile duct carcinoma.
Literature	GENERIF_SUMMARY	Novel role for shear in regulating the endocytosis of E-

Main Category	Category	Terms
		cadherin and invasiveness in metastatic esophageal cancer cells.
Literature	GENERIF_SUMMARY	PRL-3 promoted downregulation of E-cadherin.
Literature	GENERIF_SUMMARY	decreased expression of E-cadherin in upper urothelial carcinoma (UUC) in regions affected by Balkan Endemic Nephropathy tumors may be linked to tumor growth, while expression of E-cadherin in control tumors may be associated with tumor grade.
Literature	GENERIF_SUMMARY	SFK and cortactin constitute an important signaling pathway that functionally links E-cadherin adhesion and the actin cytoskeleton
Literature	GENERIF_SUMMARY	the tensile strength and the lifetime of single E-cadherin/E-cadherin bonds between parental cells are significantly lower over a wide range of loading rates
Literature	GENERIF_SUMMARY	hepatocellular carcinoma with cirrhosis was characterized by a significantly low expression of E-cadherin
Literature	GENERIF_SUMMARY	This report underlines that prophylactic gastrectomy remains the only option to eliminate the high risk for gastric cancer in CDH1 mutation carriers.
Literature	GENERIF_SUMMARY	These results suggest that alpha(5)-integrin up-regulation induced by E-cadherin loss under hypoxia has a crucial role in regulating the migration of extravillous trophoblast cells.
Literature	GENERIF_SUMMARY	e-cadherin is diminished in superficial koilocytotic cells' foci in Epidermodysplasia verruciformis
Literature	GENERIF_SUMMARY	Data show that overexpression of shrew-1 leads to preformation of an E-cadherin/EGF receptor/HER2/src-kinase/shrew-1 signaling complex and accelerated E-cadherin internalization.
Literature	GENERIF_SUMMARY	Results indicate that the expression of E-cadherin transcriptional repressors increased with malignancy in ovarian epithelial neoplasms.
Literature	GENERIF_SUMMARY	Studies show for the first time that DPAGT1 is an upstream regulator of E-cadherin N-glycosylation status and adherens junction composition and suggest that dysregulation of DPAGT1 causes disturbances in intercellular adhesion in oral cancer.
Literature	GENERIF_SUMMARY	MMP1 and MMP3, but not CDH1, IGFBP3, STK15 or VEGF, appear to play roles in familial and sporadic renal cell carcinoma
Literature	GENERIF_SUMMARY	The C/C genotype of CDH1 might be a potential risk for cervical cancer development
Literature	GENERIF_SUMMARY	Data indicate CDH1-160C>A is a risk factor for CRC, and because a high proportion of the European population are carriers of at-risk genotypes, the variant is likely to contribute substantially to the development of colorectal cancer (CRC).

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	E-cadherin, a tumor invasion suppressor in HCC, was found to be a putative gene target of miR-9. E-cadherin was up-regulated by miR-9 inhibitor.
Literature	GENERIF_SUMMARY	Mtation of E-cadherin influences Rac1 and Rho activation in opposite directions.
Literature	GENERIF_SUMMARY	Immunohistochemical analysis of E-cadherin and Zeb-1 in primary tumors confirmed that expression of the two proteins was mutually exclusive.
Literature	GENERIF_SUMMARY	cross-talk between ERalpha and EGFR could be extended to the modulation of E-cadherin expression by 4-hydroxytamoxifen and epidermal growth factor
Literature	GENERIF_SUMMARY	hMSCs caused breast cancer spheroids to become disorganized which was accompanied by a disruption of cell-cell adhesion, E-cadherin cleavage, and nuclear translocation of E-cadherin
Literature	GENERIF_SUMMARY	Using in vitro infection assays, the authors showed that Helicobacter pylori CagA interacts with E-cadherin, p120-catenin, and c-Met.
Literature	GENERIF_SUMMARY	E-cadherin may coengage KLRG1 and alpha(E)beta(7) and that KLRG1 overcomes its exceptionally weak affinity for cadherins through multipoint attachment to target cells, resulting in inhibitory signaling.
Literature	GENERIF_SUMMARY	MELF-type invasion represents a specific tumour alteration, and the reduction in hormone receptor and E-cadherin expression would be consistent with epithelial-mesenchymal transition.
Literature	GENERIF_SUMMARY	Interstitial trophoblastic cell fusion and E-cadherin immunostaining in the placental bed of normal and hypertensive pregnancies.
Literature	GENERIF_SUMMARY	Data suggest that E-cadherin-based cell-cell adhesion limits Wnt signals by promoting the activity of a junction-localized beta-catenin phosphodestruction complex, which may be relevant to tissue morphogenesis and cell fate decisions during development.
Literature	GENERIF_SUMMARY	Short-term preoperative treatment of celecoxib up-regulates the expression of E-cadherin in gastric carcinomas.
Literature	GENERIF_SUMMARY	E-cadherin and metalloproteinase-1 and -7 polymorphisms have roles in colorectal cancer
Literature	GENERIF_SUMMARY	Data show that caveolin-1 and E-cadherin closely associated at cell borders and in internalized structures upon stimulation
Literature	GENERIF_SUMMARY	Data suggest that cadherin dimerization proceeds via an induced fit mechanism where the monomers first form a tryptophan-2 independent initial encounter complex and then undergo subsequent conformational changes to form the final strand-swapped dimer.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Down reglation of E-cadherin is associated with rhabdomyosarcoma.
Literature	GENERIF_SUMMARY	increased E-cadherin and alpha-catenin expression and reduced Wnt signaling might be involved in the mechanisms of tumor suppressor function of BCSC-1.
Literature	GENERIF_SUMMARY	Expression of E-cadherin was negative in all leiomyosarcomas.
Literature	GENERIF_SUMMARY	Infection with H. pylori provokes shedding of the ectodomains of c-Met and E-cadherin. ADAM10 contributes to the shedding of c-Met and E-cadherin.
Literature	GENERIF_SUMMARY	Investigated methylation status of the CDH1 promoter in gastric cancer patients. A difference was found between the methylation level of the two alleles in some samples, but there was no mono-allelic promoter hypermethylation in any of the samples.
Literature	GENERIF_SUMMARY	The main objectives were to assess tumor proliferation and invasiveness biomarkers (Ki-67, E-cadherin) and to identify potential correlation between biomarkers and classic prognostic factors in cervical cancer.
Literature	GENERIF_SUMMARY	E-cadherin and p63 are implicated in tumoral progression and may be used as a prognostic markers.
Literature	GENERIF_SUMMARY	miR-200a appears to act as a multifunctional tumor suppressor miRNA in meningiomas through effects on the E-cadherin and Wnt/beta-catenin signaling pathways.
Literature	GENERIF_SUMMARY	the ability of CBX7 to positively regulate E-cadherin expression by interacting with HDAC2 and inhibiting its activity on the E-cadherin promoter would account for the correlation between the loss of CBX7 expression and a highly malignant phenotype
Literature	GENERIF_SUMMARY	pH 6.6 activates Src kinases, resulting in tyrosine phosphorylation of E-cadherin and p120ctn and a weakening of the association of E-cadherin with p120ctn and contributing to the instability of E-cadherin at adherens junctions
Literature	GENERIF_SUMMARY	This study identifies the E-cadherin/catenin complex as a discriminative, partly polyamine-regulated feature of IL-4/IL-13-exposed alternatively activated macrophages
Literature	GENERIF_SUMMARY	Simultaneous analyses of fascin-1 and E-cadherin expression could be more effective in evaluating the prognoses of patients with laryngeal squamous cell carcinoma.
Literature	GENERIF_SUMMARY	novel mechanism of E-cadherin gene inactivation, with CLL cells displaying a higher proportion of aberrant nonfunctional transcripts and resulting up-regulation of the wnt-beta-catenin pathway
Literature	GENERIF_SUMMARY	High methylation status of the 5' CpG island of E-cadherin gene may be one of the mechanisms in the development of gastric cardiac adenocarcinoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	aids in Wnt/beta-catenin signal transduction in invasive ductal carcinoma of breast
Literature	GENERIF_SUMMARY	In squamous cell lung cancer or lung adenocarcinomas, the abnormal cytoplasmic expression of p120ctn isoform 1 always correlates to the abnormal expression of E-cadherin.
Literature	GENERIF_SUMMARY	The expression of RKIP and E-cadherin in prostate cancer tissues was obviously lower than that in the benign prostatic hypertrophy tissues.
Literature	GENERIF_SUMMARY	CDH1 gene germ-line mutations are relatively rare in hereditary gastric cancer in China, whereas CDH1 somatic mutations and promoter methylation synergistically induce CDH1 downregulation in these patients.
Literature	GENERIF_SUMMARY	Data show that elevated Snail expression by Pdcd4 knockdown leads to downregulation of E-cadherin resulting in activating beta-catenin/Tcf-dependent transcription.
Literature	GENERIF_SUMMARY	The present findings point to a significant association between reduced E-cadherin expression and the progression and aggressivity of breast lesions.
Literature	GENERIF_SUMMARY	a decrease in CDH1, CDH13, and TIMP3 expression levels with an increase in CD44 can be used as an indicator for invasion in both ER-positive and -negative breast tumors.
Literature	GENERIF_SUMMARY	Data show that conversion of Ras-expressing keratinocytes from a premalignant to malignant state induced by decreasing E-cadherin function was associated with and required an approximately two to threefold decrease in RalA expression.
Literature	GENERIF_SUMMARY	Histologically normal tissue adjacent to tumor tissue expressing the epithelial-mesenchymal transition-inducing gene SNAI1 shows alterations in the expression of epithelial differentiation genes such as CDH1 and VDR.
Literature	GENERIF_SUMMARY	genetic variation in E-cadherin may be associated with breast cancer risk, and that this relationship may vary by menopausal status.
Literature	GENERIF_SUMMARY	Up-regulation of E-cadherin by microRNA through direct targeting of transcriptional repressors of E-cadherin, inhibits tumor progression in breast cancer cells.
Literature	GENERIF_SUMMARY	no evidence for the presence of aberrant methylation sites of E-cadherin in tumors from patients with NSCLC
Literature	GENERIF_SUMMARY	The abnormal expression of E-cadherin and P(120ctn) is closely related to the degree of differentiation, clinical stage and cervical lymph node metastasis in nasopharyngeal carcinoma.
Literature	GENERIF_SUMMARY	Data show that P-cadherin overexpression in breast cancer cells with wild-type E-cadherin promotes cell

Main Category	Category	Terms
		invasion, motility, and induces the secretion of MMP-1 and MMP-2, which then lead to P-cadherin ectodomain cleavage.
Literature	GENERIF_SUMMARY	Our results suggest that genetic instabilities of the E-cadherin gene have a role in meningioma development and progression
Literature	GENERIF_SUMMARY	The -347G-->GA promoter polymorphism in E-cadherin gene is associated with specific CRC features, and may be a prognostic factor rather than a susceptibility factor during the progression of CRC.
Literature	GENERIF_SUMMARY	The combination of the upregulation of vimentin and aberrant expression of E-cadherin/beta-catenin complexes at the tumour invasive front may provide a useful prognostic marker in oral squamous cell carcinoma.
Literature	GENERIF_SUMMARY	CDH1 new susceptibility loci for ulcerative colitis
Literature	GENERIF_SUMMARY	the extracellular domains of E- and N-cadherin determines the basic localization pattern, whereas the cytoplasmic domains modulate it thereby affects the cell adhesion activity, subapical accumulation, and the formation of adherens junction.
Literature	GENERIF_SUMMARY	Abnormality of E-cadherin expression was related to lymph node metastasis and advanced clinical stage of Her2-positive breast cancer.
Literature	GENERIF_SUMMARY	Immunohistochemistry on TMAs of 178 gastric cancers showed no correlation between COX-2 and E-cadherin expression in the conventional or early gastric cancer
Literature	GENERIF_SUMMARY	Results show that disruption of E-cadherin junctions and enhancement of cell invasion by beta-estradiol stimulation was mediated by NO-dependent activation of c-Src.
Literature	GENERIF_SUMMARY	In gastric carcinoma, up-regulation of CD147 and down-regulation of E-cadherin are in a negative correlation.
Literature	GENERIF_SUMMARY	IBH-6 and IBH-4 breast cancer cells show down-regulation of E-cadherin expression with aberrant protein localization, and up-regulation of Twist; these features can be related to their invasive/metastatic characteristics.
Literature	GENERIF_SUMMARY	CpG hypermethylation is an important mechanism of E-cad gene inactivation in ES-2 and SKOV3 ovarian carcinoma cell lines.
Literature	GENERIF_SUMMARY	Allele-specific CDH1 downregulation and hereditary diffuse gastric cancer
Literature	GENERIF_SUMMARY	Cadherin can be used to discriminate between high- and low-risk papillomavirus in urethral cytologic specimens.
Literature	GENERIF_SUMMARY	Either overexpression of a key regulator of cell cycle, Mad2, or knock down of Aurora B, an important kinase

Main Category	Category	Terms
		in mitosis, or Cdh1, a key E3 ligase in cell cycle, resulted in a significant increase of SP cells in CNE-2.
Literature	GENERIF_SUMMARY	Malignant drug resistant breast tumors are characterized by reverse relation between E-cadherin expression level and their proliferative activity
Literature	GENERIF_SUMMARY	Evaluation of Twist and E-cadherin expressions should be useful for determining tumor properties, including prognosis, in patients with esophageal squamous cell carcinoma
Literature	GENERIF_SUMMARY	molecular pathway from hypoxia to cellular transformation includes up-regulation of HIF and subsequent transcriptional induction of LOX and LOXL2, which repress E-cadherin and induce epithelial to mesenchymal transition
Literature	GENERIF_SUMMARY	The results indicate that developmentof an invasive carcinoma from an intraductal papillary mucinous neoplasms of pancreas (IPMN) is associated with a decrease in tumor cells expressing E-cadherin and beta-catenin and higher proliferative activity.
Literature	GENERIF_SUMMARY	E-cadherin methylation plays an important role in the carcinogenesis of colon carcinoma cells, and can re-express after treatment with 5-Aza-CdR.
Literature	GENERIF_SUMMARY	The TLR9, IL6, and CDH1 variants all persisted as independent risk factors for post-infectious irritable bowel syndrome (PI-IBS) when controlling for previously identified clinical risk factors.
Literature	GENERIF_SUMMARY	Differentiated prostate cell survival depended on E-cadherin and PI3K, but not keratinocyte growth factor, androgen, AR or MAPK.
Literature	GENERIF_SUMMARY	siRNA for HDAC3 reduced the cell migration with elevated E-cadherin expression
Literature	GENERIF_SUMMARY	E-cad was profiled in the NCI-60 cancer cell lines at the DNA, RNA, and protein levels using six different microarray platforms plus bisulfite sequencing.
Literature	GENERIF_SUMMARY	Listeria monocytogenes internalin interacts with E-cadherin [review]
Literature	GENERIF_SUMMARY	Gene methylation (CHFR, E-cadherin, BNIP3) in the peritoneal fluid could detect occult neoplastic cells in the peritoneum and might be a risk factor for peritoneal metastasis.
Literature	GENERIF_SUMMARY	an interaction between beta-catenin and vinculin is crucial for stabilizing E-cadherin at the cell surface
Literature	GENERIF_SUMMARY	Results indicate that cells underwent EMT exhibited overactive TGFbeta signaling and loss of expression of the CDH1, CGN, CLDN4, and KLK10 genes as a result of hypermethylation of their corresponding promoter regions.
Literature	GENERIF_SUMMARY	E-cadherin showed a reciprocal staining pattern to ILK

Main Category	Category	Terms
		in 21 of 25 cases, with only focal expression of the marker in pancreatic adenocarcinoma.
Literature	GENERIF_SUMMARY	one seminoma and three nonseminomas showed LOH of CDH1
Literature	GENERIF_SUMMARY	Data show there was a strong inverse correlation between slug and ERalpha and E-cadherin immunoreactivity in breast cancer cases.
Literature	GENERIF_SUMMARY	Hypermethylation of E-Cadherin is associated with cervical carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin and beta-catenin are differently expressed and distributed in prostate cancer cell lines with different characteristics of epithelial-mesenchymal transition.
Literature	GENERIF_SUMMARY	the AA (rather than CC) genotype of E-cadherin is associated with increased susceptibility and advanced pancreatic carcinoma in Chinese patients.
Literature	GENERIF_SUMMARY	integration of Rac1 and Rab7 activities by Armus provides an important regulatory node for E-cadherin turnover and stability of cell-cell contacts
Literature	GENERIF_SUMMARY	cadherin switching and p120(ctn) signaling as important targets of GnRH function and as novel mediators of invasiveness and tumor progression in ovarian cancer.
Literature	GENERIF_SUMMARY	High nonmembranous E-Cadherin expression was associated with advanced clinical stage, lymph node metastasis in nasopharyngeal carcinoma.
Literature	GENERIF_SUMMARY	L1 binding to ankyrin alters the regulation of neuronal branching and leads to a decrease in perisomatic synapses of GABAergic inhibitory interneurons in the developing transgenic mouse cingulate cortex.
Literature	GENERIF_SUMMARY	miR-9 miRNA downregulates E-cadherin. Expression of miR-9 is activated by MYC and MYCN, which bind to the mir-9-3 locus. Elevated expression of miR-9 may contribute to epithelial-mesenchymal transition and metastasis in some and possibly many tumours.
Literature	GENERIF_SUMMARY	The activation of Ezrin and the absence of E-cadherin contribute to the tumorigenesis and metastasis of esophageal squamous cell carcinoma.
Literature	GENERIF_SUMMARY	decrease of E-cadherin and the gain of N-cadherin gene expression are risk factors for cancer-related death in bladder cancer
Literature	GENERIF_SUMMARY	E-cadherin expression predicts precancerous lesions of the larynx.
Literature	GENERIF_SUMMARY	The results showed no differences in the expression of E-cadherin between tooth germs and solid and unicystic ameloblastomas.
Literature	GENERIF_SUMMARY	These findings indicate that independent control of E-

Main Category	Category	Terms
		cadherin expression and cell motility could be essential molecular events in p53 mutant-induced invasive phenotypes.
Literature	GENERIF_SUMMARY	Reduced E-cadherin expression may correlate to a dedifferentiated phenotype in the somatotroph pituitary adenomas.
Literature	GENERIF_SUMMARY	Kruppel-like factor 4 inhibits epithelial-to-mesenchymal transition through regulation of E-cadherin gene expression
Literature	GENERIF_SUMMARY	increased expression of Cx43 and E-cadherin may contribute to the efficient metastasis of gastric cancer to the lymph nodes.
Literature	GENERIF_SUMMARY	increased binding of DNMT3b to E-cadherin promoter region by K-Ras cause promoter hypermethylation for reduced expression of E-cadherin, leading to the decreased cell aggregation and increased metastasis of human prostate cancer cells.
Literature	GENERIF_SUMMARY	Loss of E-cadherin expression was high in gallbladder cancer(67%), while majority of the chronic cholecystitis (94%) and xanthogranulomatous cholecystitis (91%) cases retained positive E-cadherin expression.
Literature	GENERIF_SUMMARY	e-cadherin is a prognostic factor for non-small cell lung cancer
Literature	GENERIF_SUMMARY	The expression of E-cadherin in invasive ductal carcinoma and carcinoma in situ was significantly higher than that in invasive lobular carcinoma.
Literature	GENERIF_SUMMARY	provide new evidence that HAb18G/CD147-mediated cell-cell contact confers anoikis resistance in an E-cadherin-dependent manner
Literature	GENERIF_SUMMARY	Low expression of E-CD and high expression of Snail are related to the advanced stage, and poor prognosis in colorectal cancer patients.
Literature	GENERIF_SUMMARY	impaired down-regulation of E-cadherin and beta-catenin, along with Wnt/beta-catenin signaling pathway activation during the window of implantation, might be one of the potential molecular mechanisms of infertility in patients with endometriosis.
Literature	GENERIF_SUMMARY	Combined vorinostat and bortezomib therapy significantly decreased esophageal cancer epithelial-mesenchymal transition via E-cadherin up-regulation.
Literature	GENERIF_SUMMARY	ZEB1/BRG1 as a new transcriptional mechanism regulating E-cadherin expression and epithelial-to-mesenchymal transdifferentiation
Literature	GENERIF_SUMMARY	DDR1 can increase the stability of cell surface E-cadherin and promote MDCK cell aggregation, which may be mediated through the formation of DDR1/E-cadherin complexes.
Literature	GENERIF_SUMMARY	Loss of E-cadherin expression is associated with non-

Main Category	Category	Terms
		muscle-invasive bladder cancer.
Literature	GENERIF_SUMMARY	p140Cap immobilizes E-cadherin at the cell membrane and inhibits EGFR and Erk1/2 signalling, blocking scatter and proliferation of cancer cells.
Literature	GENERIF_SUMMARY	Novel heterozygous 2275G>T mutation(exon 14) of CDH1 gene detected in Iranian family with hereditary diffuse gastric cancer. The nonsense mutation generates premature stop codon(position 758)with truncated E-cadherin protein lacking cytoplasmic region.
Literature	GENERIF_SUMMARY	Elevated levels of serum E-cadherin is associated with late-stage colorectal carcinoma and familial adenomatous polyposis.
Literature	GENERIF_SUMMARY	Reduced E-cadherin or beta-catenin negative expression relates to dedifferentiation and progression of nonsmall cell lung cancer
Literature	GENERIF_SUMMARY	Different expression patterns of E-cadherin and p120-catenin proteins can be helpful to distinguish infiltrating lobular carcinoma from ductal carcinoma of the breast.
Literature	GENERIF_SUMMARY	individual cell movement, asymmetric colony expansion, Rho-associated kinase, and E-cadherin all influence embryonic stem cells clonogenicity, and provide additional guidance for improvement of clonogenic assays
Literature	GENERIF_SUMMARY	up-regulating expression of endogenous E-cadherin in breast cancer can inhibit proliferation, promote apoptosis, decrease mobility, and thus inhibit tumor growth in vivo.
Literature	GENERIF_SUMMARY	Results suggest that CDH1 gene methylation and H pylori infection are frequent events in samples from Brazilian patients with chronic gastritis and reinforces the correlation between H. pylori infection and CDH1 inactivation in early gastric tumorigenesis
Literature	GENERIF_SUMMARY	The E-cadherin-catenin complex is the factor indicative of metastasis and disease progression in gastric cancer.
Literature	GENERIF_SUMMARY	The study objective was to assess changes in the expressions of E-cadherin and alpha-, beta- and gamma-catenin proteins in pancreatic duct carcinoma in correlation with clinicopathological parameters, lymph node involvement and distant metastases.
Literature	GENERIF_SUMMARY	PAS staining instead of H&E on CDH1 mutation-positive prophylactic gastrectomy specimens may increase the detection rate of adenocarcinoma while reducing screening time.
Literature	GENERIF_SUMMARY	Stromal CD10 expression and relationship to the E-cadherin/beta-catenin complex in breast carcinoma
Literature	GENERIF_SUMMARY	Nuclear localization of E-cadherin in basal cell carcinoma is also associated with aggressive tumour

Main Category	Category	Terms
		features
Literature	GENERIF_SUMMARY	in the 123 TNBC cases, the prognosis of patients with an E-cadherin-negative expression was significantly worse than that of E-cadherin-positive patients (P=0.0265), especially for those in clinical stage II (P=0.002).
Literature	GENERIF_SUMMARY	loss of E-cad expression in gastric cancer correlated with peripheral blood micrometastasis, lymph node metastasis, TNM stage, and lymphatic invasion.
Literature	GENERIF_SUMMARY	Both P16 and CDH1 had different expression levels in tumor tissues compared to the adjacent normal tissues.
Literature	GENERIF_SUMMARY	Our study suggests that E-cadherin down-regulation may lead to enhancer of zeste homologue 2-mediated invasion and metastasis.
Literature	GENERIF_SUMMARY	the potential involvements of E-cadherin and beta-catenin in meningioma
Literature	GENERIF_SUMMARY	We suggest that COX-2 pathway reduces membranous E-cadherin expression in bladder transitional cell carcinoma
Literature	GENERIF_SUMMARY	TGF-beta1 and E-cadherin are closely associated with the metastasis of ovarian carcinoma.
Literature	GENERIF_SUMMARY	the motility of E-cadherin within, and away from, the cell surface are not exclusive or independent mechanisms and there is a fine balance between the two which when perturbed can have dramatic effects on the regulation of Adherens Junctions
Literature	GENERIF_SUMMARY	E-cadherin gene promoter hypermethylation was associated with low or absent expression of E-cadherin. Loss of E-cadherin protein may contribute to aberrant localization of beta-catenin.
Literature	GENERIF_SUMMARY	Expression of E-cadherin in endometrium and fallopian tubes from infertile women.
Literature	GENERIF_SUMMARY	Reduced expression of E-cadherin is associated with bronchioloalveolar carcinoma.
Literature	GENERIF_SUMMARY	Overexpressions of nuclear E-cadherin is associated with non-small cell lung cancer.
Literature	GENERIF_SUMMARY	Results suggests that the -160 AA genotype of human CDH1 is associated with an increased risk of gastric cancer in Oman.
Literature	GENERIF_SUMMARY	IGF2*A allele and CDH1*C allele were correlated with leiomyoma susceptibility, which may be associated with leiomyoma development.
Literature	GENERIF_SUMMARY	Analyses of CDH1 methylation suggested a significant difference between hypermethylated and non-hypermethylated samples, with a positive association between the -160 A allele and hypermethylation.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Promoter methylation of DAPK1, E-cadherin and thrombospondin-1 in de novo and therapy-related myeloid neoplasms.
Literature	GENERIF_SUMMARY	Data indicate that P-cadherin but not E-cadherin is important for maintaining adherens junctions in DU145 and MCF10A cells.
Literature	GENERIF_SUMMARY	induction of E-cadherin expression by small activating RNA leads to suppression of migration and invasion of PC3 prostate cancer cells.
Literature	GENERIF_SUMMARY	In lung carcinomas dysadherin expression seems to reflect tumour aggressiveness and may be considered a positive marker of poor prognosis when considered alone or/and in combination with down-regulation of E-cadherin.
Literature	GENERIF_SUMMARY	E-cadherin may be involved in the pathogenesis of chronic rhinosinusitis with nasal polyps and correlated with disease severity.
Literature	GENERIF_SUMMARY	The majority of patients with germline CDH1 mutation have foci of noninvasive or invasive gastric cancer by middle age.
Literature	GENERIF_SUMMARY	These results indicate that examination of CSE1L and E-cadherin distribution in colorectal epithelium glands may be valuable for evaluating the malignance of colorectal disease.
Literature	GENERIF_SUMMARY	Hypermethylation of E-cadherin is associated with recurrence of bladder cancer.
Literature	GENERIF_SUMMARY	PLEKHA7 is a cytoplasmic component of the epithelial adherens junction belt, distinct from ZO-1 and E-cadherin
Literature	GENERIF_SUMMARY	Suggest that TC21 promotes cell motility and metastasis by regulating the expression of E-cadherin and N-cadherin in hepatocellular carcinoma and is associated with tumor progression and poor prognosis in HCC.
Literature	GENERIF_SUMMARY	E-cadherin proteolyis and nuclear translocation associates with aggressive foci growth only in the peritoneal microenvironment in colorectal cancer.
Literature	GENERIF_SUMMARY	Loss of E-cadherin expression may be a marker of plasmacytoid and signet ring cell differentiation in urothelial carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin expression in nasopharyngeal carcinoma is negatively correlated with the numbers of the lymph node metastases and the metastasis distance.
Literature	GENERIF_SUMMARY	Mutational analysis of E-cadherin, in synovial sarcomas
Literature	GENERIF_SUMMARY	CDH1 gene mutation is not associated with hereditary diffuse gastric cancer.
Literature	GENERIF_SUMMARY	metalloproteinase MMP-2 and MMP-9 -to-E-cadherin

Main Category	Category	Terms
		ratio has an effect on lymphangiogenesis and lymph node metastasis in prostate cancer
Literature	GENERIF_SUMMARY	plakoglobin and E-cadherin recruit plakophilin3 to the cell border to initiate desmosome formation
Literature	GENERIF_SUMMARY	Loss of E-cadherin is associated with colorectal carcinoma.
Literature	GENERIF_SUMMARY	A negative correlation is found between Ezrin and E-cadherin expression in nasopharyngeal carcinoma tissues. Ezrin and E-cadherin are closely related to clinical staging and cervical lymph node metastases.
Literature	GENERIF_SUMMARY	These results provide evidence that E-cadherin is expressed in a proportion of invasive lobular carcinoma of the breast, however, unlike ductal carcinoma, its expression seems to be of limited significance.
Literature	GENERIF_SUMMARY	Hypermethylation of the CDH1 gene, resulting in low E-cadherin expression in endometrial carcinoma is associated with clinicopathological progress and the 5-year overall survival rate.
Literature	GENERIF_SUMMARY	rs16260 overall genotype frequencies in cleft palate only groups were significantly different with controls and rs16260 AA genotype significantly increased risk of CPO by 5.90-fold, providing 1st evidence of CDH1 genetic variation in etiology of CPO.
Literature	GENERIF_SUMMARY	ECAD inhibits Smad3/2 phosphorylation by recruiting RhoA to p120-ctn at the p120-ctn binding domain, whereas the loss of ECAD due to cadherin switching promotes the up-regulation of TGFbeta1 and its target genes, and facilitates liver fibrosis.
Literature	GENERIF_SUMMARY	Potentially pathogenic germline CDH1 mutations in women with early-onset or familial lobular breast cancer are at most infrequent.
Literature	GENERIF_SUMMARY	Data show that E-cadherin and alpha-catenin were predominantly expressed in the cell membranes, whereas beta- and gamma-catenin were found both in the cell membrane and cytoplasm.
Literature	GENERIF_SUMMARY	claudin-1 has anti-apoptotic effects, and is involved in the regulation of the expression and subcellular localization of beta-catenin and E-cadherin in MCF-7, but not T47 D cells
Literature	GENERIF_SUMMARY	CTGF downregulated the expression of E-cadherin through activation of the nuclear factor-kappaappa B (NF-kappaB) pathway.
Literature	GENERIF_SUMMARY	findings show that p53 has an important role in the progression from serous borderline ovarian tumors to invasive carcinomas. In addition, our findings suggest that downregulation of E-cadherin by the PI3K/Akt pathway contributes to this progression
Literature	GENERIF_SUMMARY	GLI1-upregulated MUC5AC interfered with the membrane localization of E-cadherin, leading to

Main Category	Category	Terms
		decreased E-cadherin-dependent cell-cell adhesion and promoting the migration and invasion of pancreatic ductal adenocarcinoma cells.
Literature	GENERIF_SUMMARY	Galpha(12) binds to p120(ctn) and modulates its phosphorylation status through a Rho-independent mechanism. Galpha(12) emerges as an important regulator of p120(ctn) function
Literature	GENERIF_SUMMARY	These data provide novel insights into the tumor-suppressor function of E-cadherin, which contributes to the suppression of c-Jun protein translation and transcriptional activity.
Literature	GENERIF_SUMMARY	Data show E-cadherin molecular dynamics in vivo following perturbation of integrin signaling by inhibiting focal adhesion kinase (FAK) and Src.
Literature	GENERIF_SUMMARY	The epigenetic change in E-Cadherin and Cyclooxygenase-2 is associated with chronic periodontitis.
Literature	GENERIF_SUMMARY	Aberrant methylation around gene promoter region may play an important part in down regulation of E-cadherin in nasopharyngeal carcinoma.
Literature	GENERIF_SUMMARY	p38 maintains E-cadherin expression by suppressing TAK1-NF-kappaB signaling, thus impeding the induction of epithelial-to-mesenchymal transition in human primary mesothelial cells.
Literature	GENERIF_SUMMARY	E-cadherin is a useful prognostic marker for ovarian clear cell adenocarcinoma patients, and paclitaxel-based chemotherapy can improve survival among patients with positive E-cadherin immunoreactivity.
Literature	GENERIF_SUMMARY	Foxo3a suppression of urothelial cancer invasiveness through Twist1, Y-box-binding protein 1, and E-cadherin regulation.
Literature	GENERIF_SUMMARY	low expression of E-cadherin contributes to the vigorous growth and transforming ability of leukaemic cells
Literature	GENERIF_SUMMARY	Dsg3, as an up-stream regulator of Src activity, helps regulate adherens junction formation through its interaction with E-cadherin
Literature	GENERIF_SUMMARY	There is no association between the CHD1 gene 3'-UTR C/T polymorphism and nephrolithiasis in our population.
Literature	GENERIF_SUMMARY	Significantly lower distribution of E-cadherin was seen in interstitial cystitis/painful bladder syndrome bladder tissue compared to control patients
Literature	GENERIF_SUMMARY	these data demonstrate that HDAC1 is a major repressive enzyme for E-cadherin expression as well as HDAC inhibitor-mediated anti-invasiveness.
Literature	GENERIF_SUMMARY	Brain metastases originating from lung adenocarcinoma and small cell lung cancer were

Main Category	Category	Terms
		significantly associated to E-cadherin genetic changes.
Literature	GENERIF_SUMMARY	Variant genotypes of the -160C>A polymorphism were associated with significantly decreased gastric cancer risk among Asians.
Literature	GENERIF_SUMMARY	E-cad could enhance adhesion and inhibit proliferation of human breast carcinoma cells through a pathway involving beta-cat and cyclin D1.
Literature	GENERIF_SUMMARY	E-cadherin appears to be a strong predictor of N stage in CRC and should be considered in pre-operative and post-operative management of colon and rectal cancer patients.
Literature	GENERIF_SUMMARY	Combined dysadherin-positive expression and E-cadherin-negative expression may be valuable information for predicting aggressive tumor behavior of differentiated-type gastric carcinoma with submucosal invasion
Literature	GENERIF_SUMMARY	Reduced levels of E-cadherin protein observed in nasopharyngeal carinoma may play an important role in invasion and metastasis.
Literature	GENERIF_SUMMARY	Data show that miR-495 expression was directly modulated by transcription factor E12/E47, and promotes oncogenesis via downregulation of E-cadherin and REDD1.
Literature	GENERIF_SUMMARY	The expression of vascular endothelial growth factor C in ovarian cancer was related to matrix metalloproteinase-2 and E-cadherin.
Literature	GENERIF_SUMMARY	Germline CDH1 mutation can be cosegregated with ILBC in the absence of DGC.
Literature	GENERIF_SUMMARY	in patients with gastric caancer, expression of E-cadherin and that of Slug were associated with tumor properties, including lymph node metastasis, stage, lymphatic invasion, and venous invasion, as well as with prognosis
Literature	GENERIF_SUMMARY	Hakai mediates E-cadherin ubiquitination and degradation triggered by Slit-Robo signaling during colorectal epithelial cell carcinogenesis.
Literature	GENERIF_SUMMARY	the loss of E-cadherin itself may contribute to dysregulated PI3K/Akt signaling through its effects on PTEN.
Literature	GENERIF_SUMMARY	Nuclear translocation of E-cadherin might contribute to ovarian folliculogenesis or granulosa/stromal cell differentiation.
Literature	GENERIF_SUMMARY	This study demonstrates that ataxin-1 occupies the promoter region of E-cadherin in vivo and that ataxin-1 activates the promoter in a CtBP2-mediated transcriptional regulation manner.
Literature	GENERIF_SUMMARY	activated STAT3 signal may associate with Twist and E-cadherin expression and mediate HCC invasiveness

Main Category	Category	Terms
		and metastasis.
Literature	GENERIF_SUMMARY	A reduction in the expression of E-cadherin was observed in vulvar squamous cell carcinoma
Literature	GENERIF_SUMMARY	Data suggest that SDF-1 down-regulates the E-cadherin/beta-catenin complex expression in HT29 cells by decreasing mRNA synthesis and increasing beta-catenin phosphorylation.
Literature	GENERIF_SUMMARY	The CDH1 gene was screened for mutations in well-characterized, Finnish, high-risk hereditary breast and/or ovarian cancer individuals.
Literature	GENERIF_SUMMARY	Kras(G12D) allele promotes metastasis in pancreatic cancer cells partly through the downregulation of E-cadherin.
Literature	GENERIF_SUMMARY	Fyn, a member of Src family tyrosine kinases, plays a critical role in mediating TGF-beta1-induced down-regulation of E-cadherin in human A549 lung cancer cells.
Literature	GENERIF_SUMMARY	Our data demonstrate that PIK3CA is a mediator of collagen I-induced down-regulation of E-cadherin.
Literature	GENERIF_SUMMARY	expression of E-cadherin was shown to be a predictor of response to trastuzumab-based treatment for HER2-overexpressing carcinomas
Literature	GENERIF_SUMMARY	CDH1 mutation are associated with signet ring cell adenocarcinoma of the stomach.
Literature	GENERIF_SUMMARY	The expression of E-cadherin was positively correlated with differentiated degree and lymphatic metastasis.
Literature	GENERIF_SUMMARY	These results suggested that the 3'-UTR +54C/T polymorphism in CDH1 may be a marker for genetic susceptibility of cancer.
Literature	GENERIF_SUMMARY	Down-regulation of E-cadherin and the up-regulation of nuclear beta-catenin expression might be crucial steps during tumour progression of tonsillar carcinomas.
Literature	GENERIF_SUMMARY	There was a positive correlation between STAT3 and vimentin expression and a negative correlation between STAT3 and E-cadherin in colon cancer.
Literature	GENERIF_SUMMARY	germline variants in the CDH1 gene contribute to a predisposition to the development of primary gastric diffuse large B-cell lymphomas.
Literature	GENERIF_SUMMARY	Single nucleotide polymorphisms in the CDH1 gene promoter is associated with development of hepatocellular carcinoma.
Literature	GENERIF_SUMMARY	These results suggest that CDH1-347 polymorphisms are associated with increased risks of oral cancer, and may be a predictive factor for tumor lymph node metastasis.
Literature	GENERIF_SUMMARY	These findings show an important role for p53 in the

Main Category	Category	Terms
		progression of serous borderline ovarian tumors to an invasive carcinoma, and suggest that downregulation of E-cadherin by DNMT1-mediated promoter methylation contributes to this process.
Literature	GENERIF_SUMMARY	High E-cadherin is associated with metastatic lymph nodes of esophageal squamous cell carcinoma.
Literature	GENERIF_SUMMARY	Cancerous tissues from 30 patients with HPSCC were examined for LOH in 4 tumor suppressor genes (TSGs) (p16, Rb, E-cadherin, and p53) at loci 9p21, 13q21, 6q22, and 17p13, respectively, using microsatellite markers amplified by polymerase chain reaction.
Literature	GENERIF_SUMMARY	Our results indicate that the cadherin switch alters through progression of ovarian high-grade serous carcinoma .
Literature	GENERIF_SUMMARY	These results indicate that PIPKIIbeta-mediated PI(4,5)P(2) signaling is important for E-cadherin upregulation and inhibition of cellular motility induced by VDR activation.
Literature	GENERIF_SUMMARY	Loss of E-cadherin is associated with triple-negative breast cancer.
Literature	GENERIF_SUMMARY	MMP20 may influence ameloblast developmental progression through hydrolysis of cadherin extracellular domains with associated release of transcription factor(s).
Literature	GENERIF_SUMMARY	Data suggest that SPARCL1, Shp2, MSH2, E-cadherin, p53, ADCY-2 and MAPK are potential prognostic markers in colorectal cancer.
Literature	GENERIF_SUMMARY	CDH1 polymorphisms are associated with epithelial E-cadherin expression contributor to airway remodelling and lung function in asthma.
Literature	GENERIF_SUMMARY	E-cadherin has a role in antimigratory and antiinvasive efficacy of silibinin in prostate cancer cells
Literature	GENERIF_SUMMARY	Among patients undergoing pancreaticoduodenectomy for pancreatic ductal adenocarcinoma, partial loss of tumoral E-cadherin expression is an independent predictor of poor outcome
Literature	GENERIF_SUMMARY	Activation of epidermal growth factor receptor promotes squamous carcinoma cell migration and invasion via inducing epithelial mesenchymal transformation-like phenotype change and matrix metallproteinase-9-mediated degradation of E-cadherin
Literature	GENERIF_SUMMARY	Data suggest that CDH1 -160C>A gene polymorphism may contribute to increased risk of gastric cancer among Caucasians[meta-analysis]
Literature	GENERIF_SUMMARY	When both E-cadherin and beta -catenin expressions were reduced, there was a significant unfavourable prognosis.
Literature	GENERIF_SUMMARY	Corticotroph tumour progression was associated with

Main Category	Category	Terms
		reduced expression of the epithelial marker E-cadherin.
Literature	GENERIF_SUMMARY	the role of the C-160A single-nucleotide polymorphism (SNP) of CDH1 in susceptibility to gastric cancer
Literature	GENERIF_SUMMARY	these findings show that epithelial-mesenchymal transition and its down-regulated expression of E-cad circumvent breast cancer dormancy in part by facilitating beta1 integrin expression necessary for metastatic outgrowth.
Literature	GENERIF_SUMMARY	elevation of E-cadherin, a cell adhesion molecule and suppression of beta-catenin, a proto-oncogene have been observed in presence of CaSR agonists whereas reverse effect has been seen in presence of CaSR antagonist
Literature	GENERIF_SUMMARY	The A allele of the CDH1 -160 C/A polymorphism confers a decreased stomach cancer risk in Asians but not in Caucasians.
Literature	GENERIF_SUMMARY	No association was found between E-cadherin immunoreactivity and nodal metastasis in oral tongue squamous cell carcinoma.
Literature	GENERIF_SUMMARY	These data suggest that loss of E-cadherin expression is associated with increased lymhogeneous metastasis of HNSCC.
Literature	GENERIF_SUMMARY	These findings suggest that hepatitis C virus core protein could play a role in hepatocellular carcinoma at least in part by altering the methylation status of CDH1 promoter in hepatocytes.
Literature	GENERIF_SUMMARY	E-cadherin expression in adenocarcinoma cells is regulated by slug.
Literature	GENERIF_SUMMARY	In the current study of patients with tongue cancer, decreased E-cadherin and increased vimentin were found in spreading tumor satellites
Literature	GENERIF_SUMMARY	Abnormal expression of E-cadherin, beta-catenin, Wnt-1 was observed.
Literature	GENERIF_SUMMARY	although genetic loss of Brca1 and E-cadherin are infrequent in breast cancer, they are down-regulated at the transcriptional level by CtBP1 expression.
Literature	GENERIF_SUMMARY	Data suggest a critical role of EZH2 in the control of cell invasionand/or metastasis by forming a co-repressor complex with HDAC1/HDAC2/Snail to repress E-cadherin.
Literature	GENERIF_SUMMARY	The results uncovered a novel function of Rack1 in maintaining the junctional homeostasis of intestinal epithelial cells by regulation of the Src- and growth factor-induced endocytosis of E-cadherin.
Literature	GENERIF_SUMMARY	H1 and PAR2 receptors enhance delivery of immune-competent cells and molecules by interrupting E-cadherin adhesion in lung epithelial cells.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Results sugggest that histone deacetylation of E-cadherin (CDH1) and downregulation of miR-373, together with the hypermethylation of CDH1 by hepatitis B virus-encoded X antigen, HBx, may be important for the understanding of HBV-related carcinogenesis.
Literature	GENERIF_SUMMARY	Suppression of E-cadherin function drives the early stages of Ras-induced squamous cell carcinoma through upregulation of FAK and Src.
Literature	GENERIF_SUMMARY	E-cadherin-mediated cell-cell contact directly regulates the Hippo signaling pathway to control cell proliferation.
Literature	GENERIF_SUMMARY	We confirmed the association between ulcerative colitis and CDH1.
Literature	GENERIF_SUMMARY	Data suggest that the TBX2/NRAGE complex protects cells against anoikis by downregulating p14ARF, representing a novel pathway for the regulation of anoikis by epithelial-to-mesenchymal transition and E-cadherin.
Literature	GENERIF_SUMMARY	E-cadherin was decreased by EGF, but increased by luteolin and quercetin.
Literature	GENERIF_SUMMARY	Cas proteins do not affect E-cadherin transcription, but rather, BCAR1 and NEDD9 signal through SRC to promote E-cadherin removal from the cell membrane and lysosomal degradation.
Literature	GENERIF_SUMMARY	Data show that CD99 combined with E-cadherin/beta-catenin and CD10 can be used as a relatively specific expression profile of SPTs.
Literature	GENERIF_SUMMARY	This is the first report of CDH1 germline truncating mutations in Japanese patients with familial gastric cancer.
Literature	GENERIF_SUMMARY	Data suggest that regulation of E-cadherin expression in proximal tubule cells involves TGFb1IRS2FoxO3a pathway
Literature	GENERIF_SUMMARY	Expression of E-cadherin is downregulated, and expression of P-cadherin is upregulated in oral mucosa of mucous membrane pemphigoid (MMP), which may be involved in the pathogenesis of MMP.
Literature	GENERIF_SUMMARY	Results suggest that decreased E-cadherin and increased MMP-2 may be associated with the capacity of GCSCs to metastasize.
Literature	GENERIF_SUMMARY	Cath-D and E-Cad essays may useful in identifying neck lymph node involvement.
Literature	GENERIF_SUMMARY	CDH1 gene promoter methylation may lead to aberrant expression of E-cadherin and beta-catenin in colonic carcinoma, and may play an important role in promoting tumor invasion.
Literature	GENERIF_SUMMARY	TGF-beta1-induced activation of Smad3 is the critical step for the uPA upregulation and E-cadherin

Main Category	Category	Terms
		downregulation
Literature	GENERIF_SUMMARY	Data show that EphB receptors interact with E-cadherin and with the metalloproteinase ADAM10 at sites of adhesion.
Literature	GENERIF_SUMMARY	E-cadherin is effectively restricted to supporting cell (SC)-SC junctions in human and murine vestibular epithelia
Literature	GENERIF_SUMMARY	Results demonstrate that the increase in E-cadherin mobility is constitutively altered by the presence of CAR at FLCARMCF7 cell junctions.
Literature	GENERIF_SUMMARY	E-cadherin,alpha-dystroglycan and beta-dystroglycan levels were decreased in the oesophageal primary tumour samples, despite the presence of normal levels of dystroglycan mRNA.
Literature	GENERIF_SUMMARY	Nuclear translocation of beta-catenin synchronized with loss of E-cadherin in oral epithelial dysplasia with a characteristic two-phase appearance
Literature	GENERIF_SUMMARY	Dietary intakes of folate, vitamins B2, B6, B12, and methionine were not associated with likelihood of promoter methylation of E- cadherin, p16, and RAR-beta2.
Literature	GENERIF_SUMMARY	study presents evidence that in the cadherin-catenin complex alpha-catenin contributes to the binding strength of another catenin, p120, to the same complex; data suggest that alpha-catenin-p120 contact within the cadherin-catenin complex can regulate cadherin trafficking
Literature	GENERIF_SUMMARY	Downregulation of E-Cadherin enhances proliferation of head and neck cancer through transcriptional regulation of epidermal growth factor receptor
Literature	GENERIF_SUMMARY	Low E-cadherin is associated with metastasis in colorectal cancer.
Literature	GENERIF_SUMMARY	Our findings suggest further investigation into the prognostic role of E-cadherin in pediatric Colorectal adenocarcinoma
Literature	GENERIF_SUMMARY	E- and N-cadherins are expressed at the surface of human beta-cells and that these adhesion molecules are involved in the maintenance of beta-cell viability
Literature	GENERIF_SUMMARY	PLGA was successfully confirmed to support parotid gland acinar cells to form more post-confluence structure by enhancing E-cadherin expression.
Literature	GENERIF_SUMMARY	Certain CDH1 gene promoter -160 C > A and -347 G > GA variants might affect the susceptibility of colorectal cancer.
Literature	GENERIF_SUMMARY	E-cadherin/beta-catenin complex and the epithelial barrier
Literature	GENERIF_SUMMARY	there are correlations between beta-catenin and TNM

Main Category	Category	Terms
		stage of colorectal cancer; there are greater amounts of beta-catenin in patients with deeper invasion of the tumor in colon layers, lymph node involvement as well as in patients with distant metastasis
Literature	GENERIF_SUMMARY	Study concludes that Smad3 regulates, at the transcriptional level, miR-200 family members, which themselves regulate ZEB1 and ZEB2, known transcriptional repressors of E-cadherin, at the posttranscriptional level in a TGF-beta-independent manner.
Literature	GENERIF_SUMMARY	p120 dynamically regulates Rho-GTPase activity at the E-cadherin complex through transient interaction with several of its up- and downstream effectors, including ROCK1
Literature	GENERIF_SUMMARY	results showed that NOTCH1 affected skin cancer cell motility by changing the expression of E-cadherin
Literature	GENERIF_SUMMARY	PIPKIgamma and phosphatidyl inositol phosphate pools at nascent E-cadherin contacts cue Exo70 targeting and orient the tethering of exocyst-associated E-cadherin
Literature	GENERIF_SUMMARY	Loss of E-cadherin is associated with invasion in esophageal cancers.
Literature	GENERIF_SUMMARY	E-cadherin is extensively expressed by oropharyngeal squamous cell carcinoma, even the non-keratinizing type, but our results suggest that cadherin expression may not be a predictor for nodal or distant metastasis in these tumors.
Literature	GENERIF_SUMMARY	Cholangiocarcinoma cell lines express E-cad and beta-cat but with different localizat- ion patterns.
Literature	GENERIF_SUMMARY	E-cadherin nuclear staining is useful in distinguishing between adult granulosa cell tumors and ovarian adenocarcinomas or carcinoid tumors
Literature	GENERIF_SUMMARY	Loss of E-CD expression is an early event in lobular neoplasia of the breast.
Literature	GENERIF_SUMMARY	The regulation and role of E-cadherin during melanoma development is the focus of this review. [review]
Literature	GENERIF_SUMMARY	The E-cadherin expression was reduced to negativity in advanced stages of gastric carcinoma.
Literature	GENERIF_SUMMARY	The combination of E-cadherin and Ki67 status might be a useful prognostic marker indicating the need for adjuvant chemotherapy in Stage II TNBC patients.
Literature	GENERIF_SUMMARY	This study finds a new mechanism that miR-9 utilizes to decrease E-cadherin expression and inhibit melanoma progression
Literature	GENERIF_SUMMARY	upregulation of connexin 43 in glioma stem cells inhibited their capacity of self-renewal, invasiveness, and tumorigenicity via modulation of E-cadherin.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Membranous E-cadherin immunostaining and cytoplasmic COX-2 expression were observed in 74.3% and 68.6% of cases respectively.
Literature	GENERIF_SUMMARY	examined integrin beta(1) , ECAD and rac1 expression in SCLC and analyzed the prognostic value of these markers
Literature	GENERIF_SUMMARY	Snail1 enhances the binding of Akt2 to the E-cadherin (CDH1) promoter and Akt2 interference prevents Snail1 repression of CDH1 gene
Literature	GENERIF_SUMMARY	CRYAB functions to suppress nasopharyngeal carcinoma progression by associating with the cadherin/catenin adherens junction and modulating the beta-catenin function.
Literature	GENERIF_SUMMARY	Loss of decorin and E-cadherin accelerated colon cancer cell growth and invasion in Dcn(-/-) mice.
Literature	GENERIF_SUMMARY	The regulation, function, and implications of E-cadherin expression in these central orchestrators of the immune system.
Literature	GENERIF_SUMMARY	The up-regulation of Ezrin and Moesin and the down-regulation of E-Cadherin were associated with the invasion and metastasis of papillary thyroid carcinoma.
Literature	GENERIF_SUMMARY	Differential expression of E-cadherin has an important role in pathogenesis of endometriosis.
Literature	GENERIF_SUMMARY	E-cadherin is necessary for localization of DLG1 but not phosphorylated MEK2 to the midbody ring during cytokinesis.
Literature	GENERIF_SUMMARY	genetic association studies in Han Chinese population
Literature	GENERIF_SUMMARY	The data demonstrate that Cdc6 acts as a molecular switch at the E-cadherin locus, linking transcriptional repression to activation of replication
Literature	GENERIF_SUMMARY	the roles of STAT3 in colorectal cancer epithelial-mesenchymal transition and in STAT3-induced down-regulation of E-cadherin
Literature	GENERIF_SUMMARY	E-cadherin could be upregulated by the suppression of ZEB1 transcriptional repressor by miRNAs in vivo
Literature	GENERIF_SUMMARY	Epigenetic regulation is a mechanism through which tumor cell colonization of metastatic sites occurs as E-cadherin-expressing cells arise from E-cadherin-deficient cells.
Literature	GENERIF_SUMMARY	Compared with conventional carcinomas, serrated adenocarcinomas showed significantly reduced membranous E-cadherin.
Literature	GENERIF_SUMMARY	CDH1 promoter methylation is associated with triple-negative breast cancer.
Literature	GENERIF_SUMMARY	The combination of a decrease of E-cadherin and an increase in vimentin might be a valuable survival

Main Category	Category	Terms
		indictor in cervical squamous cell cancer.
Literature	GENERIF_SUMMARY	E-cadherin genetic screenings performed in low-risk areas for gastric cancer identified a higher frequency of CDH1 germline mutations.
Literature	GENERIF_SUMMARY	Negative/low HER2 expression alone or combined with E-cadherin positivity is predictive of better prognosis in Tunisian patients with breast carcinoma.
Literature	GENERIF_SUMMARY	Hakai dimerization allows the formation of a phosphotyrosine-binding pocket that recognizes specific phosphorylated tyrosines and flanking acidic amino acids of Src substrates, such as E-cadherin, cortactin and DOK1.
Literature	GENERIF_SUMMARY	Data suggest that TGF-beta, TGF-betaR1, TGF-betaR2, Smad4, pSmad2/3, and E-cadherin are closely related to tumor-node-metastasis (TNM) stage of colorectal cancer (CRC).
Literature	GENERIF_SUMMARY	The reduced expression of E-cadherin and beta-catenin and EGFR over expression seems to be correlated with tumor differentiation and tumor progression than tumor invasion and tumor proliferation.
Literature	GENERIF_SUMMARY	Associations between RASSF1A and CDH1 methylation levels and clinico-pathological parameters manifested various levels of RASSF1A and CDH1 promoter methylation.
Literature	GENERIF_SUMMARY	Loss of E-cadherin is associated with liposarcoma.
Literature	GENERIF_SUMMARY	E-cadherin inhibits nuclear accumulation of Nrf2 and Nrf2-mediated gene transcription by its interaction with Nrf2; findings imply that chemoresistance of cancer cells upon the loss of E-cadherin might be associated with Nrf2
Literature	GENERIF_SUMMARY	Both membranous and cytoplasmic localization of high E-cadherin staining were associated with lymph node metastasis in nasopharyngeal carcinoma
Literature	GENERIF_SUMMARY	E-cadherin can act as a central modulator of the cell biological phenotypes and a potential prognostic marker in Colorectal cancer.
Literature	GENERIF_SUMMARY	CDH1 polymorphisms can contribute to the etiology of premalignant skin lesions in people chronically exposed to arsenic in drinking water, and that this gene may be a factor in individual susceptibility to cutaneous diseases.
Literature	GENERIF_SUMMARY	Our results suggest that endogenous PAI-1 suppresses expression of E-cadherin and differentiation in PAC cells in vitro, supporting its negative impact on tumor prognosis.
Literature	GENERIF_SUMMARY	TWIST may act upstream of E-cadherin, which can indirectly regulate the expression levels of beta-catenin in bladder cancer.
Literature	GENERIF_SUMMARY	the association between CDH1 -C160A genetic

Main Category	Category	Terms
		polymorphism and colorectal cancer susceptibility
Literature	GENERIF_SUMMARY	The modifications of E-cadherin by O-GlcNAcylation and lack of pro-region processing represent novel mechanisms for rapid regulation of cell surface transport of E-cadherin in response to intoxication.
Literature	GENERIF_SUMMARY	Snail interacted with G9a and recruited G9a and DNA methyltransferases to the E-cadherin promoter for DNA methylation.
Literature	GENERIF_SUMMARY	E-cadherin expression was present in 100% of cases of both adenocarcinomas and adenomas, with prevailing strong membranous immunoreactivity and no differences between protein expression in tumors and normal mucosa.
Literature	GENERIF_SUMMARY	Results show that the adipocytokines and glucocorticoid metabolism-related genes are overexpressed in colorectal adenocarcinomas, and expression of these genes is associated with the downregulation of E-cadherin mRNA.
Literature	GENERIF_SUMMARY	Data indicate that up-regulation of Slug was significantly correlated with a higher tumor stage and the E- to N-cadherin switch in bladder cancer cells and tissues.
Literature	GENERIF_SUMMARY	Data suggest that pulmonary metastasis is corrected with levels of Geminin, cleaved caspase-3, CD44, E-cadherin, epidermal growth factor receptor, and CD204 in cancer cells within permeated lymphatic vessels.
Literature	GENERIF_SUMMARY	E-cadherin gene polymorphism rs4783689 was marginally associated with endometriosis in the Japanese population
Literature	GENERIF_SUMMARY	by promoting EGFR internalization, reggie-1 restricts EGFR signaling involved in E-cadherin macropinocytosis and recycling and regulate adherens junction formation
Literature	GENERIF_SUMMARY	Deletions and/or microdeletions at both miR-101 genomic loci cause mature miR-101 down-regulation, subsequent EZH2 over-expression and E-cadherin dysfunction, specifically in intestinal-type gastric carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin structural models suitable to predict the impact of the majority of cancer-associated missense mutations
Literature	GENERIF_SUMMARY	High E-cadherin is associated with invasion and metastasis of gastric cancer.
Literature	GENERIF_SUMMARY	The down-regulation of E-cadherin, as modulated by the methylation of CDH1, may contribute to nerve invasion, lymphatic and distant metastasis in salivary adenoid cystic carcinoma.
Literature	GENERIF_SUMMARY	Slug is expressed markedly higher while E-cadherin markedly lower in metastatic gastrointestinal stromal tumors.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Although the expressions of E-cadherin and N-cadherin appear to be correlated with survival outcomes in bladder cancer, N-E cadherin switch may be a better predicator for postoperative recurrence and cancer-related survival.
Literature	GENERIF_SUMMARY	E-cadherin promoter hypermethylation and intake of local hot salted tea and sun-dried foods are associated with gastric tumors.
Literature	GENERIF_SUMMARY	Activation of E-cadherin induces p120-catenin dephosphorylation, as well as changes in the cadherin cytoplasmic domain.
Literature	GENERIF_SUMMARY	CDH1 methylation was found in 75 of 92 (81.5%) gastric cancer tissues, which significantly correlated with size, growth pattern, differentiation, lymphatic invasion, venous invasion, invasion depth, lymph node metastasis, distant metastasis
Literature	GENERIF_SUMMARY	E-cadherin was significantly associated with histological grade and alcohol consumption, and beta-catenin was significantly associated with nodal stage, TNM stage, and E-cadherin expression.
Literature	GENERIF_SUMMARY	Our findings suggested that the regulatory function of 4HPR on infiltration of bladder cancer cells T24 is at least partly achieved by regulating the expression of E-Cad.
Literature	GENERIF_SUMMARY	our results indicated that plasma CDH1 levels may serve as a risk marker against gastric cancer and variant genotypes of rs26160 and rs17690554 may contribute to the etiology of diffuse gastric cancer in this study.
Literature	GENERIF_SUMMARY	Growth and proliferation of SKOV-3 cells was efficiently suppressed after knocking-down of E-cadherin expression.
Literature	GENERIF_SUMMARY	The expression changes of TGF-beta1 and E-cadherin may be related to the emergence and the development of glioma.
Literature	GENERIF_SUMMARY	E-cadherin is critical for collective sheet migration and is regulated by the chemokine CXCL12 protein during restitution
Literature	GENERIF_SUMMARY	preoperative serum E-cadherin concentrations are higher in patients with less differentiated prostate cancer
Literature	GENERIF_SUMMARY	Snail interacted with Suv39H1 and recruited it to the E-cadherin promoter for transcriptional repression.
Literature	GENERIF_SUMMARY	Histone deacetylase inhibition induced the E-cadherin expression and the proper membrane localization of the E-cadherin/beta-catenin complex, leading to reduced cancer cell migration and invasion.
Literature	GENERIF_SUMMARY	alpha- and beta-catenins may be important in the early stages of phyllodes tumours development, while E-

Main Category	Category	Terms
		cadherin may be required for malignant development
Literature	GENERIF_SUMMARY	Rho-GTPase-dependent modulation of cytoskeletal function and downregulation of E-cadherin expression are identified as relevant effectors of the miR-10b-syndecan-1 axis
Literature	GENERIF_SUMMARY	Twist was expressed significantly more and E-cadherin significantly less in gastrointestinal stromal tumors (GISTs) with metastasis, and expression of both was closely related to metastasis of GISTs.
Literature	GENERIF_SUMMARY	This calpain-mediated proteolysis of E-cad generates the formation of the lymphovascular embolus and is responsible for its unique properties of increased homotypic adhesion, apoptosis resistance and budding.
Literature	GENERIF_SUMMARY	E-cadherin stimulates the synthesis of intervertebral disc matrix macromolecules aggrecan and collagen II through the induction of BMP genes and enhancement of the Smad1/5 phosphorylation.
Literature	GENERIF_SUMMARY	The M4 mutant, which contains N-terminal 1-54 amino acids and the Armadillo repeat domain, was functional in regulating E-cadherin.
Literature	GENERIF_SUMMARY	these findings identified epigenetic silencing of CDH1 in cancer cells might be a new molecular event of multidrug resistance.
Literature	GENERIF_SUMMARY	Data indicate that Rab7 siRNA knockdown causes increased E-cadherin.
Literature	GENERIF_SUMMARY	Specifically, NiCl(2) induces down-regulation of E-cadherin by reactive oxygen species generation and promoter hypermethylation
Literature	GENERIF_SUMMARY	Downregulation of E-cadherin may be valuable as risk marker for development of multiple tumours in oral cavity and for diagnosis of premalignant fields.
Literature	GENERIF_SUMMARY	Data show that Acinetobacter baumannii transposase Tnp in A549 cells induced DNA methylation of CpG regions in the promoters of E-cadherin (CDH1) gene, whereas the cytoplasmic localization of the truncated Tnp without NLSs did not induce DNA methylation.
Literature	GENERIF_SUMMARY	Investigated the expression of RKIP and E-cadherin in lung squamous cell carcinoma tissue. The rates of positive RKIP and E-cadherin mRNA expression were significantly lower in lung squamous cell carcinoma than in the surrounding normal tissues.
Literature	GENERIF_SUMMARY	Trends in the expression of E-cadherin suggests its involvement in oral carcinogenesis via Wnt pathway dysregulation.
Literature	GENERIF_SUMMARY	these results demonstrate a Fyn kinase-dependent mechanism through which IFNgamma regulates E-cadherin stability and suggest a novel mechanism of disruption of epithelial cell contact, which could contribute to perturbed epithelial barrier function.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	The aim of the present paper was to evaluate Wwox and TAZ, nuclear effectors of Hippo-related pathways, were involved in E-cadherin expression in bone metastases specimens from breast cancer.
Literature	GENERIF_SUMMARY	Approximately 38% of HDGC families have a CDHI gene mutation.
Literature	GENERIF_SUMMARY	Decreased E-cadherin expression was associated with poor survival in patients with non-small cell lung cancer, especially among Asians. [Meta-analysis]
Literature	GENERIF_SUMMARY	findings show that alpha-catulin is highly expressed in melanoma cells; expression of alpha-catulin promoted melanoma progression and occurred concomitantly with the downregulation of E-cadherin and the upregulation of expression of mesenchymal genes
Literature	GENERIF_SUMMARY	findings suggest that miR-23a regulates TGF-beta-induced epithelial-mesenchymal transition by targeting E-cadherin in lung cancer cells
Literature	GENERIF_SUMMARY	Transcription initiation arising from CDH1 intron 2 encoded a nove lCDH1a that increases gastric cancer cell invasion and angiogenesis.
Literature	GENERIF_SUMMARY	high intracellular PKCK2 activity confers anoikis resistance on esophageal cancer cells by inducing E- to N-cadherin switching.
Literature	GENERIF_SUMMARY	The expressions of E-cd and alpha-cat are significantly lower in prostate cancer than in benign prostatic hyperplasia, and they are not associated with cancerous metastasis, but negatively correlated with the PSA level in PCa patients.
Literature	GENERIF_SUMMARY	These findings establish a principal biological function of SIRT1 in the modulation of E-cadherin function.
Literature	GENERIF_SUMMARY	A meta-analysis indicated that the -160A allele of E-cadherin provides a higher risk for the development of prostate and urothelial cancers and a protective role for colorectal cancer in an ethnicity-dependent manner.
Literature	GENERIF_SUMMARY	These results provide strong evidence that the methylation status of E-cadherin gene contributes to a reduction in the expression of E-cadherin mRNA, and may play a role in the development and progression of non-small cell lung cancer.
Literature	GENERIF_SUMMARY	impact of cell-cell contact, E-cadherin and EGF receptor on the cellular radiosensitivity
Literature	GENERIF_SUMMARY	we have demonstrated an association of epithelial E-cadherin expression with stromal histologic features and disease recurrence in phyllodes tumor.
Literature	GENERIF_SUMMARY	invasiveness was significantly correlated to the loss of membranous E-cadherin, as well as to increased ASMA, S100A4, and PDGFRbeta in stromal cells.
Literature	GENERIF_SUMMARY	In the -197 A allele carrier with *1249 CC homozygote of

Main Category	Category	Terms
		IL17A, the methylation of DAPK and CDH1 increased gradually, but more rapidly, with age.
Literature	GENERIF_SUMMARY	We conclude that determination of E-cadherin expression can be used as an adjunct in selecting patients who may benefit from adjuvant chemotherapy in the presence of otherwise favorable prognostic factors.
Literature	GENERIF_SUMMARY	These findings indicate the existence of a novel E-cadherin-related mechanism by which miR-10b modulates breast cancer metastasis.
Literature	GENERIF_SUMMARY	Our results suggest that Twist was expressed significantly more and E-cadherin significantly less in osteosarcoma with metastasis
Literature	GENERIF_SUMMARY	E-cadherin cytoplasmic mutations interfere with the binding of key exocytosis-related partners
Literature	GENERIF_SUMMARY	Loss of E-cadherin is associated with hepatocellular carcinoma.
Literature	GENERIF_SUMMARY	High Slug and low E-cadherin expression in basal-like breast cancer (BLBC) correlated with histological grade, TNM stage, and lymphatic metastasis. Poor prognosis of BLBC is associated high Slug and vimentin expression and low E-cadherin levels.
Literature	GENERIF_SUMMARY	Reduced expression of E-cadherin is associated with ovarian cancer.
Literature	GENERIF_SUMMARY	There is an association between decreased E-Cadherin immunoexpression and tumor recurrence in low-grade and non-muscle invasive transitional cell carcinoma of the bladder.
Literature	GENERIF_SUMMARY	changes in the expression of Snail or E-cadherin might regulate Epithelial-mesenchymal transition development in cholangiocarcinoma
Literature	GENERIF_SUMMARY	Data indicate that cultured tubular cells (hPTECs) reorganized into patches of cells with homotypic N-cadherin or E-cadherin cell-cell adhesions.
Literature	GENERIF_SUMMARY	Recruitment of histone deacetylases by transcriptional repressors LEF-1 and Slug is responsible for E-cadherin suppression and epithelial-to-mesenchymal transition in cigarette smokers.
Literature	GENERIF_SUMMARY	E-cadherin may serve as an additional diagnostic marker for angiomyolipoma.
Literature	GENERIF_SUMMARY	CDH1 mutation is associated with gastric and breast cancer.
Literature	GENERIF_SUMMARY	CDH1 methylation-positive cervical cancer patients had a 7.8-fold risk for death and a 92.8-fold risk for relapse.
Literature	GENERIF_SUMMARY	Nanog-mediated cell migration and invasion involved its regulation of E-cadherin and FOXJ1.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Low cadherin 1 expression is associated with hepatocellular carcinoma invasion.
Literature	GENERIF_SUMMARY	Ezrin, E-cadherin and CD44V6 play an important role in the regulation of growth and meastasis of lung cancer.
Literature	GENERIF_SUMMARY	E-cadherin expression correlated negatively with PMN infiltration, compatible with the notion that E-cadherin is cleaved by PMN-derived elastase
Literature	GENERIF_SUMMARY	Studies indicate that E-cadherin and versican are involved in cancer epithelial-mesenchymal transitions and metastasis.
Literature	GENERIF_SUMMARY	the loss of CDH1/E2F4 may be associated with worse clinical and pathological findings in mammary ductal carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin expression in nevi is related to the degree of melanocytic maturation.
Literature	GENERIF_SUMMARY	The overexpression of SLUG and TWIST with down-regulation of E-cadherin was characteristic findings in hemangiopericytomas and solitary fibrous tumors, but not in meningiomas.
Literature	GENERIF_SUMMARY	SNAI1 expression was significantly correlated with lower expression of CDH1 in colorectal adenoma.
Literature	GENERIF_SUMMARY	The aim of this study is to evaluate the role of p130cas, E-cadherin, and beta-catenin expression in patients with non-small cell lung cancer.
Literature	GENERIF_SUMMARY	The drug-resistant MCF-7/Adr cells were found to be more invasive/metastatic than their parental control, possibly related to the loss of E-cadherin expression.
Literature	GENERIF_SUMMARY	Loss of SARI expression initiates epithelial-to-mesenchymal transition (EMT), which is visualized by repression of E-cadherin and up-regulation of vimentin in lung adenocarcinoma cell lines and in clinical lung adenocarcinoma specimens.
Literature	GENERIF_SUMMARY	E-cadherin promotor methylation and mutation are inversely related to motility capacity of breast cancer cells.
Literature	GENERIF_SUMMARY	E-cadherin, EMMPRIN and MMP-8 median values were significantly increased in all chronic kidney disease children, as well as in all dialyzed patients versus controls
Literature	GENERIF_SUMMARY	Key roles of ZEB1 and E-cadherin in the oncogenesis and invasiveness of lung squamous cell carcinoma.
Literature	GENERIF_SUMMARY	Enhanced expression of E-cadherin is associated with the development of cervical neoplasia and cancer.
Literature	GENERIF_SUMMARY	Interaction between alpha6 integrin and E-cadherin drives liver metastasis of colorectal cancer cells through hepatic angiopoietin-like 6.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Mapping of cancer foci showed the highest density in the anterior proximal fundus (37%) and cardia/proximal fundus (27%).
Literature	GENERIF_SUMMARY	Overexpression of NEDD9 correlated with abnormal expression of E-cadherin, beta-catenin and N-cadherin.
Literature	GENERIF_SUMMARY	After exposure to hypoxia, HIF-1alpha protein was up-regulated, both mRNA and protein levels of E-cadherin were down-regulated and MMP-2 was up-regulated, while HIF-1alpha mRNA showed no significant change
Literature	GENERIF_SUMMARY	The low expression of CD82/KAI1 and E-cadherin was closely related to the invasion, progression and metastasis of non-small cell lung cancer.
Literature	GENERIF_SUMMARY	Elevated plasma soluble E-cadherin expression was involved in the pathogenesis of PID and is useful for the diagnosis of PID.
Literature	GENERIF_SUMMARY	intermolecular interactions are required for Langerhans cell differentiation in the epidermis
Literature	GENERIF_SUMMARY	Downregulation of E-cadherin was associated with gastric cancer.
Literature	GENERIF_SUMMARY	Results show a role for E- and P-cadherin co-expression in breast cancer progression and highlight the potential benefit of targeting P-cadherin in the aggressive tumours expressing high levels of this protein.
Literature	GENERIF_SUMMARY	the E-cadherin promoter is repressed in cells expressing E6, resulting in fewer E-cadherin transcripts.
Literature	GENERIF_SUMMARY	Oncogenic B-RAF(V600E) signaling induces the T-Box3 transcriptional repressor to repress E-cadherin and enhance melanoma cell invasion.
Literature	GENERIF_SUMMARY	CDH1 missense mutations were found in 4 out of 81 (5%) patients with non-syndromic OFC. This finding opens a new pathway to reveal the molecular basis of non-syndromic OFC. Cancer risk among carriers of these mutations needs to be defined.
Literature	GENERIF_SUMMARY	The high expression of Ezrin and the low expression of E-Cadherin were related to invasion and metastasis in papillary thyroid carcinoma.
Literature	GENERIF_SUMMARY	Report immunohistochemical CDH1 expression in hyperplastic and neoplastic lesions of the prostate gland.
Literature	GENERIF_SUMMARY	The data showed that both fascin and vimentin proteins were significantly overexpressed in cholangiocarcinoma, whereas E-cadherin expression was reduced in cholangiocarcinoma compared with normal tissues.
Literature	GENERIF_SUMMARY	E-cadherin and alpha-catenin down-regulation might be associated with neoplastic transformation in laryngeal tissues and might be regarded as a risk factor for clinical recurrence.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Expression of E-cadherin is not associated with malignant progression of brainstem gliomas, unlike beta-catenin and neuronal cadherin.
Literature	GENERIF_SUMMARY	Decreased levels of E-cadherin is associated with lung cancer.
Literature	GENERIF_SUMMARY	Data confirm that E-cadherin and COX-2 expressions are related to OSF. The epigenetic changes presented in patients with chronic inflammation might demonstrate an irreversible destruction in the tissues or organs similar to the effects of cancer.
Literature	GENERIF_SUMMARY	Loss of E-cadherin may be mediated through NF-kappaB-induced Snail upregulation.
Literature	GENERIF_SUMMARY	Snail regulates levels of E-cadherin and desmoglein 2 in oral squamous cell carcinoma cells both transcriptionally and post-translationally.
Literature	GENERIF_SUMMARY	C. albicans significantly decreased E-cadherin mRNA expression & protein production in gingival epithelial cells. IFN-gamma prevented this effect.
Literature	GENERIF_SUMMARY	Dkk-1 inhibited the survival and migration of human PTC cells by regulating Wnt/beta-catenin signaling and E-cadherin expression.
Literature	GENERIF_SUMMARY	E-cadherin gene mutations are associated with gastric cancer.
Literature	GENERIF_SUMMARY	The variants of differentiated cancers with a poorer prognosis (i.e. tall cell and follicular variants of papillary cancer and widely invasive follicular cancers) present reduced intensity of E-CAD expression.
Literature	GENERIF_SUMMARY	The capacity of TFF1 to enhance prostate cancer cell migration/invasion is mediated by transcriptional repression of E-CADHERIN.
Literature	GENERIF_SUMMARY	Negative E-cadherin expression seems to predict unfavourable clinical outcome in patients with advanced-stage serous ovarian cancer
Literature	GENERIF_SUMMARY	soft tissue leiomyosarcoma patients with the gain of E-cadherin and loss of Vimentin expression represent favorable trend of survival
Literature	GENERIF_SUMMARY	WT1 binds to the E-cadherin promoter and regulates gene expression in prostate cancer.
Literature	GENERIF_SUMMARY	LSD1 expression was higher, while E-cadherin expression was lower, in colon cancer specimens of high TNM stage and with distant metastasis. Positive LSD1 expression and negative E-cadherin expression were correlated with lower overall survival.
Literature	GENERIF_SUMMARY	Loss of E-cadherin is associated with serous ovarian cancer.
Literature	GENERIF_SUMMARY	sEcad contributes to skin carcinogenesis via association with the HER/IGF-1R-family of receptors

Main Category	Category	Terms
		and subsequent activation of the MAPK and PI3K/Akt/mTOR pathways.
Literature	GENERIF_SUMMARY	the majority of cervical cancers in the study were E-cadherin positive; no correlation was found between the level of E-cadherin expression and the oxygenation status
Literature	GENERIF_SUMMARY	these results indicate that in human breast cancer loss of E-cadherin is not causal for EMT and even not a necessity.
Literature	GENERIF_SUMMARY	Increased expression of CD10 in transitional and squamous cell carcinoma (TCC/SCC) tumor and stromal cells, and decreased expression of E-cadherin in TCC tumor cells, are correlated with tumor progression, invasion, and metastasis in human bladder cancer.
Literature	GENERIF_SUMMARY	study found CDH1 somatic alterations exist in all clinical settings and histotypes of gastric cancer and associate with different survival rates
Literature	GENERIF_SUMMARY	E-cadherin is not a tumour suppressor gene in invasive ductal carcinoma of the breast.
Literature	GENERIF_SUMMARY	The expression of E-cadherin and beta-catenin in pituitary adenoma was significantly downregulated and related to subtype, invasiveness, and postoperative recurrence.
Literature	GENERIF_SUMMARY	In oral squamous cell carcinoma the probability to have or develop metastases was very low when high E-cadherin expression was found in a preoperative sample or when a low podoplanin expression was found
Literature	GENERIF_SUMMARY	E-cadherin expression is downregulated and is negatively correlated to Long non-coding RNA H19 expression in bladder cancer
Literature	GENERIF_SUMMARY	Loss of E-cadherin and aberrant beta-catenin expression could be a prognostic factor in patients with small intestinal adeno-carcinomas.
Literature	GENERIF_SUMMARY	Human c-Myc represses E-cadherin at the post-transcriptional level through miR-9.
Literature	GENERIF_SUMMARY	Data indicate that TIMP-2 promotes an anti-tumoral transcriptional profile in lung cancer including upregulation of E-cadherin.
Literature	GENERIF_SUMMARY	The aberrant high expression of beta-catenin and decreased expression of E-cadherin is associated with poor prognosis in nasopharyngeal carcinoma.
Literature	GENERIF_SUMMARY	these results suggest that E-cadherin expression in cancer cells is controlled by a balance between ZEB2 and KLF4 expression levels.
Literature	GENERIF_SUMMARY	We found upregulated miR-200a expression to increase E-cadherin and suppress the Wnt/beta-catenin pathway

Main Category	Category	Terms
		by targeting ZEB1 and ZEB2 in GA, thus delaying tumor growth in vivo.
Literature	GENERIF_SUMMARY	the CD91/IKK/NF-kappaB signaling cascade is involved in secreted HSP90alpha-induced TCF12 expression, leading to E-cadherin down-regulation and enhanced CRC cell migration/invasion
Literature	GENERIF_SUMMARY	Loss of E-cadherin is associated with ovarian cancer.
Literature	GENERIF_SUMMARY	Overexpression of delta-catenin reduces the expression of E-cadherin and alters the balance between E-cadherin and p120ctn, which in turn affects the formation of intercellular adhesions and promotes invasion and metastasis in colorectal cancer.
Literature	GENERIF_SUMMARY	alterations of their expression suggest a role of Dsg3 and gamma-catenin (additionally to E-cadherin/beta-catenin) as biomarkers of malignant transformation risk of oral dysplasia and the biological behavior (aggressiveness) of oral cancer, respectively
Literature	GENERIF_SUMMARY	Report novel CDH1 germline mutations in Chinese sporadic gastric cancer patients.
Literature	GENERIF_SUMMARY	BRAF(V600E) increases migration and invasion of thyroid cancer cells via upregulation of Snail with a concomitant decrease of its target E-cadherin.
Literature	GENERIF_SUMMARY	E-cadherin 1054del83 transcript is a frequent event in gastric cancer.
Literature	GENERIF_SUMMARY	Down-regulated LIV-1 cells showed significant inhibition of proliferation in vitro and reduction of tumor growth in vivo. Furthermore, E-cadherin expression increased in LIV-1 siRNA expressing Hep-G2.
Literature	GENERIF_SUMMARY	Studies suggest CDH1 mutation for the diagnosis of Hereditary diffuse gastric cancer.
Literature	GENERIF_SUMMARY	Aberrant expression of E-cadherin and integrin beta-1 is closely related to malignant transformation and hydatidiform mole development.
Literature	GENERIF_SUMMARY	A high membrane/cytoplasm (M/C) ratio of E-cadherin expression.
Literature	GENERIF_SUMMARY	E-cadherin alterations are associated with hereditary diffuse gastric cancer.
Literature	GENERIF_SUMMARY	Cadherin alterations are associated with inherited human diseases.
Literature	GENERIF_SUMMARY	The overexpression of EBP50 could inhibit the growth of hepatocellular carcinoma cells and promote apoptosis by modulating beta-catenin, E-cadherin.
Literature	GENERIF_SUMMARY	Data show that TGF-beta1 regulates E-cadherin and vimentin expression through KLF8 pathway.
Literature	GENERIF_SUMMARY	O-GlcNAcylation could enhance ovarian cancer cell migration and reduce the expression of E-cadherin and

Main Category	Category	Terms
		the formation of the E-cadherin/catenin complex, thereby reducing intercellular adhesion.
Literature	GENERIF_SUMMARY	Data indicate that patients with basal subtype breast cancer and high E-Cadherin expression in their tumors had a poor clinical outcome.
Literature	GENERIF_SUMMARY	HER2 is implicated in mediating the effects of epidermal growth factor (EGF) on Snail, Slug and E-cadherin expression as well as inducing invasiveness of ovarian cancer cells.
Literature	GENERIF_SUMMARY	miRNA-200c inhibited Akt signaling through its effects on E-cadherin and PTEN, resulting in the inhibition of doxorubicin resistance in breast cancer cells.
Literature	GENERIF_SUMMARY	The CDH1 -160C/A polymorphism may constitute an inheritable risk factor for breast cancer in south Indian women.
Literature	GENERIF_SUMMARY	Data indicate that genomic profiling of relapsed CDH1-mutated invasive lobular carcinoma (ILC) featured a high incidence of ERBB2 alterations.
Literature	GENERIF_SUMMARY	Loss of E-cadherin expression is associated with poor response to epidermal growth factor receptor-tyrosine kinase inhibitor treatment in lung cancer.
Literature	GENERIF_SUMMARY	results suggest that reduced expression of E-cadherin is associated with promoter methylation of E-cadherin gene, in addition to providing evidence for the aberrant nuclear localization of E-cadherin in epithelial ovarian carcinoma
Literature	GENERIF_SUMMARY	Our results establish that the E-cadherin ectodomain-specific mAb DECMA-1 inhibits Ecad(+)/HER2(+) breast cancers by hindering tumor growth and inducing apoptosis via downregulation of key oncogenic pathways
Literature	GENERIF_SUMMARY	Cx43 and E-cadherin are reduced with non-small cell lung cancer progression, and might be important biomarkers for judging the metastasis and prognosis.
Literature	GENERIF_SUMMARY	The expression of Twist is upregulated, whereas that of E-cadherin is downregulated in ovarian cancer.
Literature	GENERIF_SUMMARY	We showed that cadherin and nectin in the junctions of A431 cells and human keratinocytes are located in separate clusters.
Literature	GENERIF_SUMMARY	High E-cadherin expression is associated with lymphatic vessel invasion in abdominal metastases of breast cancer.
Literature	GENERIF_SUMMARY	The present study examined the expression of AP-2alpha, TIMP-2, MMP-2, MMP-9, and E-cadherin in severe preeclamptic placentas and normal placentas.
Literature	GENERIF_SUMMARY	GnT-III determines E-cadherin-mediated tumor suppression, and GnT-V regulates E-cadherin-mediated tumor invasion.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Data indicate that pancreatic (pro)enzymes enhanced expression of beta-catenin and E-cadherin and decreased expression of several epithelial-mesenchymal transition (EMT)-associated genes, such as Vimentin, Snail and Slug.
Literature	GENERIF_SUMMARY	CDH1 germline mutations and the hereditary diffuse gastric and lobular breast cancer syndrome
Literature	GENERIF_SUMMARY	E-cadherin expressing mononuclear histiocytes are likely precursors for multinucleated giant cells in cutaneous granulomas and may play a critical role in disease pathogenesis.
Literature	GENERIF_SUMMARY	activation of AKT plays a role in contactin-1-mediated downregulation of E-cadherin.
Literature	GENERIF_SUMMARY	ELE increased both the protein and mRNA levels of E-cadherin.
Literature	GENERIF_SUMMARY	Suggest that ZEB1 overexpression, associated with E-cadherin and miR-200s downregulation, and the expression of mesenchymal markers might enhance the metastatic potential of undifferentiated endometrial carcinomas.
Literature	GENERIF_SUMMARY	These findings support the potential value of E-cadherin for a supplementary differentiation of molecular subtypes of breast carcinoma, based on the biological significance of its capacity of expression.
Literature	GENERIF_SUMMARY	Combined detection of VM, HIF-lalpha and E-cad plays an important role in predicting the invasion, metastasis and prognosis of patients with ESCC.
Literature	GENERIF_SUMMARY	this study shows that sEcad is a novel candidate protein for drug targeting since it is increased in human and mouse HER2-positive (HER2+) breast tumors, MMTV-PyMT bodily fluids and human cell culture systems.
Literature	GENERIF_SUMMARY	Data suggest that, in tubal pregnancy, fallopian tube mucosa exhibits reduced E-cadherin expression, enhanced beta-catenin expression, and glycogen accumulation; tubal arteries exhibit neovascularization and hyperplasia in tubal pregnancy.
Literature	GENERIF_SUMMARY	High expression of GLI1 dampened expression of E-cadherin and enhanced the expression of Vimentin.
Literature	GENERIF_SUMMARY	The soluble serum E-cadherin levels in breast cancer patients undergoing preoperative systemic chemotherapy have high predictive value.
Literature	GENERIF_SUMMARY	The increased therapeutic efficacy of the PEG-IFNalpha2b plus RBV treatment could be secondary to the inhibition of claudin-1 and E-cadherin cell membrane expression.
Literature	GENERIF_SUMMARY	The abnormal expression of E-cadherin and beta-catenin proteins might be useful in distinguishing vulvar intraepithelial neoplasia from non-neoplastic vulvar

Main Category	Category	Terms
		squamous epithelium lesions in problematic cases.
Literature	GENERIF_SUMMARY	The expression of E-cadherin and beta-catenin, combined with cytological and architectural analysis, may highlight different immunophenotypes and improve classification of breast carcinomas in situ with mixed pattern.
Literature	GENERIF_SUMMARY	Loss of E-cadherin and beta-catenin with cytoplasmic ALCAM accumulation may play pivotal role in oral cancer development and progression.
Literature	GENERIF_SUMMARY	celastrol treatment restored E-cadherin promoter activity in TGF-beta1-treated cells
Literature	GENERIF_SUMMARY	Western blotting and real-time reverse transcriptase-polymerase chain reaction demonstrated that these cells had downregulated E-cadherin protein and messenger RNA levels, respectively
Literature	GENERIF_SUMMARY	methylation of E-cadherin promoter is associated with risk of lung cancer
Literature	GENERIF_SUMMARY	Data show that miR-148a downregulated vimentin expression and upregulated E-cadherin expression, suggesting that miR-148a inhibited epithelial-mesenchymal transition (EMT), and the SMAD2 gene was identified as the direct and functional target of miR-148a.
Literature	GENERIF_SUMMARY	There was a significantly positive correlation between miR-200a and E-cadherin in epithelial ovarian cancer
Literature	GENERIF_SUMMARY	We suggest that the down-regulation of E-cadherin in multicentric/multifocal breast cancer is causally connected with the worse prognosis of this tumor type.
Literature	GENERIF_SUMMARY	GLI2 represses E-cadherin gene (CDH1) expression in melanoma cells via distinct mechanisms, enhancing transcription of the EMT-activator ZEB1.
Literature	GENERIF_SUMMARY	CDH-1 expression exhibited a negative correlation with the invasiveness of colon cancer cells.
Literature	GENERIF_SUMMARY	Positive correlations between the expression of E-cadherin and NDRG2 were shown in several colon cancer cell lines as well as in colon cancer tissues.
Literature	GENERIF_SUMMARY	miR-23a inhibits E-cadherin expression in gastrointestinal cancer.
Literature	GENERIF_SUMMARY	expression and subcellular localization of p120-catenin and beta-catenin in oral carcinomas
Literature	GENERIF_SUMMARY	Suppression of CDH1 transcription is associated with the development and progression of non-small-cell lung cancer.
Literature	GENERIF_SUMMARY	FadA binds to E-cadherin, activates beta-catenin signaling, and differentially regulates the inflammatory and oncogenic responses.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	SALL4 functions to suppress CDH1 expression and to maintain cell dispersion in basal-like breast cancer.
Literature	GENERIF_SUMMARY	E-cadherin expression in squamous cells is reduced by HD-5
Literature	GENERIF_SUMMARY	Expression of E-cadherin is significantly associated with tumor location and mortality in patients with oral squamous cell carcinoma.
Literature	GENERIF_SUMMARY	Data indicate that E-cadherin was elevated , and the expression levels of N-cadherin, Vimentin and Twist1 were all decreased in oleanolic acid (OA)-treated cells.
Literature	GENERIF_SUMMARY	Our study demonstrates significantly elevated serum levels of sE-cadherin in women with systemic lupus erythematosus compared with healthy women.
Literature	GENERIF_SUMMARY	TGF-beta2 activated Smad-2/3, serine-threonine kinase (AKT), and extracellular signal-regulated kinase (ERK) signaling, and enhanced expression of beta-catenin as well as N- and OB-cadherin.
Literature	GENERIF_SUMMARY	Low level of E-cadherin expression is associated with breast cancer.
Literature	GENERIF_SUMMARY	Hypermethylation of E-cadherin is associated with gastric cancer.
Literature	GENERIF_SUMMARY	current study showed that the majority of DGC cases demonstrated concurrent CDH1 promoter methylation in tumor tissues and in non-neoplastic mucosa
Literature	GENERIF_SUMMARY	BCL2 is not involved in lobular breast carcinogenesis and is unlikely to represent an important determinant of tumor development driven by CDH1 inactivation.
Literature	GENERIF_SUMMARY	the CDH1 polymorphism is involved in the etiology of nephrolithiasis
Literature	GENERIF_SUMMARY	These results indicated a possible effect of SIRT1 and p300/CBP involved in regulating the expression of E-cadherin and MLH1, thus participating in the tumor progression of gastroesophageal junction cancer
Literature	GENERIF_SUMMARY	CDH1 hypermethylation is associated with triple-negative breast cancers.
Literature	GENERIF_SUMMARY	A novel role for ILK in glioma invasion and metastasis processes via downregulation of CDH1.
Literature	GENERIF_SUMMARY	The knockdown of flotillins dramatically affected N- and E-cadherin recruitment at CCJs in mesenchymal and epithelial cell types and perturbed CCJ integrity and functionality
Literature	GENERIF_SUMMARY	Susceptible CDH1 genotypes might modulate the development of asthma, especially for children exposed to environmental tobacco smoke.
Literature	GENERIF_SUMMARY	PP6c associates with E-cadherin in adherens junctions and is required to oppose casein kinase-1 to maintain

Main Category	Category	Terms
		cell surface localization of E-cadherin.
Literature	GENERIF_SUMMARY	No associations were noted between promoter hypermethylation of the CDH1 gene and biological features of the endometrial cancer cases.
Literature	GENERIF_SUMMARY	Epithelial-to-mesenchymal transition status is determined by calculating the ratio of vimentin to E-cadherin, whereby patients are categorized into 3 groups
Literature	GENERIF_SUMMARY	Gli-1 expression increased markedly, and was closely associated with increased Snail expression and decreased E-cadherin
Literature	GENERIF_SUMMARY	Poor prognosis for serous ovarian cancer patients with CD44(+)/E-cadherin(-) and CD44(+)/budding(+) tumor cells phenotype, has been revealed
Literature	GENERIF_SUMMARY	COX-2 likely functions upstream of NF-kappaB and regulates the expression of E-cadherin via NF-kappaB/ Snail signaling pathway in gastric cancer cells.
Literature	GENERIF_SUMMARY	filamin A generates a scaffold for organizing a signaling complex that promotes E-cadherin-mediated cell-cell adhesion and keratinocyte differentiation
Literature	GENERIF_SUMMARY	Mammosphere formation in breast carcinoma cell lines depends upon expression of E-cadherin.
Literature	GENERIF_SUMMARY	Analysis of gene expression datasets confirmed the association between low PIP4K2B and low CDH1expression.
Literature	GENERIF_SUMMARY	High E-cadherin expression was associated with increase in degree of differentiation, and decrease in lymph node involvement in laryngeal squamous cell carcinomas.
Literature	GENERIF_SUMMARY	Our findings suggest that abnormal expression of beta-catenin and E-cadherin occurred frequently in gastric carcinoma and correlated with worse clinical behaviour.
Literature	GENERIF_SUMMARY	Data suggest that E-cadherin (CDH1) screening, in addition to the identification of other risk factors, could be useful for the early detection of gastric cancer (GC) in subjects at risk.
Literature	GENERIF_SUMMARY	Loss of CDH1 induces EGFR expression via phospho-YBX1, which is activated through the AKT signaling pathway.
Literature	GENERIF_SUMMARY	Reduced E-Cadherin expression is associated with nodal metastasis in T1 esophageal squamous cell carcinoma
Literature	GENERIF_SUMMARY	Low E-Cadherin expression is associated with endometrial cancer.
Literature	GENERIF_SUMMARY	cancer-related mutations in beta-catenin can influence tumor progression by weakening the adhesion of tumor cells to one another through reduced individual Ecad/

Main Category	Category	Terms
		Ecad bond strength
Literature	GENERIF_SUMMARY	decorin-mediated inhibition of cholangiocarcinoma cell growth, migration,and invasion and promotion of cell apoptosis might be through regulation of the expression of E-cadherin in vitro.
Literature	GENERIF_SUMMARY	E-cadherin expression was the same in the cholesteatoma matrix in all samples. There were no differences in expression according to the clinical and histological characteristics of the cholesteatomas.
Literature	GENERIF_SUMMARY	Low E-Cadherin expression is a negative prognostic factor of oral squamous cell carcinoma and is likely due to the hypermethylation of CDH1 promoter.
Literature	GENERIF_SUMMARY	E-cadherin protein expression in the cecum was lower in IBS-A compared with controls and associated with longstanding symptoms. E-cadherin was further associated with abdominal pain severity in the IBS group overall, but unrelated to IBS subtype.
Literature	GENERIF_SUMMARY	Inhibition of SALL1 correlates with reduced levels of CDH1, an important contributor to epithelial-to-mesenchymal transition.
Literature	GENERIF_SUMMARY	Low expression of E-cadherin is associated with gastric carcinomas.
Literature	GENERIF_SUMMARY	Positive ZEB-1 expression and loss of E-cadherin expression are correlated with poor prognosis in hepatocellular carcinoma patients.
Literature	GENERIF_SUMMARY	E-cadherin expression was down-regulated in gastric cancer.
Literature	GENERIF_SUMMARY	A more complete study of the prognostic and predictive role of E-cadherin expression in patients with Stage I-II NSCLC will help identify a prognostically unfavorable group of patients who may be given additional treatment in the postoperative period.
Literature	GENERIF_SUMMARY	show that BMP acts prior to and independently of Cdh1 to prime pluripotent cells for mesoderm differentiation, thus helping to reinforce the block to neural differentiation
Literature	GENERIF_SUMMARY	We have shown that germline CDH1 mutations are associated with early onset of bilateral lobular carcinoma in situ with or without Invasive lobular breast cancer in women without a family history of gastric cancer.
Literature	GENERIF_SUMMARY	Deletion of beta-catenin by siRNA abrogated celecoxib-induced inhibition of MMP9 up-regulation and E-cadherin down-regulation by treatment of PGE2 in A549 cell.
Literature	GENERIF_SUMMARY	A potential important role of alpha4 in control of cell migration and/or invasion via the regulation of E-cadherin expression in bladder urothelial carcinoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Two novel CDH1 mutations were found. The c.602_603delCT mutation results in a frame-shift and a premature stop at codon 207, and is classified as pathogenic. The functional consequence of the c.1565 + 3insTT mutation was unknown and not as obvious.
Literature	GENERIF_SUMMARY	E-cadherin gene promoter hypermethylation may contribute to increased risk of bladder cancer among Asian populations.
Literature	GENERIF_SUMMARY	Studied the expression of E-cadherin, N-cadherin, TGF-beta1 and Twist protein and investigate its significance in the occurrence and development of prostate cancer.
Literature	GENERIF_SUMMARY	MTA1 promotes tumor invasion by downregulation of E-cadherin
Literature	GENERIF_SUMMARY	AKAP proteins, most likely AKAP9, maintain the bronchial epithelial barrier by regulating the E-cadherin expression at the cell membrane.
Literature	GENERIF_SUMMARY	In high grade malignant tumors (G1), CDH1 and ITGB1 gene expression was the highest, in G2 and G3 tumors the expression of both genes was gradually lowering
Literature	GENERIF_SUMMARY	GSTP1 and E-cad did not show any relevant methylation pattern in various endometrial lesions.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with laryngeal squamous cell carcinoma.
Literature	GENERIF_SUMMARY	Over-expression of miR-888 reduced the mRNA levels of E-cadherin in MCF-7 side population sphere cells.
Literature	GENERIF_SUMMARY	In conclusion NSCLC cell lines, positive for E-Cadherin,EpCAM and avb6 expression, activate normal fibroblasts through avbeta6/TGFbeta signalling in vitro, and influence both gene expression and response to therapeutic agents.
Literature	GENERIF_SUMMARY	Our meta-analysis indicates that promoter polymorphism and methylation of CDH1 gene may be involved in the development and progression of bladder cancer
Literature	GENERIF_SUMMARY	WNT/beta-catenin pathway and E-cadherin are important factors in advanced epithelial ovarian cancer.
Literature	GENERIF_SUMMARY	In increased E-cadherin expression.
Literature	GENERIF_SUMMARY	Loss of E-cadherin is associated with breast carcinoma.
Literature	GENERIF_SUMMARY	results suggest that E-cadherin loss promotes SNAI1 expression that controls the aggressiveness of prostate cancer cells
Literature	GENERIF_SUMMARY	highly sensitive rabbit E-cadherin antibody is the preferred antibody for evaluating ductal carcinomas and for distinguishing ductal versus lobular lesions, and the dual stain was superior to the single E-cadherin stain

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Loss of E-cadherin is mediated through inhibition of mTORC1 in lung cancer cells.
Literature	GENERIF_SUMMARY	the summarized analyses of these case-control studies demonstrated that the -160A of the epithelial cadherin gene exhibited no significant association with susceptibility for diffuse gastric cancer
Literature	GENERIF_SUMMARY	Extensive remodeling of the actin cytoskeleton and focal adhesions accompanies cell fusion and differentiation and appears related to alterations in E-cadherin in trophoblastic cells.
Literature	GENERIF_SUMMARY	Here, we report that infection with HCV derived from pJFH-1 replicon system that mimics natural infection elevates protein levels of DNA methyltransferase 1 and 3b to enhance DNMT activity in human hepatocytes.
Literature	GENERIF_SUMMARY	rs13689 of CDH1 gene is correlated with the syndrome differentiation of gastric cancer.
Literature	GENERIF_SUMMARY	Collectively these data unveil that FBXL19 functions as an antagonist of Rac3 by regulating its stability and regulates the TGFbeta1-induced E-cadherin down-regulation.
Literature	GENERIF_SUMMARY	CDH1 - 160C --> A promoter polymorphism and haplotypes are associated with diffuse gastric cancer.
Literature	GENERIF_SUMMARY	The expression of PRL-3, but not of E-cadherin, was associated with shorter survival of patients. PRL-3 and E-cadherin exhibit interactions in gastric cancer and are involved in the formation of lymph node metastases.
Literature	GENERIF_SUMMARY	In pancreatic adenocarcinoma cells, PAR3 knockdown enhances cell adhesion. We propose this is due to increased expression of E-cadherin, leading to a greater adhesion of free-floating cells to cells bound to the surface via integrins, particularly ITGalphav.
Literature	GENERIF_SUMMARY	MMP-2, MMP-9, and E-cadherin are expressed in the endometrium of infertile patients during the receptive phase of the natural menstrual cycle. However, there is no correlation between the expression of these molecules and the clinical IVF outcomes.
Literature	GENERIF_SUMMARY	E-cadherin and MMP-9 expression at histologically negative surgical margins shows the significance of these markers for prognostic values in oral squamous cell carcinoma patients.
Literature	GENERIF_SUMMARY	CD44 v6 and E-cad expressions had a significant correlation in the NSCLC tissue with lymphatic metastasis.
Literature	GENERIF_SUMMARY	The expression of E-cadherin in OSCC was significantly lower than the control tissues but galectin-9 expression was conversely higher.
Literature	GENERIF_SUMMARY	Ntn-1 induces MMP-12-dependent E-cad degradation via the distinct activation of PKCalpha and FAK/Fyn, which is necessary to govern the activation of ERK,

Main Category	Category	Terms
		JNK, and NF-KB in promoting motility of umbilical cord blood-derived mesenchymal stem cells.
Literature	GENERIF_SUMMARY	Lack of E-cadherin has been recorded at an early phase in carcinomas.
Literature	GENERIF_SUMMARY	The relatively slow progression of disseminated peritoneal adenomucinosis may be attributable to the smaller number of single isolated tumor cells that lack E-cadherin expression
Literature	GENERIF_SUMMARY	Low E-cadherin protein expression correlate with an aggressive, malignant phenotype in hepatocellular carcinoma.
Literature	GENERIF_SUMMARY	The disruption of cell adhesion junction by suppression of E-cadherin mediates gallotannin enhanced apoptosis in Hep G2 liver cancer cells.
Literature	GENERIF_SUMMARY	Results indicate that loss of E-cadherin expression is a consequence rather than a cause of c-erbB2-induced epithelial-mesenchymal transition.
Literature	GENERIF_SUMMARY	We found a strong association between the cancer predisposing gene CDH1 and the risk of non-syndromic cleft lip with or without cleft palate in the Polish population
Literature	GENERIF_SUMMARY	Loss of CDH1 is associated with promotes hepatocellular carcinogenesis.
Literature	GENERIF_SUMMARY	placental expression significantly increased in preeclampsia compared to normotensive pregnancies
Literature	GENERIF_SUMMARY	Data suggest that E-cadherin can act as a central modulator of tumor cell phenotype and is a potential metastasis marker in cholangiocarcinoma (CCA).
Literature	GENERIF_SUMMARY	Results showed that three polymorphisms of CDH1 were significantly associated with cancer risk, -160 C>A, -347 G>GA and 3'-UTR +54 C>T. [Meta-Analysis]
Literature	GENERIF_SUMMARY	The sialylation pattern of alpha2beta1 integrin and E-cadherin.
Literature	GENERIF_SUMMARY	E-cadherin augmented DR4/DR5 clustering and assembly of the death-inducing signaling complex, increasing caspase-8 activation in high molecular weight cell fractions.
Literature	GENERIF_SUMMARY	Downregulation of CDH-1 resulting from the epithelial-to-mesenchymal transition may be closely involved in lymph node metastasis in tongue squamous cell carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin and b-catenin immunostaining showed no prognostic value for basaloid and conventional squamous cell carcinomas
Literature	GENERIF_SUMMARY	Data suggest that macrophages contributed to the decreased expression of E-cadherin by NF-kappaB/Slug protein pathway in hepatocellular carcinomas.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Studies indicate that hypermethylation frequencies in lung cancer (LC) tissues were significantly higher than those in normal control tissues
Literature	GENERIF_SUMMARY	Shh signaling may promote invasion and metastasis of oral squamous cell carcinoma by activating MMP-9 and E-cadherin expression.
Literature	GENERIF_SUMMARY	Declining E-cadherin expression is a possible immunohistochemical predictor of patient prognosis in stage III colorectal cancer.
Literature	GENERIF_SUMMARY	In the present study, with an increasing concentration of metformin, the expression of MMP-9 was downregulated whereas that of E-cadherin was significantly upregulated.
Literature	GENERIF_SUMMARY	E-cadherin and P120 catenin cocktail immunostain can be used to differentiate ductal carcinoma in situ from lobular carcinoma in situ.
Literature	GENERIF_SUMMARY	The methylation of E-cadherin is one of the main reasons for the reduction of E-cadherin expression in ovarian cancer.
Literature	GENERIF_SUMMARY	Down-regulation of NDRG1 in gastric cancer metastatic progression was correlated to E-cadherin and MMP-9.
Literature	GENERIF_SUMMARY	Low E-cadherin expression is associated with gastrointestinal stromal tumor metastasis.
Literature	GENERIF_SUMMARY	CD133 associates with beta-catenin in early placodes, and its continued expression correlates with loss of beta-catenin and E-cadherin from the cell membrane at a time when E-cadherin transcriptional repressors Snail and Slug are not implicated.
Literature	GENERIF_SUMMARY	The loss of membranous E-cadherin with increase in cytoplasmic accumulation in differentiative layers of epithelium through the progression of dysplasia was noted along with up-regulation in VEGF expressions
Literature	GENERIF_SUMMARY	the present meta-analysis provides further evidence that promoter methylation and -160C/A polymorphism of E-cadherin gene may confer a risk to prostate cancer
Literature	GENERIF_SUMMARY	our study demonstrated that Nur77 could promote the invasion and metastasis of colorectal cancer cells through regulation of MMP-9/E-cadherin signaling
Literature	GENERIF_SUMMARY	Results showed that E-cadherin loss disrupts the organization of the cell's actin and microtubule cytoskeletons and modifies its adherence and migration characteristics but is insufficient to induce an epithelial to mesenchymal transition.
Literature	GENERIF_SUMMARY	siRNA knockdown of SMAR1 expression in these breast cancer cells results in a coordinative action of Slug-mediated repression of E-cadherin transcription.
Literature	GENERIF_SUMMARY	Reduced E-cadherin expression indicates a poor prognosis for patients with HCC (hepatocellular

Main Category	Category	Terms
		carcinoma), and it may have predictive potential for prognosis of HCC patients.[meta-analysis]
Literature	GENERIF_SUMMARY	Data demonstrated that Rab11 regulated E-cadherin expression and promoted colon cancer cell transformation.
Literature	GENERIF_SUMMARY	Loss of expression and aberrant methylation of the CDH1 gene is associated with breast cancer.
Literature	GENERIF_SUMMARY	CCND1 870G/A and CDH1 -160C/A SNPs may contribute to the etiology of colorectal cancer in South Indian population and carriers of the 870A/-160A haplotype have a higher risk for colorectal cancer susceptibility.
Literature	GENERIF_SUMMARY	The rs8049282 SNP of the E-cadherin gene may be a potential molecular marker for the development of primary infertility in northern Chinese women with ovarian endometriosis.
Literature	GENERIF_SUMMARY	Increasing evidence suggested that microRNA mediated E-cadherin expression in the head and neck cancers by directly/indirectly targeting the transcription suppressors of E-cadherin, ZEB1 and ZEB2.
Literature	GENERIF_SUMMARY	E-cadherin may directly regulate gene transcription. [Review]
Literature	GENERIF_SUMMARY	Study demonstrated that reactive oxygen species promote the migration and metastatic growth of ovarian cancer cells via upregulation of HIF-1a and LOX and E-cadherin repression.
Literature	GENERIF_SUMMARY	PLCdelta1 has tumor-suppressive functions in colorectal cancer through E-cadherin induction.
Literature	GENERIF_SUMMARY	Estrogen promotes the invasion of ovarian cancer cells via activation of the PI3K/AKT pathway, downregulation of Ecadherin and upregulation of alphaactinin4.
Literature	GENERIF_SUMMARY	we observed the coordinated loss of E-cadherin (CDH1) and transforming growth factor beta receptor II (TGFBR2) in esophageal squamous tumors.
Literature	GENERIF_SUMMARY	Zonula occludens-1, occludin and E-cadherin expression and organization in salivary glands
Literature	GENERIF_SUMMARY	Btbd7 contributes to reduced expression of E-cadherin and may be a promising cancer marker in non-small cell lung cancer.
Literature	GENERIF_SUMMARY	Data indicate that E-cadherin ubiquitination consistently increases after depletion of kinesin-like protein KIFC3 or ubiquitin-specific protease USP47.
Literature	GENERIF_SUMMARY	Abnormal CDH1 methylation occurs in high frequencies in ductal breast tumors.
Literature	GENERIF_SUMMARY	results demonstrate that adhesion of beta-cells to E- and N-cadherins is regulated by insulin secretagogues and that E- and N-cadherin engagement promotes

Main Category	Category	Terms
		stimulated insulin secretion.
Literature	GENERIF_SUMMARY	In this cohort of NPC patients, higher levels of E-cadherin and higher levels of vimentin were associated with worse outcomes
Literature	GENERIF_SUMMARY	E-cadherin and CD10 in endometrial lesions is not correlated, but reduced expression of both molecules could be critical for progression of endometrial carcinoma
Literature	GENERIF_SUMMARY	AKR1B10 is a unique enzyme involved in pancreatic carcinogenesis via modulation of the Kras-E-cadherin pathway.
Literature	GENERIF_SUMMARY	deltaEF1 associates with DNMT1 and maintains DNA methylation of the E-cadherin promoter in breast cancer cells.
Literature	GENERIF_SUMMARY	In this meta-analysis, E-cadherin expression predicts the overall survival and tumor metastasis in Asian patients with gastric cancer.
Literature	GENERIF_SUMMARY	This study focused on analyzing the immunoexpression of CK19, vimentin and E-cadherin in a number of 43 differentiated thyroid carcinomas.
Literature	GENERIF_SUMMARY	In this critical review, we describe the clinical management of CDH1 germline mutant carriers providing specific recommendations for genetic counselling, clinical criteria, surveillance and/ or prophylactic surgery.
Literature	GENERIF_SUMMARY	PDGFD, CDH1 and SLIT2 are upregulated in low grade meningiomas and schwannomas compared with healthy tissue.
Literature	GENERIF_SUMMARY	E-cadherin binding is essential for the barrier disruption with the Clostridium botulinum type B hemagglutinin complex
Literature	GENERIF_SUMMARY	Observed strong association of promoter hypermethylation of CDH1 with Gastric cancer(OR = 12.23) suggesting that epigenetic regulation of CDH1 could play a critical role in the etiology of Gastric cancer.
Literature	GENERIF_SUMMARY	NKX3.1 expression mediate beta-catenin and E-cadherin association and cell migration in prostate cells.
Literature	GENERIF_SUMMARY	low expression in hepatocellular carcinoma
Literature	GENERIF_SUMMARY	High E-cadherin expression is associated with oral squamous cell carcinoma.
Literature	GENERIF_SUMMARY	MicroRNA-9 has a role in promoting tumor metastasis via repressing E-cadherin in esophageal squamous cell carcinoma
Literature	GENERIF_SUMMARY	The results of this study support the role of E-cadherin in the pathogenesis of asthma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	high osteopontin but low E-cadherin expression can be considered as a negative, independent prognostic factor in patients with locally advanced cervical squamous cell carcinoma
Literature	GENERIF_SUMMARY	CD148 tyrosine phosphatase promotes e-cadherin cell adhesion.
Literature	GENERIF_SUMMARY	Our results suggest employing this methodology as a complementary approach to evaluate the pathogenicity of E-cadherin missense variants
Literature	GENERIF_SUMMARY	data suggests that E-cadherin regulates assembly of nectin junctions through alpha-catenin-induced remodeling of the actin cytoskeleton around the cadherin clusters.
Literature	GENERIF_SUMMARY	Octamer-binding protein 4 affects the cell biology and phenotypic transition of lung cancer cells involving beta-catenin/E-cadherin complex degradation
Literature	GENERIF_SUMMARY	The T-box transcription factor Brachyury promotes renal interstitial fibrosis by repressing E-cadherin expression
Literature	GENERIF_SUMMARY	Loss of E-cadherin expression is associated with recurrence in gastric cancer.
Literature	GENERIF_SUMMARY	Suggest that loss of PEAK1/E-cadherin may play an important role in carcinogenesis and development of gastric cancer through activating epithelial to mesenchymal transition.
Literature	GENERIF_SUMMARY	TLE1 has a role in regulating epithelial-to-mesenchymal transition in A549 cells through its repressive effect on E-cadherin
Literature	GENERIF_SUMMARY	Results suggest that, besides their role in mechanical adhesion, loss of E-cadherin expression may contribute to cancer progression by modifying pathways that regulate fundamental processes as oxidative stress, immune evasion and cell metabolism.
Literature	GENERIF_SUMMARY	The E-cadherin intracellular interactome is predominantly independent of cell-cell adhesion.
Literature	GENERIF_SUMMARY	placental level significantly reduced in placenta percreta
Literature	GENERIF_SUMMARY	epithelial cell-derived periostin exerts tumor-suppressor activities in gastric cancer through stabilizing p53 and E-cadherin proteins via the Rb/E2F1/p14(ARF)/Mdm2 signaling pathway.
Literature	GENERIF_SUMMARY	We examined allelic imbalance (AI) on loci 17p13.1 (TP53), 1p36.1 (RUNX) and 16p22 (CDHI) and microsatellite instability (MI) with BAT26 in 78 patients with gastric cancer
Literature	GENERIF_SUMMARY	ERbeta1 inhibits the migration and invasion of breast cancer cells and upregulated E-cadherin expression in a Id1-dependent manner.
Literature	GENERIF_SUMMARY	Overexpression of SPRY2 reduced EGF-induced cell

Main Category	Category	Terms
		invasion by attenuating EGF-induced E-cadherin down-regulation
Literature	GENERIF_SUMMARY	Exogenous expression of cell surface E-cadherin converts statin- sensitive cells to a partially resistant state implying that statin resistance is in part dependent on the tumor cells attaining an epithelial phenotype
Literature	GENERIF_SUMMARY	This review focuses on soluble E-cadherin in sera of patients affected by three solid cancers (breast, gastric, and colorectal cancers) and how its levels correlate or not with some cancer parameters
Literature	GENERIF_SUMMARY	self-renewal versus differentiation of human ESCs (hESCs) in response to Wnt signaling is predominantly determined by a two-layer regulatory circuit involving beta-catenin, E-cadherin, PI3K/Akt, and Slug in a time-dependent manner.
Literature	GENERIF_SUMMARY	the decrease in E-cadherin and/or increase in N-cadherin may be required for BMP4-induced migration and EMT.
Literature	GENERIF_SUMMARY	Serum soluble E-cad level is an independent prognostic factor in Asian breast cancer patients.
Literature	GENERIF_SUMMARY	Results show that telomere shortening is closely associated with severity of Helicobacter pylori-induced gastritis and CDH1 methylation status.
Literature	GENERIF_SUMMARY	Cytoskeletal regulation plays in controlling the levels of intra-junctional tension at E-cadherin junctions.
Literature	GENERIF_SUMMARY	this meta-analysis failed to confirm the association between the CDH1 C-160A polymorphism and risk of gastric cancer.
Literature	GENERIF_SUMMARY	Suggest MelanA-negative spindle-cell associated melanoma constitutes a distinct inflammatory phenotype correlated with dense infiltration of CD163 macrophages and loss of E-cadherin.
Literature	GENERIF_SUMMARY	The aim of this study was to investigate immunohistochemically the E-cadherin/N-cadherin ""switch"" and vimentin expression as markers of epithelial-mesenchymal transition process in tongue oral squamous cell carcinomas."
Literature	GENERIF_SUMMARY	The immunohistochemical expression of E-cadherin and alpha-smooth muscle actin (alpha-SMA) in 15 cases of pleomorphic adenoma of salivary glands, were investigated.
Literature	GENERIF_SUMMARY	Low CDH1 contributes to endometrial cancer progression.
Literature	GENERIF_SUMMARY	alleles of marker rs10431924 of the CDH1 gene are associated with vitiligo, especially in the presence of autoimmune comorbidities
Literature	GENERIF_SUMMARY	GPC3 and E-cadherin expressions are not independent

Main Category	Category	Terms
		prognostic factors in CRC.
Literature	GENERIF_SUMMARY	The positive expression of E-cad and anti-apoptosis Bcl-2 protein are negatively related in breast cancer.
Literature	GENERIF_SUMMARY	RhoB depletion reduces cell-cell adhesion and downregulates E-cadherin levels as well as increasing internalized E-cadherin in DU145 prostate cancer cells.
Literature	GENERIF_SUMMARY	CDH1 germline variants are present in early-onset diffuse gastric cancer patients in Chinese population, but they are mainly missense variants with unknown function which are likely associated with lymph node metastasis and survival.
Literature	GENERIF_SUMMARY	that MAD2B may play an important role in high glucose-mediated podocyte injury of diabetic nephropathy via modulation of Cdh1, cyclin B1, and Skp2 expression
Literature	GENERIF_SUMMARY	E-Cadherin expression in primary sites positively correlates with E-cadherin from LNM
Literature	GENERIF_SUMMARY	These findings indicated that increased SUZ12 promote gastric cancer cell proliferation and metastasis partly via repressing KLF2 and E-cadherin transcription.
Literature	GENERIF_SUMMARY	miR-218 and its downstream target Gli2, as well as E-cadherin, participate in the anti-invasion process
Literature	GENERIF_SUMMARY	spatial polarization of integrin and E-cadherin adhesions in a human pluripotent stem cell colony compete to recruit Rho-ROCK activated myosin II.
Literature	GENERIF_SUMMARY	C6orf106 promotes invasion in NSCLC cells. Finally, C6orf106 upregulates vimentin, and downregulates E-cadherin and P120ctn.
Literature	GENERIF_SUMMARY	Hypermethylation of Integrin alpha4 is associated with Prostate Cancer.
Literature	GENERIF_SUMMARY	The results from the current study indicate that the hypermethylation frequency of E-cadherin in NSCLC is strongly associated withnon-small cell lung carcinoma incidence and it may be an early event in carcinogenesis
Literature	GENERIF_SUMMARY	EHD2 can inhibit the metastasis of human breast cancer by regulating the epithelial-to-mesenchymal transition markers E-cadherin and N-cadherin.
Literature	GENERIF_SUMMARY	The germline missense mutant CDH1(Q16H) had a pro-invasive cell behavior.
Literature	GENERIF_SUMMARY	It was shown that both E- and N-cadherin mRNA and protein are expressed in the human renal proximal tubule.
Literature	GENERIF_SUMMARY	ABCG2 localizes to the nucleus and modulates CDH1 expression in lung cancer cells
Literature	GENERIF_SUMMARY	Negative E-cadherin expression in bone and soft tissue sarcomas was correlated with lower 5-year overall

Main Category	Category	Terms
		survival (OR = 3.831; 95% CI
Literature	GENERIF_SUMMARY	E-cadherin and CD44 may have a role in the epithelial-mesenchymal transition of bladder urothelial carcinomas
Literature	GENERIF_SUMMARY	Abnormal E-cadherin expression was correlated with more advanced gastric cancer stage, and as a consequence, with poor outcome.
Literature	GENERIF_SUMMARY	This study shows that TGF-alpha uses common and divergent molecular mediators to regulate E-cadherin expression and cell invasion.
Literature	GENERIF_SUMMARY	findings demonstrate MPP8 and SIRT1 reciprocally regulate each other's function at multiple molecular layers through physical interaction; disruption of MPP8-SIRT1 interaction de-represses E cadherin expression and reduces cell motility and invasiveness suggesting this interplay plays critical role in MPP8- and SIRT1-mediated epithelial-mesenchymal transition
Literature	GENERIF_SUMMARY	E-Cadherin under-expression is associated with hepatic carcinoma.
Literature	GENERIF_SUMMARY	We demonstrate the potential of common inherited genetic variants to inform patient outcome and show that cdh1's rs9929218 identifies approximately 8% of colorectal cancer patients with poor prognosis.
Literature	GENERIF_SUMMARY	The de-regulation of both of miR-29b/c and DNMT3A leads to the epigenetic silencing of CDH1 and contributes to the metastasis phenotype in gastric cancer.
Literature	GENERIF_SUMMARY	Review of hereditary diffuse gastric cancer associated with CDH1 gene mutation.
Literature	GENERIF_SUMMARY	MicroRNA-10b Triggers the Epithelial-Mesenchymal Transition (EMT) of Laryngeal Carcinoma Hep-2 Cells by Directly Targeting the E-cadherin.
Literature	GENERIF_SUMMARY	Nuclear factor I-C regulates E-cadherin via control of KLF4 in breast cancer
Literature	GENERIF_SUMMARY	E-cadherin is downregulated at the infiltrating front of colorectal carcinomas. The downregulation is further enhanced amongst late stage and high grade tumours compared with earlier stage and low grade tumours.
Literature	GENERIF_SUMMARY	To increased E-cadherin expression and decreased Akt phosphorylation.
Literature	GENERIF_SUMMARY	In patients with placenta accreta, extravillous trophoblasts (EVT) closer to the placental-myometrial interface demonstrated less E-CAD staining than those found deeper in the myometrium.
Literature	GENERIF_SUMMARY	expression and promoter methylation status in colon adenocarcinoma
Literature	GENERIF_SUMMARY	CDH1 hypermethylation is correlated with Helicobacter

Main Category	Category	Terms
		pylori status, indicating that it plays a more important role in the pathogenesis of Helicobacter pylori-positive gastric cancer
Literature	GENERIF_SUMMARY	The combined expression of ezrin(high) and E-cad(low) or ezrin(high) and ezrin(c) was more associated with lymph node metastasis and poor prognosis. I
Literature	GENERIF_SUMMARY	The results of the present study point to a new direction to pharmacologically accelerate corneal re-epithelialization, and should have salient clinical implication.
Literature	GENERIF_SUMMARY	Cyclin B1 could suppress the invasion and metastasis of colorectal cancer cells through regulating E-cadherin expression, which enables the development of potential intervention strategies for colorectal cancer.
Literature	GENERIF_SUMMARY	sE-cadherin performed most favorably from a large panel of serum proteins in terms of diagnostic and predictive potential in curatively treatable PCa.
Literature	GENERIF_SUMMARY	In the cascade of gastric carcinoma development, an association was found with loss of E-cadherin.
Literature	GENERIF_SUMMARY	our results revealed that activation of the P2X7 receptor by ATP promotes breast cancer cell invasion and migration, possibly via activation of the AKT pathway and regulation of E-cadherin and MMP-13 expression.
Literature	GENERIF_SUMMARY	Results suggest a regulatory function of E-cadherin that modulates Nrg1 signaling and promotes Schwann cell myelin formation
Literature	GENERIF_SUMMARY	knockdown of ILK inhibits glioma cell migration, invasion and proliferation through upregulation of E-cadherin and downregulation of cyclin D1. Our results suggest that ILK may serve as a promising therapeutic target for glioma.
Literature	GENERIF_SUMMARY	Report association between human papillomavirus and Epstein - Barr virus DNA and gene promoter methylation of RB1 and CDH1 in range of cervical lesions.
Literature	GENERIF_SUMMARY	Univariate analyses showed that AR, CDH1 and Ki-67 expression levels were significantly associated with overall survival in triple-negative breast cancer patients.
Literature	GENERIF_SUMMARY	E-cadherin expression was associated with EGFR mutation status and patient prognosis in lung adenocarcinoma patients
Literature	GENERIF_SUMMARY	E-cadherin and vimentin positive expression was associated with tumor metastasis of oral squamous cell carcinoma
Literature	GENERIF_SUMMARY	Particulate matter (PM10) downregulates E-Cadherin/ beta-Catenin expression.
Literature	GENERIF_SUMMARY	All individuals with a CDH1 mutation had a personal or family history of diffuse gastric cancer. Patients with gastric cancer and germline CDH1 mutations had

Main Category	Category	Terms
		shorter survival times than patients who met the HDGC criteria but did not have CDH1 mutations.
Literature	GENERIF_SUMMARY	Demethylase recovered CDH1 gene expression may be involved in the apoptosis process. The beta-catenin protein translocated from the nucleus and cytoplasm to the membrane result in inactivation of b-catenin signaling pathway
Literature	GENERIF_SUMMARY	Suggest ECAD expression as independent prognostic factor in colorectal carcinoma.
Literature	GENERIF_SUMMARY	We concluded that CDH1 contributes to NSCL/P with mainly rare, moderately penetrant variants, and CDH1 haploinsufficiency is the likely etiological mechanism.
Literature	GENERIF_SUMMARY	expression. E-cadherin may play a significant role in the sensitivity regulation of EGFR molecular targeting treatment. E-cadherin may provide important clues for selecting proper EGFR-TKI molecular targeting treatment
Literature	GENERIF_SUMMARY	ZIP5 knockdown inhibited the proliferation, migration and invasion of ESCC and suppressed COX2, cyclin D1 and E-cadherin expression, which led to the inhibition of cell progression in esophageal squamous cell carcinoma .
Literature	GENERIF_SUMMARY	SFN inhibited invasion by activating ERK1/2 to upregulate E-cadherin and downregulate CD44v6, thereby reducing MMP-2 expression and activity. E-cadherin is an invasion inhibitor, while CD44v6 and MMP-2 are invasion promoters
Literature	GENERIF_SUMMARY	Decreased membranous E-cadherin expression and aerogenous spread may be associated with worse disease-free survival.
Literature	GENERIF_SUMMARY	Notably, other mutations including MET, APC, CDH1, and FBXW7 were also identified in Taiwanese oral squamous cell carcinoma patients.
Literature	GENERIF_SUMMARY	the allosteric regulation of cell surface E-cadherin by p120(ctn) dephosphorylation.
Literature	GENERIF_SUMMARY	findings of our study imply that sequential staining of CK20 and E-cadherin prevents false-positive classification of BCC. Furthermore, our study demonstrated that p40 exhibits the same staining pattern in BCC, TE, and TB
Literature	GENERIF_SUMMARY	Report role of CDH1 mutations in estimates of age-associated risks of gastric and breast cancer.
Literature	GENERIF_SUMMARY	site-specific E-cadherin glycosylation modification can directly modulate E-cadherin mediated cell-cell adhesion, a key pathophysiological event in gastric cancer progression.
Literature	GENERIF_SUMMARY	There were differences in the expression of Nrf2, Keap1, p16 and E-cadherin between different ovarian carcinoma subtypes.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	CDH1 polymorphism in -347G-->GA was observed to be associated with susceptibility of pancreatic cancer
Literature	GENERIF_SUMMARY	results indicate Moesin may regulate cell motility through its interactions with MT1-MMP and E-cadherin/ p120-catenin adhesion complex and cytoplasmic expression of Moesin correlates with nodal metastasis and poor prognosis of OSCCs
Literature	GENERIF_SUMMARY	E-cadherins arrange in ordered clusters; the first demonstration of the existence of oligomeric cadherins at cell-cell contacts.
Literature	GENERIF_SUMMARY	CDH1 is reduced in endometriosis.
Literature	GENERIF_SUMMARY	intact E-Cad expression could be because of the fact that tumour cells travel as clusters and retain their expression of E-Cad
Literature	GENERIF_SUMMARY	CDH1 and IL1-beta expression dictates FAK and MAPKK-dependent cross-talk between cancer cells and human mesenchymal stem cells
Literature	GENERIF_SUMMARY	knockdown of E-cadherin decreasing MDR1 via HIF-1alpha is involved in the mechanism of multicellular resistance in colorectal cancer
Literature	GENERIF_SUMMARY	Combined therapy with cytotoxic and anti-VEGF agents was shown to result in a significant increase in the number of cases of normal membrane localization of E-cadherin as compared with control (p = 0.00043) and cytotoxic therapy-alone (p = 0.01) groups
Literature	GENERIF_SUMMARY	NKX6.1 directly enhances the mRNA level of E-cadherin by recruiting BAF155 coactivator and represses that of vimentin and N-cadherin by recruiting RBBP7 (retinoblastoma binding protein 7) corepressor.
Literature	GENERIF_SUMMARY	in the NMA cases, none of these factors was statistically significant. In a univariate analysis, neither EGFR nor E-cadherin expression showed a significant impact on disease-free or overall survival.
Literature	GENERIF_SUMMARY	These data show that MTBP aggravates the invasion and metastasis of HCC by promoting the MDM2-mediated degradation of E-cadherin.
Literature	GENERIF_SUMMARY	genetic polymorphisms of CDH1 and CTNNB1 were associated with breast cancer susceptibility and patients' prognosis
Literature	GENERIF_SUMMARY	Epithelial cells with disrupted E-cadherin gene retain the ability to form cell junctions and apico-basal polarity.
Literature	GENERIF_SUMMARY	In non-small cell lung cancer patients, the loss expression of beta-arrestin1 was frequently observed, and beta-arrestin1 expression was significantly correlated with the smoking index and E-cadherin expression, which all indicated beta-arrestin1's significant clinicopathologic role
Literature	GENERIF_SUMMARY	In patients with breast infiltrating ductal carcinoma, E-

Main Category	Category	Terms
		cadherin expression was observed conserved in most of the tumors, 92.5% of cases
Literature	GENERIF_SUMMARY	PLEKHA7 localization to adherens junctions is E-cadherin and p120 dependent.
Literature	GENERIF_SUMMARY	factor
Literature	GENERIF_SUMMARY	analysis of ARID1A, CDH1, cMET and PIK3CA expression and target-related microRNA expression in gastric cancer
Literature	GENERIF_SUMMARY	miR-200b represses esophageal squamous carcinoma cell invasion in vivo without altering the expression of E-cadherin.
Literature	GENERIF_SUMMARY	Data show that myosin II supports a stable Rho zone at the at E-cadherin junctions.
Literature	GENERIF_SUMMARY	Prophylactic laparoscopic total gastrectomy with jejunal pouch reconstruction in patients with CDH1 germline mutation is feasible with 82% of patients having intramucosal diffuse gastric signet ring cell carcinoma in the resection specimen.
Literature	GENERIF_SUMMARY	expression of E-cadherin, and p120 negatively correlated with the tumor differentiation of oral squamous cell carcinoma.
Literature	GENERIF_SUMMARY	Demonstrate that HBx-HCCR-E-cadherin regulation pathway might play an important role in HBV-induced hepatocarcinogenesis.
Literature	GENERIF_SUMMARY	The transforming growth factor beta (TGF-beta) pathway is required for E-cadherin loss to induce inhibition of Nucleotide Excision Repair.
Literature	GENERIF_SUMMARY	The CDH1 +54C/T was associated with susceptibility to endometriosis in Iranian population, and +54T allele may have a protective role in progression of endometriosis.
Literature	GENERIF_SUMMARY	Upregulation of E12/E47 by HBx ultimately and concomitant repression of E-cadherin expression led to epithelial-mesenchymal transition in human hepatocytes.
Literature	GENERIF_SUMMARY	CUGBP1 and HuR negate each other's effects in regulating E-cadherin translation by altering the recruitment of E-cadherin mRNA to PBs and play important roles in the regulation of intestinal barrier integrity.
Literature	GENERIF_SUMMARY	In Panc-1 cells, overexpression of E-cadherin activated the phosphorylation of EGFR and increased the cell sensitivity to erlotinib. In Capan1 cells, knocking down E-cadherin decreased the expression of phosphorylated EGFR, and these cells did not respond to erlotinib.
Literature	GENERIF_SUMMARY	These data suggest that targeting the IL-8/AKT1 signaling pathway and DNMT1 may provide a potential therapeutic approach for blocking NPC metastasis.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Data show that telomerase (hTERT) and ZEB1 protein form a complex, which directly binds to the E-cadherin promoter, and then inhibits E-cadherin expression and promotes epithelial-to-mesenchymal transition (EMT) in colorectal cancer cells.
Literature	GENERIF_SUMMARY	via gain-of-function analyses, DCN overexpression could inhibit renal cell carcinoma (RCC) cell proliferation and metastasis in vitro and vivo. At the mechanism level, we found that an ectopic expression of DCN significantly upregulated P21 and E-cadherin expression. Altogether, these results revealed that DCN is a tumor suppressor in RCC, and it could serve as a potential therapeutic target in patients with RCC.
Literature	GENERIF_SUMMARY	UHRF1 promotes osteosarcoma cell invasion by downregulating the expression of Ecadherin and increasing epithelialtomesenchymal transition in an Rb1dependent manner.
Literature	GENERIF_SUMMARY	Results indicate that HCV core induced epithelial-mesenchymal transition (EMT) by interacting with the transcriptional repressor complex Snail/HDAC1/2 at the E-cadherin promoter, which led to E-cadherin repression and increased invasiveness of hepatoma cells.
Literature	GENERIF_SUMMARY	Cyclin D1 also reversed EMT phenotypes with significantly increased expression of E-cadherin in KYSE-150R cells. However, cyclin D1 siRNA have no radiosensitizing effects on KYSE-150 cells, with no obvious change in EMT marker expression .Our work showed that EMT phenotypes can be reduced and the radiosensitivity of esophageal cancer cells can be enhanced by inhibiting cyclin D1.
Literature	GENERIF_SUMMARY	Downregulated Expression of E-cadherin is associated with Gastric Diseases.
Literature	GENERIF_SUMMARY	PTTG2 induces psoriasis by regulating epidermal expression of vimentin and E-cadherin.
Literature	GENERIF_SUMMARY	A significant association was found between CDH1 -160 C/A (rs16260) polymorphisms and breast cancer risk. The C/A heterozygote and combined C/A + A/A genotypes were found to be strongly associated with increased risk of breast cancer. The variant A allele also increased the breast cancer risk.
Literature	GENERIF_SUMMARY	H3K36me3 correlates with increased skipping of the final 83 base pairs of CDH1 exon 8 in gastric cancer cell lines
Literature	GENERIF_SUMMARY	Immunostaining allowed the identification of specific cell types with E-cadherin, p53, and inhibin-alpha expression in all cases.
Literature	GENERIF_SUMMARY	Loss of E-cadherin disrupts ovarian epithelial inclusion cyst formation and collective cell movement in ovarian cancer.
Literature	GENERIF_SUMMARY	G9a promotes H3K27 methylation of the E-cadherin promoter by upregulating PCL3 to increase PRC2

Main Category	Category	Terms
		promoter recruitment and by downregulating the H3K27 demethylase KDM7A to silence E-cadherin gene
Literature	GENERIF_SUMMARY	MTA1 plays an important role in Epithelial-to-mesenchymal transition (EMT) to promote metastasis via suppressing E-cadherin expression, resulting in a poor prognosis in MPM. MTA1 is a novel biomarker and indicative of a poor prognosis in MPM patients
Literature	GENERIF_SUMMARY	Data show that the expression profiles of three proteins
Literature	GENERIF_SUMMARY	downregulation of PDLIM1 in colorectal cancer samples correlated with reduced E-cadherin and membrane beta-catenin levels
Literature	GENERIF_SUMMARY	These data suggest that the number of polyploid giant cancer cells and the EMT-related proteins E-cadherin, N-cadherin, and vimentin may be valuable biomarkers to assess metastasis in patients with breast cancer.
Literature	GENERIF_SUMMARY	The role of soluble E-cad in tumorigenesis and tumor progression and its significance in clinical therapeutics. [review]
Literature	GENERIF_SUMMARY	OGT inhibited the formation of the Ecadherin/catenin complex through reducing the interaction between p120 and Ecadherin.
Literature	GENERIF_SUMMARY	The present study therefore indicated that IL-6 promoted the process of EMT in HIBECs as characterized by increased migration and invasion of HIBECs and the typical changes in mRNA and protein expression of the EMT markers E-cadherin and vimentin.
Literature	GENERIF_SUMMARY	LOC389641 promotes pancreatic ductal adenocarcinoma progression and increases cell invasion by regulating E-cadherin with the possible involvement of TNFRSF10A.
Literature	GENERIF_SUMMARY	Differential membranous E-cadherin expression, cell proliferation and O-GlcNAcylation in metastatic nodal lesion compared to primary colorectal carcinoma may play role in establishing these lesions.
Literature	GENERIF_SUMMARY	CDH1 expression is lost in 52% of colorectal carcinomas in this cohort.
Literature	GENERIF_SUMMARY	High E-cadherin concentration in presence of enhanced gamma-secretase activity is incontestably a paradoxically result, highlighting that E-cadherin loss is not a pre-requisite for Epithelial-Mesenchymal Transition.
Literature	GENERIF_SUMMARY	Calpain-mediated E-Cadherin fragmentation appears to promote intraperitoneal epithelial ovarian cancer progression.
Literature	GENERIF_SUMMARY	Evaluation of FoxC2 expression, alone or in combination with E-cadherin expression, may help to stratify non-small cell lung cancer patients for risk of disease progression, pointing to this EMT regulator as a

Main Category	Category	Terms
		potential prognostic marker
Literature	GENERIF_SUMMARY	CDH1 germline mutations in hereditary lobular breast cancer [review]
Literature	GENERIF_SUMMARY	Small interfering RNA (siRNA)-mediated knockdown of SPRY4 attenuated the AREG-induced down-regulation of E-cadherin.
Literature	GENERIF_SUMMARY	CDH1 methylation is significantly correlated with decreased CDH1 protein expression.
Literature	GENERIF_SUMMARY	findings demonstrated that reduced membranous E-cadherin and aberrant beta-catenin expression were frequent events in SRCCs of various organs, and that the altered beta-catenin expression was significantly associated with advanced disease
Literature	GENERIF_SUMMARY	As a consequence, Slug transcription is down-regulated relieving A549 cells from Slug-mediated repression of E-cadherin transcription, thereby diminishing the metastatic potential of these oncogenic Ras-expressing NSCLC cells
Literature	GENERIF_SUMMARY	Decreased expression of E-cadherin and Caveolin-1 plays a role in the progression of gastric cancer.
Literature	GENERIF_SUMMARY	E-cadherin expression is down-regulated in colon cancer cells through inactivation of UTX.
Literature	GENERIF_SUMMARY	findings define a major role for Grhl3 in the induction of migration and invasion by the downregulation of E-cadherin in cancer cells
Literature	GENERIF_SUMMARY	Mechanistic investigations showed that LINC01133 could interact with EZH2, LSD1 and recruit them to KLF2, P21 or E-cadherin promoter regions to repress their transcription.
Literature	GENERIF_SUMMARY	Findings indicated that HIF-2alpha promotes epithelial-mesenchymal transition in pancreatic cancer by regulating Twist2 binding to the promoter of E-cadherin.
Literature	GENERIF_SUMMARY	These metastatic tumors revealed no detectable expression of CK8/18, E-cadherin, VCAM-1, and ICAM-1
Literature	GENERIF_SUMMARY	EZH2 is recruited to the E-cadherin promoter by the long non-coding RNA, MALAT-1 (metastasis associated in lung adenocarcinoma transcript 1), where it represses E-cadherin expression
Literature	GENERIF_SUMMARY	Collectively, using tetraspanin CD9 in tandem with E-cadherin as a biomarker in renal cell carcinoma will help to not only distinguish between types, but also predict the metastatic potential of RCC.
Literature	GENERIF_SUMMARY	Results suggest that epigenetic aberration of the e-cadherin gene (CDH1) may occur in the endometrium of patients with fibroids, which may be associated with E-cadherin protein expression in endometrial tissue.
Literature	GENERIF_SUMMARY	Findings indicate that promoter hypermethylation plays

Main Category	Category	Terms
		a significant role in switching off E-cadherin gene expression in malignant melanoma and suggest that loss of E-cadherin may contribute to increased cell proliferation, invasiveness, ulcerative process, metastasis, and reduced disease-free/overall survival.
Literature	GENERIF_SUMMARY	EZH2 mediates the regulation of E-cadherin expression by S100A4 and controls the proliferation and migration of gastric cancer cells.
Literature	GENERIF_SUMMARY	The levels of circadian protein Per2 were significantly increased and E-cadherin was significantly decreased in the tissue of human esophageal cancer with metastasis as compared with non-metastatic esophageal cancer.
Literature	GENERIF_SUMMARY	Truncating somatic alterations in the CDH1 gene occur in 84% of plasmacytoid carcinomas and are specific to this histologic variant.
Literature	GENERIF_SUMMARY	Data show that the nuclear HDAC3 and cytoplasmic CDH1 have independent prognostic value in pancreatic cancer and provide targets for prognostic therapeutics.
Literature	GENERIF_SUMMARY	Data show that grape seed proanthocyanidins could inhibit cell proliferation, invasion and migration of A549 cells, which might be related to the down-regulation of epidermal growth factor receptor and N-cadherin and the up-regulation of E-cadherin.
Literature	GENERIF_SUMMARY	Data suggest that microRNA miR-101 may inhibit MDA-MB-231 breast cancer cell proliferation and migration by repressing DNA methyltransferase 3a (DNMT3a) expression and up-regulating E-cadherin expression.
Literature	GENERIF_SUMMARY	cancer stem cells amount in human lung adenocarcinoma cell line A549 depends on E-cadherin expression.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with Metastasis in Colorectal Cancer.
Literature	GENERIF_SUMMARY	Mutant BRAF represses E-cadherin expression, implicating a catalytic role for BRAF in epithelial-mesenchymal transition.
Literature	GENERIF_SUMMARY	Active ADAM10 promotes the carcinogenesis, invasion, metastasis and proliferation of ESCC and controls invasion and metastasis at least in part through the shedding of E-cadherin activity.
Literature	GENERIF_SUMMARY	The epithelial cell membrane protein E-cadherin physically binds to PKD1 which leads to a subcellular redistribution of PKD1.
Literature	GENERIF_SUMMARY	Study of E-cadherin mediated, intercellular binding kinetics, provide quantitative evidence for confinement-enabled cadherin interactions within intercellular junctions.
Literature	GENERIF_SUMMARY	Data indicate that E-cadherin and caspase-3 were targets of miR-421, which was up-regulated by HIF-1alpha.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	TWIST2 regulates epithelial-mesenchymal transition by depriving the epithelial cell phenotype of E-cadherin and endowing the mesenchymal cell phenotype with Vimentin, which may be involved in the progression and prognosis of ovarian cancer.
Literature	GENERIF_SUMMARY	overexpressed miR-544a may be a promoter for breast cancer migration and invasion by targeting CDH1.
Literature	GENERIF_SUMMARY	CDH1 promoter methylation can be a potential biomarker in ovarian cancer risk prediction. [meta-analysis]
Literature	GENERIF_SUMMARY	A recursively partitioned mixture model of all 3 somatic alteration burden measures resulted in classes of tumors whose epigenetic and genetic burden profile were associated with the PAM50 subtype and mutations in TP53, PIK3CA, and CDH1
Literature	GENERIF_SUMMARY	expression correlated with laryngeal cancer survival
Literature	GENERIF_SUMMARY	Amla extract (Emblica officinalis, AE) decreases the gene and protein expression of IGF1R, a target of miR-375, and SNAIL1, a transcription factor that represses E-cadherin expression.
Literature	GENERIF_SUMMARY	miR23b is key in promoting NPC cell proliferation, migration and invasion through targeting Ecadherin.
Literature	GENERIF_SUMMARY	E-cadherin expression abnormalities in colorectal cancer are rather qualitative than quantitative.
Literature	GENERIF_SUMMARY	Expression of CDH1 and CD24 was transcriptionally upregulated by direct binding of HOXA5 to their promoter sequences as demonstrated by luciferase and ChIP analyses
Literature	GENERIF_SUMMARY	aberrant Notch1 is linked to hepatocellular carcinoma development, tumor recurrence and invasion, which might be mediated, at least in part, through the Notch1-E-Cadherin pathway
Literature	GENERIF_SUMMARY	The combined biomarkers E-cadherin, membranous epidermal growth factor receptor (EGFR) and vimentin show a stronger prognostic value for and disease-free survival than any of the single biomarkers.
Literature	GENERIF_SUMMARY	Slug-upregulated miR-221 promotes breast cancer progression via reducing E-cadherin expression.
Literature	GENERIF_SUMMARY	PAF stimulation dose-dependently promoted the invasion, migration and growth of prostate cancer cells in vitro, while knockdow our findings demonstrate that PAFR can activate ERK1/2 pathway, and subsequently increase MMP-3 expression and decrease E-cadherin expression
Literature	GENERIF_SUMMARY	cortactin binds to E-cadherin, and that a posttranslational modification of cortactin, RhoA-induced phosphorylation by protein kinase D1 (PKD1; also known as PRKD1) at S298, impairs adherens junction assembly and supports their dissolution.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Findings unveil a negative role for PKC-delta in cell-cell adhesion through phosphorylation of E-cadherin.
Literature	GENERIF_SUMMARY	this study shows that the expression and the immunoreactivity of E-cadherin is decreased in the nasal epithelium of patients with allergic rhinitis
Literature	GENERIF_SUMMARY	While Claudin 1 and 4 expression and activities of small GTPases were not affected, gamma-Secretase-mediated cleavage of E-Cadherin was attenuated by Leda-1/Pianp.
Literature	GENERIF_SUMMARY	EHD2 expression, along with the epithelial marker E-cadherin, was markedly reduced in tumor tissues than in adjacent noncancerous tissues. Molecular data indicated that EHD2 inhibited migration and invasion of hepatocellular carcinoma probably by interacting with E-cadherin.
Literature	GENERIF_SUMMARY	Thrombomodulin up-regulated E-cadherin but down-regulated N-cadherin expression, resulting in reversal of epithelial-mesenchymal transition (EMT) in the lung cancer cells.
Literature	GENERIF_SUMMARY	analysis of a novel SMAD-independent pathway linking enhanced activin B signaling to reduced E-cadherin expression and increased migration in type II endometrial cancer
Literature	GENERIF_SUMMARY	findings suggest that E-cadherin and P-cadherin, but not N-cadherin staining, might be useful in diagnosing CIN and for predicting prognosis in patients with early-stage Cervical Squamous Cell Carcinoma.
Literature	GENERIF_SUMMARY	we sequenced the entire 6.5-kb coding region of the CDH1 gene. Among the oral cleft patients, four nonsynonymous single nucleotide variants (SNVs), c.1235T>C (p.V412A), c.1273G>A (p.V425I), c.1565C>T (p.T522I), and c.1888C>G (p.L630V), were identified. The frequency of nonsynonymous SNVs on the CDH1 gene in oral cleft patients (4/80) was significantly higher than that in the control group (1/138)
Literature	GENERIF_SUMMARY	There was no significant difference in E-cadherin and CD44v6 expression among the partial, complete, and invasive hydatidiform moles patient groups.
Literature	GENERIF_SUMMARY	Infiltrating lobular carcinoma (ILC)of the breast represents a biologically distinct subset of breast cancer, a biology defined by specific genetic aberrations in E-cadherin, the high prevalence of ER-positive disease, the relatively low frequency of HER2-positive disease, and specific mutational events revealed by deep sequencing of ILC genomes
Literature	GENERIF_SUMMARY	our findings suggest that Dsg3 functions as a key in cell-cell adhesion through at least a mechanism of regulating E-cadherin membrane trafficking
Literature	GENERIF_SUMMARY	Analysis highlights significant association between CHD1 rs9929218 polymorphism and CRC susceptibility.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	CDH1 -160C>A Polymorphism is not associated with the Risk of Gastrointestinal Cancer.
Literature	GENERIF_SUMMARY	High CDH1 expression is associated with metastasis occurrence in Prostate Cancer.
Literature	GENERIF_SUMMARY	data suggest that syndecan-2 induces extracellular shedding of E-cadherin and supports the acquisition of a fibroblast-like morphology by regulating MMP-7 expression in a colon cancer cell line
Literature	GENERIF_SUMMARY	Progressive loss of e-cadherin/alpha-catenin expression is associated with an aggressive phenotype (low differentiation, increased metastatic activity/ advanced stage) in thyroid carcinomas.
Literature	GENERIF_SUMMARY	dynamic interplay among E-cadherin, tight junctions, and epithelial-to-mesenchymal transition exists and mediates an important function in hepatitis C virus entry
Literature	GENERIF_SUMMARY	The CDH1 downregulation acted as an independent prognostic biomarker in cytogenetically normal acute myeloid leukemia (CN-AML) patients.
Literature	GENERIF_SUMMARY	CDH1 -160C/A polymorphism is associated with breast cancer.
Literature	GENERIF_SUMMARY	Hypermethylation Status of E-Cadherin Gene is associated with Gastric Cancer.
Literature	GENERIF_SUMMARY	Data indicate that CDH11, ICAM1 and CLDN3 were overexpressed in tumors when compared to normal esophagus, normal gastric and non-dysplastic Barrett's.
Literature	GENERIF_SUMMARY	Therefore, EGF is suggested to induce E-cadherin down-regulation at the transcriptional level through the MEK/ ERK pathway, which might result in, at least in part, the induction of cellular morphological changes and cell migration in LoVo cells.
Literature	GENERIF_SUMMARY	Data show that gene silence of ezrin inhibits the proliferation and invasion of prostate cancer PC-3 cells, meanwhile the level of E-cadherin is upregulated and N-cadherin is downregulated.
Literature	GENERIF_SUMMARY	The important role of disturbances in E-cadherin-beta-catenin interactions in the mechanisms of malignant transformation of follicular epithelium.
Literature	GENERIF_SUMMARY	human pluripotent stem cells are subject to matrix metalloproteinase-dependent E-cadherin ectodomain shedding.
Literature	GENERIF_SUMMARY	E-cadherin is transcriptionally downregulated in gastric cancer tissues comparing to adjacent tissue controls.
Literature	GENERIF_SUMMARY	class I HDACs (HDAC1, 2, 3 and 8) play a major role in regulating extracellular matrix and Epithelial-mesenchymal transition, whereas class IIa HDACs (HDAC4 and 5) are less effective.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	DNA methylation level was negatively correlated with mRNA expression of the E-Cadherin gene. Methylation in tumor tissues was significantly higher than that in tumor adjacent tissues (P
Literature	GENERIF_SUMMARY	A novel regulator, myoferlin, of ADAM12 is discovered in HeLa cells and this protein increases ADAM12 expression level, stability, and its enzymatic activity, leading to the reduction of its substrate, E-cadherin, which plays important roles in the regulation of cell adhesion and tumor metastasis.
Literature	GENERIF_SUMMARY	Serum sE-cadherin levels maybe an important indicator of the pathogenesis of progression of HBV-related disorders
Literature	GENERIF_SUMMARY	26-year-old female and her father, a 54-year-old male, were found to be carriers of the CDH-1 mutation
Literature	GENERIF_SUMMARY	CDH1 expression in high grade papillary urothelial carcinomas predicted disease-free survival and overall survival.
Literature	GENERIF_SUMMARY	DNMT1 down regulation mediated upregulation of CDH1 (hereafter, DNMT1/CDH1 loop) in presence of ectopic-excess of miR-152 prevents migration of cancer cells.
Literature	GENERIF_SUMMARY	GDF8 promotes ovarian cancer cell migration via ALK4/5-SMAD2/3-E-cadherin signaling.
Literature	GENERIF_SUMMARY	The Hippo signaling pathway was significantly associated with ER-negative breast cancer (pathway level P = 0.02). Gene-based analyses revealed that CDH1 was responsible for the pathway association (P
Literature	GENERIF_SUMMARY	IGF-II-mediated loss of E-cadherin is central in developing hepatomegaly in mice and abnormal cell growth in the hepatoma cell line
Literature	GENERIF_SUMMARY	Loss of CDH1 expression is associated with prostate and liver cancer.
Literature	GENERIF_SUMMARY	The expression level of E-cadherin in liver cirrhotic tissues were significantly lower than that in chronic hepatitis tissues.
Literature	GENERIF_SUMMARY	E-cadherin inhibits beta-catenin in the context of disruption of the APC-destruction complex, and that this function is also EC1 domain dependent. Both binding functions of E-cadherin may be required for its tumour suppressor activity.
Literature	GENERIF_SUMMARY	PARP3 controls of TGFbeta-induced epithelial mesenchymal transformation and acquisition of stem-like cell features by stimulation transglutaminase 2/ SNAI1 signaling.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with lung metastasis in melanoma.
Literature	GENERIF_SUMMARY	Bmi-1 promotes the invasion and migration of colon

Main Category	Category	Terms
		cancer stem cells through the downregulation of E-cadherin.
Literature	GENERIF_SUMMARY	Decreased expression of CDH1 or CTNNB1 in the cell membranes of cancer cells is associated with poor survival of patients with esophageal cancer.
Literature	GENERIF_SUMMARY	alpha-catenin is activated during the initial adhesion assembly process involving filopodia retraction-mediated nucleation and clustering of E-cadherin; activated alpha-catenin continues to be present in the active conformation while E-cadherin clusters are dynamically organized into peripheral and lateral assemblies. Further, alpha-catenin does not require continuous force to maintain its open conformation.
Literature	GENERIF_SUMMARY	the loss of MUC16 and E-cadherin expression resulted in the formation of more compact spheroids. In addition, our data describe an unusual link between E-cadherin expression and less compact spheroids. Our data also emphasize the role of MUC16 and b1 integrin in Epithelial ovarian cancer (EOC) spheroid formation.
Literature	GENERIF_SUMMARY	AXIN1 is expressed in meningioma along with APC and E-cadherin
Literature	GENERIF_SUMMARY	studies indicate Bit1 is an inhibitor of EMT and metastasis in lung cancer and hence can serve as a molecular target in curbing lung cancer aggressiveness
Literature	GENERIF_SUMMARY	Cleavage of e-cadherin, as evidenced by the presence of c-terminal fragments and n-terminal fragments on biopsy, contributes to defective barrier function in neosquamous epithelium.
Literature	GENERIF_SUMMARY	The CDH1 -160C/A polymorphism does not seem to significantly influence the expression of E-cadherin in PA and ACC of salivary glands.
Literature	GENERIF_SUMMARY	Low expression of CDH1 is associated with invasion in non-small cell lung cancer.
Literature	GENERIF_SUMMARY	This study reports the identification of a novel human E-cadherin variant mRNA produced by alternative splicing.
Literature	GENERIF_SUMMARY	Low expression of E-cadherin is associated with bladder cancer.
Literature	GENERIF_SUMMARY	ADH1B and CDH1 polymorphisms predict prognosis in male patients with non-metastatic laryngeal cancer
Literature	GENERIF_SUMMARY	Anxa5 mediates the in vitro malignant behaviours of murine hepatocarcinoma Hca-F cells via ERK2/c-Jun/p-c-Jun(Ser73) and ERK2/E-cadherin pathways.
Literature	GENERIF_SUMMARY	we concluded that the CDH1 -472delA and -160C>A polymorphisms are not associated with diffuse (DGC) or intestinal (IGC) types of gastric cancer in patients from western Mexico.
Literature	GENERIF_SUMMARY	Neutrophil elastase has the capacity to cleave E-cad

Main Category	Category	Terms
		and interfere with its cell-cell adhesion function in acutely injured lung epithelium.
Literature	GENERIF_SUMMARY	Results show that CDH1 expression is significantly downregulated in human hepatocellular carcinoma (HCC) tissues and inversely correlates with RARgamma expression. Also, the study shows that CDH1 expression is under the regulation of RARgamma.
Literature	GENERIF_SUMMARY	-347GA/GA and -160A/A genotypes may be prognostic markers that can help to identify patients at increased risk of invasive/metastatic cancer in northern China
Literature	GENERIF_SUMMARY	E-cadherin and beta-catenin expression provides discriminative prognostic power independent of conventional pathologic factors, thus further reinforcing the important role of cell adhesion molecules in the process of tumor metastasis, especially in triple-negative breast cancer (TNBC).
Literature	GENERIF_SUMMARY	Expression of E-cadherin was significantly associated with the diffuse-type familial EOGCs and early stage, whereas expression of ERalpha might have little role in EOGC tumorigenesis.
Literature	GENERIF_SUMMARY	CDH4 may play a negative role in the growth and metastasis of salivary adenoid cystic carcinoma via co-expression with E-cadherin.
Literature	GENERIF_SUMMARY	Mutations in CDH1 gene is associated with breast cancer.
Literature	GENERIF_SUMMARY	Endometrial expression of HOXA10 and E-cadherin are reduced in women with recurrent implantation failure and recurrent miscarriage.
Literature	GENERIF_SUMMARY	Subsequent mechanistic investigation indicated that cadherin 1 (CDH1)/beta-catenin signaling and Notch-1/Akt signaling were the downstream pathways of miR-92a-3p in glioma cells and glioma stem-like cells, respectively
Literature	GENERIF_SUMMARY	Intralobular breast carcinoma patients with tumors carrying both CDH1 and ERBB2 mutations have a worse prognosis.
Literature	GENERIF_SUMMARY	Study shows that the protein level of E-CAD is indirectly regulated by MIR-193a in non-small cell lung cancers through downregulation of WT1.
Literature	GENERIF_SUMMARY	Sprouty2 inhibits amphiregulin-induced down-regulation of E-cadherin and cell invasion in human ovarian cancer cells.
Literature	GENERIF_SUMMARY	E-cadherin-160 genotype represents a valid risk factor for high-grade squamous intraepithelial lesions and cervical intraepithelial neoplasia.
Literature	GENERIF_SUMMARY	E-cadherin genetic variants control disease severity and progression and could be a marker of disease outcome.
Literature	GENERIF_SUMMARY	Radixin knockdown suppresses the metastasis of

Main Category	Category	Terms
		SGC-7901 cells in vitro by up-regulation of E-cadherin. The NF-kappaB/snail pathway contributes to the regulation of E-cadherin in response to depletion of radixin.
Literature	GENERIF_SUMMARY	the decrease of CDH1 transcript, corresponding to the canonical protein, and the presence of CDH1a, corresponding to an alternative isoform, are likely to perturb E-cadherin-mediated signaling and cell-cell adhesion, thus contributing to intestinal-type gastric carcinogenesis
Literature	GENERIF_SUMMARY	Reduced E-cadherin expression is commonly reported in esophageal adenocarcinoma and independently predicts neoplasm metastasis and poor patient survival.
Literature	GENERIF_SUMMARY	PI3Kbeta associated with N-cadherin and PIK3CB depletion selectively disrupted N-cadherin cell adhesions in PTEN-mutant bladder carcinoma.
Literature	GENERIF_SUMMARY	HIF-1alpha expression was upregulated in the vasculogenic mimicry-positive CRC cell line HCT-116 and thereby affected the expression of the EMT-related markers Claudin-4, E-cadherin (E-cd) and Vimentin(VIM).
Literature	GENERIF_SUMMARY	Results provide evidence that Cdh1 is a major contributor to E2F1 degradation during epidermal keratinocyte differentiation.
Literature	GENERIF_SUMMARY	eIF4E promoted cholangiocarcinoma cell metastasis by up-regulating the expression of VEGF-C, MMP-2 and suppressing E-cadherin expression.
Literature	GENERIF_SUMMARY	ESE3 as a negative regulator of pancreatic ductal adenocarcinomas (PDAC) progression and metastasis by enforcing E-cadherin upregulation.
Literature	GENERIF_SUMMARY	There was no association between the CDH1 rs334558 polymorphism and breast cancer. Aberrant promoter methylation profile was found in breast cancer patients and it was related with early cancer stages. CDH1 down-regulation was significantly associated with metastasis and promoter methylation.
Literature	GENERIF_SUMMARY	The data suggest that reduced expression of E-cadherin and over-expression of Slug, Snail, and TGF-beta induce epithelial to mesenchymal transition in ameloblastoma.
Literature	GENERIF_SUMMARY	The results show how E-cadherin instructs the assembly of the LGN/NuMA complex at cell-cell contacts, and define a mechanism that couples cell division orientation to intercellular adhesion.
Literature	GENERIF_SUMMARY	only residual disease and negative E-cadherin immunoexpression were useful independent predictors of unfavourable clinical outcome and shorter overall survival in ovarian serous carcinomas
Literature	GENERIF_SUMMARY	These results suggest that stabilization of delta-catenin by Hakai is dependent on Src.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Slug expression was inhibited and E-cadherin expression was increased in the Slug siRNA-transfected anaplastic thyroid tumors.
Literature	GENERIF_SUMMARY	Data show that paired box gene 5 (B-cell lineage specific activator) protein (Pax-5) induces E-cadherin expression in breast cancer cells.
Literature	GENERIF_SUMMARY	High RAS-associated protein RAB5 expression correlated with the presence of lymphatic invasion and venous invasion and low E-cadherin expression.
Literature	GENERIF_SUMMARY	Results show that interleukin-10 (IL-10) positively regulated the expression of E-cadherin by downregulation of microRNA miR-9 in leukemia cells.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with colon cancer.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with recurrence in pancreatic head cancer.
Literature	GENERIF_SUMMARY	Data indicate that ellagic aicd (EA) down-regulates the expression of COX-2, NF-kappa B, vimentin and up-regulates the expression of E-cadherin in in pancreatic carcinom PANC-1 cells.
Literature	GENERIF_SUMMARY	Study demonstrate that CdGAP acts as a novel co-transcriptional repressor with Zeb2 to suppress E-cadherin expression in breast cancer cells.
Literature	GENERIF_SUMMARY	Our study demonstrates that the CDH1 polymorphisms were significantly associated with the risk of nonsyndromic cleft lip with or without cleft palate in the northern Chinese Han population
Literature	GENERIF_SUMMARY	AnxA5 2D-network mediates E-cadherin mobility in the plasmalemma that triggers human trophoblasts aggregation and thereby cell fusion.
Literature	GENERIF_SUMMARY	Data suggest ATM and E-cadherin expression as potential supportive predictive markers for EGFR-targeted therapy.
Literature	GENERIF_SUMMARY	lnc AGAP2-AS1 upregulated by SP1 plays an important role in gastric cancer development and progression by suppressing P21 and E-cadherin
Literature	GENERIF_SUMMARY	High p120-catenin expression is associated with skin cancer.
Literature	GENERIF_SUMMARY	Impairment of E-cadherin/N-cadherin adhesion abrogates the ability of Cancer-associated fibroblasts to guide collective cell migration and blocks cancer cell invasion.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with gastric cancer.
Literature	GENERIF_SUMMARY	ANP inhibits TGF-beta1-induced EMT in 16HBE-14o and A549 cells through cGMP/PKG signaling, by which it targets TGF-beta1/Smad3 via attenuating phosphorylation of Smad3. These findings suggest the potential of ANP in the treatment on pulmonary

Main Category	Category	Terms
		diseases with airway remodeling.
Literature	GENERIF_SUMMARY	Data indicate a positive correlation between metastasis-associated protein 1 (MTA1) and microRNA miR-22, supporting their inhibitory effect on E-cadherin expression.
Literature	GENERIF_SUMMARY	Data suggest expression of ECAD (E-cadherin) in bladder cancer cell lines correlates with presence of intercellular junctions and level of PAK5 (p21-activated kinase PAK5 protein) expression; endogenous PAK5 is localized to intercellular junctions and interacts with ECAD; depletion of PAK5 expression reduces integrity of intercellular junctions; PAK5 mRNA levels are reduced in bladder cancer compared with normal controls.
Literature	GENERIF_SUMMARY	At the molecular level, transcription of the adherens junction protein E-cadherin is upregulated on nicotinic acid addition, leading to accumulation of E-cadherin protein at the cell-cell boundary. This can be attributed to nicotinic acid's ability to facilitate the ubiquitination and degradation of Snail1, a transcription factor that represses E-cadherin expression.
Literature	GENERIF_SUMMARY	miR-203/ERK1/2/Slug/E-cadherin signaling pathway plays an essential role on SGC7901 cell invasion and motility. miR-203 can be novel modalities to prevent peritoneal metastasis of invasive cancers such as gastric cancer.
Literature	GENERIF_SUMMARY	total loss of expression in 55% of undifferentiated/ dedifferentiated endometrial carcinomas
Literature	GENERIF_SUMMARY	current study demonstrates that hypoxia reduces E-cadherin expression and increases cell migration, regardless of the methylation profile; additionally, no differences in E-cadherin methylation patterns were observed among oral mucosa, potential malignant oral lesion and oral squamous cell carcinoma
Literature	GENERIF_SUMMARY	The extracellular domain can be defined as a colorectal cancer 5FU resistance marker
Literature	GENERIF_SUMMARY	The BC cells showed the coexpression of E- and P-cadherins, as well as release of the molecules b- and p120-catenins into the cytoplasm of tumor cells, which leads to the activation of intracellular mechanisms for changing the structure of the cytoskeleton and the level of proliferation
Literature	GENERIF_SUMMARY	E-cadherin expression was increased by transfection of p300 small interfering RNA in a dose-dependent manner.. There was a correlation between Snail and p300 expressions in lung cancer. Moreover, p300 acetylates Snail both in vivo and in vitro, and K187 may be involved in this modification.
Literature	GENERIF_SUMMARY	JianPi JieDu Recipe could inhibit TGF-beta-induced epithelial-to-mesenchymal transition in colorectal cancer through TGF-beta/Smad mediated Snail/E-cadherin expression

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	In 11 BCD patients from eight families, we identified five CDH1 deleterious missense mutations and three CTNND1 truncating mutations.
Literature	GENERIF_SUMMARY	E-cadherin expression in breast cancers is highly comparable between briefly fixed specimens and the corresponding traditionally fixed resection specimens.
Literature	GENERIF_SUMMARY	Combining the targets E-cadherin, epithelial membrane antigen (EMA), human epidermal growth receptor type 2 (Her2/neu), carcinoembryonic antigen (CEA) resulted in nearly 100% detection of ductal ovarian metastases, whereas the combination of EMA, Her2/neu and epithelial cell adhesion molecule (EpCAM) was most suitable to detect lobular ovarian metastases.
Literature	GENERIF_SUMMARY	mDia1 was recruited to the zonula adherens (ZA) of established Caco-2 monolayers in response to E-cadherin and RhoA.
Literature	GENERIF_SUMMARY	ILK aids trophoblast syncytialization via downregulation of CDH1, perhaps through an ILK-PARP1-SNAI1 pathway
Literature	GENERIF_SUMMARY	We silenced Srx by short hairpin RNA in HeLa and SiHa cells. Diminished Srx expression upregulated E-cadherin expression. The cell invasion and migration activity in the ShSrx group were obviously decreased in HeLa and SiHa cells.
Literature	GENERIF_SUMMARY	Study demonstrated that down-regulation of E-cadherin contributes to epithelial-to-mesenchymal transition-mediated chemo-resistance of prostate cancer.
Literature	GENERIF_SUMMARY	integrin signaling potentiates transforming growth factor-beta 1 (TGF-beta1) dependent down-regulation of E-Cadherin expression - Important implications for epithelial to mesenchymal transition (EMT) in renal cell carcinoma
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with Non Small-cell Lung Cancer.
Literature	GENERIF_SUMMARY	protocadherin 7 may play important roles in the invasion and metastasis of gastric cancer, and protocadherin 7 could suppress cell migration and invasion through E-cadherin inhibition.
Literature	GENERIF_SUMMARY	The expression of Bcl-2 and E cadherin immunopositivity was associated positively with tumor grade, high T category and histopathological grades. The results of this study points to the significance of cell proliferation and invasion as a major determinant of prognosis in OSCC.
Literature	GENERIF_SUMMARY	House dust mite exposure on human bronchial epithelial cells (16HBE) increased the expression of TSLP and E-cadherin dysfunction by PI3K/Akt signaling pathway. Exposure of 16HBE to TSLP and HDM resulted in aberrant cleavage and redistribution of E-cadherin.
Literature	GENERIF_SUMMARY	A correlation was found between stage, tumor

Main Category	Category	Terms
		differentiation grade, risk for relapse or progression of disease, and the impaired expression of different EMT markers
Literature	GENERIF_SUMMARY	CDH1 hypermethylation may play a key role in the carcinogenesis of HCC
Literature	GENERIF_SUMMARY	Data suggest that epithelial-mesenchymal transition (EMT) is regulated by ZMYND8 (receptor for activated protein kinase C) which selectively activates gene promoters of CLDN1 (claudin 1) and CDH1 (E-cadherin) in breast cancer cells; thus, the presence of ZMYND8 could be implicated in maintaining epithelial phenotype of cells; ZMYND8 regulates invasion/migration of breast cancer cells.
Literature	GENERIF_SUMMARY	Positive expression of matrix metalloproteinase-9 and vascular endothelial growth factor and negative expression of E-cadherin were malignant markers for gastric cancer. Positive expression of vascular endothelial growth factor and negative expression of E-cadherin were associated with L-OHP resistance.
Literature	GENERIF_SUMMARY	select gene expression alterations in c.1380delA CDH1 SB.mhdgc-1 cells, are reported.
Literature	GENERIF_SUMMARY	MEG3/miR-421/E-cadherin regulatory axis may be a novel therapeutic target for breast cancer.
Literature	GENERIF_SUMMARY	While there is a notion that the epithelial-mesenchymal transition zinc finger E-box-binding homeobox 1-E-Cadherin cascade is initiated by zinc finger E-box-binding homeobox 1, our results are contrary and outline other possible pathways influencing the regulation of E-Cadherin.
Literature	GENERIF_SUMMARY	The results showed that, despite being a missense mutation, c.977T>A (p.I326N) is a new germline pathogenic mutation of the CDH1 gene, not described to date.
Literature	GENERIF_SUMMARY	Loss of TET1 expression facilitates colon cancer cell migration via H3K27me3-mediated repression of E-cadherin expression.
Literature	GENERIF_SUMMARY	the blockage of SPHK1 activity to attenuate autophagy may be a promising strategy for the prevention and treatment of patocellular carcinoma
Literature	GENERIF_SUMMARY	Higher proportions of early-onset Diffuse Gastric Cancers (DGCs) contained somatic mutations in CDH1 or TGFBR1 compared with late-onset DGCs. A smaller proportion of early-onset DGCs contained somatic mutations in RHOA. CDH1 alterations, but not RHOA mutations, were associated with shorter survival times. Female predominance in early-onset DGC may be related to relatively high rates of somatic CDH1 and TGFBR1 mutations.
Literature	GENERIF_SUMMARY	73.0% of the evaluated tumors showed abnormal E-cadherin expression in IHC.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	High expression of CDH1 is associated with colorectal cancer.
Literature	GENERIF_SUMMARY	MEG3 acts as a ceRNA to regulate expression of E-cadherin and FOXO1 by competitively binding miR-9 and may be used as a potential biomarker in predicting ESCC patients' progression and prognosis
Literature	GENERIF_SUMMARY	Here, the authors use a quantitative Forster resonance energy transfer (FRET) approach to show that epithelial cadherin forms constitutive dimers and that these dimers exist independently of the actin cytoskeleton or cytoplasmic proteins.
Literature	GENERIF_SUMMARY	All substances, nilotinib, dasatinib, erlotinib and gefitinib have a significant impact on beta-catenin and E-cadherin expression in both HPV16-positive and HPV16-negative cells in vitro. Hence, alterations of beta-catenin and E-cadherin could provide novel insights for future targeted therapies of head and neck SCC
Literature	GENERIF_SUMMARY	Application of force to E-cadherin stimulates liver kinase B1 to activate AMP-activated protein kinase, a master regulator of energy homeostasis. LKB1 recruits AMPK to the E-cadherin mechanotransduction complex, thereby stimulating actomyosin contractility, glucose uptake and ATP production.
Literature	GENERIF_SUMMARY	Heterozygosity and mutant homozygosity as well as the combination of both TP53 Arg72Pro and CDH1 rs16260 polymorphisms are responsible to increase the risk of colorectal cancer development in Bangladeshi population.
Literature	GENERIF_SUMMARY	E-cadherin silencing relies on the formation of a complex between the paRNA and microRNA-guided Argonaute 1 that, together, recruit SUV39H1 and induce repressive chromatin modifications in the gene promoter
Literature	GENERIF_SUMMARY	DcR3 treatment caused HepG2 cell cytoskeleton remodeling, inhibited E-cadherin expression, and promoted cell migration.
Literature	GENERIF_SUMMARY	E-cadherin promoter methylation in Epstein-Barr virus-associated oral squamous cell carcinomas
Literature	GENERIF_SUMMARY	VIM is downregulated and E-cad is upregulated in T1 stage non-small-cell lung cancer, suggesting that a mesenchymal-epithelial transition may take place in the early-stage of tumor development
Literature	GENERIF_SUMMARY	PCNA and E-cadherin may have roles in prognosis of patients with gastric cancer
Literature	GENERIF_SUMMARY	Results show that mesenchymal N-cadherin-expressing (Ncad+) cells and MCAs invade much more efficiently than E-cadherin-expressing (Ecad+) cells. Data emphasize the role of Ncad in intraperitoneal seeding of EOC and provide the rationale for future studies targeting Ncad in preclinical models of epithelial

Main Category	Category	Terms
		ovarian cancer metastasis.
Literature	GENERIF_SUMMARY	our data suggest that miR-1285-5p functions as a tumor promoter in the development of non-small-cell lung carcinoma by targeting Smad4 and CDH1, indicating a novel therapeutic strategy for non-small-cell lung carcinoma patients.
Literature	GENERIF_SUMMARY	we report that E-cadherin can mediate UVR-induced melanin transfer
Literature	GENERIF_SUMMARY	Studies indicate that Grainyhead-like transcription factor 2 (GRHL2) controls the expression of E-cadherin (CDH1) required for adherens junctions and possibly regulates the expression of claudin-4 (CLDN4) in tight junctions.
Literature	GENERIF_SUMMARY	mechanistic experiments revealed that ADAM10-RNAi resulted in an increase in E-cadherin and a decrease in N-cadherin and vimentin expression. Our study implies that high expression of ADAM10 promotes the proliferation and migration of hypopharyngeal squamous cell carcinoma (HSCC). These findings may help to provide a method for treatment of HSCC
Literature	GENERIF_SUMMARY	Real-time PCR showed E-cad mRNA decreased in SCC25 while increased in RAW 264.7 of the indirect cell co-culture model, and immunofluoresence (IF) observed the evident switch of E-cad staining from SCC25 to RAW 264.7
Literature	GENERIF_SUMMARY	Findings show that the expression of E-cadherin in endometrial carcinoma is significantly related to GalNAc-T6 expression but it is not associated with better prognosis.
Literature	GENERIF_SUMMARY	A transition in the stiffness of E-cadherin homotypic interactions regulates actin and membrane dynamics during initial stages of cell-cell adhesion.
Literature	GENERIF_SUMMARY	FDG uptake on PET was associated with negative E-cadherin expression in patients with lung adenocarcinoma.
Literature	GENERIF_SUMMARY	overexpression of SPRY4-IT1 increased E-cadherin and decreased N-cadherin and vimentin protein levels, indicating that SPRY4-IT1 may regulate ovarian cancer cell metastasis through the inhibition of epithelial-mesenchymal transition.
Literature	GENERIF_SUMMARY	E-cadherin re-expression and subsequent induction of NF-kappaB signaling likely enhance energy production and cell proliferation.
Literature	GENERIF_SUMMARY	reduction of Alex3 correlates with the development of non-small cell lung cancer and predicts adverse clinical outcome of non-small cell lung cancer patients. The effect of Alex3 on inhibiting invasion and migration may attribute to upregulation of E-cadherin expression through AKT-Slug pathway inactivation.
Literature	GENERIF_SUMMARY	Reduced expression of E-cadherin in clear cell renal

Main Category	Category	Terms
		carcinoma
Literature	GENERIF_SUMMARY	E-cadherin expression promotes growth of primary breast tumors and conceivably the formation of metastases, supporting a role for MET in metastasis. E-cadherin needs to be reevaluated as a tumor suppressor.
Literature	GENERIF_SUMMARY	Early-stage invasive ductal carcinomas or invasive non-lobular breast cancers with the triple-negative breast cancer (TMBC) molecular phenotype have a higher risk for the loss of E-cad expression than do tumors with non-TNBC molecular phenotypes.[meta-analysis]
Literature	GENERIF_SUMMARY	Depletion of Brg1 improves the integrity of airway epithelium in asthma by regulating E-cadherin expression in lung epithelial cells. Knockdown of Brg1 increased the cell proliferation and migration by human bronchial epithelial 16HBE cells.
Literature	GENERIF_SUMMARY	Findings uncover a new regulatory network in RCC involving metastasis-promoting miR-720 that directly targets expression of key metastasis-suppressing proteins E-cadherin and alphaE-catenin complex.
Literature	GENERIF_SUMMARY	High Twist-1 expression, aberrant E-cadherin and high EZH2 expression in primary prostate cancer are considered as adverse prognostic markers of an aggressive tumor with high metastatic potential.
Literature	GENERIF_SUMMARY	Immunostaining revealed that NKA was co-localized with E-cadherin in the glands of the gastric epithelium. Both NKA activity and alpha1-isoform protein expression were reduced in the study group (P
Literature	GENERIF_SUMMARY	ZIP6 deficiency disturbs intracellular Zn2+ homeostasis, leading to increased cell survival in hypoxia and reduced E-cadherin expression, indicating that decreased ZIP6 expression is strongly associated with resistance to hypoxia.
Literature	GENERIF_SUMMARY	There was also lower expression of the genes CADM1 and CDH1 in melanomas greater than 1.0mm thick compared to melanomas less than 1.0mm
Literature	GENERIF_SUMMARY	Results identified a weak expression of E-cadherin in the majority of patients with glioma. Also, E-cadherin, as a representative EMT marker, seems to have limited prognostic value in glioma.
Literature	GENERIF_SUMMARY	down-regulation of USP48 increases E-cadherin expression and epithelial barrier integrity through reducing TRAF2 stability
Literature	GENERIF_SUMMARY	patients with both abnormal expression of p0071 and E-cadherin (cytoplasmic expression) had a statistically significant shorter survival than the patients without both abnormal expression (P
Literature	GENERIF_SUMMARY	Taken together, these findings demonstrated that Twist was upregulated in high invasion and metastasis cell lines as well as invasive ductal carcinoma (IDC) tissues

Main Category	Category	Terms
		companioned with downregulated expression of E-cadherin and ER, which provides important clues for the deeper study of breast cancer
Literature	GENERIF_SUMMARY	N-cadherin can be used as a potential biomarker for early diagnosis of OSCC (oral squamous cell carcinoma). However, the N-cadherin expression did not show any correlation with the histological grade of OSCC (oral squamous cell carcinoma).
Literature	GENERIF_SUMMARY	Epithelial marker E-cadherin is remarkably reduced on the surface of cervical adenocarcinoma cells and squamous cell carcinoma cells.
Literature	GENERIF_SUMMARY	Study demonstrated that E-cadherin is a determinant molecule associated with ovarian cancer (OC) progression, dissemination and aggressiveness; showed that total and membranous E-cadherin expression levels is a specific and sensitive marker to differentiate advanced- from early-stage tumors and serous tumors from other histological types.
Literature	GENERIF_SUMMARY	Data suggest that ECAD, STAT3, Bak, and Bcl-xL are expressed in affected endometrial tissues of women with endometrioid adenocarcinoma depending on neoplasm staging and cell differentiation. This study was conducted using immunohistochemistry of surgically resected tissues. (STAT3 = signal transducer and activator of transcription 3 protein; Bak = pro-apoptotic protein BAK; Bcl-xL = BCL2 associated agonist of cell death)
Literature	GENERIF_SUMMARY	CDH1 single-nucleotide polymorphisms role in susceptibility to non-small cell lung cancer
Literature	GENERIF_SUMMARY	Negative correlation between HOTAIR and E-cadherin was found in gastric cancer tissues and cell lines, and HOTAIR contributed to the regulation of E-cadherin through binding to EZH2 with the E-cadherin promoter.
Literature	GENERIF_SUMMARY	Our results explored that autocrine production of human growth hormone enhances cellular proliferation by the activation of nuclear factor-kappaB (65 kDa) and downregulation of E-cadherin expression.
Literature	GENERIF_SUMMARY	Results show that E-cadherin/beta-catenin complex is disrupted by ICAT promoting epithelial-mesenchymal transition of cervical cancer cells. .
Literature	GENERIF_SUMMARY	The expression of E-cad and VEGF and their relationship with clinical features of triple-negative breast cancer (TNBC) suggest that Uygur patients might have different prognostic factors as compared with Han patients. The expression of E-cad was negatively correlated with lymph node metastasis, stage, and histological grade.
Literature	GENERIF_SUMMARY	Results strongly indicate a protective role of allele A of the SNP CHD1 -160C>A against urinary calculi formation in children.
Literature	GENERIF_SUMMARY	In conditioned media of PIBF-deficient JEG-3 cells,

Main Category	Category	Terms
		primary lung and ovarian cancer cells MMP9 activity was reduced. PIBF silencing resulted in increased E-cadherin expression, suggesting that by down regulating E-cadherin expression, PIBF might interfere with the cell-cell adhesion mechanisms and by increasing MMP activity induced extracellular matrix degradation, facilitates the invasion of tumor cells.
Literature	GENERIF_SUMMARY	TGF-beta1 induced epithelial-mesenchymal transition in non-small cell lung cancer cells by upregulating miR-9 and downregulating miR-9's target, E-cadherin.
Literature	GENERIF_SUMMARY	The presence of E-cadherin decreases cortical contractility during mitosis through a signaling cascade leading to multipolar divisions, and its knockout promotes clustering and survival of cells with multiple centrosomes.
Literature	GENERIF_SUMMARY	The - 73A > C CDH1 promoter variation may lead to differences in the overall survival of sporadic gastric carcinoma patients and allele-specific repressions of CDH1.
Literature	GENERIF_SUMMARY	CDH1 methylation may play a role in the initiation and progression of salivary carcinoma ex pleomorphic adenoma
Literature	GENERIF_SUMMARY	Results show that E-cadherin expression levels were negatively regulated by 90K via ubiquitination-mediated proteasomal degradation in a cell density-dependent manner.
Literature	GENERIF_SUMMARY	Studies categorize cadherin 1 (CDH1) variants, as neutral or deleterious.
Literature	GENERIF_SUMMARY	CDH1 promoter methylation may be correlated with cervical cancer carcinogenesis, especially for Caucasians. It was associated with histological subtypes
Literature	GENERIF_SUMMARY	Twist, E-cadherin, and N-cadherin protein were differently expressed in endometrioid adenocarcinoma tissues and in normal endometrium which indicates their potential function for endometrioid adenocarcinoma development.
Literature	GENERIF_SUMMARY	The aim of our study was to analyze the immunohistochemical expression of beta-catenin, E-cadherin and Snail, depending on clinico-morphological aspects of the laryngeal squamous cell carcinomas. Results revealed variable E-cadherin, beta-catenin and Snail expression, depending on differentiation degree and tumor stage.
Literature	GENERIF_SUMMARY	We consider that the presence of long-standing diabetes mellitus has a negative impact on the prognosis of pancreatic ductal carcinoma patients which may be relevant to a high frequency of promoter methylation of CDH1
Literature	GENERIF_SUMMARY	Overall, hypoxia-induced activation of Twist/miR-214/E-cadherin axis is involved in the EMT of TECs, and anti-

Main Category	Category	Terms
		miR-214 may be an attractive strategy to ameliorate the progression of renal fibrosis.
Literature	GENERIF_SUMMARY	The CDH1 -160C/A polymorphism was significantly associated with CRC risk.
Literature	GENERIF_SUMMARY	The aim of this study was to improve knowledge of the HER2 and the E-cadherin alterations in the context of gastric cancer to characterize subtypes of patients that could better benefit from targeted therapy.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with Hepatocellular Carcinoma.
Literature	GENERIF_SUMMARY	found that E-cadherin, N-cadherin and fibronetin are involved in CHD4-mediated epithelial-mesenchymal transition
Literature	GENERIF_SUMMARY	Upregulated miR-373 levels and simultaneously downregulated levels of CD44 and E-cadherin were noted in this study.
Literature	GENERIF_SUMMARY	the dysregulation of TET2/E-cadherin/beta-catenin regulatory loop is a critical oncogenic event in HCC progression
Literature	GENERIF_SUMMARY	We illustrate the approach using immunohistochemical measurements of the epithelial-mesenchymal transition marker E-cadherin in a set of colorectal primary tumors from a population-based prospective cohort in North Carolina
Literature	GENERIF_SUMMARY	Activation of estrogen receptor beta (ERbeta) regulates the expression of N-cadherin, E-cadherin and beta-catenin in androgen-independent prostate cancer cells.
Literature	GENERIF_SUMMARY	CDH1 plays an essential role in epithelial cell adherence. Causative of blepharocheilodontic syndrome CDH1 mutations impair the cell adhesion function of the cadherin-catenin complex in a dominant-negative manner.
Literature	GENERIF_SUMMARY	High UTX expression is independently associated with a better prognosis in patients with esophageal squamous cell carcinoma (ESCC) and downregulation of UTX increases ESCC cell growth and decreases E-cadherin expression. Our results suggest that UTX may be a novel therapeutic target for patients with ESCC.
Literature	GENERIF_SUMMARY	Upon identification of a homogenous downregulated profile for miR-205-5p in colon adenocarcinoma patients, functional studies demonstrated that experimental upregulation of this sequence is able to significantly raise the levels of E-cadherin through direct inhibition of ZEB1.
Literature	GENERIF_SUMMARY	E-cadherin expression is not significantly linked to metastatic disease in pancreatic ductal adenocarcinoma.
Literature	GENERIF_SUMMARY	In the former, p53 binds to the CDH1 (encoding E-cadherin) locus to antagonize EZH2-mediated H3K27

Main Category	Category	Terms
		trimethylation (H3K27me3) to maintain high levels of acetylation of H3K27 (H3K27ac).
Literature	GENERIF_SUMMARY	SIRT1 induces the epithelial-mesenchymal transition (EMT) by accelerating E-cadherin degradation via autophagy and facilitates melanoma metastasis.
Literature	GENERIF_SUMMARY	Plasma sE-cadherin levels and sE-cadherin/sVE-cadherin ratios are potential biomarkers for COPD.
Literature	GENERIF_SUMMARY	Together, these data suggest that the S18-2 protein induces epithelial to mesenchymal cell transition through the TWIST2/E-cadherin signalling and, consequently, CXCR4-mediated migration of prostate cancer cells.
Literature	GENERIF_SUMMARY	Investigated twist family bHLH transcription factor 1 (Twist), snail family transcriptional repressor 1 (Snail) and E-cadherin in prognosis, recurrence-free survival and overall survival in completely resected N0 NSCLC patients. Found high expression of Twist and Snail and low expression of E-cadherin were associated with unfavourable prognosis.
Literature	GENERIF_SUMMARY	UHRF1 controls both CDH1 and antisense-directed non coding RNA RCDH1-AS by directly binding this bidirectional promoter region.
Literature	GENERIF_SUMMARY	Smad4 could be considered as a central component of EMT transition in human colorectal cancer that combines with transcriptional factors to reduce E-cadherin and alter the expression of the epithelial phenotype.
Literature	GENERIF_SUMMARY	Results showed that decreased expression level of E-cadherin was strongly associated with malignant transformation of the cervical epithelium and the histological progression of cervical squamous cell carcinoma.
Literature	GENERIF_SUMMARY	EGFR-mediated kinase cascade controls the force-dependent recruitment of vinculin to stressed E-cadherin complexes - a key early signature of cadherin-based mechanotransduction.
Literature	GENERIF_SUMMARY	Both at the primary site and metastatic lymph nodes of invasive breast cancer samples, E-cadherin-positive tumors have worse disease-free and overall survival. E-cadherin and vimentin proteins are colocalized within the same tumor cells in a human breast cancer specimen.
Literature	GENERIF_SUMMARY	There were significant associations between CDH1 promoter hypermethylation and Oral Cancer risk.
Literature	GENERIF_SUMMARY	E-cadherin expression was preserved in 10 (21.28%) of the 47 NSCLCs immunostained with anti-E-cadherin antibody and reduced/absent in 37 of the 47 (78.72%) NSCLCs studied. E-cadherin plays a major role in the intercellular adhesion.
Literature	GENERIF_SUMMARY	Somatic variants in the CDH1 gene are associated with

Main Category	Category	Terms
		Diffuse Gastric Cancer.
Literature	GENERIF_SUMMARY	detection of Ezrin and E-cadherin expression in cervical smears, could be a potential prognostic marker for identifying cervical lesions with high-risk of progression to invasive cervical cancer, and may help on the selection of an appropriate therapy or avoid unnecessary treatment
Literature	GENERIF_SUMMARY	Study results demonstrate that germline CDH1 mutations are a significant contributor to the high frequency of diffuse gastric cancer in New Zealand Maori.
Literature	GENERIF_SUMMARY	Lobular breast cancer (LBC) is a component of the Hereditary Diffuse Gastric Cancer (HDGC) syndrome as a sporadic or familial setting. and novel E-cadherin germline mutations have recently been detected in women affected by LBCs without history for HDGC
Literature	GENERIF_SUMMARY	Up-regulation of H19 in bladder cancer tissues is correlated with clinical stage or metastasis of cancer. By cell transfection to suppress H19 expression in bladder cancer cells, E-cadherin expression is up-regulated, thus weakening metastatic potency of cancer cells.
Literature	GENERIF_SUMMARY	High CDH1 expression is associated with the Pathogenesis of Adamantinomatous Craniopharyngiomas.
Literature	GENERIF_SUMMARY	Using single-molecule localization microscopy, we show that pAJs in these cells reach more than 1 mum in length and consist of several cadherin clusters with crystal-like density interspersed within sparser cadherin regions. Notably, extrajunctional cadherin appears to be monomeric, and its density is almost four orders of magnitude less than observed in the pAJ regions.
Literature	GENERIF_SUMMARY	E-cadherin low expression is associated with lobular histology, tumor characteristics and menopausal hormone use, with no evidence of an association with breast cancer specific survival. These data support loss of E-cadherin expression as an important marker of tumor subtypes.
Literature	GENERIF_SUMMARY	High glucose enhances the formation of EZH2/Snail/HDAC1 complex in the nucleus, which in turn causes E-cadherin repression.
Literature	GENERIF_SUMMARY	Reciprocal loop of hypoxia-inducible factor-1alpha and metastasis-associated protein 2 contributes to the progression of pancreatic adenocarcinoma by suppressing E-cadherin transcription and promoting epithelial mesenchymal transformation.
Literature	GENERIF_SUMMARY	E-cadherin and N-cadherin staining scores could be merged into a cadherin (CDH) logistic index, which showed better discrimination than E-cadherin or N-cadherin alone.
Literature	GENERIF_SUMMARY	these results indicate that increased alpha-actinin-1

Main Category	Category	Terms
		expression destabilizes E-cadherin-based adhesions, which is likely to promote the migratory potential of breast cancer cells. Furthermore, our results identify a-actinin-1 as a candidate prognostic biomarker in basal-like breast cancer.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with metastasis in breast cancer.
Literature	GENERIF_SUMMARY	the HDAC inhibitors augmented both Ecadherin and vimentin expression and their effects varied in different cholangiocarcinoma cell lines. Therefore, the clinical use of HDAC inhibitors in biliary cancer should be considered cautiously
Literature	GENERIF_SUMMARY	CDH1 germline missense variant segregates with gastric cancer in three unrelated Spanish families.
Literature	GENERIF_SUMMARY	Six2 is negatively correlated with good prognosis and decreases 5-FU sensitivity via suppressing E-cadherin expression in HCC cells.
Literature	GENERIF_SUMMARY	alpha-catenin represents a bona fide tumor suppressor for the development of lobular-type breast cancer and as such provides an alternative event to E-cadherin inactivation, adherens junction dysfunction, and subsequent constitutive actomyosin contraction.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with Multicellular Spheroids in Breast Cancer during Simulated Microgravity.
Literature	GENERIF_SUMMARY	pathogenic variants are described in four genes encoding components of the p120-catenin complex (CTNND1, PLEKHA7, PLEKHA5) and an epithelial splicing regulator (ESRP2), in addition to the known Cleft lip/Palate-associated gene, CDH1, which encodes E-cadherin.
Literature	GENERIF_SUMMARY	E-Cadherin and epithelial syndecan-1 were more highly expressed in intraluminal/luminal unicystic ameloblastoma than in mural unicystic ameloblastoma and solid/multicystic ameloblastoma, whereas the stromal expression of syndecan-1 was higher in mural unicystic ameloblastoma and solid/multicystic ameloblastoma.
Literature	GENERIF_SUMMARY	Overexpression of KLF6-SV1 is associated with young patients, and loss of E-cadherin suggests that this variant correlated with the aggressiveness of nasopharyngeal carcinoma.
Literature	GENERIF_SUMMARY	restoring the expression of MIST1 reverses the EMT and reduces the tumorigenicity of pancreatic cancer cells partly via the Snail/E-cadherin pathway
Literature	GENERIF_SUMMARY	miR-219-5p promotes tumor growth and metastasis of HCC by regulating CDH1 and can serve as a prognostic marker for HCC patients.
Literature	GENERIF_SUMMARY	High CDH1 expression is associated with invasive breast carcinoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Soluble E-cadherin (sE-cad) (an 80-kDa soluble form), which is highly expressed in the malignant ascites of ovarian cancer patients, is a potent inducer of angiogenesis. In addition to ectodomain shedding, we provide further evidence that sE-cad is abundantly released in the form of exosomes.
Literature	GENERIF_SUMMARY	The disassociation of the beta-catenin/E-cadherin complex in the osteoblast membrane under stretch loading and the subsequent translocation of beta-catenin into the nucleus may be an intrinsic mechanical signal transduction mechanism.
Literature	GENERIF_SUMMARY	CHIP depletion inhibited cell growth, migration and invasion potential of colorectal cancer cells, accompanied by downregulation of MAPK and AKT signaling activities and upregulation of E-cadherin.
Literature	GENERIF_SUMMARY	NEDD9, E-cadherin and gamma-catenin proteins have roles in pancreatic ductal adenocarcinoma
Literature	GENERIF_SUMMARY	the definition of a specific protocol for CDH1 genetic screening and ongoing coordinated management of patients with Hereditary lobular breast cancer is crucial for the effective surveillance and early detection of lobular breast cancer.
Literature	GENERIF_SUMMARY	we present a diverse set of data indicating that the loss of E-cadherin enhances IGF1R pathway activity and sensitivity to IGF1R/InsR therapy.
Literature	GENERIF_SUMMARY	E-cadherin methylation is strongly associated with the incidence of nasopharyngeal cancer (review and meta-analysis).
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with pancreatic cancer.
Literature	GENERIF_SUMMARY	LncRNA RP11-789C1.1 inhibited EMT in GC through the RP11-789C1.1/miR-5003/E-cadherin axis, which could be a promising therapeutic target for Gastric Cancer.
Literature	GENERIF_SUMMARY	The authors demonstrate that Ecad interacts with isoform 2 of Dsg via a conserved Leu-175 on the Ecad cis binding interface and facilitates desmosome assembly.
Literature	GENERIF_SUMMARY	Low E-cadherin immunoexpression among pT3b prostate carcinoma patients indicates those with poorer survival and might benefit from specific therapy.
Literature	GENERIF_SUMMARY	Aberrant expression of E-cad may be a promising prognostic signature for HNSCC, especially when it is used with other prognostic markers.
Literature	GENERIF_SUMMARY	CDH1 variants are significantly reverse associated with mutation of EGFR tyrosine kinase;CDH1 variants might contribute to pathological development in lung adenocarcinoma
Literature	GENERIF_SUMMARY	When ANXA5 expression increased, cell proliferation was inhibited by regulating the expression of bcl-2 and

Main Category	Category	Terms
		bax while cell metastasis was suppressed by regulating E-cadherin and MMP-9 expression.
Literature	GENERIF_SUMMARY	study provides evidence for genetic polymorphisms of the adherent junction component cadherin gene and the association of its haplotypes with leukoaraiosis
Literature	GENERIF_SUMMARY	Silent gonadotroph tumours presented high immunoreactive score (IRS) for N-cadherin and low IRS for the extracellular domain of E-cadherin. A substantial proportion of the tumours presented nuclear staining for the intracellular domain of E-cadherin, accompanied by a reduced membranous expression of the intracellular domain of E-cadherin.
Literature	GENERIF_SUMMARY	Study constructed the negative correlation between miR-217 and CDH1 level in GC patients and cells; unveiled part of the miR-217 function during the pathogenesis of GC.
Literature	GENERIF_SUMMARY	Findings provide evidence that p.L13_L15del variant affects a conserved region of the CDH1 signal peptide core which is essential for E-cadherin synthesis and delivery. Failure in this checkpoint leads to loss of protein expression and function, and ultimately to Hereditary Diffuse Gastric Cancer.
Literature	GENERIF_SUMMARY	hnRNP H/F are important for maintenance and differentiation of embryonic stem cells and that this at least in part reflects a switch in TCF3 alternative splicing that leads to repression of CDH1/E-cadherin.
Literature	GENERIF_SUMMARY	These findings suggested that PHF8 played an oncogenic role in facilitating FIP200-dependent autophagic degradation of E-cadherin, EMT and metastasis in hepatocellular carcinoma (HCC). PHF8 might be a promising target for prevention, treatment and prognostic prediction of HCC.
Literature	GENERIF_SUMMARY	In the present study, we demonstrated that miR711mediated downregulation of CD44 expression inhibited EMT of gastric cancer cells in vitro and in vivo by downregulating vimentin protein expression and upregulating Ecadherin protein expression through transfection, qRTPCR and western blotting
Literature	GENERIF_SUMMARY	Results found that NRBE3 down-regulates E-cadherin expression by suppressing its transcription which expression is negatively correlated with NRBE3 expression in breast cancer tissues and cell lines.
Literature	GENERIF_SUMMARY	E-cadherin expression on multiple myeloma cells activates tumor-promoting properties in plasmacytoid DCs.
Literature	GENERIF_SUMMARY	The progression of the endometrioid carcinoma may occur in the setting of various molecular changes, in particular, with decreased expression of E-cadherin and b-catenin and high expression of vimentin, or in the absence of vimentin, utilizing other mechanisms of regulation of proliferative and metastatic potential.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	It is a tumor suppressor gene and can be used as a biomarker to assess and evaluate the progression and prognosis of oral dysplastic lesions and oral squamous cell carcinoma.
Literature	GENERIF_SUMMARY	We examined effects of induced mosaic knockdown of molecular regulators of cortical tension (ROCK1) and cell-cell adhesion (CDH1) with CRISPR interference. Mosaic knockdown of ROCK1 or CDH1 resulted in differential patterning within hiPSC colonies due to cellular self-organization, while retaining an epithelial pluripotent phenotype
Literature	GENERIF_SUMMARY	Review of the literature revealed a significant enrichment of CDH1 mutations within the extracellular cadherin repeat domains in cleft lip/palate/hereditary diffuse gastric cancer (CLP/HDGC) families in comparison to CDH1 mutations associated with HDGC only.
Literature	GENERIF_SUMMARY	The frequency of E-cadherin promoter hypermethylation in CT is significantly higher than in AC, indicating that promoter hypermethylation of E-cadherin plays an important role in non-small cell lung cancer carcinogenesis.
Literature	GENERIF_SUMMARY	Findings reveal a novel mechanism of tumor progression through posttranslational modification of E-cadherin.
Literature	GENERIF_SUMMARY	EZH2-mediated epigenetic suppression of EphB3 inhibits gastric cancer proliferation and metastasis by affecting E-cadherin and vimentin expression.
Literature	GENERIF_SUMMARY	The analysis of chemosensitivity and E-cadherin expression in colorectal cell lines showed that the absence of E-cadherin in HCT116 cells correlated with their higher chemosensitivity, whereas HT29 cells with marked membrane E-cadherin expression were more resistant to all drugs tested
Literature	GENERIF_SUMMARY	decreased in Fallopian tube epithelium in tubal pregnancy
Literature	GENERIF_SUMMARY	Study identifies an epigenetic regulation of Ecadherin expression by the methylation and acetylation of H3K27 in gastric neoplasm.
Literature	GENERIF_SUMMARY	Hereditary diffuse gastric cancer (HDGC) is a recently reported hereditary cancer syndrome and represents 1-3% of gastric adenocarcinoma cases. The CDH1 gene codes for E-cadherin which is located on the short arm of chromosome 16. CDH1 mutation induces cell disruption, facilitating infiltration into adjacent tissues.
Literature	GENERIF_SUMMARY	the E-cadherin and beta-catenin immunoexpressions in different variants of papillary thyroid carcinoma were assessed.
Literature	GENERIF_SUMMARY	Results found that E-CAD expression is epigenetically suppressed by SNAIL2 during the metastasis of colorectal cancer.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	CDH1 rs5030625 GAGA genotype is associated with Familial Breast Cancer risk.
Literature	GENERIF_SUMMARY	Calreticulin regulates syncytialization by ensuring appropriate control of both the synthesis and transportation of E-cadherin.
Literature	GENERIF_SUMMARY	Helicobacter pylori plays an important role in the development and progression of gastric carcinoma through silencing of CDH1, RUNX3, p21WAF and p27 expression
Literature	GENERIF_SUMMARY	results of the present study suggest the potential involvement of Twist and E-cadherin in the modulation of events related to worse prognoses in lip squamous cell carcinoma
Literature	GENERIF_SUMMARY	Ecadherin expression levels were regulated by the manifold, and UPF1, a potential tumor suppressor, may promote the EMT process in Huh7 HCC cells. The findings of the present study suggested that UPF1 expression levels affected the EMT process by targeting Ecadherin, Ncadherin, Vimentin and Twist.
Literature	GENERIF_SUMMARY	While no pathogenic somatic mutations were found that could explain the diffuse histology of gastric cancer in DGC and MGC, methylation was the most common somatic inactivation event of the CDH1 gene, and LOH was rare. The previously unreported c.-137C > A variant modify the CDH1 gene expression since it alters the binding sites for transcription factors.
Literature	GENERIF_SUMMARY	A comprehensive overview of the diseases most frequently associated with CDH1 germline alterations, as well as the type and functional consequences of the mutations found in each setting (review)
Literature	GENERIF_SUMMARY	Type and location of CDH1 germline variants may help to identify families at increased risk for concomitant cancers that might benefit from individualised surveillance and intervention strategies.
Literature	GENERIF_SUMMARY	We conclude that E-cadherin loss in invasive ductal carcinoma is likely a consequence of genomic instability occurring during carcinogenesis.
Literature	GENERIF_SUMMARY	The low endometrial expression of beta-catenin, E-cadherin and K-cadherin were associated to fertility-related problems, such as primary intertility and recurrent pregnancy loss.
Literature	GENERIF_SUMMARY	CBX7 and PRMT1 contribute to regulate E-cadherin expression through several mechanisms.
Literature	GENERIF_SUMMARY	the ability of an inflammatory protease, neutrophil elastase, to process the adherens junction protein E-cadherin to generate short peptide fragments with effects on epithelial function, is reported.
Literature	GENERIF_SUMMARY	The Immunohistochemical Pattern of Epithelial-Mesenchymal Transition Markers In Endometrial Carcinoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	IL-6 secreted from varicella zoster virus-infected primary human perineurial cells facilitated changes in E- and N-cadherin expression and cell migration, reminiscent of an epithelial-to-mesenchymal cell transition, potentially contributing to loss of perineurial cell barrier integrity and viral spread.
Literature	GENERIF_SUMMARY	Our results indicate that the immunohistochemical detection of E-cadherin might be useful in categorizing acromegaly patients based on the response to Somatostatin analogues (SSAs)
Literature	GENERIF_SUMMARY	BMP4 increases E-cadherin and claudin expression in glioblastoma through activation of SMAD signaling, thereby suppressing tumor cell invasion
Literature	GENERIF_SUMMARY	E-cadherin and beta-Catenin may have roles in initial lymph node metastasis of renal cell carcinoma
Literature	GENERIF_SUMMARY	E-cadherin can restrict aggressive behavior in sarcomas.
Literature	GENERIF_SUMMARY	CDH1 -160C/A and Exo1 K589E polymorphisms are associated with increased susceptibility to prostate cancer in Bangladeshi population.
Literature	GENERIF_SUMMARY	Study results indicate that CDH1-160 C>A polymorphism does not contribute to the genetic susceptibility of colorectal cancer (CRC) and the polymorphism may not be a direct effect on the progression of the disease in Turkish CRC patients.
Literature	GENERIF_SUMMARY	Study confirmed the association between CDH1 C/T polymorphisms and the risk of developing vitiligo. A relationship was found between CDH1 CC genotype and late age of onset, clinical type, the absence of autoimmune comorbidities and there family history. However, CDH1 TT genotype was found to be increased significantly in patients with autoimmune comorbidities.
Literature	GENERIF_SUMMARY	No unequivocal pathogenic germline CDH1 variants were identified in Brazilian early age onset gastric cancer patients
Literature	GENERIF_SUMMARY	PRPS2 promotes the migration and invasion of cultured colorectal cancer cells through regulating the activity of MMP-9 and the expression of E-cadherin.
Literature	GENERIF_SUMMARY	The expression of E-cadherin in breast cancer positively correlated with FOXA1 (P = 0.028), whereas negative association was seen between the expression of Twist1 and FOXA1 (P = 0.016).
Literature	GENERIF_SUMMARY	Reduced membranous expression of E-cadherin and increased cytoplasmic E-cadherin expression predict poor survival in metastatic colorectal cancer.
Literature	GENERIF_SUMMARY	MiR-9 induces EMT by targeting CDH1, and activates MAPK/ERK and Wnt/beta-catenin signal pathways, thus promoting human synovial sarcoma tumorigenesis.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	circFNC3B increased CD44 expression, which was associated with cell adhesion, via the formation of a ternary complex of circFNDC3B-IGF2BP3-CD44 mRNA.
Literature	GENERIF_SUMMARY	Study observed increased protein levels of E-cadherin in extracellular vesicles in breast cancer; furthermore, Tspan8 and p120-catenin were co-immunoprecipitated, indicating that they may interact with each other.
Literature	GENERIF_SUMMARY	Comparison of CDH1 Gene Hypermethylation Status in Blood and Serum among Gastric Cancer Patients.
Literature	GENERIF_SUMMARY	LATS degradation by APC/C(Cdh1) represents a previously unappreciated and evolutionarily conserved layer of Hippo signaling regulation.
Literature	GENERIF_SUMMARY	TRERNA1 promotes cell metastasis and the invasion of HCC via the recruitment of EHMT2 and/or the EHMT2/SNAI1 complex to suppress CDH1.
Literature	GENERIF_SUMMARY	This results in elevated levels of APC/CCDH1 substrates, including CDC20. We also demonstrate that small-molecule inhibition of APC/CCDH1/CDC20 can increase mitotic abnormalities and reduce cancer stem-like cells (CSC) viability.
Literature	GENERIF_SUMMARY	LLGL1 is coexpressed with E-cadherin; loss of expression of either protein is associated with diffuse gastric cancer and peritoneal metastases
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with Epithelial-Mesenchymal Transition in Breast Cancer.
Literature	GENERIF_SUMMARY	ALDOLASE A regulates invasion of bladder cancer cells via E-cadherin-EGFR signaling.
Literature	GENERIF_SUMMARY	Multivariate analysis indicated that RUNX3 and CDH1 promoter methylation were independent prognostic factors for overall survival (P = 0.03 and 0.04; respectively), while DAPK1, RUNX3 and MT1G PM were independent prognostic factors for disease free survival.
Literature	GENERIF_SUMMARY	Serum level of the epithelial marker E-cadherin could be considered as reliable biomarker of renal injury and is associated with the degree of renal damage in type 2 diabetic patients.
Literature	GENERIF_SUMMARY	E-cadherin and p120ctn protein expression are lost in hidradenitis suppurativa lesions.
Literature	GENERIF_SUMMARY	E-cadherin-downregulated cells were morphologically like mesenchymal cells, and the number of CD133(+) - and CD44(+) -cells (CSC-like cells) increased. These cells can be used as stable models to study cancer cells and screening of antitumor therapeutics.
Literature	GENERIF_SUMMARY	MicroRNA-572 targets CDH1 to promote metastasis of Wilms' tumor.
Literature	GENERIF_SUMMARY	these data demonstrate that cell cohesion is maintained differently in meibomian gland cells and

Main Category	Category	Terms
		indicate that Ecad is important for meibomian gland function.
Literature	GENERIF_SUMMARY	Studied E-cadherin - petidomimetic interactions using nuclear magnetic resonance spectroscopy and molecular docking simulation.
Literature	GENERIF_SUMMARY	Kaiso plays a role in regulating intestinal epithelial cell integrity and function
Literature	GENERIF_SUMMARY	these data indicate that E-cadherin is a target gene of HOXC8 and that the loss of E-cadherin promotes the growth and migration of NSCLC.
Literature	GENERIF_SUMMARY	Our results indicate that tight junctions are compromised in benign prostatic hyperplasia and loss of E-cadherin is potentially an important underlying mechanism
Literature	GENERIF_SUMMARY	Optimal Timing of Total Gastrectomy to Prevent Diffuse Gastric Cancer in Individuals With Pathogenic Variants in CDH1.
Literature	GENERIF_SUMMARY	HSPD1 repressed E-cadherin expression to promote cell invasion and migration for poor prognosis in oral squamous cell carcinoma.
Literature	GENERIF_SUMMARY	activin A downregulates E-cadherin expression by upregulating SLUG and SNAIL expression via SMAD2/3-SMAD4-dependent signaling.
Literature	GENERIF_SUMMARY	Investigated the role of Wnt11 in neuroendocrine (NE) differentiation, cell proliferation, and epithelial-to-mesenchymal transition in small-cell lung cancer (SCLC). Ascl1 plays an important role in regulating the Wnt signaling pathway and is one of the driver molecules of Wnt11 in SCLC. Wnt11 controlled NE differentiation, cell proliferation, and E-cadherin expression under the regulation of Ascl1 in SCLC cell lines.
Literature	GENERIF_SUMMARY	clinical criteria for CDH1 testing should be widened, including breast cancer families only, and a consideration for delayed prophylactic gastrectomy/ surveillance should be evaluated.
Literature	GENERIF_SUMMARY	SARNP, a participant in mRNA splicing and export, negatively regulates E-cadherin expression via interaction with pinin.
Literature	GENERIF_SUMMARY	Long non-coding RNA H19 is responsible for the progression of lung adenocarcinoma by mediating methylation-dependent repression of CDH1 promoter.
Literature	GENERIF_SUMMARY	Low CDH1 expression is associated with migration and invasion in renal cell carcinoma.
Literature	GENERIF_SUMMARY	silencing of OGT in HT29 cells upregulates E-cadherin (a major actor of epithelial-to-mesenchymal transition) and changes its glycosylation. On the other hand, OGT silencing perturbs biosynthesis of glycosphingolipids resulting in a decrease in gangliosides and an increase

Main Category	Category	Terms
		in globosides.
Literature	GENERIF_SUMMARY	MiR-9 accelerates epithelial-mesenchymal transition of ovarian cancer cells via inhibiting e-cadherin.
Literature	GENERIF_SUMMARY	The expression level of GATA3 in the experimental group was significantly higher than that in the control group (p
Literature	GENERIF_SUMMARY	Soluble E-cadherin is associated with oxidative stress in patients with chronic HBV infection.
Literature	GENERIF_SUMMARY	High podoplanin and low E-cadherin levels correlate with better prognosis in adenoid cystic carcinoma.
Literature	GENERIF_SUMMARY	Detailed cell fractionation, immunoprecipitation, microscopy, and immunohistochemical analyses demonstrated that CDCP1 promotes translocation of the key regulators of Wnt signaling, b-catenin, and E-cadherin, to the nucleus. Of functional importance, disruption of CDCP1 reduces nuclear localized, chromatin-associated b-catenin and nuclear localized E-cadherin, increases sequestration of these proteins in cell membranes
Literature	GENERIF_SUMMARY	APC/Cdh1 targets PECAM-1 for ubiquitination and degradation in endothelial cells.
Literature	GENERIF_SUMMARY	Results strongly indicate the link between E-cadherin (CDH1) and Intestinal-type gastric cancer (IGC).
Literature	GENERIF_SUMMARY	GADD45G Interacts with E-cadherin to Suppress the Migration and Invasion of Esophageal Squamous Cell Carcinoma.
Literature	GENERIF_SUMMARY	LINC00978 promotes the progression of hepatocellular carcinoma by regulating EZH2-mediated silencing of p21 and E-cadherin expression.
Literature	GENERIF_SUMMARY	The 107-residue intrinsically disordered N-terminal segment of beta-catenin forms a flexible ""tongue"" that is inserted into alpha-catenin and participates in the assembly of the alpha-catenin*beta-catenin*epithelial cadherin complex."
Literature	GENERIF_SUMMARY	The epithelial cadherin -160C/A polymorphism is associated with decreased risk of colorectal cancer
Literature	GENERIF_SUMMARY	High CDH1 methylation is associated with Epithelial Ovarian Cancer.
Literature	GENERIF_SUMMARY	The essential role of E-cadherin in maintaining prostatic epithelial integrity and organization was demonstrated using organoid culture of human prostate cancer cell lines, LNCaP and LAPC4.
Literature	GENERIF_SUMMARY	miR-25 mediates metastasis and epithelial-mesenchymal-transition in human esophageal squamous cell carcinoma via regulation of E-cadherin signaling.
Literature	GENERIF_SUMMARY	TBC1D2b suppresses E-cadherin internalization, thus

Main Category	Category	Terms
		hindering cancer cell invasion and metastasis.
Literature	GENERIF_SUMMARY	miR-219 regulates liver cancer stem cell expansion via E-cadherin pathway.
Literature	GENERIF_SUMMARY	Analysis of data from two clinical trials and literature meta-analysis revealed that colorectal cancer-risk loci related CDH1 single nucleotide polymorphisms rs12597188 and rs9939049 influence colorectal cancer patient survival, and rs12597188 also associated with poor response to therapy.
Literature	GENERIF_SUMMARY	Glutathione S-transferase omega 2 regulates cell growth and the expression of E-cadherin via post-transcriptional down-regulation of beta-catenin in human esophageal squamous cells.
Literature	GENERIF_SUMMARY	Regulation of cell-cell adhesion in prostate cancer cells by microRNA-96 through upregulation of E-Cadherin and EpCAM.
Literature	GENERIF_SUMMARY	S100P is a molecular determinant of E-cadherin function in gastric cancer.
Literature	GENERIF_SUMMARY	Crosstalk between miR-215 and epithelial-mesenchymal transition specific markers (E-cadherin and N-cadherin) in different stages of chronic HCV Infection.
Literature	GENERIF_SUMMARY	E-cadherin expression is regulated by thyroid hormones in the squamous cell carcinomas.
Literature	GENERIF_SUMMARY	Methylation of tumour suppressor genes RUNX3, RASSF1A and E-Cadherin in HCV-related liver cirrhosis and hepatocellular carcinoma.
Literature	GENERIF_SUMMARY	Engagement of Robo1 by Slit2 induces formation of a trimeric complex consisting of Src-Robo1-E-cadherin for E-cadherin phosphorylation and epithelial-mesenchymal transition.
Literature	GENERIF_SUMMARY	Diagnostic accuracy of E-cadherin for malignanteffusions
Literature	GENERIF_SUMMARY	The ultrastructures of adherent junctions, gap junctions, and tight junctions could be observed in mouse RPE by TEM. E-cadherin, ZO-1, and connexin 43 were colocalized and physically bound to each other. The knockdown of one of these three proteins led to downregulation of the other two proteins and compromised epithelial function.
Literature	GENERIF_SUMMARY	Study showed that in Ishikawa endometrial carcinoma cells, downregulation of PTEN was associated with the expression of the CDH1 and CDH2 genes and upregulated expression of the cell membrane glycoprotein, CD133, which are associated with epithelial-mesenchymal transition (EMT) in malignancy.
Literature	GENERIF_SUMMARY	An association between CDH1 loss of function alterations and high Gleason grade was found in prostatic adenocarcinoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Clinical relevance of somatic mutations in main driver genes detected in gastric cancer patients by next-generation DNA sequencing.
Literature	GENERIF_SUMMARY	Silencing of FOXA2 decreases E-cadherin expression and is associated with lymph node metastasis in oral cancer.
Literature	GENERIF_SUMMARY	BST2 regulates interferon gamma-dependent decrease in invasion of HTR-8/SVneo cells via STAT1 and AKT signaling pathways and expression of E-cadherin.
Literature	GENERIF_SUMMARY	Complete histopathologic examination of risk reduction gastrectomy specimens for CDH1 germline mutation
Literature	GENERIF_SUMMARY	Reduced E-cadherin expression as a prognostic factor in non-muscle-invasive bladder cancer
Literature	GENERIF_SUMMARY	Expression of E- Cadherin and Levels of Dysplasia in Oral Leukoplakia - A Prospective Cohort Study.
Literature	GENERIF_SUMMARY	CDH1 gene mutation in early-onset, colorectal signet-ring cell carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin Plays a Role in Hepatitis B Virus Entry Through Affecting Glycosylated Sodium-Taurocholate Cotransporting Polypeptide Distribution.
Literature	GENERIF_SUMMARY	Expression of E-cadherin, Ki-67, and p53 in urinary bladder cancer in relation to progression, survival, and recurrence.
Literature	GENERIF_SUMMARY	Using this method, the CD45-negative cells were characterized into three Circulating tumor cells (CTCs) subsets based on protein expression levels
Literature	GENERIF_SUMMARY	NLK interacts with 1433zeta to restore the expression of Ecadherin.
Literature	GENERIF_SUMMARY	CDH1 mutations in gastric cancers are not associated with family history.
Literature	GENERIF_SUMMARY	THE STUDY OF THE EXPRESSION OF CDH1, KI67, P53 AND HER2 IN DIFFUSE GASTRIC CARCINOMA.
Literature	GENERIF_SUMMARY	MicroRNA 345 (miR345) regulates KISS1-E-cadherin functional interaction in breast cancer brain metastases.
Literature	GENERIF_SUMMARY	CDH1 Mutation Distribution and Type Suggests Genetic Differences between the Etiology of Orofacial Clefting and Gastric Cancer.
Literature	GENERIF_SUMMARY	Hispanic/Latino Patients with Gastric Adenocarcinoma Have Distinct Molecular Profiles Including a High Rate of Germline CDH1 Variants.
Literature	GENERIF_SUMMARY	WNT1, a target of miR-34a, promotes cervical squamous cell carcinoma proliferation and invasion by induction of an E-P cadherin switch via the WNT/beta-catenin pathway.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	CDH1-related blepharocheilodontic syndrome is associated with diffuse gastric cancer risk.
Literature	GENERIF_SUMMARY	Integrin linked kinase regulates endosomal recycling of N-cadherin in melanoma cells.
Literature	GENERIF_SUMMARY	CDH1 and SNAI1 are regulated by E7 from human papillomavirus types 16 and 18.
Literature	GENERIF_SUMMARY	Twist1 induces chromosomal instability (CIN) in colorectal cancer cells.
Literature	GENERIF_SUMMARY	Aberrant Expression of E-cadherin, N-cadherin, and P-cadherin in Clear Cell Renal Cell Carcinoma
Literature	GENERIF_SUMMARY	The loss of expression of E-cadherin may be a simple biological factor to predict lymph nodes metastasis in patients with undifferentiated early gastric cancer.
Literature	GENERIF_SUMMARY	Case report
Literature	GENERIF_SUMMARY	Germline mutations of the E-cadherin gene (CDH1) in early onset gastric cancer.
Literature	GENERIF_SUMMARY	FSH Levels Are Related to E-cadherin Expression and Subcellular Location in Nonfunctioning Pituitary Tumors.
Literature	GENERIF_SUMMARY	MicroRNA-9 regulates the proliferation, migration and invasion of human glioma cells by targeting CDH1.
Literature	GENERIF_SUMMARY	Elevated soluble E-cadherin during the epithelial-mesenchymal transition process and as a diagnostic marker in colorectal cancer.
Literature	GENERIF_SUMMARY	Application of proximity ligation assays to identify potential plasma biomarkers in oral cavity cancer patients
Literature	GENERIF_SUMMARY	E-cadherin to P-cadherin switching in lobular breast cancer with tubular elements.
Literature	GENERIF_SUMMARY	E-cadherin immunohistochemical expression in invasive lobular carcinoma of the breast
Literature	GENERIF_SUMMARY	Proper E-cadherin membrane location in colon requires Dab2 and it modifies by inflammation and cancer.
Literature	GENERIF_SUMMARY	CDCA2 promotes proliferation and migration of melanoma by upregulating CCAD1.
Literature	GENERIF_SUMMARY	TRIM29 mediates lung squamous cell carcinoma cell metastasis by regulating autophagic degradation of E-cadherin.
Literature	GENERIF_SUMMARY	p190A RhoGAP induces CDH1 expression and cooperates with E-cadherin to activate LATS kinases and suppress tumor cell growth.
Literature	GENERIF_SUMMARY	Family's History Based on the CDH1 Germline Variant (c.360delG) and a Suspected Hereditary Gastric Cancer Form.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Invasive plasmacytoid urothelial carcinoma
Literature	GENERIF_SUMMARY	Impact of E-cadherin and its transcription regulators on assessing epithelial-mesenchymal transition in chronic hepatitis C virus infection.
Literature	GENERIF_SUMMARY	MK615 Suppresses Hypoxia Tolerance by Up-regulation of E-cadherin in Colorectal Cancer Cells With Mutant KRAS.
Literature	GENERIF_SUMMARY	Overexpression of miR-21-5p in colorectal cancer cells promotes self-assembly of E-cadherin-dependent multicellular tumor spheroids.
Literature	GENERIF_SUMMARY	Hereditary diffuse CDH1 positive gastric cancer. Prophylactic gastrectomy?"", trans ""Cancer gastrico difuso hereditario CDH1 positivo inverted question markgastrectomia profilactica?"
Literature	GENERIF_SUMMARY	Expression and Role of E-Cadherin, beta-Catenin, and Vimentin in Human Papillomavirus-Positive and Human Papillomavirus-Negative Oropharyngeal Squamous Cell Carcinoma.
Literature	GENERIF_SUMMARY	LncRNA UCA1 maintains the low-tumorigenic and nonmetastatic status by stabilizing E-cadherin in primary prostate cancer cells.
Literature	GENERIF_SUMMARY	Overview on new progress of hereditary diffuse gastric cancer with CDH1 variants.
Literature	GENERIF_SUMMARY	Radiation-induced H3K9 tri-methylation in E-cadherin promoter during lung EMT
Literature	GENERIF_SUMMARY	Cadherin clusters stabilized by a combination of specific and nonspecific cis-interactions.
Literature	GENERIF_SUMMARY	First report of a Mexican family with mutation in the CDH1 gene.
Literature	GENERIF_SUMMARY	Wilms' tumor 1 (WT1) promotes ovarian cancer progression by regulating E-cadherin and ERK1/2 signaling.
Literature	GENERIF_SUMMARY	SUMO E3 ligase CBX4 regulates hTERT-mediated transcription of CDH1 and promotes breast cancer cell migration and invasion.
Literature	GENERIF_SUMMARY	Potential use of serum insulin-like growth factor 1 and E-cadherin as biomarkers of colorectal cancer.
Literature	GENERIF_SUMMARY	SUMOylation of HSP27 regulates PKM2 to promote esophageal squamous cell carcinoma progression.
Literature	GENERIF_SUMMARY	Identification of Significant Genes and Pathways in Acute Pancreatitis via Bioinformatical Analysis.
Literature	GENERIF_SUMMARY	Golgi phosphoprotein-3 (GOLPH3) promote metastasis of nasopharyngeal carcinoma through regulating E-cadherin.
Literature	GENERIF_SUMMARY	Differential Expression of E-Cadherin and P-Cadherin in

Main Category	Category	Terms
		Breast Cancer Molecular Subtypes.
Literature	GENERIF_SUMMARY	Indications for Total Gastrectomy in CDH1 Mutation Carriers and Outcomes of Risk-Reducing Minimally Invasive and Open Gastrectomies.
Literature	GENERIF_SUMMARY	The circular RNA circPTK2 inhibits EMT in hepatocellular carcinoma by acting as a ceRNA and sponging miR-92a to upregulate E-cadherin.
Literature	GENERIF_SUMMARY	Prognostic value of PSMA, c-MET and E-cadherin in salivary duct carcinoma.
Literature	GENERIF_SUMMARY	Epithelial Distribution of E-Cadherin, p63, and Mitotic Figures in ApoTome Images to Determine the Oncogenic Potentiality of Oral Submucous Fibrosis.
Literature	GENERIF_SUMMARY	Genes regulating membrane-associated E-cadherin and proliferation in adenomatous polyposis coli mutant colon cancer cells
Literature	GENERIF_SUMMARY	Inactivation of homeodomain-interacting protein kinase 2 promotes oral squamous cell carcinoma metastasis through inhibition of P53-dependent E-cadherin expression.
Literature	GENERIF_SUMMARY	Decorin deficiency promotes epithelial-mesenchymal transition and colon cancer metastasis.
Literature	GENERIF_SUMMARY	ZEB1 and oncogenic Ras constitute a regulatory switch for stimulus-dependent E-cadherin downregulation.
Literature	GENERIF_SUMMARY	CD82 Suppresses ADAM17-Dependent E-Cadherin Cleavage and Cell Migration in Prostate Cancer.
Literature	GENERIF_SUMMARY	Differential expression of Oct-4, CD44, and E-cadherin in eutopic and ectopic endometrium in ovarian endometriomas and their correlations with clinicopathological variables.
Literature	GENERIF_SUMMARY	Mapping transmembrane binding partners for E-cadherin ectodomains.
Literature	GENERIF_SUMMARY	Hereditary Diffuse Gastric Cancer Syndrome and the Role of CDH1
Literature	GENERIF_SUMMARY	Comparative analysis of the AIB1 interactome in breast cancer reveals MTA2 as a repressive partner which silences E-Cadherin to promote EMT and associates with a pro-metastatic phenotype.
Literature	GENERIF_SUMMARY	""""Switch of E-Cadherin to N-Cadherin expression in different molecular subtypes of breast invasive duct carcinomas and its correlation with clinicopathological features"""".
Literature	GENERIF_SUMMARY	Downregulation of E-cadherin in pluripotent stem cells triggers partial EMT.
Literature	GENERIF_SUMMARY	Predictive and prognostic significance of tumour subtype, SSTR1-5 and e-cadherin expression in a well-defined cohort of patients with acromegaly.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Tumor-suppressor function of Beclin 1 in breast cancer cells requires E-cadherin.
Literature	GENERIF_SUMMARY	Chloride Intracellular Channel Protein 1 (CLIC1), E-cadherin and P-cadherin Define Distinct Subclasses of HER2, Luminal B and Triple-negative Breast Cancer.
Literature	GENERIF_SUMMARY	PLEKHA7, an Apical Adherens Junction Protein, Suppresses Inflammatory Breast Cancer in the Context of High E-Cadherin and p120-Catenin Expression.
Literature	GENERIF_SUMMARY	PET-CT radiomics by integrating primary tumor and peritumoral areas predicts E-cadherin expression and correlates with pelvic lymph node metastasis in early-stage cervical cancer.
Literature	GENERIF_SUMMARY	Salivary Duct Carcinoma With Rhabdoid Features-No or Aberrant Expression of E-cadherin and Genetic Changes in CDH1
Literature	GENERIF_SUMMARY	LncRNA SNHG1 and RNA binding protein hnRNPL form a complex and coregulate CDH1 to boost the growth and metastasis of prostate cancer.
Literature	GENERIF_SUMMARY	Resolving pathogenicity classification for the CDH1 c.[715G>A] (p.Gly239Arg) Variant.
Literature	GENERIF_SUMMARY	Variation rs9929218 and risk of the colorectal Cancer and adenomas
Literature	GENERIF_SUMMARY	TMEM52B suppression promotes cancer cell survival and invasion through modulating E-cadherin stability and EGFR activity.
Literature	GENERIF_SUMMARY	Long noncoding RNA LOC284454 promotes hepatocellular carcinoma cell invasion and migration by inhibiting Ecadherin expression.
Literature	GENERIF_SUMMARY	CDH1 variants leading to gastric cancer risk management decision-making experiences in emerging adults
Literature	GENERIF_SUMMARY	Loss of lncRNA SNHG8 promotes epithelial-mesenchymal transition by destabilizing CDH1 mRNA.
Literature	GENERIF_SUMMARY	Type 2 human papillomavirus E7 attenuates E-cadherin expression in human keratinocytes.
Literature	GENERIF_SUMMARY	Role of SPARC in the epithelial-mesenchymal transition induced by PTHrP in human colon cancer cells.
Literature	GENERIF_SUMMARY	Assessment of mismatch repair deficiency, CDX2, beta-catenin and E-cadherin expression in colon cancer
Literature	GENERIF_SUMMARY	E-cadherin Gene (CDH1) Expression in Low Grade Astrocytoma.
Literature	GENERIF_SUMMARY	CDH1 pathogenic variants and cancer risk in an unselected patient population.
Literature	GENERIF_SUMMARY	CDH1 Gene rs1801552 C/T Polymorphism Increases Susceptibility to Esophageal Squamous Cell Carcinoma

Main Category	Category	Terms
		but Not to Gastric Cardiac Adenocarcinoma.
Literature	GENERIF_SUMMARY	Rare germline variants in the E-cadherin gene CDH1 are associated with the risk of brain tumors of neuroepithelial and epithelial origin.
Literature	GENERIF_SUMMARY	Modulation of the EMT/MET Process by E-Cadherin in Airway Epithelia Stress Injury.
Literature	GENERIF_SUMMARY	Indolent feature of Helicobacter pylori-uninfected intramucosal signet ring cell carcinomas with CDH1 mutations.
Literature	GENERIF_SUMMARY	Phenotypic Heterogeneity and Metastasis of Breast Cancer Cells.
Literature	GENERIF_SUMMARY	E-cadherin and N-cadherin Immunohistochemical Expression in Proliferating Urothelial Lesions
Literature	GENERIF_SUMMARY	MicroRNA4545p promotes breast cancer progression by inducing epithelialmesenchymal transition via targeting the FoxJ2/Ecadherin axis.
Literature	GENERIF_SUMMARY	CDH1 germline mutations in families with hereditary lobular breast cancer.
Literature	GENERIF_SUMMARY	Oral Squamous Cell Carcinoma Bone Invasion
Literature	GENERIF_SUMMARY	A Keratin 7 and E-Cadherin Signature Is Highly Predictive of Tubo-Ovarian High-Grade Serous Carcinoma Prognosis.
Literature	GENERIF_SUMMARY	E-Cadherin Expression and Blunted Interferon Response in Blastic Plasmacytoid Dendritic Cell Neoplasm.
Literature	GENERIF_SUMMARY	E-cadherin (CDH1) gene -160C/A polymorphism and the risk of colorectal cancer
Literature	GENERIF_SUMMARY	Knockout of FosB gene changes drug sensitivity and invasion activity via the regulation of Bcl-2, E-cadherin, beta-catenin, and vimentin expression.
Literature	GENERIF_SUMMARY	Mixed ductal-lobular carcinoma
Literature	GENERIF_SUMMARY	Inhibiting the redox function of APE1 suppresses cervical cancer metastasis via disengagement of ZEB1 from E-cadherin in EMT.
Literature	GENERIF_SUMMARY	hsa-miR-199b-3p Prevents the Epithelial-Mesenchymal Transition and Dysfunction of the Renal Tubule by Regulating E-cadherin through Targeting KDM6A in Diabetic Nephropathy.
Literature	GENERIF_SUMMARY	Potential value of circulatory microRNA10b gene expression and its target E-cadherin as a prognostic and metastatic prediction marker for breast cancer.
Literature	GENERIF_SUMMARY	Association of E-cadherin & vimentin expression with clinicopathological parameters in lingual squamous cell carcinomas & their role in incomplete epithelial mesenchymal transition.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Tumorigenic mesenchymal clusters are less sensitive to moderate osmotic stresses due to low amounts of junctional E-cadherin.
Literature	GENERIF_SUMMARY	ARID1A regulates E-cadherin expression in colorectal cancer cells
Literature	GENERIF_SUMMARY	Differential impact of clinicopathological risk factors within the 2 largest ProMisE molecular subgroups of endometrial carcinoma.
Literature	GENERIF_SUMMARY	E-Cadherin and Angiopoietin-2 as Potential Biomarkers for Colorectal Cancer With Peritoneal Carcinomatosis.
Literature	GENERIF_SUMMARY	Collagen deposition correlates with loss of E-cadherin and increased p63 expression in dysplastic conditions of oral submucous fibrosis.
Literature	GENERIF_SUMMARY	Integrin beta1 orchestrates the abnormal cell-matrix attachment and invasive behaviour of E-cadherin dysfunctional cells.
Literature	GENERIF_SUMMARY	The role of E-Cadherin expression in primary site of breast cancer.
Literature	GENERIF_SUMMARY	CDH1 germline mutations in a Chinese cohort with hereditary diffuse gastric cancer.
Literature	GENERIF_SUMMARY	HSPB8 promoted intrahepatic cholangiocarcinoma progression by enhancing epithelial-mesenchymal transition and autophagy.
Literature	GENERIF_SUMMARY	Association Between Hereditary Lobular Breast Cancer Due to CDH1 Variants and Gastric Cancer Risk.
Literature	GENERIF_SUMMARY	Expression of N-Terminal-Deficient E-Cadherin Protein in Invasive Lobular Carcinoma of the Breast.
Literature	GENERIF_SUMMARY	Prognostic Significance of Wnt1, Wnt2, E-Cadherin, and beta-catenin Expression in Operable Non-small Cell Lung Cancer.
Literature	GENERIF_SUMMARY	C-Mannosylated tryptophan-containing WSPW peptide binds to actinin-4 and alters E-cadherin subcellular localization in lung epithelial-like A549 cells.
Literature	GENERIF_SUMMARY	Epigenetic Regulation of CDH1 Is Altered after HOXB7-Silencing in MDA-MB-468 Triple-Negative Breast Cancer Cells.
Literature	GENERIF_SUMMARY	MYSM1 inhibits human colorectal cancer tumorigenesis by activating miR-200 family members/ CDH1 and blocking PI3K/AKT signaling.
Literature	GENERIF_SUMMARY	Modulation of E-Cadherin and N-Cadherin by ovarian steroids and embryonic stimuli.
Literature	GENERIF_SUMMARY	Prognostic value of E-Cadherin and its tumor suppressor role in Saudi women with advanced epithelial ovarian cancer.
Literature	GENERIF_SUMMARY	Transcriptional repression of E-cadherin in nickel-

Main Category	Category	Terms
		exposed lung epithelial cells mediated by loss of Sp1 binding at the promoter.
Literature	GENERIF_SUMMARY	Circular RNA circHERC4 as a novel oncogenic driver to promote tumor metastasis via the miR-556-5p/CTBP2/E-cadherin axis in colorectal cancer.
Literature	GENERIF_SUMMARY	Upregulation of homeobox D10 expression suppresses invasion and migration of clear cell renal cell carcinoma through targeting of E-cadherin.
Literature	GENERIF_SUMMARY	Air pollution-regulated E-cadherin mediates contact inhibition of proliferation via the hippo signaling pathways in emphysema.
Literature	GENERIF_SUMMARY	Crosstalk between E-Cadherin/beta-Catenin and NF-kappaB Signaling Pathways
Literature	GENERIF_SUMMARY	E-cadherin is a robust prognostic biomarker in colorectal cancer and low expression is associated with sensitivity to inhibitors of topoisomerase, aurora, and HSP90 in preclinical models.
Literature	GENERIF_SUMMARY	Extracellular domains of E-cadherin determine key mechanical phenotypes of an epithelium through cell- and non-cell-autonomous outside-in signaling.
Literature	GENERIF_SUMMARY	CDH1 germline variants are enriched in patients with colorectal cancer, gastric cancer, and breast cancer.
Literature	GENERIF_SUMMARY	CDH1 germline mutations in healthy individuals from families with the hereditary diffuse gastric cancer syndrome.
Literature	GENERIF_SUMMARY	Emerging Adults Carrying a CDH1 Pathogenic or Likely Pathogenic Variant Face Diet and Lifestyle Challenges after Total Gastrectomy.
Literature	GENERIF_SUMMARY	PKD-dependent PARP12-catalyzed mono-ADP-ribosylation of Golgin-97 is required for E-cadherin transport from Golgi to plasma membrane.
Literature	GENERIF_SUMMARY	Characterization of the first microRNA in human CDH1 that affects cell cycle and apoptosis and indicates breast cancers progression.
Literature	GENERIF_SUMMARY	TdIF1-LSD1 Axis Regulates Epithelial-Mesenchymal Transition and Metastasis via Histone Demethylation of E-Cadherin Promoter in Lung Cancer.
Literature	GENERIF_SUMMARY	E-cadherin and aquaporin 1 co-expression analysis in hepatocellular carcinoma
Literature	GENERIF_SUMMARY	HOXA10 improves endometrial receptivity by upregulating E-cadherindagger.
Literature	GENERIF_SUMMARY	Muscle Transcriptomics Shows Overexpression of Cadherin 1 in Inclusion Body Myositis.
Literature	GENERIF_SUMMARY	Expression of HER2/neu in gastric adenocarcinoma and its correlation with serum HER2/neu level and E-cadherin expression.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Pseudomonas aeruginosa Triggered Exosomal Release of ADAM10 Mediates Proteolytic Cleavage in Trans.
Literature	GENERIF_SUMMARY	E-cadherin, fibronectin and Slug immunoexpression in non-melanoma skin cancers.
Literature	GENERIF_SUMMARY	Pleiotropic cancer manifestations of germline CDH1 mutations
Literature	GENERIF_SUMMARY	TMEM139 prevents NSCLC metastasis by inhibiting lysosomal degradation of E-cadherin.
Literature	GENERIF_SUMMARY	CDH1 Germline Variants in a Tunisian Cohort with Hereditary Diffuse Gastric Carcinoma.
Literature	GENERIF_SUMMARY	Global distribution of prophylactic total gastrectomy in E-cadherin (CDH1) mutations.
Literature	GENERIF_SUMMARY	miR-888 and E-Cadherin Levels in Ductal Carcinoma Breast Cancer as Possible Cancer-Related Markers.
Literature	GENERIF_SUMMARY	Mechanical Cues, E-Cadherin Expression and Cell """"Sociality"""" Are Crucial Crossroads in Determining Pancreatic Ductal Adenocarcinoma Cells Behavior."
Literature	GENERIF_SUMMARY	Loss of E-Cadherin Expression Correlates With Ki-67 in Head and Neck Squamous Cell Carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin to N-cadherin switching in the TGF-beta1 mediated retinal pigment epithelial to mesenchymal transition.
Literature	GENERIF_SUMMARY	Dysregulation of E-cadherin in pulmonary cell damage related with COPD contributes to emphysema.
Literature	GENERIF_SUMMARY	Control of CDH1/E-Cadherin Gene Expression and Release of a Soluble Form of E-Cadherin in SARS-CoV-2 Infected Caco-2 Intestinal Cells
Literature	GENERIF_SUMMARY	Negative E-cadherin expression on bone marrow myeloma cell membranes is associated with extramedullary disease.
Literature	GENERIF_SUMMARY	Combination of AKT1 and CDH1 mutations predicts primary resistance to immunotherapy in dMMR/MSI-H gastrointestinal cancer.
Literature	GENERIF_SUMMARY	Epigenetic silencing of E-cadherin gene induced by lncRNA MALAT-1 in acute myeloid leukaemia.
Literature	GENERIF_SUMMARY	The Potential Role of CDH1 as an Oncogene Combined With Related miRNAs and Their Diagnostic Value in Breast Cancer.
Literature	GENERIF_SUMMARY	Barrier proteins and eosinophilic esophagitis in children
Literature	GENERIF_SUMMARY	KRAS mutation in intrahepatic cholangiocarcinoma
Literature	GENERIF_SUMMARY	Kaiso phosphorylation at threonine 606 leads to its accumulation in the cytoplasm, reducing its transcriptional repression of the tumour suppressor CDH1.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Detection of mRNAs of Ribosomal Protein L15 and E-Cadherin in Living Circulating Tumor Cells at Single Cell Resolution To Study Tumor Heterogeneity.
Literature	GENERIF_SUMMARY	Papillary thyroid cancer with Hashimoto's thyroiditis attenuates the tumour aggressiveness through the up-regulation of E-cadherin and TGF-beta expression.
Literature	GENERIF_SUMMARY	CDH1 mutations recurrence and global clustering in genetically tested families with hereditary diffuse gastric cancer syndrome
Literature	GENERIF_SUMMARY	Expression of HnRNP A1, ZEB1, and E-cadherin in Hepatocellular carcinoma and their impact on patients' prognosis and survival.
Literature	GENERIF_SUMMARY	p120 regulates E-cadherin expression in nasal epithelial cells in chronic rhinosinusitis.
Literature	GENERIF_SUMMARY	Familial and hereditary gastric cancer, an overview.
Literature	GENERIF_SUMMARY	Constitutive Occurrence of E
Literature	GENERIF_SUMMARY	Short and long-term outcomes of prophylactic total gastrectomy in 54 consecutive individuals with germline pathogenic mutations in the CDH1 gene.
Literature	GENERIF_SUMMARY	MICAL2 contributes to gastric cancer cell migration via Cdc42-dependent activation of E-cadherin/beta-catenin signaling pathway.
Literature	GENERIF_SUMMARY	CDH1 overexpression predicts bladder cancer from early stage and inversely correlates with immune infiltration.
Literature	GENERIF_SUMMARY	Investigation of the expression levels of CDH1, FHIT, PTEN, and TTPAL genes in colorectal tumors.
Literature	GENERIF_SUMMARY	[Association of CDH1, FANCB and APC Gene Polymorphisms with Lung Cancer Susceptibility in Chinese Population].
Literature	GENERIF_SUMMARY	Combined high NEDD9 expression and E-cadherin loss correlate with poor clinical outcome in gastric cancer.
Literature	GENERIF_SUMMARY	Human antigen R affects the migration and invasion of human lung cancer A549 cells via regulating E-cadherin suppressor Snail.
Literature	GENERIF_SUMMARY	Somatic Mutational Landscape in Mexican Patients
Literature	GENERIF_SUMMARY	Association between CDH1 methylation and esophageal cancer risk
Literature	GENERIF_SUMMARY	Immunohistochemical evaluation of E-cadherin in oral epithelial dysplasia and squamous cell carcinoma.
Literature	GENERIF_SUMMARY	[Association of genomic instability of CDH1 gene with clinicopathological characteristics of gastric cancer].
Literature	GENERIF_SUMMARY	ARID1A downregulation promotes cell proliferation and migration of colon cancer via VIM activation and CDH1

Main Category	Category	Terms
		suppression.
Literature	GENERIF_SUMMARY	Genotype-first approach to identify associations between CDH1 germline variants and cancer phenotypes
Literature	GENERIF_SUMMARY	Soluble E-cadherin promotes invasiveness of neoplastic cells in salivary gland carcinoma ex pleomorphic adenoma.
Literature	GENERIF_SUMMARY	GAS5 attenuates the malignant progression of glioma stem-like cells by promoting E-cadherin.
Literature	GENERIF_SUMMARY	Clinical and prognostic features of E-cadherin in adenocarcinoma of the esophagogastric junction patients.
Literature	GENERIF_SUMMARY	The Identification of Large Rearrangements Involving Intron 2 of the CDH1 Gene in BRCA1/2 Negative and Breast Cancer Susceptibility.
Literature	GENERIF_SUMMARY	Soluble E-cadherin
Literature	GENERIF_SUMMARY	Low expression of E-Cadherin and CDH1 variants associated with diffuse gastric cancer.
Literature	GENERIF_SUMMARY	[Influence of E-cadherin methylation on prognosis in children with acute lymphoblastic leukemia].
Literature	GENERIF_SUMMARY	Scribble and E-cadherin cooperate to control symmetric daughter cell positioning by multiple mechanisms.
Literature	GENERIF_SUMMARY	Stabilization of E-cadherin adhesions by COX-2/GSK3beta signaling is a targetable pathway in metastatic breast cancer.
Literature	GENERIF_SUMMARY	Succinylation of CTBP1 mediated by KAT2A suppresses its inhibitory activity on the transcription of CDH1 to promote the progression of prostate cancer.
Literature	GENERIF_SUMMARY	Association of ATM, CDH1 and TP53 genes polymorphisms with familial breast cancer in patients of Khyber Pakhtunkhwa, Pakistan.
Literature	GENERIF_SUMMARY	Lysine-specific Demethylase 6A Upregulates Cadherin-1 and Accelerates Gastric Cancer Growth.
Literature	GENERIF_SUMMARY	Exosomal circSPIRE1 mediates glycosylation of E-cadherin to suppress metastasis of renal cell carcinoma.
Literature	GENERIF_SUMMARY	The E3 ubiquitin ligase MIB1 suppresses breast cancer cell migration through regulating CTNND1 protein level.
Literature	GENERIF_SUMMARY	Neural crest E-cadherin loss drives cleft lip/palate by epigenetic modulation via pro-inflammatory gene-environment interaction.
Literature	GENERIF_SUMMARY	Combined loss of CDH1 and downstream regulatory sequences drive early-onset diffuse gastric cancer and increase penetrance of hereditary diffuse gastric cancer.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	E-cadherin and aquaporin-3 are downregulated in wound edges of human chronic wounds.
Literature	GENERIF_SUMMARY	Modulation of the E-cadherin in human cells infected in vitro with Coxiella burnetii.
Literature	GENERIF_SUMMARY	miR-25 promotes cell migration and invasion by inhibiting the expression of CDH1 in rectal carcinoma.
Literature	GENERIF_SUMMARY	E-cadherin expression in the tumor microenvironment of advanced epidermal growth factor receptor-mutant lung adenocarcinoma and the association with prognosis.
Literature	GENERIF_SUMMARY	Association of quantitative analysis of intratumoral reduced E-cadherin expression with lymph node metastasis and prognosis in patients with breast cancer.
Literature	GENERIF_SUMMARY	Trophoblast cell surface antigen-2 phosphorylation triggered by binding of galectin-3 drives metastasis through down-regulation of E-cadherin.
Literature	GENERIF_SUMMARY	CDH1 hypermethylation
Literature	GENERIF_SUMMARY	CDH1 overexpression sensitizes TRAIL resistant breast cancer cells towards rhTRAIL induced apoptosis.
Literature	GENERIF_SUMMARY	Expression of cytotoxic T lymphocyte-associated antigen 4, CD44, and E-cadherin in the microenvironment of breast carcinomas.
Literature	GENERIF_SUMMARY	Analysis of E-cadherin (CDH1) Gene Polymorphism and Its Association with Cervical Cancer Risk in Bangladeshi Women.
Literature	GENERIF_SUMMARY	Germline CDH1 variants in hereditary diffuse gastric cancer syndrome with focus on younger women.
Literature	GENERIF_SUMMARY	Correlation between E-cadherin/beta-catenin, Vimentin expression, clinicopathologic features and drug resistance prediction in naive prostate cancer
Literature	GENERIF_SUMMARY	Early Immune Changes Support Signet Ring Cell Dormancy in CDH1-Driven Hereditary Diffuse Gastric Carcinogenesis.
Literature	GENERIF_SUMMARY	Expression of CD105 but not of E-cadherin is associated with malignancy recurrence and disease-free interval in laryngeal cancer in men.
Literature	GENERIF_SUMMARY	PATHOLOGICAL SIGNIFICANCE OF CDH1/E-CADHERIN GERMLINE SEQUENCE VARIANTS IN BREAST CANCER PATIENTS.
Literature	GENERIF_SUMMARY	The chromatin architectural regulator SND1 mediates metastasis in triple-negative breast cancer by promoting CDH1 gene methylation.
Literature	GENERIF_SUMMARY	Dynamic interactions between E-cadherin and Ankyrin-G mediate epithelial cell polarity maintenance.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Exosomal miR-205-5p Improves Endometrial Receptivity by Upregulating E-Cadherin Expression through ZEB1 Inhibition.
Literature	GENERIF_SUMMARY	MGRN1 depletion promotes intercellular adhesion in melanoma by upregulation of E-cadherin and inhibition of CDC42.
Literature	GENERIF_SUMMARY	PLK1 and its substrate MISP facilitate intrahepatic cholangiocarcinoma progression by promoting lymphatic invasion and impairing E-cadherin adherens junctions.
Literature	GENERIF_SUMMARY	Prostaglandin E2 Exposure Disrupts E-Cadherin/ Caveolin-1-Mediated Tumor Suppression to Favor Caveolin-1-Enhanced Migration, Invasion, and Metastasis in Melanoma Models.
Literature	GENERIF_SUMMARY	Host-Cell-Dependent Roles of E-Cadherin in Serratia Invasion.
Literature	GENERIF_SUMMARY	Tag, you're it
Literature	GENERIF_SUMMARY	LINC MIR503HG Controls SC-beta Cell Differentiation and Insulin Production by Targeting CDH1 and HES1.
Literature	GENERIF_SUMMARY	Oxidative Stress Promotes Liver Cancer Metastasis via RNF25-Mediated E-Cadherin Protein Degradation.
Literature	GENERIF_SUMMARY	Increased CpG methylation at the CDH1 locus in inflamed ileal mucosa of patients with Crohn disease.
Literature	GENERIF_SUMMARY	The clinical significance of SNAIL, TWIST, and E-Cadherin expression in gastric mesentery tumor deposits of advanced gastric cancer.
Literature	GENERIF_SUMMARY	E-cadherin loss drives diffuse-type gastric tumorigenesis via EZH2-mediated reprogramming.
Literature	GENERIF_SUMMARY	Cell division-dependent dissemination following E-cadherin loss underlies initiation of diffuse-type gastric cancer.
Literature	GENERIF_SUMMARY	RAC1-mediated integrin alpha-6 expression in E-cadherin-deficient gastric cancer cells promotes interactions with the stroma and peritoneal dissemination.
Literature	GENERIF_SUMMARY	Expression of CD44, PCNA and E-cadherin in pterygium tissues.
Literature	GENERIF_SUMMARY	CDH1 Genotype Exploration in Women With Hereditary Lobular Breast Cancer Phenotype.
Literature	GENERIF_SUMMARY	CDH1 methylation analysis in invasive lobular breast carcinomas with and without gene mutation.
Literature	GENERIF_SUMMARY	E-cadherin expression and gene expression profiles in corticotroph pituitary neuroendocrine tumor subtypes.
Literature	GENERIF_SUMMARY	A novel noncanonical function for IRF6 in the recycling of E-cadherin.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Associations between CDH1 gene polymorphisms and the risk of gastric cancer
Literature	GENERIF_SUMMARY	Biomarkers of epithelial-mesenchymal transition
Literature	GENERIF_SUMMARY	Molecular Crosstalk Between Adherens Junction Proteins, E-cadherin and Nectin-4.
Literature	GENERIF_SUMMARY	IL-8 promotes lens capsular residual cells migration by down-regulates expression of E-cadherin and ZO-1 via the CXCR1/2-NF-kappaB-RhoA signal pathway.
Literature	GENERIF_SUMMARY	Human RB and c-Myc activate expression of the murine E-cadherin gene in epithelial cells through interaction with transcription factor AP-2
Pathways	KEGG_PATHWAY	Rap1 signaling pathway
Pathways	KEGG_PATHWAY	Apelin signaling pathway
Pathways	KEGG_PATHWAY	Hippo signaling pathway
Pathways	KEGG_PATHWAY	Cell adhesion molecules
Pathways	KEGG_PATHWAY	IgSF CAM signaling
Pathways	KEGG_PATHWAY	Adherens junction
Pathways	KEGG_PATHWAY	Bacterial invasion of epithelial cells
Pathways	KEGG_PATHWAY	Pathways in cancer
Pathways	KEGG_PATHWAY	Endometrial cancer
Pathways	KEGG_PATHWAY	Thyroid cancer
Pathways	KEGG_PATHWAY	Melanoma
Pathways	KEGG_PATHWAY	Bladder cancer
Pathways	KEGG_PATHWAY	Gastric cancer
Pathways	REACTOME_PATHWAY	Apoptosis
Pathways	REACTOME_PATHWAY	Apoptotic cleavage of cellular proteins
Pathways	REACTOME_PATHWAY	Developmental Biology
Pathways	REACTOME_PATHWAY	Adaptive Immune System
Pathways	REACTOME_PATHWAY	Degradation of the extracellular matrix
Pathways	REACTOME_PATHWAY	Extracellular matrix organization
Pathways	REACTOME_PATHWAY	Cell-Cell communication
Pathways	REACTOME_PATHWAY	Signal Transduction
Pathways	REACTOME_PATHWAY	Disease
Pathways	REACTOME_PATHWAY	Immune System
Pathways	REACTOME_PATHWAY	Signaling by Rho GTPases

Main Category	Category	Terms
Pathways	REACTOME_PATHWAY	RHO GTPase Effectors
Pathways	REACTOME_PATHWAY	Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell
Pathways	REACTOME_PATHWAY	Integrin cell surface interactions
Pathways	REACTOME_PATHWAY	Apoptotic cleavage of cell adhesion proteins
Pathways	REACTOME_PATHWAY	Adherens junctions interactions
Pathways	REACTOME_PATHWAY	Cell-cell junction organization
Pathways	REACTOME_PATHWAY	Cell junction organization
Pathways	REACTOME_PATHWAY	Programmed Cell Death
Pathways	REACTOME_PATHWAY	RHO GTPases activate IQGAPs
Pathways	REACTOME_PATHWAY	Infectious disease
Pathways	REACTOME_PATHWAY	Apoptotic execution phase
Pathways	REACTOME_PATHWAY	Listeria monocytogenes entry into host cells
Pathways	REACTOME_PATHWAY	InlA-mediated entry of Listeria monocytogenes into host cells
Pathways	REACTOME_PATHWAY	Signaling by Rho GTPases, Miro GTPases and RHOBTB3
Pathways	REACTOME_PATHWAY	MITF-M-regulated melanocyte development
Pathways	REACTOME_PATHWAY	Developmental Cell Lineages
Pathways	REACTOME_PATHWAY	Developmental Cell Lineages of the Integumentary System
Pathways	REACTOME_PATHWAY	Gastrulation
Pathways	REACTOME_PATHWAY	Regulation of Homotypic Cell-Cell Adhesion
Pathways	REACTOME_PATHWAY	Regulation of CDH1 Expression and Function
Pathways	REACTOME_PATHWAY	Regulation of Expression and Function of Type I Classical Cadherins
Pathways	REACTOME_PATHWAY	Regulation of CDH1 Function
Pathways	REACTOME_PATHWAY	Regulation of CDH1 mRNA translation by microRNAs
Pathways	REACTOME_PATHWAY	Degradation of CDH1
Pathways	REACTOME_PATHWAY	Regulation of CDH1 posttranslational processing and trafficking to plasma membrane
Pathways	REACTOME_PATHWAY	Formation of definitive endoderm
Pathways	REACTOME_PATHWAY	Bacterial Infection Pathways
Pathways	REACTOME_PATHWAY	Transcriptional and post-translational regulation of MITF-M expression and activity

Main Category	Category	Terms
Pathways	REACTOME_PATHWAY	MITF-M-dependent gene expression
Pathways	REACTOME_PATHWAY	Developmental Lineages of the Mammary Gland
Pathways	REACTOME_PATHWAY	Regulation of MITF-M-dependent genes involved in extracellular matrix, focal adhesion and epithelial-to-mesenchymal transition
Pathways	REACTOME_PATHWAY	Developmental Lineage of Mammary Gland Luminal Epithelial Cells
Pathways	REACTOME_PATHWAY	Developmental Lineage of Mammary Gland Myoepithelial Cells
Pathways	REACTOME_PATHWAY	Developmental Lineage of Mammary Stem Cells
Pathways	WIKIPATHWAYS	Integrated breast cancer pathway
Pathways	WIKIPATHWAYS	Neural crest differentiation
Pathways	WIKIPATHWAYS	Pathogenic Escherichia coli infection
Pathways	WIKIPATHWAYS	Mammary gland development embryonic development stage 1 of 4
Pathways	WIKIPATHWAYS	Mammary gland development involution stage 4 of 4
Pathways	WIKIPATHWAYS	Bladder cancer
Pathways	WIKIPATHWAYS	Hair follicle development organogenesis stage 2 of 3
Pathways	WIKIPATHWAYS	Pre implantation embryo
Pathways	WIKIPATHWAYS	TGF beta signaling in thyroid cells for epithelial mesenchymal transition
Pathways	WIKIPATHWAYS	H19 Rb E2F1 and CDK beta catenin in colorectal cancer
Pathways	WIKIPATHWAYS	Endometrial cancer
Pathways	WIKIPATHWAYS	Epithelial to mesenchymal transition in colorectal cancer
Pathways	WIKIPATHWAYS	Mechanoregulation and pathology of YAP TAZ via Hippo and non Hippo mechanisms
Pathways	WIKIPATHWAYS	Hippo signaling regulation
Pathways	WIKIPATHWAYS	Hippo Merlin signaling dysregulation
Pathways	WIKIPATHWAYS	Gastrin signaling
Pathways	WIKIPATHWAYS	Melanoma
Pathways	WIKIPATHWAYS	CAMKK2 pathway
Pathways	WIKIPATHWAYS	Pleural mesothelioma
Pathways	WIKIPATHWAYS	TROP2 regulatory signaling
Pathways	WIKIPATHWAYS	10q11 21q11 23 copy number variation syndrome
Pathways	WIKIPATHWAYS	Macrophage stimulating protein MSP signaling

Main Category	Category	Terms
Pathways	WIKIPATHWAYS	Cell lineage map for neuronal differentiation
Pathways	WIKIPATHWAYS	Cancer pathways
Pathways	WIKIPATHWAYS	Hallmark of cancer metastasis and epithelial to mesenchymal transition
Protein Domains	INTERPRO	Cadherin_Y-type_LIR
Protein Domains	INTERPRO	Cadherin-like_dom
Protein Domains	INTERPRO	Cadherin_pro_dom
Protein Domains	INTERPRO	Cadherin-like_sf
Protein Domains	INTERPRO	Cadherin_CS
Protein Domains	INTERPRO	Catenin-bd_sf
Protein Domains	INTERPRO	Cadherin
Protein Domains	SMART	CA
Protein Domains	SMART	Cadherin_pro
Protein Domains	UP_KW_DOMAIN	Repeat
Protein Domains	UP_KW_DOMAIN	Signal
Protein Domains	UP_KW_DOMAIN	Transmembrane
Protein Domains	UP_KW_DOMAIN	Transmembrane helix