Main Category	Category	Terms
Functional Annotations	UP_KW_BIOLOGICAL_PROCESS	Cell cycle
Functional Annotations	UP_KW_CELLULAR_COMPONENT	Chromosome
Functional Annotations	UP_KW_CELLULAR_COMPONENT	Nucleus
Functional Annotations	UP_KW_MOLECULAR_FUNCTION	DNA-binding
Functional Annotations	UP_KW_PTM	Acetylation
Functional Annotations	UP_KW_PTM	Phosphoprotein
Functional Annotations	UP_KW_PTM	Ubl conjugation
Functional Annotations	UP_KW_PTM	Isopeptide bond
Functional Annotations	UP_SEQ_FEATURE	COMPBIAS:Basic and acidic residues
Functional Annotations	UP_SEQ_FEATURE	COMPBIAS:Basic residues
Functional Annotations	UP_SEQ_FEATURE	COMPBIAS:Polar residues
Functional Annotations	UP_SEQ_FEATURE	CROSSLNK:Glycyl lysine isopeptide (Lys-Gly) (interchain with G-Cte
Functional Annotations	UP_SEQ_FEATURE	CROSSLNK:Glycyl lysine isopeptide (Lys-Gly) (interchain with G-Cte alternate
Functional Annotations	UP_SEQ_FEATURE	CROSSLNK:Glycyl lysine isopeptide (Lys-Gly) (interchain with G-Cte
Functional Annotations	UP_SEQ_FEATURE	CROSSLNK:Glycyl lysine isopeptide (Lys-Gly) (interchain with G-Cte alternate
Functional Annotations	UP_SEQ_FEATURE	DOMAIN:FHA
Functional Annotations	UP_SEQ_FEATURE	DOMAIN:PP1-binding
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:KRRRVSFGGHLRPELFDENLPPNTPLKRGEAPTKRKSLVM QPQPSGKQESGSEIHVEVKAQSLVISPPAPSPRKTPVASDQRRRSCKT VPKRGGRKSGNLPSKRVSISRSQHDILQMICSKRRSGASEANLIVAKSV TQTKVIK->AAAAVSFGGHLAPELFDENLPPNTPLAAGEAPTAAASLVM AEQPQPSGAQESGSEIHVEVAAQSLVISPPAPSPAATPVASDQAAASC TEVPAAGGAASGNLPSAAVSISASQHDILQMICSAAASGASEANLIVA AQTQTAVIA: Abolished ability to undergo liquid-liquid phase separ
Functional Annotations	UP_SEQ_FEATURE	MUTAGEN:VSF->ASA: Abolihed binding with PPP1CC; induces a sl MKI67 enrichment on chromosomes.

Main Category	Category	Terms
Functional Annotations	UP_SEQ_FEATURE	REGION:16 X 122 AA approximate repeats
Functional Annotations	UP_SEQ_FEATURE	REGION:Disordered
Functional Annotations	UP_SEQ_FEATURE	REGION:Positively charged patch (CP)
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 1
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 10
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 11
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 12
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 13
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 14
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 15
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 16
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 2
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 3
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 4
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 5
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 6
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 7
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 8
Functional Annotations	UP_SEQ_FEATURE	REPEAT:K167R 9
Gene Ontology	GOTERM_BP_DIRECT	chromosome segregation
Gene Ontology	GOTERM_BP_DIRECT	regulation of mitotic nuclear division

Main Category	Category	Terms
Gene Ontology	GOTERM_BP_DIRECT	cell population proliferation
Gene Ontology	GOTERM_BP_DIRECT	regulation of chromatin organization
Gene Ontology	GOTERM_CC_DIRECT	condensed chromosome
Gene Ontology	GOTERM_CC_DIRECT	nucleus
Gene Ontology	GOTERM_CC_DIRECT	nucleoplasm
Gene Ontology	GOTERM_CC_DIRECT	chromosome
Gene Ontology	GOTERM_CC_DIRECT	nucleolus
Gene Ontology	GOTERM_CC_DIRECT	membrane
Gene Ontology	GOTERM_MF_DIRECT	nucleotide binding
Gene Ontology	GOTERM_MF_DIRECT	DNA binding
Gene Ontology	GOTERM_MF_DIRECT	RNA binding
Gene Ontology	GOTERM_MF_DIRECT	protein binding
Gene Ontology	GOTERM_MF_DIRECT	ATP binding
Gene Ontology	GOTERM_MF_DIRECT	molecular condensate scaffold activity
General Annotations	CHROMOSOME	10
General Annotations	CYTOGENETIC_LOCATION	10q26.2
General Annotations	ENTREZ_GENE_SUMMARY	Enables RNA binding activity and molecular condensate scaffold a in chromosome segregation and regulation of mitotic nuclear divis chromosome; nucleolus; and nucleoplasm. Is active in condensed Implicated in several diseases, including Crohn's disease; colorecta endocrine gland cancer (multiple); graft-versus-host disease; and h immunodeficiency virus infectious disease. Biomarker of several d including Barrett's esophagus; autoimmune disease of musculosk (multiple); endocrine gland cancer (multiple); gastrointestinal syste (multiple); and lung cancer (multiple). [provided by Alliance of Gen Jul 2025]
General Annotations	UP_COMMENT_ALTERNATIVE_PRODUCTS	Event=Alternative splicing; Named isoforms=2; Name=Long; IsoI Sequence=Displayed; Name=Short; IsoId=P46013-2; Sequence=V

Main Category	Category	Terms
General Annotations	UP_COMMENT_BIOTECHNOLOGY	Widely used as a marker to assess cell proliferation, as it is detect nucleus of proliferating cells only (PubMed:21960707, PubMed:63 cancer research field for example, MKI67 is the most widely used r comparing proliferation between tumor samples (PubMed:219607 PubMed:26680267). {ECO:0000269|PubMed:6339421, ECO:00003 PubMed:21960707, ECO:0000303|PubMed:26680267}.
General Annotations	UP_COMMENT_DEVELOPMENTAL_STAGE	Expression occurs preferentially during late G1, S, G2 and M phase cycle, while in cells in G0 phase the antigen cannot be detected (at (PubMed:6206131). Present at highest level in G2 phase and durin protein level). In interphase, forms fiber-like structures in fibrillarin- regions surrounding nucleoli (PubMed:2674163, PubMed:8799815 {ECO:0000269|PubMed:2674163, ECO:0000269|PubMed:6206131 PubMed:8799815}.
General Annotations	UP_COMMENT_DOMAIN	The positively charged patch (CP) region mediates liquid-liquid pha {ECO:0000269|PubMed:39153474}.
General Annotations	UP_COMMENT_FUNCTION	Protein that associates with the surface of mitotic chromosomes a as a chromosome repellent during early mitosis and chromosome during late mitosis (PubMed:27362226, PubMed:32879492, PubM PubMed:39153474). Required to maintain individual mitotic chrom dispersed in the cytoplasm following nuclear envelope disassembl (PubMed:27362226). During early mitosis, relocalizes from nucleo chromosome surface where it forms extended brush structures th substantial fraction of the chromosome surface (PubMed:273622 brush structure prevents chromosomes from collapsing into a sing mass by forming a steric and electrostatic charge barrier: the prote net electrical charge and acts as a surfactant, dispersing chromos enabling independent chromosome motility (PubMed:27362226). anaphase, the MKI67 brush structure collapses and MKI67 switche chromosome repellent to a chromosome attractant to promote ch clustering and facilitate the exclusion of large cytoplasmic particle future nuclear space (PubMed:32879492, PubMed:39153474). Me dephosphorylation during mitotic exit and simultaneous exposure basic patch induce the RNA-dependent formation of a liquid- like c phase on the chromosome surface, promoting coalescence of nei chromosome surfaces and clustering of chromosomes (PubMed:3 Binds premature ribosomal RNAs during anaphase; promoting liqu separation (PubMed:28935370, PubMed:39153474). Binds DNA, w for supercoiled DNA and AT-rich DNA (PubMed:10878551). Does n the internal structure of mitotic chromosomes (By similarity). May chromatin organization; it is however unclear whether it plays a dir chromatin organization or whether it is an indirect consequence of mitotic chromosome (PubMed:24867636). {ECO:0000250|UniProt ECO:0000269|PubMed:10878551, ECO:0000269|PubMed:2486763 ECO:0000269|PubMed:27362226, ECO:0000269|PubMed:2893537 ECO:0000269|PubMed:32879492, ECO:0000269|PubMed:3551370 ECO:0000269|PubMed:39153474}.
General Annotations	UP_COMMENT_INTERACTION	P46013; Q9NS87: KIF15; NbExp=3; IntAct=EBI-876367, EBI-712159 Q9BYG3: NIFK; NbExp=3; IntAct=EBI-876367, EBI-2561019;
General Annotations	UP_COMMENT_PTM	Hyperphosphorylated by CDK1 in mitosis; hyperphosphorylatiom p undergoing liquid-liquid phase separation (PubMed:10502411, Pub PubMed:25012651, PubMed:39153474). Dephosphorylated by PP onset of anaphase (PubMed:25012651). Dephosphorylated by pro phosphatase 2A (PP2A) at the onset of anaphase (By similarity). Dephosphorylation by protein phosphatase 2A (PP2A) and simulta of the positively charged patch (CP) during mitotic exit induce the formation of a liquid-like condensed phase on the chromosome su

Main Category	Category	Terms
		(PubMed:39153474). {ECO:0000250|UniProtKB:E9PVX6, ECO:0000 PubMed:10502411, ECO:0000269|PubMed:10653604, ECO:00002 PubMed:25012651, ECO:0000269|PubMed:39153474}.
General Annotations	UP_COMMENT_PTM	Ubiquitinated by the APC/C complex after neuronal progenitors ex brain development, leading to clearance from constitutive heteroch {ECO:0000269|PubMed:34942119}.
General Annotations	UP_COMMENT_SUBCELLULAR_LOCATION	Chromosome {ECO:0000269|PubMed:15896774, ECO:0000269| PubMed:22002106, ECO:0000269|PubMed:27362226, ECO:00002 PubMed:32879492, ECO:0000269|PubMed:35513709, ECO:00002 PubMed:39153474, ECO:0000269|PubMed:9510506}. Nucleus {EC PubMed:10878551, ECO:0000269|PubMed:22002106}. Nucleus, n {ECO:0000269|PubMed:10878551, ECO:0000269|PubMed:220021 ECO:0000269|PubMed:2674163, ECO:0000269|PubMed:8799815} early mitosis, relocalizes from nucleoli to the surface of the mitoti the perichromosomal layer, and covers a substantial fraction of the chromosome surface (PubMed:27362226). Associates with satelli phase (PubMed:9510506). Binds tightly to chromatin in interphase binding decreases in mitosis when it associates with the surface c condensed chromosomes (PubMed:15896774, PubMed:22002106 Predominantly localized in the G1 phase in the perinucleolar regior phases it is also detected throughout the nuclear interior, being pre localized in the nuclear matrix (PubMed:22002106). {ECO:000026 PubMed:15896774, ECO:0000269|PubMed:22002106, ECO:00002 PubMed:27362226}.
General Annotations	UP_COMMENT_SUBCELLULAR_LOCATION	Nucleus {ECO:0000256|ARBA:ARBA00004123}.
General Annotations	UP_COMMENT_SUBUNIT	Interacts with KIF15 (PubMed:10878014). Interacts (via the FHA d NIFK (PubMed:11342549, PubMed:14659764, PubMed:16244663) PPP1CC (PubMed:24867636, PubMed:25012651). Component of least composed of ZNF335, HCFC1, CCAR2, EMSY, MKI67, RBBP5, WDR5; the complex is formed as a result of interactions between c a nuclear receptor-mediated transcription complex and a histone m complex (PubMed:19131338). Interacts with ZNF335 (PubMed:19 PubMed:23178126). {ECO:0000269|PubMed:10878014, ECO:0000 PubMed:11342549, ECO:0000269|PubMed:14659764, ECO:00002 PubMed:16244663, ECO:0000269|PubMed:19131338, ECO:00002 PubMed:23178126, ECO:0000269|PubMed:24867636, ECO:00002 PubMed:25012651}.
Identifiers	ENTREZ_GENE_ID	4288
Identifiers	ENSEMBL_GENE_ID	ENSG00000148773
Identifiers	UNIPROT_ID	Q70L74_HUMAN
Identifiers	UNIPROT_ID	Q70L72_HUMAN
Identifiers	UNIPROT_ID	Q70L73_HUMAN
Identifiers	UNIPROT_ID	KI67_HUMAN
Interactions	UP_KW_LIGAND	ATP-binding
Interactions	UP_KW_LIGAND	Nucleotide-binding
Literature	GENERIF_SUMMARY	There is no significant correlation between Ki-67 labelling index an anterior pituitary adenomas, even if this index can be considered a in the determination of the infiltrative behavior of these tumors.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Not significantly correlated with lymph node metastasis of breast i carcinoma.
Literature	GENERIF_SUMMARY	Use of tissue microarray technique shows immunohistochemical t heterogeneity in expression levels of Ki-67 in malignant fibrous his
Literature	GENERIF_SUMMARY	Ki-67 antigen overexpressioin is an early event in esophageal carci useful biomarkers for early detection.
Literature	GENERIF_SUMMARY	patients that progressed to breast cancer showed significantly hig expression in their HUT (hyperplasia of usual type)foci compared v
Literature	GENERIF_SUMMARY	overexpression may be marker of radioresistance in head and nec
Literature	GENERIF_SUMMARY	Results indicate that loss of RAR-beta expression and accumulatio Ki67 proteins may serve as biomarkers for early identification of es cancer in the high-risk populations.
Literature	GENERIF_SUMMARY	The combined evaluation of p27Kip1 and Ki-67 expression provide information on overall survival of ovarian carcinoma patients.
Literature	GENERIF_SUMMARY	presence in CNS tumors correlates with histological type and grad
Literature	GENERIF_SUMMARY	This marker is masked to MIB-1 staining after expression of its tar
Literature	GENERIF_SUMMARY	p53 and Ki-67 expression in epithelial gastric dysplasia and in gast
Literature	GENERIF_SUMMARY	Increase of CD44s, MMP-9, and Ki-67 were involved in the growth a invasion of osteosarcoma.
Literature	GENERIF_SUMMARY	Data suggest that marsupial chmadrin and human Ki-67 induce ch compaction across species, possibly via the interaction of its LR d heterochromatin protein 1.
Literature	GENERIF_SUMMARY	No relationship was observed between Ki-67 staining & the E-cadh status. The presence of E-cadherin mutations does not correlate w staining.
Literature	GENERIF_SUMMARY	increased p53 and Ki-67 expression in varying grades of urothelial bladder
Literature	GENERIF_SUMMARY	Ki-67 proved to be a better prognostic marker, as compared to met large intestine adenocarcinomas
Literature	GENERIF_SUMMARY	Ki-67 antigen reactivity was significantly higher in the patient with carcinoma remission, compared with the control group.
Literature	GENERIF_SUMMARY	a biological marker in Langerhans cell histiocytosis
Literature	GENERIF_SUMMARY	distribution of staining pattern for Ki-67 paralleled the staining pat betaE12
Literature	GENERIF_SUMMARY	Detected in nuclei of neoplastic cells and in nuclei of epithelial cell colonic mucosa.
Literature	GENERIF_SUMMARY	nmr analysis of the solution structure of the FHA domain of human mapping of the binding surface for NIFK binding
Literature	GENERIF_SUMMARY	compared the expression of Ki-67 between primary breast tumors to regional lymph nodes; estimated the relationships between Ki-6 anatomoclinical features of the breast cancer
Literature	GENERIF_SUMMARY	The expression of Ki67 and p53 in various forms of leukoplakia po

Main Category	Category	Terms
		increasing instability of the genome in parallel with the severity of
Literature	GENERIF_SUMMARY	Mutations in the cDNA of the proliferation marker Ki-67 protein is a cervix, colonic and lung neoplasms
Literature	GENERIF_SUMMARY	study expands the immunophenotype of granular cell tumor (S100 gene product 9.5, and inhibin-alpha) regardless of location and sup origin
Literature	GENERIF_SUMMARY	Ki-67 may have a role in distant metastasis of radiotherapy-treated neoplasms
Literature	GENERIF_SUMMARY	Ki-67 LI (labeling index) is a useful prognosticator for PENs (pancr neoplasms).
Literature	GENERIF_SUMMARY	Ki-67 is a factor of poor prognosis for survival in NSCLC.
Literature	GENERIF_SUMMARY	ER alpha and Ki67 might be involved in distinct pathological and m features during breast cancer development
Literature	GENERIF_SUMMARY	Expression correlates with intensity of apoptosis in colorectal ade
Literature	GENERIF_SUMMARY	Expression of Ki-67, PCNA proteins were closely connected with th stomach tumour malignancy.
Literature	GENERIF_SUMMARY	Evaluation of expression of PCNA and Ki-67 may provide an additi about the progression of uveal melanoma.
Literature	GENERIF_SUMMARY	Ki-67 and hTERT have roles in meningioma cell proliferation, but o role in disease recurrence
Literature	GENERIF_SUMMARY	a useful msarkser in patients with lung adenocarcinoma.
Literature	GENERIF_SUMMARY	Ki-67 by itself is a less reliable marker of dysplasia.
Literature	GENERIF_SUMMARY	Altered expression is associated with therapy failure and death in multiple types of cancer.
Literature	GENERIF_SUMMARY	Study suggests that the clinical characteristics such as visual field recurrence are correlated with the high Ki-67 labeling index.
Literature	GENERIF_SUMMARY	Proliferative activity of 13 biopsy specimens of Kaposi sarcoma w immunohistochemically using the monoclonal antibody to KI-67.
Literature	GENERIF_SUMMARY	Aberrant proliferative patterns with Ki67 staining are not useful in reactive epithelia from low-grade dysplasia, but may prove useful i of high-grade dysplasia.
Literature	GENERIF_SUMMARY	p27(kip1) and Ki-67 have roles in post-surgical disease progressio cancer
Literature	GENERIF_SUMMARY	ability of the gef gene to down regulate Ki-67 expression and induc a breast cancer cell line
Literature	GENERIF_SUMMARY	ChIP assays provided evidence for a physical association of pKi-67 chromatin of the promoter and transcribed region of the rRNA gen data strongly suggest a role for pKi-67 in the early steps of rRNA s
Literature	GENERIF_SUMMARY	liver expression in HBV and HCV cirrhosis and hepatocellular carci associated with dysplastic cell changes and tumour grade
Literature	GENERIF_SUMMARY	Ki 67 usefulness in studying cellular proliferation, and underline th lymphocytic leukemia (CLL) with polymorphic cytology are more p

Main Category	Category	Terms
		typical CLL.
Literature	GENERIF_SUMMARY	To investigate the prognostic and predictive relevance of p53 prote antigen, MMP-2 and MMP-9 in patients with transitional cell carcin the upper urinary tract
Literature	GENERIF_SUMMARY	The expression rate of Ki-67 was relatively high in small cell lung c any prognostic significance.
Literature	GENERIF_SUMMARY	Increased expression of Ki-67 protein suggests an increased cellul which is involved in the process of occurrence and progression of dysplasia in the flat mucosa of colon.
Literature	GENERIF_SUMMARY	Expression of iNOS and Ki-67 protein in hemangioma is significant that in vascular malformation.
Literature	GENERIF_SUMMARY	Ki-67/podoplanin-immunoreactivity was useful to identify prolifera vessels.
Literature	GENERIF_SUMMARY	The Ki-67 Antigen expression was associated with the highest leve proliferation, suggesting a synergistic effect.
Literature	GENERIF_SUMMARY	Ki67 stain alone is a powerful tool for distinguishing benign from m proliferations within the selected groups.
Literature	GENERIF_SUMMARY	The combination of MCM2, geminin and Ki67 could represent a va the understanding of hepatocellular carcinoma progression in cirrh
Literature	GENERIF_SUMMARY	Ki67 was an independent prognosticator if nodal status, tumour si histological grade were included in the analysis but not if analysed standardized mitotic index.
Literature	GENERIF_SUMMARY	changes in COX-2 and Ki-67 expression peaked during the prolifera fell during the late luteal phase in endometrial polyps and disease-
Literature	GENERIF_SUMMARY	Overexpression of Ki67 and VEGF were related to poor prognosis c cell carcinoma of the larynx.
Literature	GENERIF_SUMMARY	In univariate analyses, low Ki-67 proliferation index (P = 0.045) and immunoreactivity (P = 0.04) were associated to patient's overall su
Literature	GENERIF_SUMMARY	Ki-67 did not display apparent prognostic value in NSCLC in the stu
Literature	GENERIF_SUMMARY	Height of Ki-67 staining within the epithelium increased significant and moderate and between mild and severe dysplasia.
Literature	GENERIF_SUMMARY	Based on these results it is suggested that Ki-67 is a reliable micro to predict metastatic potential of rectal carcinoids.
Literature	GENERIF_SUMMARY	analysis of Ki67 and thymidylate synthase expression in primary tu compared with metastatic nodes in breast cancer patients
Literature	GENERIF_SUMMARY	In an analysis of breast cancer-specific survival up to 5 years, high and negative ER as well as aneuploidy of the tumour were significa factors.
Literature	GENERIF_SUMMARY	p53 and Ki-67 were expressed with increasing frequency, and bcl-2 mdm-2 with decreasing frequency in thyroid carcinoma progressio cyclin D1 were expressed in
Literature	GENERIF_SUMMARY	The expression of STAT3 was positively related to that of Survivin nasal NK/T cell lymphoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	This study shows that histologic features are reliable in identifying tumors that have a higher proliferative potential as shown by highe immunoreactivity for Ki-67 and phosphorylated histone H3 (PHH3)
Literature	GENERIF_SUMMARY	The basal-like subtype was associated with elevated Ki-67 index.
Literature	GENERIF_SUMMARY	Analysis of p27 and Ki-67 expression levels might indicate that low activity of desmoid fibroblasts is connected with another mechani one, in which p27 takes part.
Literature	GENERIF_SUMMARY	MUM1 expression dichotomises follicular lymphoma (FL) into low- CD10+/Bcl-6+/MUM1-/Ki-67low phenotype, and high-grade FL of C weak/MUM1+/ Ki-67high phenotype
Literature	GENERIF_SUMMARY	No siatistically significant value in assocation with survival or even in head and neck squamous cell carcinoma.
Literature	GENERIF_SUMMARY	Ki-67 index and PIK3CA amplification were strong predictors for an associated death in ovarian cancer patients.
Literature	GENERIF_SUMMARY	The present study showed elevated expression of topo-I, Ki-67, and patients with uterine carcinosarcoma suggesting sensitivity to top drug treatment.
Literature	GENERIF_SUMMARY	There are significant differences between Ki-67 indices and CCND1 among melanocytic lesions.
Literature	GENERIF_SUMMARY	When MKI67 immunohistochemistry is used for diagnostic purpos suspected hyaline trabecular tumor of the thyroid, only MIB-1 clone temperature should be used.
Literature	GENERIF_SUMMARY	Expression of MKI67 as a proliferation marker was studied, using M
Literature	GENERIF_SUMMARY	crucial role of pKi-67 in RNA polymerase I-dependent nucleolar rRN
Literature	GENERIF_SUMMARY	p16, p53 and Ki-67 expressions were all negative in normal cervica but all positive in cervical intraepithelial neoplasia.
Literature	GENERIF_SUMMARY	The Ki67+ proliferation index was inversely correlated with desmop plakophilin-1 during cervical squamous cell carcinogenesis
Literature	GENERIF_SUMMARY	Ki-67 and coagulative tumor necrosis both should be evaluated wh outcome predictions for patients with clear cell renal cell carcinom
Literature	GENERIF_SUMMARY	findings suggest Ki67 expression as a prognostic marker for lympl metastasis in penile squamous carcinoma
Literature	GENERIF_SUMMARY	Expression rates of EGFR and Ki-67 in squamous cell carcinoma o and paranasal sinuses were significantly higher than that in nasal i papilloma and nasal polyps.
Literature	GENERIF_SUMMARY	Ki-67 expression is predictive of prognosis, and our prognostic mo become a useful tool for predicting prognosis in patients with stag NK/T-cell lymphoma, nasal type
Literature	GENERIF_SUMMARY	Ki-67 antibody and cyclin D1 overexpression seem to parallel each supporting the concept that cyclin D1 serves as a cell cycle activat cell carcinoma of the penis.
Literature	GENERIF_SUMMARY	p53 expression may have a role in the carcinogenesis of squamou carcinoma whereas Bcl-2 expression has no role. Ki-67 expression used in determining the aggressiveness of CIN lesions

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Ki-67 levels were not helpful in identifying patients who would ben neoadjuvant treatment of rectal cancer.
Literature	GENERIF_SUMMARY	p53 and Ki-67, but not bcl-2, cyclin D1 or HER-2 may have roles in t tumor genesis in non-small cell lung carcinoma
Literature	GENERIF_SUMMARY	Ki-67 represents an independent prognostic predictor in stage III o
Literature	GENERIF_SUMMARY	Expression of CCND1 is correlated with Ki-67 expression.
Literature	GENERIF_SUMMARY	These results suggest that papillary thyroid carcinomas with BRAF mutation are more aggressive than those with wildtype BRAF.
Literature	GENERIF_SUMMARY	in colorectal cancer patients c-reactive protein levels and Ki67 (pro related Ki-67 antigen) expression were associated with poorer can survival; Ki67 labelling index and C-reactive protein were correlated
Literature	GENERIF_SUMMARY	MIB-1(Ki-67) positive tumor cells were maximally aggregated at th an excised specimen and can provide better information about the biological nature in benign meningiomas
Literature	GENERIF_SUMMARY	high cyclin expression may contribute to deregulation of the cell cy soft tissue tumors
Literature	GENERIF_SUMMARY	No significant association was observed between FHIT and Ki-67 e malignant pleural mesothelioma
Literature	GENERIF_SUMMARY	Ki-67 index remains an important prognostic marker in the era of a therapy in advanced mantle cell lymphoma.
Literature	GENERIF_SUMMARY	expression of p53 and Ki-67 antigen is helpful to predict tumor rec prognosis in skull base chordomas
Literature	GENERIF_SUMMARY	Aim of the studies included detection and subcellular localization proteins (NS3, NS5A and C) in liver biopsies from adults (n=19) wit lasting hepatitis C as related to hepatocyte proliferative activity.
Literature	GENERIF_SUMMARY	In ER-alpha-negative tumors, we detected a correlation between ER expression and high levels of Ki-67.
Literature	GENERIF_SUMMARY	Ki-67- and p53-immunostained cells were mainly located in the sup p63-positive cells were found throughout the lining cystic epitheliu
Literature	GENERIF_SUMMARY	Only FIGO stage, cyclin D1, p27 (Kip1) and Ki-67 are independent p factors that might help in predicting outcome of cervical cancer pa
Literature	GENERIF_SUMMARY	The combination of Ki-67 and PCNA, specific proliferative markers improve the accuracy of early diagnosis of prostatic cancer.
Literature	GENERIF_SUMMARY	Non-smoking and non-drinking patients with squamous cell carcin same risk for developing multiple tumors as their smoking and drin counterparts without an increased expression of Ki-67.
Literature	GENERIF_SUMMARY	Conclude that the Ki-67 and p53 labeling indices are useful additio discriminating atypical from benign or anaplastic meningiomas.
Literature	GENERIF_SUMMARY	Ki-67 labeling index was an independent predictor of progression (analysis, P
Literature	GENERIF_SUMMARY	DPPIV-expression and enzyme activity, Ki-67 antigen and K16 are s upregulated in the centre and inner margin of the psoriatic lesion c clinically uninvolved skin and the healthy volunteers skin.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Ki-67 and PCNA are related to proliferation activities of tumor cells neuromas.
Literature	GENERIF_SUMMARY	The role of p53, followed by Ki-67, as predictive factors.
Literature	GENERIF_SUMMARY	Ki67 expression was higher in planoepithelial metaplasia and eosi infiltrations.
Literature	GENERIF_SUMMARY	Expression of Ki-67 in cervical intraepithelial neoplasia and invasiv cell carcinoma of the uterine cervix
Literature	GENERIF_SUMMARY	Report relationship of Ki67, TP53, MDM-2 and BCL-2 expressions v and WHO/ISUP grades, tumor category and overall patient survival tumors of the bladder.
Literature	GENERIF_SUMMARY	Post-chemotherapy Ki67 is a strong predictor of outcome for breas patients not achieving a pathological complete response
Literature	GENERIF_SUMMARY	The pattern and extent of Ki-67 are useful parameters for identifyin Barrett esophagus.
Literature	GENERIF_SUMMARY	Ki-67 expression is likely to be associated with tumour recurrence prognosis in jaw bone ES in adult patients.
Literature	GENERIF_SUMMARY	Inhibition of renal cancer cell growth in vitro and in vivo with oncol armed short hairpin RNA targeting Ki-67 encoding mRNA
Literature	GENERIF_SUMMARY	Ki-67 and bcl-2 expression was correlated with tumor grade, and th tumor grade, the higher the Ki-67 and bcl-2 expression.
Literature	GENERIF_SUMMARY	Significance of the expression of nucleolar argyrophilic proteins ar in the evaluation of cell proliferative activity and in the prediction o lung cancer
Literature	GENERIF_SUMMARY	Ki-67 LI merits further study as a possible biomarker of early prost behaviour.
Literature	GENERIF_SUMMARY	Ki-67 expression level is a potentially useful clinical marker for pre recurrence in surgically treated stage I oral tongue squamous cell
Literature	GENERIF_SUMMARY	low Ki-67 expression also remained as an adverse prognostic facto independent of non-GCB phenotype and bcl-2 expression
Literature	GENERIF_SUMMARY	proliferative marker Ki-67 was analyzed in 11 nasal NK/T-cell lymp immunohistochemical methods
Literature	GENERIF_SUMMARY	Ki-67 plays a role in breast carcinoma cell proliferation & invasion. Ki-67 is essential for cell growth & tumorigenicity.
Literature	GENERIF_SUMMARY	In this immunohistochemical study, 80% of all vulvar intra-epithelia were positive of MIB1 protein.
Literature	GENERIF_SUMMARY	Overexpression of Ki67 was negatively correlated with cell differen gastric carcinoma.
Literature	GENERIF_SUMMARY	analysis of the expression of fatty acid synthase, Ki-67 and p53 in carcinomas of the larynx
Literature	GENERIF_SUMMARY	Its expression is prognostic parameter in breast carcinomas.
Literature	GENERIF_SUMMARY	significant interrelationships of the investigated molecular alterati clinicopathological characteristics in diffuse gliomas, which suppo role of molecular markers (MGMT, p53, Ki-67)in the diagnostic ass

Main Category	Category	Terms
		these neoplasms
Literature	GENERIF_SUMMARY	Human eccrine sweat glands express CK7, CK8, CK14, CK18, CK19 Ki67, p63, EGF and EGFR. In skin, CEA can be used as a specific im marker of sweat glands.
Literature	GENERIF_SUMMARY	Expression of p53, BCL-2 and Ki-67 was upregulated in clinically lo cancer compared with benign prostate tissue, with no alteration in expression.
Literature	GENERIF_SUMMARY	peripheral blood lymphocyte CD25 and Ki67 have roles in renal cel
Literature	GENERIF_SUMMARY	Overexpression of Bmi-1 and Ki67 protein are significantly correlat tumorigenesis, metastasis and prognosis of colorectal carcinoma.
Literature	GENERIF_SUMMARY	Clinical outcome of bladder urothelial tumours showed a significar with MIB1 LI and molecular grade.
Literature	GENERIF_SUMMARY	Correlation between poor survival and altered expression rates of I malignancies indicate a biological role in salivary cancer pathogen
Literature	GENERIF_SUMMARY	Tumors with lower Ki67 expression were more sensitive to neoadju than that with higher expression of Ki67.
Literature	GENERIF_SUMMARY	Analysis of Ki-67 expression in aggressive fibromatosis (desmoid reported.
Literature	GENERIF_SUMMARY	Trends suggested down-modulation of cyclooxygenase-2 and Ki-67 tissues, increased pAKT-Ser473 expression, and an inverse relation PGE(2) and BCL2 expression.
Literature	GENERIF_SUMMARY	MCM7 is a more reliable marker than MCM2 or Ki-67 in lung adenc
Literature	GENERIF_SUMMARY	Clonal status was associated with the expression rate of Ki-67 to a which may be useful in assessing susceptibility to gastric carcinor
Literature	GENERIF_SUMMARY	The expression of EGFR, Ki67, p16 in middle ear cholesteatoma wa higher compared with controls.
Literature	GENERIF_SUMMARY	Data show that Cav-1 expression in tumor tissues was correlated v Ki-67 and p53 expression.
Literature	GENERIF_SUMMARY	high expression is associated with post-operative recurrence of lur adenocarcinomas
Literature	GENERIF_SUMMARY	Standard uptake value of 18F-FDG PET-CT is linearly correlated wit expression of GLUT1, but not with the expression of Ki67.
Literature	GENERIF_SUMMARY	Immunohistochemistry detected inhibin-alpha, -betaA and -betaB, glycodelin A in invasive trophoblastic mole.
Literature	GENERIF_SUMMARY	Membranous/cytoplasmic Ki67 identifies a group of breast carcino be important to consider separately in prognostic and predictive st
Literature	GENERIF_SUMMARY	All multicentric and nodular basal cell carcinomas displayed a pron Srcasm staining in association with a marked increase in Ki-67 sta
Literature	GENERIF_SUMMARY	Ki67 immunoexpression was significantly correlated with raised tra hepatocellular carcinoma
Literature	GENERIF_SUMMARY	Concurrent expression of Ki67 and CD34 may be an important cha involved in the progression of colorectal cancer.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	For routine clinical purposes, grading appears to add only limited i about the prognosis in comparison with Ki-67 expression
Literature	GENERIF_SUMMARY	Results indicated that Ki-67 evaluation in routine EUS-FNA cytolog can be used as a potential prognostic marker in pancreatic NETs.
Literature	GENERIF_SUMMARY	The present results suggest that the decrease in syndecan-1 expre increase in the Ki-67 index observed in ameloblastic carcinomas is with its higher aggressiveness as compared to the rare desmoplas peripheral ameloblastomas.
Literature	GENERIF_SUMMARY	Data show that the over-expression of hTERT, c-myc and Ki-67 may in the progress of hepatocellular carcinoma developing.
Literature	GENERIF_SUMMARY	Expression of ER, progesterone receptor, and HER2 proteins and th appear to distinguish luminal A from luminal B breast cancer subty
Literature	GENERIF_SUMMARY	Immunodetection of HER2/neu receptor correlated positively with and PR immunolabelling. The Ki-67 proliferation index correlated o ERalpha immunopositivity.
Literature	GENERIF_SUMMARY	The expressions of p53, p21 and Ki67 proteins are closely related pathogenesis and progression of nasal NK/T-cell lymphoma.
Literature	GENERIF_SUMMARY	The expression of Ki67 in squamous carcinoma of cervix was not the major clinic pathological parameters, and there was no obviou between the expression of RhoC gene and Ki67 antigen.
Literature	GENERIF_SUMMARY	Combined MDM2 and Ki-67 expression levels could be considered stratification of patients with prostate cancer in clinical trials.
Literature	GENERIF_SUMMARY	The staining pattern for p16 in desmoplastic melanomas and Spitz conjunction with the histopathologic features, S100 staining, Ki67 index and clinical scenario may aid in the difficult differential diagn
Literature	GENERIF_SUMMARY	The results of this study showed that MIB-1 LI is a rapid and cost-e modality for predicting tumor grade in oligodendrogliomas.
Literature	GENERIF_SUMMARY	Both Ki-67 and c-kit immunoreactivities were explored and compar normal breast tissues and subsequently documented cancers.
Literature	GENERIF_SUMMARY	Studies indicate that estrogen receptor-positive tumors with increa proliferation, evaluated by Ki-67, may be selectively responsive to a inhibitor therapy.
Literature	GENERIF_SUMMARY	An immunohistochemical and molecular study of the BCL2, MKI67 genes in Synovial sarcoma (SS) was performed. The high level of E correlated with a high level of CCND1 mRNA, but not with MKI67 m
Literature	GENERIF_SUMMARY	Positive expression of p16, Ki-67, and ProEx C and negative expres seem to be related to HPV-16 infection
Literature	GENERIF_SUMMARY	P63, followed by Ki-67, showed better correlation with cancer prog p53.
Literature	GENERIF_SUMMARY	The main objectives were to assess tumor proliferation and invasi biomarkers (Ki-67, E-cadherin) and to identify potential correlation biomarkers and classic prognostic factors in cervical cancer.
Literature	GENERIF_SUMMARY	High mitotic index (MI) and high MIB-1 expression were found.
Literature	GENERIF_SUMMARY	increased EZH2 protein expression and Ki-67 expression in >40% c

Main Category	Category	Terms
		cells are both associated with an improved relapse-free survival in stages II and III, but not so in rectal cancer.
Literature	GENERIF_SUMMARY	Ki67 appears to be a time-varying biomarker of risk of breast canc with atypical hyperplasia
Literature	GENERIF_SUMMARY	the quantitative and qualitative (pattern of distribution) evaluation expressions of Ki67, P27 and P53 may be helpful in differentiating premalignant epidermal lesions, particularly in unsatisfactory or fra specimens
Literature	GENERIF_SUMMARY	Endothelial proliferation assessed on Ki-67 combined with the lack expression is often associated with invasive cervical carcinoma.
Literature	GENERIF_SUMMARY	MCM3 may be a more reliable proliferation marker than Ki-67 in ac growth of tumor and evaluating tumor aggressiveness of papillary carcinoma.
Literature	GENERIF_SUMMARY	Genes previously shown to be of relevance in high-grade gliomas i Ki-67, tp53, and O6-methylguanine-DNA methyltransferase(MGMT) are not significant biomarkers of outcome in pilocytic astrocytoma
Literature	GENERIF_SUMMARY	Data show that the CIS surrounding morphologically normal urothe high frequency of abnormal DNA content, clear aneuploid cells, p5 protein expression, and a proliferative status underlying a field carc
Literature	GENERIF_SUMMARY	Report correlation of the standardized uptake value in FDG-PET wi expression level of Ki-67 antigen in resected non-small cell lung ca
Literature	GENERIF_SUMMARY	Ki67 is a prognostic factor in node-negative breast cancer
Literature	GENERIF_SUMMARY	This study evaluates the association of Forkhead-box protein A1 (F GATA-binding protein 3 (GATA3) expressions with Oncotype DX rec scores in 77 cases of patients with ER-positive node-negative brea
Literature	GENERIF_SUMMARY	An immunochemical study of the expressions of Ki-67, estrogen re progesterone receptor, and p53 in 9 cases of symplastic uterine lei performed.
Literature	GENERIF_SUMMARY	Toker cells and mammary Paget cells differ substantially in their ex c-erbB-2 and Ki-67.
Literature	GENERIF_SUMMARY	patients with Gleason scale > 8 prostate cancer were characterized nuclear expression of p53, p16INK4a and Ki-67, and also by overex cytoplasmic Bcl-2.
Literature	GENERIF_SUMMARY	Studies indicate that Ki67 at baseline is a highly prognostic factor cancer.
Literature	GENERIF_SUMMARY	Studies indicate that Ki67 at baseline is a highly prognostic factor cancer.
Literature	GENERIF_SUMMARY	Data show that a positive correlation was observed between the S PET at the biopsy site and the Ki-67 proliferation index (MIB-1 labe
Literature	GENERIF_SUMMARY	The expression of p53, c-erbB-2 and Ki67 proteins in 20 simple inte metaplasia (SIM), 27 atypical intestinal metaplasia (AIM), and 37 g carcinomas showed significant differences between SIM and gast
Literature	GENERIF_SUMMARY	CyclinB1, FHIT and Ki-67 may play significant roles in the occurren evolution of gastric carcinoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	The Ki-67 Labeling Index in the malignant GIST group was higher t benign group, in the p53-positive cases was higher than that in the cases, in the C-kit-positive group was higher than that in the C-kit-n
Literature	GENERIF_SUMMARY	Data show that SOX9 expression was associated with increasing G and higher Ki67 staining in prostate cancer samples.
Literature	GENERIF_SUMMARY	The clinical trial of Kim et al. shows an effect of the cyclooxygenas inhibitor celecoxib at a high dose on Ki-67 expression in the norma epithelia of current and former smokers.
Literature	GENERIF_SUMMARY	no significant correlation with disease-free survival in nasopharyng
Literature	GENERIF_SUMMARY	There is no statistically significant association between the expres and overall survival or relapse of patients with laryngeal cancer. Nu staining was statistically significantly correlated to Ki-67 expressio
Literature	GENERIF_SUMMARY	MIB1 overexpression is associated with urothelial carcinomas of t tract.
Literature	GENERIF_SUMMARY	There is a significant statistical difference between immunohistoc expression of p16(INK4a) and Ki-67 in HPV-associated dysplastic dysplastic vulvar lesions.
Literature	GENERIF_SUMMARY	MLH1, MSH2 and MSH6 expression in initial glioblastoma lesions associated with the Ki67 antigen proliferation index.
Literature	GENERIF_SUMMARY	Increased expression of Ki-67 was associated with the presence o metastasis, advanced stages of disease, tumors occurring in the fl and moderately/well-differentiated oral squamous cell carcinoma.
Literature	GENERIF_SUMMARY	High MIB-1 expression is associated with gallbladder carcinoma.
Literature	GENERIF_SUMMARY	The expressions of P53, P21 and Ki67 proteins are closely related pathogenesis and progression of nasal NK/T-cell lymphoma.
Literature	GENERIF_SUMMARY	Prognostic value of of the expression Ki-67, CD31, and VEGF in somatotropinomas
Literature	GENERIF_SUMMARY	expression of CD147 was positively correlated with matrix metallo Ki-67 labeling index, or both in invasive pituitary adenomas (P
Literature	GENERIF_SUMMARY	Ki-67 was a stronger and more robust prognostic indicator than mi this series of nodular melanoma.
Literature	GENERIF_SUMMARY	overexpression in esophageal mucosa correlates with the severity
Literature	GENERIF_SUMMARY	High MIB1 is associated with high-grade ovarian serous carcinoma
Literature	GENERIF_SUMMARY	High levels of HER2 and Ki-67 or TK1 expression correlate with the tumor grades and tumor recurrence in meningiomas.
Literature	GENERIF_SUMMARY	A review of literature is carried out to correlate the role of Ki-67 lab its correlation with clinical findings, tumor size, invasiveness, recur adenoma subtype, adenoma doubling time, and pituitary carcinom addressed.
Literature	GENERIF_SUMMARY	CD133 expression positively correlates with that of Ki-67 in gastric adenocarcinoma.
Literature	GENERIF_SUMMARY	Endometrial adenocarcinomas show micro-anatomical variations i expression and this is often inversely correlated with CK7 immuno

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Ki-67, bcl-2 and caspase-3 positive cells changed in a temporally a restricted manner coincident with upper jaw differentiation
Literature	GENERIF_SUMMARY	Patients with a higher expression of Ki67 has a better tumor respo chemotherapy.
Literature	GENERIF_SUMMARY	High KI-67 is associated with endometrial cancer.
Literature	GENERIF_SUMMARY	MCM-2 expression did not show significant associations with any clinicopathological parameters, while Ki-67 expression was merely associated with tumor size. Neither MCM-2 nor Ki-67 expression s prognostic value.
Literature	GENERIF_SUMMARY	intracellular Ki67 expression provides a specific, quantitative and r measure of antigen-specific T cell proliferation in vitro.
Literature	GENERIF_SUMMARY	Most cases that were p53 positive had an increased proliferation a determined with Ki67.
Literature	GENERIF_SUMMARY	MIB-1 showed a weak expression in the basement membrane of th epithelium and a total absence in the lamina propria deep layer
Literature	GENERIF_SUMMARY	The high expression of CAIX in non-small cell lung cancer tissues i with VEGF level but not with Ki67 level.
Literature	GENERIF_SUMMARY	The positive intensity of Ki67 protein expression in primary ovarian significantly higher than in metastatic ovarian cancers.
Literature	GENERIF_SUMMARY	There was no correlation of Ki-67 expression and tumor stage or g and neck squamous cell carcinoma.
Literature	GENERIF_SUMMARY	We conclude that Ki-67 labeling detects anal HPV-related changes degree of sensitivity and specificity.
Literature	GENERIF_SUMMARY	Data show that the antibody TuBB-9 targeting a physiologically act pKi-67 but not with MIB-1, directed to a different epitope.
Literature	GENERIF_SUMMARY	Overexpression of KI-67 is associated with neuroendocrine carcino
Literature	GENERIF_SUMMARY	Data show that the prognostic abilities of TOP2A and MIB-1 were s better in ERG(-) patients.
Literature	GENERIF_SUMMARY	Negative EGFR, high Ki-67 proliferation index, and positive nm23-H be significantly predictive of a pathological complete response in b patients.
Literature	GENERIF_SUMMARY	study demonstrated a positive correlation between expressions of MCM-2 in ductal breast cancers; most intense expression of these was demonstrated in G3 grade cancers
Literature	GENERIF_SUMMARY	Ki67 after NAC, but not before NAC, is prognostic in breast cancer
Literature	GENERIF_SUMMARY	FOXP1 expression negatively correlated with Ki67 expression in cl cell carcinomas (p = .036).
Literature	GENERIF_SUMMARY	Data show that colocalization of CAIX, a membrane-bound hypoxi Ki-67, a nuclear proliferation marker were well analyzed with param of immunohistochemistry.
Literature	GENERIF_SUMMARY	The aim of this study was to investigate the correlation between th EZH 2 with p53 and Ki-67 expression and other clinicopathological primary breast carcinomas.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	A prognostic index in skin melanoma through the combination of r metalloproteinase-2, Ki67, and p53.
Literature	GENERIF_SUMMARY	In hormone-independent breast cancer cases negative for CerbB-2 expressions were significantly higher than that in hormone-depend negative cases.
Literature	GENERIF_SUMMARY	Triple negative breast cancer with high Ki-67 was associated with a aggressive clinical feature despite a higher pathologic complete re
Literature	GENERIF_SUMMARY	reduced immunostaining of PCNA and Ki67 and increased immun and p57 in the dexamethasone-induced intrauterine growth restric development compared to control placental development
Literature	GENERIF_SUMMARY	combined evaluation of RET and Ki-67 could act as an adjuvant pro useful for ameliorating the initial risk stratification of patients with thyroid carcinoma
Literature	GENERIF_SUMMARY	The purpose was to correlate markers expression of apoptosis (p5 and cell proliferation (Ki-67 and PCNA) with histological indicators severity.
Literature	GENERIF_SUMMARY	Nondysplastic leukoplakias without dual p53p16INK4a and triple p53p16INK4aKi-67 aberration seem to be harmless. Nondysplastic with these aberrations are at risk of progression, with very promisi predictive value.
Literature	GENERIF_SUMMARY	lack of Ki-67 proteins expression which may explain commonly kn activity of desmoid tumour cells
Literature	GENERIF_SUMMARY	findings indicate that miR-519d targets the MKi67 transcript and su cell growth, suggesting that miR-519d has a tumor suppressive rol HCC pathogenesis
Literature	GENERIF_SUMMARY	we investigated atypical tubal metaplasia in regard to its immunoe p53, Ki-67, and human telomerase reverse transcriptase (TERT), ar clinical outcome
Literature	GENERIF_SUMMARY	Both semiscale and fullscale quantitative analyses of Ki67/MART1 valuable diagnostic tools to distinguish melanomas and nevi with a of certainty.
Literature	GENERIF_SUMMARY	p53 inhibited Ki-67 promoter activity via p53- and Sp1-dependent p
Literature	GENERIF_SUMMARY	Findings define a MKI67, TOP2A, and E2F1 tri-marker proliferation possible utility for improved prognostication and treatment stratifi prostate cancer.
Literature	GENERIF_SUMMARY	For women
Literature	GENERIF_SUMMARY	Multiple sclerosis patients treated with interferon (IFN)-beta alone combination with low-dose prednisolone display increased propor natural killer (NK) cell subsets in the active phase of the cell cycle
Literature	GENERIF_SUMMARY	The aim of this work is to study the distribution of CD105 positive proliferation rate of endothelial cells from liver metastasis of colon based on double immunostaining CD105/Ki67.
Literature	GENERIF_SUMMARY	The prognostic value of the correlations between the expression o progesterone receptors, Ki67 expression and the histological stage the degree of myometrial and vascular invasion, was investigated.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	The study included a total of 28 cases of urothelial carcinomas, wh analyzed histopathologically and immunohistochemically using an c-erbB2 and Ki67 antibodies.
Literature	GENERIF_SUMMARY	Results indicate that carbonic anhydrase IX (CA IX) expression wa associated with patient age, T stage, grade, lymphovascular invasi expression, but was associated with p53 expression.
Literature	GENERIF_SUMMARY	p53 overexpression and increased Ki-67 proliferation index were m gastric pit dysplasia compared with adjacent gastric mucosa.
Literature	GENERIF_SUMMARY	The differential expression of DUSP6 correlated with Ki-67 index, s DUSP6 plays an important role in cancer resistance in different sul small cell lung carcinoma.
Literature	GENERIF_SUMMARY	Studies indicate that 48 biomarkers have been identified for ccRCC CD44, CA9, p53, Ki67 and PCNA have shown prognostic value, and VEGF-R2 are predictors of survival of pRCC patients.
Literature	GENERIF_SUMMARY	The expression levels of PCNA, Ki-67, Bcl-2 and mP53 are significa in the pterygium samples as compared to the normal conjunctiva s
Literature	GENERIF_SUMMARY	MIB-1 and p53 immunohistochemistry do not correlate with aggres the most common type of giant pituitary adenoma.
Literature	GENERIF_SUMMARY	immunohistochemical expressions of cyclin D1, p16(INK4a) and K analyzed in a total of 127 cases showing multinucleated giant cells
Literature	GENERIF_SUMMARY	Data suggest that there is no correlation between Ki-67 LI or cytok distribution pattern and MGMT immunoreactivity in GH-secreting p adenomas. These data may have implications in drug resistance re treatment with temozolomide.
Literature	GENERIF_SUMMARY	In grade 3 samples of CIN (Cervixal Intreapithelial Neoplasia), expr separately or combined with thymidine phosphorylase, seemed to progression to invasive cancer. Ki-67 levels were shown to correlat CIN grade.
Literature	GENERIF_SUMMARY	Data indicate that assessment of Ki67 in 200 tumor cells showed a average proliferation difference of 3.9% between core biopsies and samples with the core biopsies being the more proliferative sampl
Literature	GENERIF_SUMMARY	analysis of Aurora A and B kinases, Ki-67 and the estrogen and pro receptors from an endometriosis tissue microarray model
Literature	GENERIF_SUMMARY	Ki-67 labeling index after video-assisted thoracoscopic surgery seg was a prognostic factor of disease-free survival in stage I non-sma cancer.
Literature	GENERIF_SUMMARY	Parathyroid adenomas come into existence as a result of neoplast of PCNA/Ki-67 positive chief cells, especially vacuolated chief cell
Literature	GENERIF_SUMMARY	The Ki-67 index predicts progression-free survival of patients with differentiated neuroendocrine tumors.
Literature	GENERIF_SUMMARY	High Ki-67 expression correlates with morphologic features of agg and the expression pattern of Ki-67 correspond to the expected cy pattern in normal endometrium.
Literature	GENERIF_SUMMARY	The expressions of p16, Ki-67, and L1 proteins and human papillor were investigated as potential molecular markers for the diagnosis transforming potential of low cervical intraepithelial neoplasia

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	This study suggests that Ki67 is an independent predictor of poor patients with gastrointestinal stromal tumors.
Literature	GENERIF_SUMMARY	Proliferation marker Ki-67 is an independent prognostic indicator i patients.
Literature	GENERIF_SUMMARY	Low expression of Ki-67 was correlated with multiple lesions and h of KI-67 may be related with tumor across the midline in patients v glioblastoma multiforme.
Literature	GENERIF_SUMMARY	High-grade tumors overall showed significantly higher claudin-4 ar significantly lower claudin-7 expression when compared with low-g
Literature	GENERIF_SUMMARY	it was demonstrated that the region from -223 to +12 nt could drive transcription of the Ki-67 gene, and the Sp1 binding site is essentia transcriptional regulation of the Ki-67 gene.
Literature	GENERIF_SUMMARY	The combination of E-cadherin and Ki67 status might be a useful p marker indicating the need for adjuvant chemotherapy in Stage II T
Literature	GENERIF_SUMMARY	Ki-67 antigen,which reflects cell proliferation,was overexpressed in Barrett's epithelium samples.The finding of strong Ki-67 antigen ex long segments of metaplastic epithelium may be related to increas potential.
Literature	GENERIF_SUMMARY	The high-grade serous carcinomas had a significantly higher expre p53 and Ki-67, while low-grade carcinomas had a significantly high level of Pax2.
Literature	GENERIF_SUMMARY	Ki-67 staining percentage may represent a marker to select hepato carcinoma patients for liver transplantation and to follow posttrans recurrence.
Literature	GENERIF_SUMMARY	Ki-67 and p53 staining may be helpful adjuncts in the differential d Necrotizing sialometaplasia from squamous cell carcinoma in app setting.
Literature	GENERIF_SUMMARY	Ki-67, but not proliferating cell nuclear antigen, has a role in poor d survival in patients with adenocarcinoma of the lung
Literature	GENERIF_SUMMARY	Ki67 was a significant predictor of disease progression and should in determining appropriate treatments and planning follow-up for p neuroendocrine tumors.
Literature	GENERIF_SUMMARY	Ki-67 overexpression (Ki-67 >/= 10%) was identified as an indepen factor for distant-metastatic-free survival (DFS) and overall surviva univariate and multivariate analysis.
Literature	GENERIF_SUMMARY	Report apoptosis along with BCL-2 and Ki-67 expression in individu intestinal metaplasia, chronic active gastritis, and normal gastric m
Literature	GENERIF_SUMMARY	There was a significant relation between CD10 expression as well labelling index and recurrence of ameloblastoma
Literature	GENERIF_SUMMARY	Increased Ki-67 and bcl-2 expression plus density of new vessels a prognostic value for the disease outcome in medulloblastoma pati
Literature	GENERIF_SUMMARY	In anaplastic oligodendroglial tumours the Ki67 index has a strong impact on univariable analysis.
Literature	GENERIF_SUMMARY	Low MGMT expression and high Ki-67 expression were more frequ oligodendrogliomas than in astrocytomas and mixed tumors.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Women with a HER2/neu/neu+/Ki67+ ductal carcinoma in situ (DC higher risk of developing DCIS local recurrence after breast-conser
Literature	GENERIF_SUMMARY	Ki-67 expression was higher in patients with high-grade endometri
Literature	GENERIF_SUMMARY	analysis of P53, p16 and Ki67 immunoexpression in oral squamou
Literature	GENERIF_SUMMARY	analysis of P53, Bcl-2 and Ki67 immunoexpression in follicular soli ameloblastomas
Literature	GENERIF_SUMMARY	Determination of the expression of Ki-67 in an attempt to establish and/or associations with clinical characteristics of patients with n pituitary adenomas.
Literature	GENERIF_SUMMARY	HIF-2a was correlated with ABCG2 expression, histology-grade and expression in breast invasive ductal carcinoma
Literature	GENERIF_SUMMARY	High MIB1 proliferation index trended toward worse response to a radiotherapy in gliomas.
Literature	GENERIF_SUMMARY	Osteopontin expression seems to be related to Ki-67 (proliferation CD34 expression (angiogenesis marker) confirming its role in the p psoriasis.
Literature	GENERIF_SUMMARY	lower levels of Ki-67 and PCNA expression were measured in cellu than in leiomyosarcomas
Literature	GENERIF_SUMMARY	Ki-67 staining, a reflection of tumor-proliferation activity, was signi to the FDG uptake value in lymphoma lesions.
Literature	GENERIF_SUMMARY	Higher expression of Smac and Ki-67 appear to play a role in the p pancreatic cancer. Combined detection of these proteins may imp prognostic evaluation of this disease.
Literature	GENERIF_SUMMARY	The expression of MIB-1 was statistically elevated in higher-grade patients 25 years or younger and in tumors located in extremities a lesions.
Literature	GENERIF_SUMMARY	Scoring of the percentage of Ki67 expressing cell nuclei is diagnos and reproducible in terms of delineating flat lesions of the bladder
Literature	GENERIF_SUMMARY	These findings do not provide best evidence for the routine use of prognosticate overall short-term survival in patients with pulmonar tumors.
Literature	GENERIF_SUMMARY	Stromal Ki-67 expression was statistically correlated with epithelia epithelial AR, and stromal ER-beta expression.
Literature	GENERIF_SUMMARY	upregulated CENP-H and Ki67 levels are significantly associated w relapse-free survival in hypopharyngeal squamous cell carcinoma factors may be predictors of a relapsing phenotype in HSSC cases
Literature	GENERIF_SUMMARY	It is thought that Ki-67 and AgNOR might be helpful as a prognosti recurrences of keratocystic odontogenic tumors.
Literature	GENERIF_SUMMARY	Data show that the high sensitivity of p16/Ki-67 with significantly i specificity compared with HPV testing makes p16/Ki-67 a viable o grade squamous intraepithelial lesion (LSIL) triage.
Literature	GENERIF_SUMMARY	MIB-1 labeling index predominantly affected the pattern of failure i patients treated with a multimodal approach

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with response to therapy in liv malignancies.
Literature	GENERIF_SUMMARY	The results of this study indicated that the higher MCM-2 expressi the KCOT might be related to the inflammation and this protein mi sensitive to inflammation.
Literature	GENERIF_SUMMARY	The combination of Galectin-3 and PTTG is complementary as a d biomarker for patients with papillary carcinoma.
Literature	GENERIF_SUMMARY	Over-expression of Ki-67 proliferation protein was found in 62% of carcinoma cases.
Literature	GENERIF_SUMMARY	High Ki67 expression in the primary tumor is an independent adve factor in metastatic breast cancer.
Literature	GENERIF_SUMMARY	Results suggest that rigorous comparisons of measurement appro be applied in a tumour-type and treatment-specific manner to enha application of Ki67 assessment.
Literature	GENERIF_SUMMARY	FDG uptake in malignant melanoma is determined by GLUT-1 and HK-2 and Ki-67 play no role in FDG uptake of malignant melanoma
Literature	GENERIF_SUMMARY	The Ki-67 proliferative index was high in 4 of 15 (27%) carcinomas adenomas, and none of the hyperplasia or normal tissues.
Literature	GENERIF_SUMMARY	Although there is no difference in overall survival between primary with less than or at least 10% Ki-67 expression, there is significant survival when Ki-67 expression in lymph nodes is less than 10%.
Literature	GENERIF_SUMMARY	Ki-67 expression revealed pituitary adenomas with progression ha proliferation index of 3.66 +/- 3.00%, which was significantly highe without progression of 1.89 +/- 1.25% (mean +/- SD, n = 27) (p
Literature	GENERIF_SUMMARY	Tumor sizes and high Ki67 PI in the invasive front appear to be imp prognostic tools for head and neck squamous cell carcinoma.
Literature	GENERIF_SUMMARY	The data show that p16(INK4a) /Ki-67 co-expression allows most to be detected initially and in the follow-up period
Literature	GENERIF_SUMMARY	We quantified the spatial distribution of Ki67-expressing cells with epithelium in a large cohort of 613 lung biopsy sections
Literature	GENERIF_SUMMARY	Digital image analysis and manual counting are the acceptable sta assessment in neuroendocrine tumors of the gastroenteropancrea
Literature	GENERIF_SUMMARY	Decreasing expressions of HIF-1alpha, Bcl-2 and Ki-67 were obser chemoradiotherapy of recel cancer, whereas GLUT-1 overall was u
Literature	GENERIF_SUMMARY	A moderate correlation between the stage of endometriosis and th Ki-67 staining was found.
Literature	GENERIF_SUMMARY	A positive correlation between MT and Ki-67 expression was obse studied cases but was even stronger in the metatypic subtype of b carcinoma.
Literature	GENERIF_SUMMARY	High Ki67 protein expression was an adverse prognostic factor for risk early breast cancer patients
Literature	GENERIF_SUMMARY	In untreated patients, Ki67, TOP2A, and RacGAP1 are significant a prognostic markers.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	A predictive impact of ki-67 to treatment in non-small cell lung can unclear. [review]
Literature	GENERIF_SUMMARY	The Ki67 marker was more important factor than P53 protein in pr breast cancer patients in Iran.
Literature	GENERIF_SUMMARY	Disease free survival was significantly different between patients w was reduced and those not reduced for Luminal B, triple negative a negative breast cancer.
Literature	GENERIF_SUMMARY	Ki67 overexpression is associated with central nervous system me operable breast cancer.
Literature	GENERIF_SUMMARY	Assessment of the Ki67 index remains a useful prognostic tool if a performed in the relapse situation.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with proliferation in salivary gl
Literature	GENERIF_SUMMARY	Data indicate that higher number of intratumoural FOXP3+ Tregs w with a more advanced tumour stage and higher Ki67 labelling inde
Literature	GENERIF_SUMMARY	KI67 antigen is significantly higher in young breast cancer patients ones
Literature	GENERIF_SUMMARY	Polyps in postmenopause have a high COX-2 expression that is hig malignant polyps than in benign polyps; there was no difference in expression between malignant polyps and premalignant/malignan
Literature	GENERIF_SUMMARY	Livin expression was positively correlated with the proliferation ma was negatively correlated with caspase-3 expression in human am carcinoma patients.
Literature	GENERIF_SUMMARY	Ki-67 may be a good marker of biological aggressiveness useful in patients for more aggressive treatment in prostate cancer
Literature	GENERIF_SUMMARY	Ki-67 presents a significant prognostic factor for gastrointestinal s recurrence.
Literature	GENERIF_SUMMARY	distribution pattern of Ki67 may be a new independent prognostic cancer
Literature	GENERIF_SUMMARY	Ki-67 overexpression was a positive predictor of survival in colorec metastases.
Literature	GENERIF_SUMMARY	Ki67 gene up-regulation was noted at the donor site wound healing
Literature	GENERIF_SUMMARY	In multivariate analysis, Ki67 remained the only significant prognos subgroups of ER positive HER2 negative early-stage (pT1-pT2, pN0
Literature	GENERIF_SUMMARY	The assessment of syndecan-1 and Ki-67 expression in skin biops tool for differentiating keratoacanthoma and squamous cell carcin
Literature	GENERIF_SUMMARY	Phosphohistone-H3 and Ki-67 immunostaining in cutaneous pilar l leiomyosarcoma (atypical intradermal smooth muscle neoplasm).
Literature	GENERIF_SUMMARY	Grade according to Ki-67 was a better prognostic marker than mito metastatic pancreatic and midgut neuroendocrine neoplasms.
Literature	GENERIF_SUMMARY	The results predict that an increased level of MIB-1 (Ki-67) express tissue, points to a less favourable outcome for ovarian cancer pati
Literature	GENERIF_SUMMARY	In patients with breast cancer neoplasm grade and expression of E Lys27H3, vimentin, and KI67 can be used to predict BRCA1 germlin

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Ki-67 protein may be regulated by specific miRNA-mRNA interactic contribute to the proliferation of glioma cells.
Literature	GENERIF_SUMMARY	Ki-67 was associated with common histopathological parameters, to be an independent prognostic parameter for DFS and OS in brea patients.
Literature	GENERIF_SUMMARY	Carbonic anhydrase II and Ki-67 are significant prognostic factors gastrointestinal stromal tumors.
Literature	GENERIF_SUMMARY	p53 and Ki-67 markers are associated with cell proliferation in chil Eosinophilic esophagitis but do not represent a premalignant (dysp condition of the esophagus
Literature	GENERIF_SUMMARY	In HMG tissues, Ki-67-positive cells were scattered in both the gray matter, with a significantly higher Ki-67 labeling index in the white compared with gray matter.
Literature	GENERIF_SUMMARY	Using Ki-67, Bcl-2, p53, and Cox-2 markers, we managed to disting disordered proliferative/benign hyperplastic endometrium endome grade endometrial carcinoma.
Literature	GENERIF_SUMMARY	Through phenotype analysis, non-hepatitis A virus-specific antibod cells (ASCs) are found to be Ki-67lowCD138highCD31highCD38hig with hepatitis A virus-specific ASCs.
Literature	GENERIF_SUMMARY	Ki-67 and p53 might have influence on meningioma development a
Literature	GENERIF_SUMMARY	Ki-67 and matrix metalloproteinase-9 expression in the follicular cy odontogenic tumor, and ameloblastoma
Literature	GENERIF_SUMMARY	Posttreatment Ki67 levels provide prognostic information for patie hormone-receptor-positive breast cancer and residual disease afte chemotherapy.
Literature	GENERIF_SUMMARY	Report prognostic value of Ki-67 determination in fine needle aspir tumors.
Literature	GENERIF_SUMMARY	Immunohistochemical expression of Ki67 was not associated with of invasive breast cancers.
Literature	GENERIF_SUMMARY	Ki67 levels were altered after neoadjuvant chemotherapy in breast
Literature	GENERIF_SUMMARY	Expression of Ki67 was correlated with grade and stage in bladder
Literature	GENERIF_SUMMARY	Ki-67 overexpression was detected in the epithelial component in (42%) Carcinosarcoma cases, and in the mesenchymal componen (36%) Carcinosarcoma neoplasms.
Literature	GENERIF_SUMMARY	MIB-1 is an important clinical marker in astrocytomas. Nevertheles shortcomings make it difficult to standardize a proliferation index purposes.
Literature	GENERIF_SUMMARY	cell cycle related proteins PCNA, Ki67, cyclin D3, p27 and p57 were both normal and diabetic human term placentas.
Literature	GENERIF_SUMMARY	Ki67 was significantly linked to prognosis in uni- and multivariate a complete cohort and in HR-positive, but not triple-negative breast c
Literature	GENERIF_SUMMARY	We demonstrate that axillary lymph node involvement and high exp Ki-67 are associated with poorer prognosis.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Report Ki67 expression in gastrointestinal stromal tumors.
Literature	GENERIF_SUMMARY	in oral squamous cell carcinoma cohort, below-median Ki67 and to sCAIX expression (Ki67(lo)sCAIX(hi)) were associated with signific disease-specific survival in univariate and multivariate analyses
Literature	GENERIF_SUMMARY	Ki-67 staining is stronger than PHH3, making 'hot spots' easier to i Ki-67 is more ideal than PHH3(Phosphohistone H3) for staining N in tumors with borderline grades.
Literature	GENERIF_SUMMARY	Tonsil surface epithelium is a useful control for monitoring Ki-67 st Achieving optimal staining results could minimize variations in Ki-6 differences in the staining methods used by different laboratories.
Literature	GENERIF_SUMMARY	Ki67 index, TP53 status, or PR negativity can be used to segregate HER2-negative tumors into prognostically meaningful subgroups w different clinical outcomes.
Literature	GENERIF_SUMMARY	Ki-67 and p53 expression rates were statistically significantly corre grade of mammary phyllodes tumors.
Literature	GENERIF_SUMMARY	A significant positive correlation was found between age and PCN values (P
Literature	GENERIF_SUMMARY	p16/Ki-67 immunostaining demonstrated better accuracy than hur papillomavirus for detecting cervical intraepithelial neoplasia 2 or l patients with atypical squamous cells
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with uterine cervical neoplasm
Literature	GENERIF_SUMMARY	The integration of the immunocytochemical score of Ki-67 and p53 cytological score resulted in a final score that was also diagnostica
Literature	GENERIF_SUMMARY	Suggest that Ki67 labeling be performed on all pancreatic neuroen in addition to mitotic rate determination to define more accurately and prognosis.
Literature	GENERIF_SUMMARY	p16(INK4a)/Ki-67 and L1 capsid protein immunostaining and HR-H residual LBCS diagnosed with ASC-H or LSIL-H are useful objective predicting CIN2+.
Literature	GENERIF_SUMMARY	The imunohistochemical staining for topoisomerase IIalpha expres useful as additional marker for mucosal dysplasia degree determin
Literature	GENERIF_SUMMARY	Studied cohort of breast cancer patients that received neoadjuvant between 2002 and 2010. Ki-67 expression was measured using qu immunofluorescence automated quantitative analysis (AQUA) tech
Literature	GENERIF_SUMMARY	Ki-67 and p53 expression varies in morphologically normal tubal ep depending on age and BRCA1 mutation status.
Literature	GENERIF_SUMMARY	Apparent diffusion coefficient values of prostate cancer could refle proliferative activity and differentiated degree of prostate cancer.
Literature	GENERIF_SUMMARY	High Ki67 expression is predictive of poor prognosis and of resista tamoxifen therapy in postmenopausal BC. We recommend conside one of the risk factors that guide adjuvant treatment decisions.
Literature	GENERIF_SUMMARY	HPV genotyping, p16/Ki-67 dual staining and koilocytic morpholog in the prediction of clinical outcome in women initially diagnosed w squamous intraepithelial lesion cytology.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Immunofluorescence expression (IFE) of cell cycle regulators p16i in association with Ki-67, p53 in cell cultures, obtained from differe cervical intra-epithelial lesions, was assessed.
Literature	GENERIF_SUMMARY	The aim of the present study was to evaluate the prognostic value proliferation factors mitotic activity index (MAI), phosphohistone H cyclin B1, cyclin A and Ki67, alone and in combinations.
Literature	GENERIF_SUMMARY	Significant relationship between Ki-67 proliferation index and CDC2 index in astrocytomas.
Literature	GENERIF_SUMMARY	KB cell Ki-67 mRNA significantly increased after doxorubicin treatr
Literature	GENERIF_SUMMARY	Data indicate that Ki-67 positivity is prognostic for predicting non-r bladder cancer (NMIBC) recurrence and progression.
Literature	GENERIF_SUMMARY	CISH would help to distinguish episomal from integrated HPV, in o conclusions regarding the prognosis of the lesion, while p16(INK4a stain approach would confer a high level of standardization to the procedure
Literature	GENERIF_SUMMARY	importance of temporal-spatial expression of Ki-67 for the thickeni cranial spinal cord lateral wall
Literature	GENERIF_SUMMARY	It is a tumor marker for proliferation and invasion.
Literature	GENERIF_SUMMARY	Ki67 immunostaining provides prognostic information beyond trad assessed clinicopathological variables in node-negative breast car
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with breast cancer.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with breast cancer.
Literature	GENERIF_SUMMARY	In residual pituitary adenomas with a Ki-67 Li above 2.2%, regrowth expected, and these tumors may require shorter intervals of follow resonance imaging (MRI) and/or early adjuvant therapy.
Literature	GENERIF_SUMMARY	Each 1% increase in Ki-67 expression was associated with a 12% i prostate cancer-specific death.
Literature	GENERIF_SUMMARY	Data suggest that Ki-67 status was an independent prognostic fac metastasis free survival (MFS) and overall survival in breast cance 1-3 positive nodes.
Literature	GENERIF_SUMMARY	The automatic Ki-67 score is very accurate compared with patholo estimation.
Literature	GENERIF_SUMMARY	Expression of beta-catenin, Ki67 and Her-2/Neu was studied in gas protein expression of beta-catenin, Ki67 and Her-2/Neu had close with the clinical pathologic characteristics of gastric cancer.
Literature	GENERIF_SUMMARY	Ki-67 overexpression suggests that cytotoxic agents are effective
Literature	GENERIF_SUMMARY	Ki67 index, MVD and HIF-1 alpha index have promising prognostic and should be validated in further studies.
Literature	GENERIF_SUMMARY	Investigated expression of a pathological biomarker, Ki-67, in addit the usefulness of (18)F-FDG PET-CT in the pretreatment evaluation epithelial neoplasms.
Literature	GENERIF_SUMMARY	Data observed that MIB-1 is a reliable independent prognostic indi negative infiltrating ductal carcinoma in this study.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	the prognostic significance of Ki-67 expression varied in different s lymphoma and in DLBCL and MCL after the introduction of rituxim valuable for clinical decision-making and individual prognostic eva
Literature	GENERIF_SUMMARY	Ki67 expression increases in gliomas.
Literature	GENERIF_SUMMARY	It has found a statistically significant correlation between the expr markers and WHO grade and recurrence.
Literature	GENERIF_SUMMARY	Ki-67 is an independent prognostic marker for patients with epithe carcinoma
Literature	GENERIF_SUMMARY	p16/Ki-67 dual-stained cytology may provide a valuable novel appr pregnant women with persistent or progressing cervical intraepith disease
Literature	GENERIF_SUMMARY	Data provide methodology to achieve accurate immunohistochem labelling index (Ki67 LI) reflects proliferative activity estimation by image analysis (DIA) approach.
Literature	GENERIF_SUMMARY	Establishment of Ki67, p53 and CD34 expression in human tooth b stages of odontogenetic development.
Literature	GENERIF_SUMMARY	The significant positive correlations were observed between Bcl-2 Ki-67 antigen in sinonasal inverted papilloma and sinonasal carcin and between Bcl-2 protein and CD34 antigen in the chronic rhinosi
Literature	GENERIF_SUMMARY	This study attests the reproducibility of Ki67-LI of pNETs whether c cytology or histology with a very good inter-observer correlation
Literature	GENERIF_SUMMARY	data shows that the Ki-67 value predicts the response to neoadjuv chemotherapy as a function of the molecular subtype
Literature	GENERIF_SUMMARY	confirms that the deregulated cell cycle, as demonstrated by p16/k labelling, is independent of the oncogenic action of hr-papillomavir grade urothelial proliferations
Literature	GENERIF_SUMMARY	Ki-67 was expressed in 13 (65%) tumors, and negatively correlated prognosis, histologic type and stage of Neuroblastoma
Literature	GENERIF_SUMMARY	Basal cell carcinoma exhibited a wide variation of Ki67 proliferativ ranging from 1% to 61%, among the histological subtypes
Literature	GENERIF_SUMMARY	PCNA and Ki67 staining intensities significantly increased in villou significantly decreased in basal plates of Preeclampsia group
Literature	GENERIF_SUMMARY	Ki-67 is a protein phosphatase 1-interacting protein that organises chromosome periphery.
Literature	GENERIF_SUMMARY	In triple negative breast cancer, Ki67 expression was correlated wi margin and intensive posterior echo with spotty flow imaging by ul
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with dysplastic areas of large hyperplastic polyps.
Literature	GENERIF_SUMMARY	Report counting strategy which acknowledges small highly prolifer of Ki67 expression in breast neoplasms.
Literature	GENERIF_SUMMARY	The level of Ki-67 may be considered one of the valuable biomarke cancer patients with respect to process and recurrence.
Literature	GENERIF_SUMMARY	Ki67 index is an effective complementation of modified NIH criteri

Main Category	Category	Terms
		the prognosis of gastrintestinal stromal tumors, and Ki67 index>8 unfavorable factor for imatinib adjuvant therapy.
Literature	GENERIF_SUMMARY	Notch-1 and Ki-67 expression may act as promising predictors for and prognosis of Kimura's disease.
Literature	GENERIF_SUMMARY	Ki-67 may be a useful marker for discriminating benign from borde p16 and pRb may be a useful combination of markers for distingui benign PTs in core biopsy specimen.
Literature	GENERIF_SUMMARY	Although no obvious defects in the progression of mitosis were ob timing of dephosphorylation of the mutant Ki67 in anaphase was c indicating that Ki67 itself is one of the substrates of PP1gamma-K
Literature	GENERIF_SUMMARY	Letter
Literature	GENERIF_SUMMARY	this is a first study showing the correlation of Ki-67, PCNA interact might lead to malignant transformation.
Literature	GENERIF_SUMMARY	Expression of miR-34a was lower in metastases than in primary tu inversely correlated with expression of cyclin D1 and Ki-67
Literature	GENERIF_SUMMARY	high Ki-67 labeling index is predictive sign of a poor prognosis in w resistant prolactinoma.
Literature	GENERIF_SUMMARY	Ki-67 may be a useful diagnostic adjunct in irradiated breast
Literature	GENERIF_SUMMARY	In multiple pathology specimens of primary neuroendorine tumors metastases, Ki67 expression varied over the course of the disease
Literature	GENERIF_SUMMARY	elevated protein expression of p53, cyclin D1 and Ki-67 in the seco malignancy is an indicator of more aggressive malignant behavior secondary tumor
Literature	GENERIF_SUMMARY	In uterine cervical lesions, Ki-67 expression was observed in high g carcinoma
Literature	GENERIF_SUMMARY	In this study, PR negativity was a prognostic factor for early relaps HER2-negative breast cancer, while a high Ki-67 index suggested a early relapse.
Literature	GENERIF_SUMMARY	Data suggest Ki-67 labeling index (immunochemical detection of M most useful prognostic marker for thyroid follicular cell tumors an objective parameter to stratify patient prognosis postoperatively. [
Literature	GENERIF_SUMMARY	Our data showed that Ki-67 is an effective prognostic indicator of s ENKTL patients. This prognostic index may be helpful in identifying patients with ENKTL.
Literature	GENERIF_SUMMARY	an association between the vimentin and ki-67 expression levels a of spread of metastatic renal cell carcinoma was identified
Literature	GENERIF_SUMMARY	Ki67 was highly expressed in five cases of primary mucosal malign while the expression in the other five cases was low. There was no between the Ki67 expression and prognosis.
Literature	GENERIF_SUMMARY	The results emphasize the potential of Ki-67 and p53 as biomarker carcinogenesis in oral squamous cell carcinoma.
Literature	GENERIF_SUMMARY	This study demonstrates that, Ki-67 LI serves as an important prog in human astrocytomas.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	No significant association was observed between Ki-67 expressior tumor thickness, Clark level, or lymph node involvement.
Literature	GENERIF_SUMMARY	Ki-67 index is a useful diagnostic marker for neuroendocrine tumo
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with gastric cancer.
Literature	GENERIF_SUMMARY	Ki67 index is a valuable IHC marker to distinguish indolent from ag lymphomas especially in small needle biopsies where exact typing possible.
Literature	GENERIF_SUMMARY	Circulating tumor cells expression of ER, BCL-2, HER2, and Ki67 ca reproducibly measured with high analytical validity using the CellS
Literature	GENERIF_SUMMARY	it is not possible to relate the increased Ki67 index of villous troph last trimester in preeclampsia with one cause
Literature	GENERIF_SUMMARY	-67 proliferation index levels were higher in patients with legal abo patients with placenta percreta or noncomplicated cesarean delive
Literature	GENERIF_SUMMARY	Suggest that Ki67 expression in HR-negative breast cancer may im assessment of pathological response after neoadjuvant chemothe
Literature	GENERIF_SUMMARY	High Ki67 Expression is associated with Oropharyngeal Squamous Carcinoma.
Literature	GENERIF_SUMMARY	Computerized image analysis of the Ki-67 index in MCL is an attra to semiquantitative estimation and can be introduced easily in a ro diagnostic setting for risk stratification in MCL.
Literature	GENERIF_SUMMARY	We found that Ki-67 expression in pathologic BM is a novel signific parameter of worse prognosis in DLBCL patients with BMI in the ri
Literature	GENERIF_SUMMARY	increased expression of Cyclin D1 and Ki-67 in patients with endor carcinoma relative to proliferative endometrium and simple hyperp D1 expression had a positive correlation with Ki-67 expression.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with prostate carcinomas.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with hypopharyngeal cancers.
Literature	GENERIF_SUMMARY	No associations were found between Ki67 expression and histolog breast neoplasms.
Literature	GENERIF_SUMMARY	The present study shows that high Ki-67 LI is a risk factor for poor elderly age group and non-GCB subtype in patients with DLBCL tre CHOP.
Literature	GENERIF_SUMMARY	Elevated Ki67 and geminin expression distinguish a fraction of me carcinoma with worse prognosis.
Literature	GENERIF_SUMMARY	Ki-67 and PgR expression were independently associated with the patients with ER+ and HER2- breast cancer
Literature	GENERIF_SUMMARY	Results show that patients with Ki67 higher than median had signi median overall survival than those with low Ki67 suggesting its pro epithelioid malignant pleural mesothelioma.
Literature	GENERIF_SUMMARY	The combined Ki67/Bcl2 phenotyping was useful in luminal cance prognostication. It also refined prognostication in intermediate gro and stage 2 cancers) of luminal cancers; and aided in further class luminal A cancers.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	In ACC, the proliferation marker Ki-67 was associated with significa overall and disease-free survival.
Literature	GENERIF_SUMMARY	Visual assessment of Ki67 in breast neoplasms by a 5-grade scale easy, and reliable with acceptably low interobserver and intraobser
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with glioma.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with recurrence in invasive du cancer.
Literature	GENERIF_SUMMARY	Immunohistochemistry analysis shows that co-expression of p16, COX-2 is associated with basal phenotype in high-grade ductal car of the breast.
Literature	GENERIF_SUMMARY	Ki-67 is a strong predictor of central nervous system relapse in pat mantle cell lymphoma.
Literature	GENERIF_SUMMARY	High expression of ki67 is associated with papillary thyroid microc
Literature	GENERIF_SUMMARY	High KI-67 antigen expression is associated with Malignant Perito Mesothelioma.
Literature	GENERIF_SUMMARY	Neoadjuvant chemotherapy reduces Ki67 expression in breast can
Literature	GENERIF_SUMMARY	Studied the prognostic value of Ki-67, p27, and p53 immunoreactiv gastric cancer.
Literature	GENERIF_SUMMARY	Ki-67/MIB-1 had a prognostic value for overall survival in cervical c
Literature	GENERIF_SUMMARY	There was a statistically significant difference between well-differe endocrine neoplasm (Grade I), and well-differentiated endocrine ca (WDEC, Grade II) and PDEC for Ki-67 >20% (p
Literature	GENERIF_SUMMARY	negative-ER ,negative-PR ,positive-p53 and higher Ki-67 expression to a higher mtDNA copy ratio. In addition, advanced T-status (p=0.0 negative-HER-2/neu expression (p=0.061) were associated with m mutations
Literature	GENERIF_SUMMARY	Among women with SA, increased Ki-67 expression in either the SA normal background terminal duct lobular units carried an approxim increased odds of subsequent breast cancer as compared to lowe expression.
Literature	GENERIF_SUMMARY	It was indicated, that index of maturation correlates with level of b cells, containing Philadelphia chromosome, so as with percentage Bcl-2+, Pgp-170+ and Ki67+ cells in bone marrow of CML patients.
Literature	GENERIF_SUMMARY	Visual assessment of Ki67 labeling index at 10 % intervals is a can standard method in breast cancer clinical practice.
Literature	GENERIF_SUMMARY	Treg cell level was related to Ki-67 expression and immune status lung cancer patients.
Literature	GENERIF_SUMMARY	Ki-67 index is a useful parameter for more objective grading and ca prognostic value to determine the biological potential of preneopla lesions.
Literature	GENERIF_SUMMARY	Evaluating p16 and Ki-67 in combination would be significant in pr recurrence in pTa and pT1 urothelial carcinomas.
Literature	GENERIF_SUMMARY	Univariate analyses showed that AR, CDH1 and Ki-67 expression le

Main Category	Category	Terms
		significantly associated with overall survival in triple-negative brea patients.
Literature	GENERIF_SUMMARY	Ki67 expression varied in breast cancer patients according to men phase.
Literature	GENERIF_SUMMARY	Suggest Ki-67 proliferative index as an important prognostic facto recurrence and death in prostate cancer post prostatectomy.
Literature	GENERIF_SUMMARY	AR is not associated with the proliferation marker Ki-67, but is ass lower 21-gene recurrence score in estrogen receptor positive/HER breast cancer.
Literature	GENERIF_SUMMARY	Suggest that Ki67 expression correlates with prognosis in gastroin tumors.
Literature	GENERIF_SUMMARY	The present work aims to investigate the relationship between the AEG-1(astrocyte elevated gene-1), b-FGF(basic-fibroblast growth fa catenin, Ki-67, TNF-alpha (tumor necrosis factor-alfa) other progno in DC (Ductal Carcinomas) and ductal intraepithelial neoplasm. We relationship between these factors.
Literature	GENERIF_SUMMARY	High ki-67 expression is associated with poor Response to chemo Patients with Nasal Natural Killer/T-cell Lymphoma.
Literature	GENERIF_SUMMARY	Suggest MIB-1 index may be used as a surrogate for the mitosis-ka index in neuroblastoma.
Literature	GENERIF_SUMMARY	elevated Ki-67 expression was associated with a poorer outcome i patients, particularly in early-stage Asian ADC patients.
Literature	GENERIF_SUMMARY	Data suggest that tumor expression of Ki67 and GRIM19 (cell deat protein GRIM19) correlate with malignancy in thyroid Hurthle cell (variable expression of Ki67 and GRIM19 may helped differentiate H from HC adenoma.
Literature	GENERIF_SUMMARY	high expression of VEGF and Ki-67 were independent poor progno overall survival in adenoid cystic carcinoma
Literature	GENERIF_SUMMARY	Region of interest size is a critically important parameter for semia quantification by computer-assisted image analysis in breast canc
Literature	GENERIF_SUMMARY	Suggest that clinical course of KI67/Her2 positive breast cancer w by hormone receptor status rather than Ki67 expression.
Literature	GENERIF_SUMMARY	Both p16/Ki-67 double staining and the hr-HPV DNA test had simil in predicting high-grade cervical intraepithelial neoplasia among w US
Literature	GENERIF_SUMMARY	Overexpression of p53 and Ki-67 could be used to discriminate low subtype in breast cancer.
Literature	GENERIF_SUMMARY	Suggest that combination of Bcl-2 and Ki67 to detect gastric canc effective.
Literature	GENERIF_SUMMARY	Our data support Ki67 evaluation to estimate non-small-cell lung c patients' prognosis, particularly for adenocarcinoma.
Literature	GENERIF_SUMMARY	High Ki67 expression in colorectal cancer is associated with good outcome.
Literature	GENERIF_SUMMARY	Her2 and Ki67 expression predicts recurrence of ductal carcinoma

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Classification based on IDH1/2 mutation status and Ki-67 express be more convenient for clinical application and guide personalized malignant gliomas.
Literature	GENERIF_SUMMARY	The expression of Ki67 and FasL is closely related to the occurren development of cervical carcinoma.
Literature	GENERIF_SUMMARY	Tissue handling of surgical specimen is critical for assessing the K surgical removed breast cancer.
Literature	GENERIF_SUMMARY	COX-2 and Ki-67 are abundantly expressed in hepatolithiasis and b carcinoma tissues and may play an important role in the disease o progression, and metastasis
Literature	GENERIF_SUMMARY	the expression of PSMB4 was significantly increased in the human ovarian cancer)tissues and was correlated with Ki-67.
Literature	GENERIF_SUMMARY	Ki-67 Expression in CRC Lymph Node Metastasis Does Not Predict
Literature	GENERIF_SUMMARY	High expression of Ki-67 is associated with a higher tumor grade a mortality.
Literature	GENERIF_SUMMARY	We observed an association of higher Ki-67 expression with Gleas National Comprehensive Cancer Network risk (P=0.013) and PSA r
Literature	GENERIF_SUMMARY	High-grade cell proliferation, as measured by Ki-67 labeling index, s correlated with an tumor infiltration pattern that indicated a more a squamous cell carcinoma.
Literature	GENERIF_SUMMARY	we propose that PELP1/Ki-67 double high expression in tumors is prognostic factor for predicting a poor outcome for patients with T breast cancer
Literature	GENERIF_SUMMARY	Intraobserver reproducibility ranged from very good to perfect for b Our results suggested that specimen-specific cut-off value should both scoring methods
Literature	GENERIF_SUMMARY	Findings indicate that high Ki67 antigen/ SATB1 protein ratio was a predictor of worse overall survival (OS), while Ki67 and SATB1 did r statistical significance as single predictors.
Literature	GENERIF_SUMMARY	oligonucleotide-mediated treatment could be more effective when towards KI-67 and BCL2. This could be accomplished by dual mon targeting KI-67 and BCL2
Literature	GENERIF_SUMMARY	EZH2 is highly expressed in pituitary adenomas, positively associa expression, and associated with proliferation.
Literature	GENERIF_SUMMARY	Ki-67 index >/= 4% was found significantly associated with the sur pseudopapillary tumor of the pancreas.
Literature	GENERIF_SUMMARY	No significant relationship observed between Ki-67 immunostainin intraephitelial neoplasia grade in pregnancy
Literature	GENERIF_SUMMARY	Genomic Grade Index or centrally reviewed Ki67 significantly impro clinicopathological models to determine distant recurrence of brea
Literature	GENERIF_SUMMARY	he expression pattern of ERCC2 was correlated with the expressio Ki-67 and a significant correlation (r = 0.230, p
Literature	GENERIF_SUMMARY	correlation between TGFb1 and pSmad2 as well as between HtrA1 the very significant increase of Ki67 in stromal compartment of eu

Main Category	Category	Terms
		endometrium suggest a possible involvement of HtrA1 in the path endometriosis.
Literature	GENERIF_SUMMARY	These results indicated there might be a tight correlation among M hepatocellular carcinoma prognosis
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with breast cancer.
Literature	GENERIF_SUMMARY	the markers Ets-1 and APAF-1 relative to p53, Ki-67 and PTEN expr adenomas/polyps, were investigated.
Literature	GENERIF_SUMMARY	The investigation of p53 and Ki-67 expression is useful in identifyi proliferative forms of salivary tumors, with aggressive potential of
Literature	GENERIF_SUMMARY	The use of basal Ki67 status as a predictive factor of chemotherap facilitate the identification of a patient subpopulation with high pro achieving pathological complete response when treated with prim chemotherapy, and thus with a potentially favorable long-term prog
Literature	GENERIF_SUMMARY	Analysis revealed linearly independent intrinsic factors of the Ki67 immunohistochemistry variance
Literature	GENERIF_SUMMARY	We also found that Fibulin-5 reduces the level of expression of Ki-6 protein associated with cell proliferation. Moreover, reduction in Fi expression corresponds to an increase of Ki-67 detection in breast
Literature	GENERIF_SUMMARY	High Ki-67 LI were associated with retroperitoneal soft tissue sarc
Literature	GENERIF_SUMMARY	Our results suggest that the ADC values on DW-MRI may be used a measurement of cell proliferation and hypoxia in RC. J. Magn. Res 2016;44
Literature	GENERIF_SUMMARY	Analysis of vimentin and Ki67 expression can straightforwardly be CTCs from patients with mCRPC. Poorer survival outcomes were c vimentin- and Ki67-positive CTC patients.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with Triple Negative Breast Ca
Literature	GENERIF_SUMMARY	Using the Ki-67 index is superior to using cytology and growth patt prognostic factors in Mantle-cell lymphoma.
Literature	GENERIF_SUMMARY	The addition of p16/Ki-67 to HPV DNA testing leads to a more acc stratification of CIN in women presenting with minor cytological ab
Literature	GENERIF_SUMMARY	Altogether, these results suggest that Ki-67 expression in prolifera spatially organises heterochromatin, thereby controlling gene expr
Literature	GENERIF_SUMMARY	Ki-67 expression showed a substantial concordance between biop specimens. Extremely discordant Ki-67 levels may be associated v tumor biology.
Literature	GENERIF_SUMMARY	Infiltrating Lobular Breast Carcinoma of non-lactating women show 0.009), lymph node involvement (p = 0.051) and distant metastasis They were also more proliferative tumors measured by Ki-67.
Literature	GENERIF_SUMMARY	The expression of Ki-67 protein is associated with the expression o pterygium.
Literature	GENERIF_SUMMARY	The simultaneous determination of the chemokine receptor CXCR4 proliferative activity (Ki-67 expression) can define disease prognos relapse-survival rates in patients with Stage II colon cancer after ra treatment

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Report reliability of manual and digital image analysis of Ki67 labe adrenocortical carcinoma.
Literature	GENERIF_SUMMARY	High Ki67 expression was frequently observed in resected T3 gast adenocarcinoma and was a significant prognostic factor for poor
Literature	GENERIF_SUMMARY	Ki-67 proliferation index (P = 0.027) proved to be independent prog
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with soft tissue sarcomas.
Literature	GENERIF_SUMMARY	MIB-1 confirms that a small proportion (19%) of congenital choled malformation has marked epithelial proliferation but no clinical co be identified.
Literature	GENERIF_SUMMARY	In localized PC treated by radical prostatectomy, higher Ki67 PI ass clinical grade automated algorithm is strongly associated with a hi SVI and ECE and greater probability of recurrence.
Literature	GENERIF_SUMMARY	p16/Ki-67 dual-stained cytology may serve as a more objective alt cytology for triage of high-risk HPV-positive women.
Literature	GENERIF_SUMMARY	The endometrial expression of PR and Ki67 along with serum CA1 development of lymph node metastasis in endometrial cancer.
Literature	GENERIF_SUMMARY	TOP2A and Ki-67 antibodies may be used in combination for cervi screening in immunocytochemistry assays.
Literature	GENERIF_SUMMARY	Combination of Ki-67 expression and CT value determination was classification of differentiation and metastatic or proliferative pote peripheral lung adenocarcinoma.
Literature	GENERIF_SUMMARY	higher Ki-67 expression was related to the tumour invasiveness in adenomas.
Literature	GENERIF_SUMMARY	Data show that comparing with Ki-67 and TOP2A, RacGAP1 allowe prognostic statement.
Literature	GENERIF_SUMMARY	High Ki-67 Expression is associated with Mutiple Myeloma.
Literature	GENERIF_SUMMARY	Ki-67 in tumor tissue is probably a more reliable marker than serun CYFRA 21-1 in predicting the clinical course of esophageal squam carcinoma
Literature	GENERIF_SUMMARY	This study showed that the overexpressions of Ki67, RacGAP1, an the prognosis of female breast cancer patients adversely
Literature	GENERIF_SUMMARY	High Ki67 expression is only present in 6.8% of CaP patients and is reduced survival and increased risk of metastasis, independent of score and D'Amico risk category. DLX2 is a novel marker of increas risk found in 73% patients and 8.2% showed co-expression with a
Literature	GENERIF_SUMMARY	elucidation of a biomechanical role of the mitotic chromosome pe mammalian cells and suggests that Ki-67 can function as ""surfact intracellular compartmentalization"
Literature	GENERIF_SUMMARY	Ki-67 cell expression over 50% in CIN2 biopsies and high-grade sq intraepithelial lesion (HSIL) in the previous cytology were statistica with CIN3 results in the cone
Literature	GENERIF_SUMMARY	Response to chemotherapy in neuroendocrine tumors increases w but Ki-67 alone is an unreliable means to select patients for chem

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Nested multiplex polymerase chain reactions to detect the presenc types, as well as immunohistochemical reactions were performed detection of cell cycle-related proteins p16, p27, p53 and Ki-67. Th all analyzed cell cycle-related proteins was increased in HPV-infect
Literature	GENERIF_SUMMARY	MCM4 and MCM7 expression is significantly correlated with Ki-67, cyclin E expression in esophageal adenocarcinoma, squamous cel precancerous lesions.
Literature	GENERIF_SUMMARY	Proliferative markers-mitotic count and Ki67 index-have limited val recurrence or metastasis in congenital mesoblastic nephromas wi component
Literature	GENERIF_SUMMARY	In non-germinal center B-cell subtype primary central nervous syst patients, high Ki-67 index was an adverse independent prognostic could be used to stratify patients for more suitable management.
Literature	GENERIF_SUMMARY	High grade ductal intraepithelial neoplasia lesions show haphazard while low grade ductal intraepithelial neoplasia lesions show basal Ki67 staining in the proliferating epithelial cells.
Literature	GENERIF_SUMMARY	High Ki67 expression is associated with Cervical Squamous Intrae Lesions and Cancer.
Literature	GENERIF_SUMMARY	Innon-muscle-invasive bladder cancer expression of ESR1, ERBB2 significantly different between stage and grade
Literature	GENERIF_SUMMARY	The aim of this study is to analyze the immunoexpression of Ki67, PCNA markers in epithelial remnants of dental follicles of impacte identify a possible correlation between the immunoexpression of t dentigerous cysts and keratocystic odontogenic tumors.
Literature	GENERIF_SUMMARY	Ki-67 antigen gene expression correlated with overall survival in pa malignant cutaneous melanoma.
Literature	GENERIF_SUMMARY	Here the authors show how Ki-67 and RepoMan form mitotic exit p by recruiting PP1, how they distinguish between distinct PP1 isofo the assembly of these two holoenzymes are dynamically regulated kinase during mitosis.
Literature	GENERIF_SUMMARY	Optimal tissue sampling for immunohistochemical Ki67 biomarker dependent on the heterogeneity of the tissue under study.
Literature	GENERIF_SUMMARY	p16/Ki-67 co-expressions associated strongly with high-risk huma papillomavirus persistence, especially with HPV16/18, and could b a suitable biomarker for cervical cancer screening
Literature	GENERIF_SUMMARY	the overexpression of USP7 might promote cell proliferation by de Ki-67 protein
Literature	GENERIF_SUMMARY	experiments presented confirmed that Ki67 plays a role in the form maintenance of mitotic chromosome architecture
Literature	GENERIF_SUMMARY	CD44(+)ALDH1(+)Ki-67(-) tumor cells may favor distant metastasi may have a crucial role for tumor progression, treatment resistanc metastatic ability of breast cancer stem cells
Literature	GENERIF_SUMMARY	The positive rates of Ki-67 expression was 85.42% (41/48) in the c acuminata tissues, 61.11% (11/18) in the controls, respectively.
Literature	GENERIF_SUMMARY	Describe hot-spot selection algorithm with an extended context-ba whole slide images and hot-spot gradual extinction algorithm for a

Main Category	Category	Terms
		staining in meningiomas.
Literature	GENERIF_SUMMARY	Findings from this large-scale multicentre analysis with centrally g automated KI67 scores show strong evidence in support of a prog automated KI67 scoring in breast cancer.
Literature	GENERIF_SUMMARY	the expression of p53, p16 and Ki67 in 91 cases of cutaneous squa carcinoma and its precursors, was investigated.
Literature	GENERIF_SUMMARY	Anillin (ANLN) expression in tumor cells is correlated to poor progn cancer patients, independent of Ki-67 or tumor size.
Literature	GENERIF_SUMMARY	Using p53 and/or Ki-67 in addition to cytology increases the specif penalising the sensitivity
Literature	GENERIF_SUMMARY	high Ki-67 expression was associated with poor survival in patients tract urinary carcinoma.
Literature	GENERIF_SUMMARY	For patients with ER+/HER2- breast cancer, three distinct risk patte levels were confirmed according to the 2015 St Gallen consensus. with clearly low or high Ki67-LI, straightforward clinical decisions c but for patients with intermediate Ki67-LI, other factors might prov information.
Literature	GENERIF_SUMMARY	High Ki-67 before NAC was a predictor for pCR in neoadjuvant sett cancer patients.
Literature	GENERIF_SUMMARY	Ki-67 before and after NAC, as well as the change of Ki-67 before a might be prognostic factors for OS and DFS for breast cancer patie
Literature	GENERIF_SUMMARY	the combination of the TERT promoter/BRAFV600E mutations and promising marker to predict recurrence of PTC.
Literature	GENERIF_SUMMARY	There are considerable differences between the different Ki-67 ant capacity to detect proliferating tumor cells and to separate low- an breast cancer patient groups.
Literature	GENERIF_SUMMARY	Neoadjuvant therapy in microsatellite-stable colorectal carcinoma concomitant loss of MSH6 and Ki-67 expression.
Literature	GENERIF_SUMMARY	Tumor size, lymph node involvement, family history, Ki-67 and p53 independent variables associated with either DFS or OS. Triple-neg cancer (TNBC) patients with positive p53 or Ki-67 high index or far cancer have a significant association with worse prognosis. This s that p53, Ki-67 and family history are useful prognostic markers in
Literature	GENERIF_SUMMARY	Differences in Ki67 expressions between pre- and post-neoadjuvar specimens might predict early recurrence of breast cancer.
Literature	GENERIF_SUMMARY	high Ki67/BCL2 index significantly correlated with advanced stage intestinal type, high histologic grade, and lymphatic and perineural gastric adenocarcinoma
Literature	GENERIF_SUMMARY	Studies indicate that high Ki-67 expression was correlated with po and advanced clinicopathological features, and it could serve as a disease management [Review].
Literature	GENERIF_SUMMARY	Expression of topoisomerase IIalpha and mitosin was significantly recurrent meningioma than in non-recurrent meningioma (P
Literature	GENERIF_SUMMARY	In leukoplakia, the expression of survivin associated with that of ki the assumption that all these lesions are potentially malignant.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	High Ki-67 expression Correlates with Breast Cancer.
Literature	GENERIF_SUMMARY	IHC based post-Ki67 levels may have distinct predictive power con naive IHC Ki67.
Literature	GENERIF_SUMMARY	High Ki-67 immunohistochemical expression levels in distant meta were independently associated with poorer overall survival outcom of recurrence lesion in breast cancer patients.
Literature	GENERIF_SUMMARY	Ten international pathology institutions participated in this study a messenger RNA expression levels of ERBB2, ESR1, PGR, and MKI6 centrally and locally extracted RNA from formalin-fixed, paraffin-en cancer specimens with the MammaTyper(R) test. Samples were m repeatedly on different days within the local laboratories, and repro assessed by means of variance comp...
Literature	GENERIF_SUMMARY	Upregulation of RAD51 and overexpression of Ki67 may be associa progression of thyroid cancer.
Literature	GENERIF_SUMMARY	the combination therapy upregulated the expression level of p-p53 decreased Ki-67 expression in vivo
Literature	GENERIF_SUMMARY	Immunochemical detection of the protein p16INK4a in cervical cyt combination with Ki-67.
Literature	GENERIF_SUMMARY	Ki-67 expression in ureteroscopic biopsy specimens is potentially clinical decision making for patients with suspected upper urinary carcinoma.
Literature	GENERIF_SUMMARY	A high Ki-67 LI correlated significantly with a worse prognosis in ga (GC)patients. Further cumulative studies for the optimal cutoff valu LI are needed before application in clinical practice.
Literature	GENERIF_SUMMARY	Ki-67 expression level failed to have a markedly significant impact patients with extensive-stage small cell lung cancer
Literature	GENERIF_SUMMARY	All cases of DF showed much higher Ki67 proliferation index (P = 0 with increased mitotic figures both on H&E and with anti-PHH3
Literature	GENERIF_SUMMARY	We performed immunohistochemistry for Ki67, p16INK4a, and WN HPs (hyperplastic polyps), sessile serrated adenomas/polyps (SSA traditional serrated adenomas (TSAs) .The distribution of Ki67 and positive cells in TSAs was different from that in HPs and SSA/Ps.
Literature	GENERIF_SUMMARY	Results suggest that Ki-67 expression can be a good prognostic bi early gastric cancer but not for advanced gastric cancer. Furtherm expression is a better prognostic marker in gastric cancers with we histology.
Literature	GENERIF_SUMMARY	Study shows that a Ki67 increase occurs in a significant proportion with entero-pancreatic neuroendocrine neoplasms at time of disea particularly in those with pancreatic origin.
Literature	GENERIF_SUMMARY	High Ki-67 expression in localized PCa is a factor of poor prognosi cancer
Literature	GENERIF_SUMMARY	Grading of pancreatic neuroendocrine tumors is currently based on and Ki67 proliferation index. The results suggest that Phosphohist effective marker for determining mitotic activity and can be used a Ki67.
Literature	GENERIF_SUMMARY	though TK1 expression was an independent prognostic factor for r

Main Category	Category	Terms
		for survival, TK1 is a more informative expression than Ki-67 for LI overall survival rates. Thus, when TK1 is combined with MDACC gr staging and lymph node metastasis, IHC determination of TK1 exp improve the overall prediction of prognosis in patients with ovarian
Literature	GENERIF_SUMMARY	Variants near TTN and CCDC8 were associated with KI67 expressi rs2288563 and rs2562832 in TTN are potential biomarkers for the clinical outcomes in hepatitis B-related hepatocellular carcinoma p
Literature	GENERIF_SUMMARY	The different values of the cycling nuclear area major dimension m connected with the biological behaviour of the three examined gro the endometrial epithelial cells may follow a Ki-67 increase pathwa the relatively stable pathway which the rapidly proliferating adenoc may use.
Literature	GENERIF_SUMMARY	Age-associated expression of the proliferation and immature neuro MKI67 and DCX, respectively, was unrelated, suggesting that neuro associated processes are independently altered at these points in development from stem cell to neuron.
Literature	GENERIF_SUMMARY	PD-L1, Ki-67, and p53 staining individually had significant prognost patients with stage II and III colorectal cancer
Literature	GENERIF_SUMMARY	The expression of alpha-enolase, Ki67 and p53 in pancreatic cance normal tissues were evaluated by IHC using the corresponding pri on the commercial tissue arrays
Literature	GENERIF_SUMMARY	Studies suggest that Ki-67 acts as an organiser of chromosome pe [Review].
Literature	GENERIF_SUMMARY	Prognosis of luminal breast carcinoma can be predicted using Ki6 continuous variable and a standard cut-off value of 14%. Informati specimen type used to determine ki67 should be recorded in the p report
Literature	GENERIF_SUMMARY	Ki67 depletion results in the dissociation of both pre-ribosomal RN nucleolar proteins from the perichromosomal layer (PCL), which in Ki67 is required for the PCL accumulation of pre-ribosomal RNAs, nucleolar proteins are associated.
Literature	GENERIF_SUMMARY	suggest that upregulation of NG2/CSPG4 rather than changes in C expression is associated with low overall survival in glioblastoma patients, supporting NG2/CSPG4 as a potential prognostic marker
Literature	GENERIF_SUMMARY	The results demonstrate that high expression of Ki-67 contributes resistance and acts a poor prognosis indicator in patients with loc nasopharyngeal carcinoma.
Literature	GENERIF_SUMMARY	Adjuvant chemotherapy was 9% less likely to be recommended by multidisciplinary board when using the current criteria compared w combination of the St. Gallen criteria and Ki67 and uPA/PAI-1 statu Taken together, our data show discordance among markers in iden of recurrence, even though each marker may prove to be independ
Literature	GENERIF_SUMMARY	Ki67 expression in gastric carcinoma is directly correlated with the and depth of invasion.
Literature	GENERIF_SUMMARY	Data show there was no trend to higher Ki-67 antigen in metastatic pancreatic neuroendocrine tumors (NETs).
Literature	GENERIF_SUMMARY	Suggest that different mitotic count or Ki-67 index may have differ values in different stages of small intestinal neuroendocrine tumor

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Data suggest that Ki-67 antigen proliferative index has important li hhosphohistone H3 (PHH3) is an alternative proliferative marker.
Literature	GENERIF_SUMMARY	both the value and the level of Ki-67 expression were correlated po normalized iodine concentration (NIC) values (r=0.344, P=0.002 ar P=0.026); HIF-1alpha expression was correlated positively with the the RC (r=0.598, P
Literature	GENERIF_SUMMARY	Expression of Ki-67, p53 and K-ras in the intestinal mucosa was de immunohistochemistry and quantitative real-time polymerase chai PCR), respectively. 1.03% patients (n = 7) underwent polypectomy to be the early stage of Colorectal cancer (CRC). Histological analy expression of p53 and K-ras can guarantee to screen the early stag
Literature	GENERIF_SUMMARY	According to immunohistochemistry and immunoblot analysis, the levels of cyclin D1, cyclin E, pRb, and Ki67 in psoriasis lesions decr treatment and were similar with those in the normal group
Literature	GENERIF_SUMMARY	lost BMAL1 and Ki-67 overexpression are associated with poor ove nasopharyngeal carcinoma patients
Literature	GENERIF_SUMMARY	High Ki67, EZH2, and SMYD3 immunoexpression, adjusted for star clinicopathological parameters, independently predicts outcome in prostate cancer, at diagnosis.
Literature	GENERIF_SUMMARY	TP53 expression (>50% positive cells) has shorter time to treatme poor overall survival independent of both the Mantle Cell Lymphom Prognostic Index score and Ki67 index.
Literature	GENERIF_SUMMARY	apparent diffusion coefficient was correlated with PSA, Gleason sc HIF-1alpha and VEGF expression in patients with Prostate cancer.
Literature	GENERIF_SUMMARY	An overview of functional roles of Ki-67 across the cell cycle has b along with recent experiments that clarify its role in regulating cell progression in human cells. (Review)
Literature	GENERIF_SUMMARY	Meta-analysis highlights Ki-67 expression as a predictor of breast carcinoma in situ recurrence; nevertheless, additional adjusted stu adequate follow-up periods, stemming from various world regions needed on this topic.[meta-analysis]
Literature	GENERIF_SUMMARY	In human adenocarcinoma tissues, PFKFB3 and Ki67 protein levels to the clinical characteristics and overall survival
Literature	GENERIF_SUMMARY	Ki-67 and TOPO 2A expression correlated with tumour size and tur invasiveness in somatotropinomas.
Literature	GENERIF_SUMMARY	Low Ki-67 expression is associated with glioblastoma.
Literature	GENERIF_SUMMARY	DIA using VDS is an accurate method to determine the Ki67 prolife breast cancer, as an alternative to manual scoring of whole section practice.
Literature	GENERIF_SUMMARY	Data suggest that Ki-67 is a strong prognostic factor for overall su disease-free survival (DFS) and should be included in all pancreati neuroendocrine tumor pathology.
Literature	GENERIF_SUMMARY	KI-67 expression correlates with SATB1 expression in non-small ce carcinoma.
Literature	GENERIF_SUMMARY	Ki-67 may be of diagnostic value in distinguishing between partial hydatidiform moles

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	In ACTH-secreting pituitary tumors, Ki-67 was expressed in 7 of 28 tumors and 8 of 27 nonrecurrent tumors, and there was no stainin pituitary samples. It was expressed predominantly in the nucleus c cells. There was no significant difference between Ki67 expressior nonrecurrent group and the recurrent group.
Literature	GENERIF_SUMMARY	For the first time coexpression of CD44 and Ki-67 on particulate th neoplasms of the colon is shown and the potential reasons for this are discussed.
Literature	GENERIF_SUMMARY	Aggressive adjuvant systemic treatments should be considered fo HER2-rich and triple-negative subtype who exhibit tumor shrinkage have high levels of Ki67.
Literature	GENERIF_SUMMARY	Ki-67 and condensins, which localize to the external surface and th of mitotic chromosomes, have independent yet cooperative functio supporting the structural integrity of mitotic chromosomes.
Literature	GENERIF_SUMMARY	Data suggest that Ki-67 index and survivin may be useful biomarke cancer with preoperative chemoradiotherapy.
Literature	GENERIF_SUMMARY	High Ki67 expression in the index prostate cancer lesion is an inde predictor of biochemical recurrence in patients with positive surgic following radical prostatectomy.
Literature	GENERIF_SUMMARY	For HR-positive, HER2-negative breast cancer patients with 0-1 noc chemotherapy use declined significantly after the adoption of the 2 PgR status and Ki-67 were useful for chemotherapy decision-maki without the 21-gene assay.
Literature	GENERIF_SUMMARY	Dual p16 and Ki-67 expression can be used in cervical screening o women.
Literature	GENERIF_SUMMARY	Human Endometriosis Tissue Microarray Reveals Site-specific Exp Estrogen Receptors, Progesterone Receptor, and Ki67.
Literature	GENERIF_SUMMARY	PR [progesterone receptor] immunoreactive score and Ki-67 index within WHO Grade I tumours and overlapped considerably with Gra PR expression reduced and Ki-67 index increased with increasing V 0.000). A moderate inverse correlation was found between Ki-67 in score (R = -0.7371).
Literature	GENERIF_SUMMARY	NMR diffusion tensor imaging has a high application value in the d brain glioma. The expressions of Ki-67 and VEGF are related to the grade of glioma, which can be used as biological indexes for the di glioma.
Literature	GENERIF_SUMMARY	This study demonstrated that p16/Ki-67 dual staining represents a method for cervical cancer screening. Application of this method c reduction of unnecessary colposcopy referrals and misdiagnosis.
Literature	GENERIF_SUMMARY	Overexpression of Ki-67 is associated with recurrence in Pancreati Neuroendocrine Tumors.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with papillary thyroid carcinon
Literature	GENERIF_SUMMARY	Ki-67-a proliferation marker is easily identified and provides compa information. In contrast to poor reproducibility of mitotic counts, K high agreement between pathologists; is more reproducible; adds value to the MBR grading system and correlates well with other cli parameters.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	p16/Ki-67 dual staining can increase the efficiency of screening m cervical cancer.
Literature	GENERIF_SUMMARY	BDNF and Tau as biomarkers of severity in multiple sclerosis
Literature	GENERIF_SUMMARY	he mean value was 12.33 +/- 21.77%. ADC correlated with weak st significant with expression of Ki-67 (p = - 0.202, p
Literature	GENERIF_SUMMARY	The aim of this study was to investigate the expression of p16 and proteins in regard to expression of the Ki-67 antigen and available clinicopathological data (i.a. receptor status, staging and grading
Literature	GENERIF_SUMMARY	Ki67 LI, CD105, and alpha-SMA expression showed significant diff between normal, low-risk oral epithelial dysplasia and high-risk ora dysplasia. These findings further support that features such as inc cell layer hyperplasia, abnormal superficial mitosis, increased nucl ratio, and hyperchromasia could be transformation-relevant displa
Literature	GENERIF_SUMMARY	Studied predictive value of using mitotic count vs. Ki67 score for p response to tamoxifen in postmenopausal breast cancer patients; count is a better selection marker for reduced tamoxifen benefit th
Literature	GENERIF_SUMMARY	positive correlations were found between higher Ki-67 expression differentiation,larger tumor size and higher pathologic stages, lym metastasis positivity, and advanced TNM stages.
Literature	GENERIF_SUMMARY	RAGE, EGFR and Ki-67 were immunohistochemicalyl studied for th in biopsy specimens from primary breast tumors
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with Central Giant Cell Granul
Literature	GENERIF_SUMMARY	Our results identified FGFR3(high)/Ki67(high) papillary pTa tumors with poor prognosis and encourage histological grading as high gr
Literature	GENERIF_SUMMARY	Overexpression of HER2 was observed in 48% (n = 12) of gallblad adenocarcinomas, 58% (n = 7) of high-grade dysplasia, 47% (n = 8) dysplasia, and 74% (n = 25) of intestinal metaplasia. Ki-67 index in non-linear fashion as the precursor lesions progress toward malig markers might be used as a prognostic biomarker in early neoplas transformation of gallbladder epithelium.
Literature	GENERIF_SUMMARY	We show that it is possible to approximate the true Ki67 Index acc detecting individual nuclei and also statically demonstrate the wea commonly adopted approaches that use both tumor and non-tum together while compensating for the latter with higher order appro
Literature	GENERIF_SUMMARY	Advanced cervical cancer with a high Ki-67 expression responds b cisplatin-based chemotherapy.
Literature	GENERIF_SUMMARY	Data suggest that Ki-67 labeling index (LI) grading of endoscopic u guided fine needle aspiration biopsy (EUS-FNAB) specimens shoul applied in clinical practice because of the possibility of grading un with grade 2 to 3 pancreatic neuroendocrine tumors (PanNET) cas
Literature	GENERIF_SUMMARY	p16/Ki-67 dual immunostaining had comparable sensitivity and im specificity in screening high-grade cervical intraepithelial neoplasn when compared with hrHPV detection. Further studies may be ben assess the efficacy of this novel biomarker, which can be potential of the initial screening assays.
Literature	GENERIF_SUMMARY	Our study shows that tumour grade in patients with PanNENs can determined by assessing the Ki-67 index in EUS-FNA specimens b

Main Category	Category	Terms
		2017 World Health Organisation classification and grading system
Literature	GENERIF_SUMMARY	These results show that higher Ki67 expression in fibroblast like ce front indicated better clinical outcomes for oral squamous cell car patients.
Literature	GENERIF_SUMMARY	This preliminary study showed, evidently for the first time, that mul haphazardly distributed clusters of proliferating cells are present in the Ki-67 proliferation marker only labels progenitor daughter cells stem cells, each MIB+ cluster in each NDC must have been produc stem cell.
Literature	GENERIF_SUMMARY	Oral and Extraoral Intermediate Tumors
Literature	GENERIF_SUMMARY	This is the first study to show that the Ki-67 LI in metachronous ax node metastases(ALNM) is a prognostic factor for patients with m axillary lymph node (ALN) recurrence of breast cancer.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with recurrence in papillary ur cancer.
Literature	GENERIF_SUMMARY	Apparent diffusion coefficient (parameters were correlated with K index; the parameters may help indicate p53 expression.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with Oral Squamous Cell Carc
Literature	GENERIF_SUMMARY	securin and Ki67 are expressed in invasive breast carcinoma
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with Pancreatic Neuroendocri Recurrence.
Literature	GENERIF_SUMMARY	Ki-67 may be useful diagnostic markers for carcinoma ex-pleomor
Literature	GENERIF_SUMMARY	High value of Ki-67 is associated with Breast Carcinomas.
Literature	GENERIF_SUMMARY	The cytokine interleukin-15 promotes the induction of Ki67 on NK
Literature	GENERIF_SUMMARY	Expression of Cyclin E and Ki-67 is elevated in gestational trophob
Literature	GENERIF_SUMMARY	There is a strong correlation with PHH-3 and Ki-67 and mitosis ind and grade II meningiomas
Literature	GENERIF_SUMMARY	A well-established marker of tumor proliferation, Ki-67 can indepen prognosis of patients with operable breast cancer, and a reduction to neoadjuvant chemotherapy has been associated with favorable
Literature	GENERIF_SUMMARY	High Ki-67 expression significantly predicts poor overall survival ar survival in colorectal cancer patients (meta-analysis).
Literature	GENERIF_SUMMARY	Head and neck squamous cell carcinoma (HNSCC) are fast-growin which treatment must not be delayed. Clinical tumour growth rates by EGFR, KI67 and CD44 expression.
Literature	GENERIF_SUMMARY	This meta-analysis indicate that p16 staining and p16/Ki-67 stainin specific for CIN2+/CIN3+ than high-risk human papillomavirus (hrH testing [Meta-analysis].
Literature	GENERIF_SUMMARY	Ki-67, MCM-3, and MCM-7, but not MCM-5 are reliable proliferative markers in discerning benign and malignant adrenocortical tumors
Literature	GENERIF_SUMMARY	Study suggests that both positive CD44v9 and high Ki-67 expressio associated with poor prognosis in early gastric cancer, and the cor these markers provides better prognostic stratification than the sir

Main Category	Category	Terms
		them.
Literature	GENERIF_SUMMARY	he only marker clearly related to 2-year survival is Ki-67, which is sl good prognostic factor in the multivariate analysis (hazard ratio = confidence interval = 0.27-0.90). Therefore, in a prospective cohort cancer patients, only Ki-67 is a marker of good prognosis in short-t
Literature	GENERIF_SUMMARY	Our study shows that Ki-67 is a continuous marker which is extrem pathologist variation. Even if only 50% of variance is attributable to based proliferation (ICC = 0.5), this information is sufficient to obta significant differences in clinical cohorts.
Literature	GENERIF_SUMMARY	Positive correlations were demonstrated between the expression l and the Ki-67 antigen. In vitro studies have confirmed that the expr MCM5 is elevated in cancer cells. MCM5 protein may be used as a marker of cancer cell proliferation in laryngeal squamous cell canc
Literature	GENERIF_SUMMARY	Our results have shown that high levels of TGF-alpha and Ki-67 are with shorter progression-free survival in patients with chordoma. W confirmed the role of Ki-67 as a functional molecular marker of po
Literature	GENERIF_SUMMARY	ble prognoThe nuclear protein, Ki-67, is associated with cell prolife which may be an indicator of tumour aggressiveness.
Literature	GENERIF_SUMMARY	A proliferation marker protein Ki-67 (Ki-67) index of 5% is better tha stratifying G1 and G2 pancreatic neuroendocrine tumors (pNETs).
Literature	GENERIF_SUMMARY	Expression of PCNA, Ki-67 and COX-2 in breast cancer based on D information.
Literature	GENERIF_SUMMARY	Ki-67, p53 and BCL-2 Expressions and their Association with Clinic Histopathology of Breast Cancer among Women in Tanzania.
Literature	GENERIF_SUMMARY	The findings suggest that the Ki67 proliferation index (PI) and tum lymphocytes (TILs) could be used as prognostic markers, as a pot alternative to the Hyams grade.
Literature	GENERIF_SUMMARY	More than 20% of AR-positive cytoplasmic staining of Oral squamo carcinoma (OSCC) epithelium was significantly associated with nu in the epithelium and increased AR levels in the stroma. In metasta patients, Ki-67 was significantly higher than in non-metastatic OSC
Literature	GENERIF_SUMMARY	multiplex immunofluorescence-based evaluation of Ki-67 expressi tumour and immune cells represents a novel prognostic approach
Literature	GENERIF_SUMMARY	Ki-67 levels correlate with neoplasm histologic stage and grade an term recurrence in pulmonary carcinoid tumors.
Literature	GENERIF_SUMMARY	Proliferation Tumour Marker Network (PTM-NET) for the identifica region in Ki67 stained breast cancer whole slide images.
Literature	GENERIF_SUMMARY	This study is the first to demonstrate disproportionally high Ki-67 l in a large number of cribriform-morular variant-papillary thyroid ca despite the fact that these cases had favorable prognoses.
Literature	GENERIF_SUMMARY	The expressions of ER, PR, HER-2 and Ki-67 in metastatic lesions a with the prognosis of breast cancer patients.Their expression disc between primary and metastatic lesions can guide the treatment a risks of recurrence and prognosis.
Literature	GENERIF_SUMMARY	We have determined in this study that p53, Ki-67 and laminin may l adjuvant immunohistochemical markers in differential diagnosis o

Main Category	Category	Terms
		Keratoacanthoma and well-differentiated squamous cell carcinom
Literature	GENERIF_SUMMARY	It is recommended to monitor positivity rate of ER, PR, Ki67 and He periodically to comply with quality assurance requirements.
Literature	GENERIF_SUMMARY	Evaluation of T cell activation with Ki67 expression.
Literature	GENERIF_SUMMARY	high Ki-67 expression was significantly related with adverse pathol finally with worse biochemical recurrence-free survival.
Literature	GENERIF_SUMMARY	High Ki-67 expression is associated with lymph node metastasis i Cancer.
Literature	GENERIF_SUMMARY	Hot spots in fetal human penile and clitoral development.
Literature	GENERIF_SUMMARY	Gene expression profiling revealed MCM3 to be a better marker tha prognosis of invasive ductal breast carcinoma patients.
Literature	GENERIF_SUMMARY	Expression of Ki-67 and Betaeta-Catenin in Pseudoepitheliomatous and Squamous Cell Carcinoma in Oral Mucosal Biopsies
Literature	GENERIF_SUMMARY	Immunoexpression of estrogen receptor-beta and progesterone re prostate adenocarcinoma, does it inhibit neoplastic proliferation a
Literature	GENERIF_SUMMARY	levels negatively influenced survival in meningiomas [review]
Literature	GENERIF_SUMMARY	Immunohistochemical staining patterns of Ki-67 and p53 in florid r urothelial atypia and urothelial carcinoma in situ demonstrate sign
Literature	GENERIF_SUMMARY	Expression of E-cadherin, Ki-67, and p53 in urinary bladder cancer progression, survival, and recurrence.
Literature	GENERIF_SUMMARY	THE STUDY OF THE EXPRESSION OF CDH1, KI67, P53 AND HER2 GASTRIC CARCINOMA.
Literature	GENERIF_SUMMARY	Ki67 is a risk factor in the prognosis of prostate cancer
Literature	GENERIF_SUMMARY	Association between Ki-67 Labeling index and Histopathological G Glioma in Indonesian Population.
Literature	GENERIF_SUMMARY	High Expression of miR-34a Associated with Less Aggressive Can Not with Survival in Breast Cancer.
Literature	GENERIF_SUMMARY	Ki-67 and Clinical Correlations in Patients with Resistant Prolactin
Literature	GENERIF_SUMMARY	Factors influencing agreement of breast cancer luminal molecular Ki67 labeling index between core needle biopsy and surgical resec
Literature	GENERIF_SUMMARY	Ki67 in endocrine neoplasms
Literature	GENERIF_SUMMARY	Correlation between Skip N2 Metastases and SUVmax, Long Diam and Ki67 Expression in Patients with Non-Small-Cell Lung Cancer.
Literature	GENERIF_SUMMARY	STMN1 and MKI67 Are Upregulated in Uterine Leiomyosarcoma a Biomarkers for its Diagnosis.
Literature	GENERIF_SUMMARY	Clinical complete responders experienced a greater overall variatio compared with partial responders and patients with progressive/s (66.7% vs 30.7%, p = 0.009). High responders showed a better out responders in terms of both disease-free survival (p = 0.009) and o (p = 0.002).
Literature	GENERIF_SUMMARY	Correlation Between p16/Ki-67 Expression and the Grade of Cervic

Main Category	Category	Terms
		Intraepithelial Neoplasias.
Literature	GENERIF_SUMMARY	Stage-dependent Increase in Expression of miR-155 and Ki-67 and Tumour-associated Inflammatory Cells in Folliculotropic Mycosis F
Literature	GENERIF_SUMMARY	Expression of Ki-67 and CD34 on blood and bone marrow cells of C with different response to imatinib and nilotinib therapy.
Literature	GENERIF_SUMMARY	Ki67, CD105, and alpha-SMA Expression Supports Biological Distin Squamous Cell Carcinoma Arising in the Background of Oral Subm
Literature	GENERIF_SUMMARY	Relationships of expressions of P16 and Ki-67 with tumor markers of cervical cancer patients.
Literature	GENERIF_SUMMARY	Recurrence of Solitary Fibrous Tumor/Hemangiopericytoma Could by Ki-67 Regardless of Its Origin.
Literature	GENERIF_SUMMARY	Is Ki-67 index overexpression in IDH wild type glioblastoma a predi Progression Free survival? A clinical and Molecular analytic investi
Literature	GENERIF_SUMMARY	Chromosome clustering by Ki-67 excludes cytoplasm during nucle
Literature	GENERIF_SUMMARY	Impact of Ki-67 Labeling Index on Prognostic Significance of the C Response Score in Women With Tubo-ovarian Cancer Treated With Chemotherapy.
Literature	GENERIF_SUMMARY	P16, Ki67, P53, and WT1 Expression in Uterine Smooth Muscle Tur
Literature	GENERIF_SUMMARY	Curative Surgery and Ki-67 Value Rather Than Tumor Differentiatio Survival of Patients With High-grade Neuroendocrine Neoplasms.
Literature	GENERIF_SUMMARY	Ki67 and P53 Expression in Relation to Clinicopathological Feature Tumour of the Breast.
Literature	GENERIF_SUMMARY	The Relationship of GATA3 and Ki-67 With Histopathological Progn Parameters, Locoregional Recurrence and Disease-free Survival in Carcinoma of the Breast.
Literature	GENERIF_SUMMARY	Validation of Clinical Treatment Score post-5 years (CTS5) risk stra premenopausal breast cancer patients and Ki-67 labelling index.
Literature	GENERIF_SUMMARY	Expression and significance of Her2 and Ki-67 in gastric adenocar distant metastasis
Literature	GENERIF_SUMMARY	Expression of COX-2, p16, and Ki67 in the range from normal breas breast cancer.
Literature	GENERIF_SUMMARY	Long-term outcome and prognostic value of Ki67 after perioperati therapy in postmenopausal women with hormone-sensitive early b (POETIC)
Literature	GENERIF_SUMMARY	Correlation between immunohistochemical expression of Ki-67and aggressiveness of urinary bladder urothelial carcinoma.
Literature	GENERIF_SUMMARY	Efficacy and safety of an adsorbent and anti-oxidative vaginal gel o on high-risk HPV, and on p16/Ki-67
Literature	GENERIF_SUMMARY	Digital Quantification of Ki-67 and PHH3 in the Classification of Ut Muscle Tumors.
Literature	GENERIF_SUMMARY	[Detection and value of serum antigen KI-67 (ki67) in clinical diagn cancer patients].

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	MCM3 proliferative index is worthier over Ki-67 in the characterizat gland tumors.
Literature	GENERIF_SUMMARY	Predictive Ki-67 Proliferation Index of Cervical Squamous Cell Carc on IVIM-DWI Combined with Texture Features.
Literature	GENERIF_SUMMARY	Should the solid variant of odontogenic keratocyst and keratoamel classified as the same entity? A clinicopathological analysis of nin review of the literature.
Literature	GENERIF_SUMMARY	Role of WT1, B-cell lymphoma 2, Ki-67 (Mib1), and Her2/Neu as dia prognostic immunomarkers in ovarian serous and endometroid ca
Literature	GENERIF_SUMMARY	Expression of p53 and Ki-67 proteins in patients with increasing se duration of pterygium.
Literature	GENERIF_SUMMARY	Analysis of the association of BRAFV600E mutation and Ki-67 ove with clinical and pathological characteristics in papillary thyroid ca
Literature	GENERIF_SUMMARY	Different Stages of Quiescence, Senescence, and Cell Stress Ident Molecular Algorithm Based on the Expression of Ki67, RPS6, and E Galactosidase Activity.
Literature	GENERIF_SUMMARY	To Ki or Not to Ki
Literature	GENERIF_SUMMARY	Ki67 does not predict recurrence for low-grade appendiceal mucin with peritoneal dissemination after cytoreductive surgery and HIPE
Literature	GENERIF_SUMMARY	Correlationship between Ki67, VEGF, and p53 and Hepatocellular C Recurrence in Liver Transplant Patients.
Literature	GENERIF_SUMMARY	Clinical significance of circulating tumour cells and Ki-67 in renal c
Literature	GENERIF_SUMMARY	Gata-3 and KI-67 expression in correlation with molecular subtypes cancer.
Literature	GENERIF_SUMMARY	S1P Lyase Regulates Intestinal Stem Cell Quiescence via Ki-67 and
Literature	GENERIF_SUMMARY	Association of p16 and Ki-67 with Risk of Recurrence in Previously Cervical High-Grade Squamous Intraepithelial Lesions.
Literature	GENERIF_SUMMARY	Prognostic and clinicopathologic value of ki-67 and profilin 1 immunohistochemical expression in primary pT1 urothelial bladde
Literature	GENERIF_SUMMARY	Autophagy activity in cholangiocarcinoma is associated with anato localization of the tumor.
Literature	GENERIF_SUMMARY	Prognostic Values of MicroRNA-21 and Ki-67 in Diffuse Large B-Ce Patients
Literature	GENERIF_SUMMARY	Role of Ki-67, MRE11, and PD-L1 as Predictive Biomarkers for Recu in Muscle-invasive Bladder Cancer.
Literature	GENERIF_SUMMARY	Characterisation of anal intraepithelial neoplasia and anal cancer i men by immunohistochemical markers p16, Ki-67, HPV-E4 and DN markers.
Literature	GENERIF_SUMMARY	Diagnostic utility and prognostic significance of the Ki-67 labeling large B-cell lymphoma transformed from follicular lymphoma
Literature	GENERIF_SUMMARY	The chromatin-binding domain of Ki-67 together with p53 protects chromosomes from mitotic damage.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Cavernous Malformations to Be Investigated for Familiarity
Literature	GENERIF_SUMMARY	Ki67 as Proliferative Marker in Patients with Early Breast Cancer ar Association with Clinicopathological Factors.
Literature	GENERIF_SUMMARY	Ki67 as a prognostic marker in uterine leiomyosarcoma
Literature	GENERIF_SUMMARY	Correlation analysis of Ki67 and CK7 expression with clinical chara prognosis of postoperative cervical adenocarcinoma patients.
Literature	GENERIF_SUMMARY	Visualization of endogenous p27 and Ki67 reveals the importance driven metabolic switch in promoting survival of quiescent cancer
Literature	GENERIF_SUMMARY	Diffusive Ki67 and vimentin are associated with worse recurrence- upper tract urothelial carcinoma
Literature	GENERIF_SUMMARY	The relationship between the apparent diffusion coefficient and th proliferation index in intracranial solitary fibrous tumor/hemangiop
Literature	GENERIF_SUMMARY	Comparison of Different Molecular Subtypes with 14% Ki-67 Cut-of Breast Cancer Patients of Pakistan- An Indication of Poor Prognos
Literature	GENERIF_SUMMARY	Evaluation of Ki-67 as a Prognostic Marker in Diffuse Large B-Cell Single-Center Retrospective Cohort Study.
Literature	GENERIF_SUMMARY	Prognostic Value of Copine 1 in Patients With Renal Cell Carcinom
Literature	GENERIF_SUMMARY	Clinicopathological Analysis of Non-B Non-C Hepatocellular Carcin on Cellular Proliferation.
Literature	GENERIF_SUMMARY	Apparent Diffusion Coefficient Histogram Analysis for the Preopera of Ki-67 Expression in Pituitary Macroadenoma.
Literature	GENERIF_SUMMARY	Immunoexpression of PTEN, HER2/neu, and Ki-67 in endoscopic g carcinoma biopsies; their correlation with histological subtypes an survival.
Literature	GENERIF_SUMMARY	Integration of the Ki-67 proliferation index into the Ogata score imp diagnostic sensitivity for low-grade myelodysplastic syndromes.
Literature	GENERIF_SUMMARY	Utility of KI-67 as a prognostic biomarker in pulmonary neuroendoc
Literature	GENERIF_SUMMARY	Clinical significance of CD34, podoplanin and Ki-67 expression in p locally advanced squamous cell cervical carcinoma.
Literature	GENERIF_SUMMARY	Significance of Ki67 expression in endometriosis for infertility.
Literature	GENERIF_SUMMARY	Prognostic Utility of the Ki-67 Labeling Index in Follicular Thyroid T
Literature	GENERIF_SUMMARY	Symptom-related Ki-67 expression in endometrial polyps.
Literature	GENERIF_SUMMARY	Noninvasive Method for Predicting the Expression of Ki67 and Pro Small-Cell Lung Cancer Patients
Literature	GENERIF_SUMMARY	Loss of E-Cadherin Expression Correlates With Ki-67 in Head and N Cell Carcinoma.
Literature	GENERIF_SUMMARY	The value of Ki67 for the diagnosis of LSIL and the problems of p1 diagnosis of HSIL.
Literature	GENERIF_SUMMARY	Can we use Ki67 expression to predict prostate cancer aggressive
Literature	GENERIF_SUMMARY	Validation of absolutely quantitated Ki67 and cyclinD1 protein leve

Main Category	Category	Terms
		of Luminal-like breast cancer patients.
Literature	GENERIF_SUMMARY	Clinical significance and interrelations of PD-L1 expression, Ki-67 i molecular alterations in sporadic medullary thyroid carcinoma fron population.
Literature	GENERIF_SUMMARY	Correlation of the Ki67 Working Group prognostic risk categories v Oncotype DX Recurrence Score in early breast cancer.
Literature	GENERIF_SUMMARY	Correlation of H3K9ac and H4K12ac With Cell Proliferation Marker Leukoplakia
Literature	GENERIF_SUMMARY	Does the Overall Survival of the Resectable Periampullary Carcinor with High Expression of p53 and ki67?
Literature	GENERIF_SUMMARY	Human herpesvirus 6A and 6B and polyomavirus JC and BK infect carcinoma and their relationship with p53, p16INK4a, Ki-67, and nu kappa B expression.
Literature	GENERIF_SUMMARY	Association between epidemiological and clinico-pathological feat cancer with prognosis, family history, Ki-67 proliferation index and Tunisian breast cancer patients.
Literature	GENERIF_SUMMARY	Cyclin-dependent kinase inhibitor p21 and proliferative marker ki67 carcinoma.
Literature	GENERIF_SUMMARY	A Consensus Diagnosis Utilizing Surface KI-67 Expression as an A in Low-Grade Dysplasia Helps Identify Patients at High Risk of Pro High-Grade Dysplasia and Esophaegal Adenocarcinoma.
Literature	GENERIF_SUMMARY	Lung carcinoid tumours
Literature	GENERIF_SUMMARY	High rate of Ki-67 increase in entero-pancreatic NET relapses after curative intent.
Literature	GENERIF_SUMMARY	Circulating tumor cells correlating with Ki-67 predicts the prognosi cancer patients.
Literature	GENERIF_SUMMARY	Practice patterns for reporting digestive system neuroendocrine ne
Literature	GENERIF_SUMMARY	The relationships of OSBPL3 expression with KI-67 expression and mutations in CRC
Literature	GENERIF_SUMMARY	Mesenchymal circulating tumor cells and Ki67
Literature	GENERIF_SUMMARY	Expression of Markers Ki-67, Nestin, VEGF, CD34 and Apoptosis in Healthy Lung Tissue with Non-Changed and Metaplastic Bronchial
Literature	GENERIF_SUMMARY	Evaluation of variable p16 immunostaining patterns, Ki-67 indices in cervical SILs and squamous cell carcinomas
Literature	GENERIF_SUMMARY	C/EBPbeta expression decreases in cervical cancer and leads to tu
Literature	GENERIF_SUMMARY	CCDC86 is a novel Ki-67-interacting protein important for cell divis
Literature	GENERIF_SUMMARY	[Features of expression of GLUT-1 and Ki-67 proteins in various co mixed variants of ameloblastoma].
Literature	GENERIF_SUMMARY	Identifying key genes related to inflammasome in severe COVID-19 on a joint model with random forest and artificial neural network.
Literature	GENERIF_SUMMARY	Clinicopathological Value Of Epidermal Growth Factor Receptor (E Expression In Colorectal Adenoma And Adenocarcinoma.

Main Category	Category	Terms
Literature	GENERIF_SUMMARY	Identification of MKI67, TPR , and TCHH Mutations as Prognostic I Patients With Defective Mismatch Repair Colon Cancer Stage II/III
Literature	GENERIF_SUMMARY	Ki67 and prostate specific antigen are prognostic in metastatic ho prostate cancer.
Literature	GENERIF_SUMMARY	The expression of p53, ki67 and COX-2 in erosive-type oral lichen p
Literature	GENERIF_SUMMARY	Effects of different KRAS mutants and Ki67 expression on diagnos prognosis in lung adenocarcinoma.
Literature	GENERIF_SUMMARY	Ki-67 is necessary during DNA replication for fork protection and g
Literature	GENERIF_SUMMARY	Expression of cytokeratin 7/20 and Ki67 in Helicobacter pylori-ass and intestinal metaplasia.
Literature	GENERIF_SUMMARY	Relationship between p16/ki67 immunoscores and PAX1/ZNF582 status in precancerous and cancerous cervical lesions in high-risk women.
Pathways	WIKIPATHWAYS	Primary focal segmental glomerulosclerosis FSGS
Protein Domains	INTERPRO	FHA_dom
Protein Domains	INTERPRO	SMAD_FHA_dom_sf
Protein Domains	INTERPRO	KI67R
Protein Domains	INTERPRO	PP1-bd
Protein Domains	SMART	FHA
Protein Domains	SMART	K167R
Protein Domains	UP_KW_DOMAIN	Repeat