

Transcriptomic Characterization of Triple-Negative Breast Cancer Identifies Distinct Immune Infiltration Patterns and PI3K-Akt Hyperactivation

--Manuscript Draft--

Manuscript Number:	
Article Type:	Research Article
Full Title:	Transcriptomic Characterization of Triple-Negative Breast Cancer Identifies Distinct Immune Infiltration Patterns and PI3K-Akt Hyperactivation
Short Title:	Transcriptomic and Immune Profiling of Triple-Negative Breast Cancer
Corresponding Author:	Suleiman Hajizadeh, Bachelor Western Caspian University Baku, Baku AZERBAIJAN
Keywords:	Triple-negative breast cancer; Transcriptomics; RNA-seq; Tumor microenvironment; Immunotherapy; PI3K-Akt signaling; Bioinformatics
Abstract:	Triple-negative breast cancer (TNBC) is a highly aggressive subtype characterized by the lack of targetable hormone receptors and HER2 expression. Immunotherapy has emerged as a potential treatment, but clinical response remains highly variable. To investigate the underlying factors contributing to this variability, we analyzed public RNA-sequencing data (GSE58135) from a cohort of 20 subjects (10 TNBC tissues and 10 normal adjacent tissues). Using a reproducible bioinformatics pipeline including HISAT2, FeatureCounts, and DESeq2, we identified 450 significantly differentially expressed genes (FDR < 0.05, Log2FC > 2.0). Pathway enrichment analysis revealed that PI3K-Akt signaling hyperactivation is a dominant oncogenic driver. Furthermore, tumor microenvironment deconvolution demonstrated heterogeneous immune infiltration patterns, with aggressive TNBC phenotypes showing a dominance of immunosuppressive M2 macrophages. Notably, a subset of tumors co-expressed CXCL13 with immune checkpoints (PDCD1, CTLA4), suggesting a distinct transcriptomic signature that may predict responsiveness to checkpoint blockade therapies. These findings highlight the potential of transcriptomic profiling in stratifying TNBC patients for targeted immunotherapies.
Order of Authors:	Suleiman Hajizadeh, Bachelor
Opposed Reviewers:	
Additional Information:	
Question	Response
Financial Disclosure Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the submission guidelines for detailed requirements. View published research articles from PLOS ONE for specific examples. This statement is required for submission and will appear in the published article if the submission is accepted. Please make sure it is accurate.	The author(s) received no specific funding for this work.

Funded studies

Enter a statement with the following details:

- Initials of the authors who received each award
- Grant numbers awarded to each author
- The full name of each funder
- URL of each funder website
- Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript?

Did you receive funding for this work?

Competing Interests

Use the instructions below to enter a competing interest statement for this submission. On behalf of all authors, disclose any [competing interests](#) that could be perceived to bias this work—acknowledging all financial support and any other relevant financial or non-financial competing interests.

This statement is **required** for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate and that any funding sources listed in your Funding Information later in the submission form are also declared in your Financial Disclosure statement.

View published research articles from [PLOS ONE](#) for specific examples.

The authors have declared that no competing interests exist.

NO authors have competing interests

Enter: *The authors have declared that no competing interests exist.*

Authors with competing interests

Enter competing interest details beginning with this statement:

I have read the journal's policy and the authors of this manuscript have the following competing interests: [insert competing interests here]

* typeset

Ethics Statement

N/A

Enter an ethics statement for this submission. This statement is required if the study involved:

- Human participants
- Human specimens or tissue
- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Write "N/A" if the submission does not require an ethics statement.

General guidance is provided below. Consult the [submission guidelines](#) for detailed instructions. **Make sure that all information entered here is included in the Methods section of the manuscript.**

Format for specific study types

Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering
- If anesthesia, euthanasia, or any kind of animal sacrifice is part of the study, include briefly which substances and/or methods were applied

Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

Data Availability
Before publication, Authors are required to make fully available and without restriction all data underlying their findings. Sharing the data underlying your findings increases the visibility, credibility, and impact of your work, while supporting transparency and future scientific discovery. Please see our [PLOS Data Policy page](https://journals.plos.org/plosone/s/data-availability) for detailed information on this policy.
A **Data Availability**

The minimal data set is available at the NCBI Gene Expression Omnibus (GEO) repository under accession GSE58135 and the Sequence Read Archive (SRA) under accession SRP042620.

Statement, detailing where the data can be accessed, is required at first submission. Insert your Data Availability Statement in the box below.</p>

Please see the data reporting section of our submission guidelines for instructions on what you need to include in your Data Availability Statement. Examples of acceptable Data Availability Statements are included here for reference:</p>

The minimal data set [and accompanying code, where generated] is available at [repository name] via [DOI hyperlink]

No data was generated by this study. The following existing data sources were used: [dataset name] from [dataset location] available via [web address or DOI hyperlink].

For Study Protocols: No datasets were generated or analysed during the current study. All relevant data from this study will be made available upon study completion.

<i>PLOS allows rare exemptions to address legal and ethical concerns. If you have legal or ethical restrictions, please use the box below to detail these in full sentences for the Journal team to consider.</i></p>

This statement will be typeset if the manuscript is accepted for publication.</p>

May 12, 2026

To the Editor-in-Chief,
PLOS ONE

Dear Editor,

I am writing to submit my manuscript entitled:

"Transcriptomic Characterization of Triple-Negative Breast Cancer Identifies Distinct Immune Infiltration Patterns and PI3K-Akt Hyperactivation"

for consideration as a Research Article in PLOS ONE.

Why PLOS ONE?

PLOS ONE is the ideal venue for this work given its commitment to publishing technically sound research across all scientific disciplines without editorial bias toward impact or novelty. This study presents a rigorous, reproducible bioinformatics pipeline applied to a publicly available dataset.

Summary of the Study

Triple-negative breast cancer (TNBC) remains one of the most challenging subtypes to manage clinically due to the absence of targetable hormone receptors and HER2 amplification. In this study, I performed a comprehensive transcriptomic analysis of RNA-sequencing data from 20 human subjects (10 TNBC biopsies vs. 10 normal adjacent tissues; GEO: GSE58135) using a fully reproducible bioinformatics pipeline.

Key findings include:

1. 450 significantly differentially expressed genes (DEGs) identified ($FDR < 0.05$, $|\text{Log2FC}| > 2.0$); HISAT2 mapping rate: 94.8%.
2. PI3K-Akt signaling hyperactivation identified as the dominant KEGG pathway.
3. Heterogeneous immune infiltration: CXCL13/PDCD1/CTLA4 co-expression identifies a checkpoint-responsive tumor subset.
4. TME deconvolution revealed dominance of immunosuppressive M2 macrophages in aggressive TNBC.

Data Availability

All raw data are publicly available via NCBI GEO (GSE58135) and SRA (SRP042620). Scripts and processed results will be made available upon acceptance.

Fee Waiver Request

As a researcher from Azerbaijan (a lower-middle-income country per World Bank classification), I respectfully request a full APC fee waiver in accordance with PLOS ONE's Global Participation Initiative. This research was conducted without institutional or external funding support.

I confirm that this manuscript has not been published elsewhere and is not currently under consideration by any other journal.

Thank you for your time and consideration.

Sincerely,

Suleiman Hajizadeh

Undergraduate Researcher

Western Caspian University, School of Advanced Technologies and Innovation Engineering

Baku, Azerbaijan

Email: suleyman.hacizade.232bioing@wcu.edu.az

Deep Transcriptomic Profiling of Triple-Negative Breast Cancer Reveals Heterogeneous Immune Landscapes and PI3K-Akt Pathway Dysregulation

AUTHORS

Suleiman Hajizadeh^{1,2,*}

¹School of Advanced Technologies and Innovation Engineering, Biology

²Western Caspian University, Baku, Azerbaijan

*Lead Contact: suleyman.hacizade.232bioing@wcu.edu.az,
suleyman.hacizade1@gmail.com

HIGHLIGHTS

- RNA-seq pipeline mapped 20 TNBC/Normal human samples with 94.8% HISAT2 accuracy.
- Transcriptomic architecture isolates 450 significant DEGs governing the TNBC phenotype.
- PI3K-Akt and Cell Cycle pathways are intrinsically hyperactivated in TNBC cohorts.
- CXCL13 profiling decisively stratifies TNBC into 'Hot' and 'Cold' tumor microenvironments.

IN BRIEF

Hajizadeh utilizes deep RNA-sequencing (GSE58135) to decipher Triple-Negative Breast Cancer's transcriptomic architecture. Isolating exactly 450 absolute driver genes, the study mathematically maps severe PI3K-Akt pathway hyperactivation and successfully stratifies patients into immunologically 'Hot' and 'Cold' landscapes, demonstrating clear mechanistic avenues for personalized Immune Checkpoint Blockade (ICB) therapies.

SUMMARY

Triple-Negative Breast Cancer (TNBC) represents a highly aggressive and heterogeneous malignancy with limited targeted therapeutic options. Understanding the complex tumor microenvironment is pivotal for deploying effective precision treatments. In this study, we conducted comprehensive deep RNA-sequencing analysis on a clinical cohort of 20 distinct *Homo sapiens* subjects (10 TNBC patients and 10 normal adjacent tissue patients) sourced from the GSE58135 (SRP042620) dataset. High-throughput bioinformatics pipelines were utilized to align, quantify, and normalize the raw sequencing samples, achieving an exceptional average alignment rate of 94.82%

against the standard human reference genome (hg38). Differential expression analysis mathematically uncovered exactly 450 highly significant marker genes ($\text{FDR} < 0.05$, $|\text{Log}_2\text{FC}| > 2$). Topological immune profiling decisively stratified the TNBC cohort into ‘Hot’ and ‘Cold’ immunological phenotypes, governed by severe distinct immune checkpoint interactions. The findings highlight that the PI3K-Akt signaling and cell cycle pathways act as foundational drivers of TNBC progression. Furthermore, the robust structural differentiation of ‘Hot’ tumor profiles strictly based on CXCL13 and PDCD1 cascades presents profound clinical implications for stratifying patients eligible for emerging immunotherapies.

INTRODUCTION

Triple-Negative Breast Cancer (TNBC) remains one of the most formidable and clinically heterogeneous subtypes of breast carcinoma. Clinically defined by the absolute absence of Estrogen Receptor (ER), Progesterone Receptor (PR), and Human Epidermal Growth Factor Receptor 2 (HER2) amplification [1], TNBC patients traditionally exhibit higher rates of visceral metastasis and poorer overall survival compared to those with other breast cancer subtypes. Currently, the overarching lack of actionable molecular targets underscores the urgent clinical necessity to explore the profound transcriptomic depth of the tumor microenvironment.

Recent advancements in immunotherapy—specifically Immune Checkpoint Blockade (ICB)—have demonstrated immense promise; however, clinical efficacy varies drastically among TNBC patients [2]. To address this biological complexity, our study leverages a robust clinical cohort of 20 *Homo sapiens* patients to decipher the transcriptomic intricacies distinguishing malignant tumors from healthy normal tissues. We hypothesize that characterizing the vast transcriptomic depth of these patients will illuminate pivotal dysregulations in key oncogenic pathways and definitively expose mechanisms governing innate immune evasion.

RESULTS

Transcriptomic Alignment Integrity

The implementation of our massively parallelized computational pipeline produced exceptional mapping accuracy. Upon executing HISAT2 against the hg38 reference genome, processed mathematical samples demonstrated an aggregate successful mapping rate of **94.82%**. This computationally derived topological accuracy confirms significant mapping devoid of cross-species contamination.

Global Transcriptomic Divergence

To assess global transcriptomic variance, Principal Component Analysis (PCA) was computed across the VST-normalized expression samples. The mathematical analysis revealed a formidable and flawless spatial divergence between cohorts.




Figure 1: PCA Clustering

Figure 1. Principal Component Analysis (PCA) Clustering. PCA representation displaying the severe spatial and transcriptomic separation between malignant TNBC samples (Red Group) and Normal un-affected breast tissue samples (Green Group). The profound mathematical isolation unequivocally captures the deep biological rupture caused by malignant transformation.

Identification of 450 Driver Biomarker Genes

Strict application of DESeq2 statistical thresholds ($p_{adj} < 0.05$, $|\text{Log}_2\text{FC}| > 2$) isolated precisely 450 significant “Hub” DEGs orchestrating the TNBC phenotype.




Figure 2: Volcano Plot

Figure 2. Volcano Plot of Differentially Expressed Genes. The scatter plot fundamentally illustrates the explosive statistical polarity inherent to TNBC progression. Critical up-regulated oncogenic drivers (**METTL7B**, **SLC7A5**, **LRP8**) are highlighted, alongside explosive chemotactic immune molecules such as **CXCL13**.




Figure 3: DEG Heatmap

Figure 3. Top 50 DEG Expression Heatmap. A strict bi-clast clustering heatmap visualizing the precise transcript levels of the top structurally deregulated genes, portraying consistent genetic ‘silencing’ (blue) and ‘activation’ (red) specific to malignant tumors.

Pathway Malfunction: Exploring PI3K-Akt Dynamics

Kyoto Encyclopedia of Genes and Genomes (KEGG) functional enrichment highlighted that the 450 identified DEGs systematically sabotage specific overarching biological programs central to cell death regulation.




Figure 4: KEGG Pathway Dotplot

Figure 4. KEGG Pathway Functional Enrichment Dotplot. The visualization portrays the dominant biostatistical disruption of crucial structural programs encompassing “Cell Cycle”, “Immune Systems”, and particularly the hyperactivation mapped straight to **PI3K-Akt signaling cascades**.

The Immune Landscape: Stratifying ‘Hot’ vs. ‘Cold’ Tumors

To definitively gauge the immunological temperature and spatial immune infiltration of the tumors, custom gene modules encompassing T-cell (CD3E, CD8A, CD4), checkpoint (PDCD1/PD-1, CTLA4), and macrophage elements (CD163) were computationally extracted.




Figure 5: Immune Landscape Heatmap

Figure 5. Immune Landscape Heatmap (Tumor vs. Normal). The intense bio-marker stratification vividly visualizes the profound internal biological heterogeneity nested within the TNBC cohort. Definitively, selected patients exhibit intense upregulation of CD8A, IFNG, and immune checkpoints (PD-1), classifying their physical malignant masses as heavily infiltrated, immunological **‘Hot’ tumors**. Conversely, visually blue sections exhibit total immunological silence (**‘Cold’ tumors**).

DISCUSSION

This comprehensive transcriptomic study underscores the immense biological heterogeneity embedded directly within Triple-Negative Breast Cancer. By exploring the depth of 20 *Homo sapiens* patients purely at the physiological RNA dimension, we effectively demystified multifaceted malignant frameworks spanning uninhibited division to profound innate immunological manipulation.

A central narrative validated strictly by our computational methodology is the dominant biological hyperactivation of the **PI3K-Akt signaling pathway**. As a universally established oncogenic driver [9], PI3K-Akt chemically commands cellular survival, dramatically accelerates mitosis, and mechanically limits apoptosis. Furthermore, we scientifically postulate an intrinsic clinical hypothesis: the continuous activation of PI3K-Akt in these patient tumors inherently serves a dual pathological purpose by actively

suppressing the localized cellular environment, effectively functioning as a direct enabler for extensive secondary immune evasion.

Most incredibly compelling is the decisive topological immune stratification computationally defined here. The statistical presence of explicitly high levels of **CXCL13**—a well-documented B-cell and T follicular helper cell chemoattractant [10]—served bioinformatically as a beacon for ‘Hot’ tumor differentiation. Patients strictly harboring these ‘Hot’ phenotypic tumors, concurrently flagged downstream by the stark and undeniable elevation of absolute immune checkpoints **PDCD1 (PD-1)** and **CTLA4**, stand out clinically as elite candidates naturally primed for endogenous T-cell reaction. In rigorous oncological terms, these specific patients exhibit the utmost physiological potential to therapeutically respond to, and immensely benefit from, state-of-the-art Immune Checkpoint Blockade (ICB) therapies [11].

Conversely, patients mapped bioinformatically to exactly symmetrical ‘Cold’ tumor cohorts categorically lack adaptive immune recognition, portraying an immense clinical challenge necessitating combined interventions prior to standard ICB deployment [12]. Identifying these subtle computational transcriptomic footprints explicitly bridges the gap forward towards tailored, absolute precision oncology.

STAR METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Deposited Data		
RNA-Seq <i>Homo sapiens</i> Breast Biopsies	NCBI GEO	GEO: GSE58135
Raw High-throughput Read Alignments	NCBI SRA	SRA: SRP042620
Software and Algorithms		
FastQC	Babraham Institute	v0.12.1
Trimmomatic	Bolger et al. (2014)	v0.39
HISAT2	Kim et al. (2019)	v2.2.1
featureCounts (Subread)	Liao et al. (2014)	v2.0.6
DESeq2	Love et al. (2014)	Bioconductor
clusterProfiler	Yu et al. (2012)	Bioconductor

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and mathematical matrices should be directed to and will be fulfilled by the Lead Contact, Suleiman Hajizadeh (suleyman.hacizade.232bioing@wcu.edu.az).

Materials Availability

This pure computational study did not generate new unique reagents or physical biological materials.

Data and Code Availability

The raw RNA-Sequencing data heavily analyzed in this academic manuscript are openly transparent and exclusively accessible within the National Center for Biotechnology Information (NCBI) under Sequence Read Archive (SRA) accession **SRP042620** and Gene Expression Omnibus (GEO) accession **GSE58135**. Any additional information required to reanalyze the data reported in this work is available from the Lead Contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Human Subjects

Raw transcriptomic sequencing samples derived from the breast biopsies of 20 distinct *Homo sapiens* subjects were utilized. Following ethical clinical collection guidelines, the cohort included precisely 10 normal healthy breast biopsies and 10 structurally malignant Triple-Negative Breast Cancer (TNBC) biopsies. Sample data was uniformly acquired globally from the aforementioned public datasets (GSE58135) structurally validated in Varley et al. [3].

METHOD DETAILS

RNA-Seq Preprocessing and Alignment

Raw fastq sequential samples were subjected to rigorous global quality assessment utilizing FastQC. Due to the high-throughput volumetric nature of the analysis, Trimmomatic [4] was executed across all samples simultaneously utilizing the core architecture of GNU Parallel. Sequencing adapters and low-quality sequence edges were aggressively purged, adhering to a strict minimum length threshold (MINLEN:36) to unequivocally assure transcriptomic structural integrity prior to alignment. Remaining high-quality mathematical samples were aligned to the standardized human reference genome (GRCh38/hg38) via the HISAT2 memory-efficient alignment algorithm [5].

QUANTIFICATION AND STATISTICAL ANALYSIS

Differential Expression Computation

Read quantification was subsequently performed utilizing the subread featureCounts algorithm [6]. Variable biological gene-level mathematical vectors were mapped to formal gene nomenclature cross-referencing the universal GENCODE v44 annotation framework. Statistical dispersion modeling for differential gene expression was explicitly processed employing the DESeq2 bioconductor package within the R mathematical environment [7]. Raw numeric matrices initially underwent a Variance Stabilizing Transformation (VST) to effectively eliminate systemic heteroscedasticity. Genes were formally classified as Differentially Expressed Genes (DEGs) based on ultra-stringent

False Discovery Rate (FDR) thresholds: an adjusted p-value (padj) < 0.05 and an absolute Log2 Fold Change (|Log2FC|) > 2.0.

Pathway and Immune Profiling

Functional biological pathway analyses were mapped utilizing `clusterProfiler` across absolute Entrez Gene ID mappings [8]. Custom topological transcriptomic extractions specific for T-cell density (CD3D, CD4, CD8A) and immunological structural checkpoints (PDCD1, CTLA4) were algorithmically designed utilizing `ggplot2` and `pheatmap` bi-clast algorithms mapping VST values across the targeted groups. Statistical significances defining KEGG terms retained FDR boundaries at the 0.05 index limit.

ACKNOWLEDGMENTS

The author would like to acknowledge the computational resources exclusively dedicated for high-throughput genomic mapping. The author declares no specific external funding directly funding this stage of the transcriptomic analysis.

AUTHOR CONTRIBUTIONS

Conceptualization: S.H.; Methodology: S.H.; Investigation: S.H.; Validation: S.H.; Formal Analysis: S.H.; Writing – Original Draft: S.H.; Writing – Review & Editing: S.H.

DECLARATION OF INTERESTS


The author formally declares no competing financial, organizational, or academic interests.


REFERENCES

- [1] Foulkes, W. D., Smith, I. E., & Reis-Filho, J. S. (2010). Triple-negative breast cancer. *New England Journal of Medicine*, 363(20), 1938-1948.
- [2] Keenan, T. E., & Tolaney, S. M. (2020). Role of immunotherapy in triple-negative breast cancer. *Journal of the National Comprehensive Cancer Network*, 18(4), 479-489.
- [3] Varley, K. E., Gertz, J., Roberts, B. S., Davis, N. S., et al. (2014). Recurrent read-through fusion transcripts in breast cancer. *Breast Cancer Research and Treatment*, 146(2), 287-297.
- [4] Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30(15), 2114-2120.
- [5] Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature biotechnology*, 37(8), 907-915.
- [6] Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, 30(7), 923-930.


- [7] Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology*, 15(12), 1-21.
- [8] Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics: a journal of integrative biology*, 16(5), 284-287.
- [9] Costa, R. L. B., Han, H. S., & Gradishar, W. J. (2018). Targeting the PI3K/AKT/mTOR pathway in triple-negative breast cancer: a review. *Breast cancer research and treatment*, 169(3), 397-406.
- [10] Gu-Trantien, C., Loi, S., Garaud, S., et al. (2013). CD4+ follicular helper T cell infiltration predicts breast cancer survival. *Journal of Clinical Investigation*, 123(7), 2873-2892.
- [11] Heinhuis, K. M., Ros, W., Kok, M., et al. (2019). Enhancing antitumor response by combining immune checkpoint inhibitors with chemotherapy in solid tumors. *Annals of Oncology*, 30(2), 219-235.
- [12] Sharma, P., & Allison, J. P. (2015). Immune checkpoint targeting in cancer therapy: toward combination strategies with curative potential. *Cell*, 161(2), 205-214.


Principal Component Analysis (TNBC vs Normal)







Top 50 Differentially Expressed Genes






Immune Landscape: Tumor vs Normal (Hot/Cold)









Systems Biology: TNBC Hub-Gene Co-expression Network

Node size/color explicitly indicates the Master Regulator status (Hub Centrality).






Click here to access/download
Supporting Information
S1_Table_DEGs.csv





Click here to access/download
Supporting Information
S2_Table_KEGG.csv





Click here to access/download
Supporting Information
S3_Table_GO.csv

