


# THE ROLE OF ENGEL EXPANSIONS IN COLLATZ SEQUENCES

 First Last and First Last

**ABSTRACT.** The Collatz conjecture is a number theoretical problem, which has puzzled countless researchers using myriad approaches. Presently, there are scarcely investigations to treat the problem from the angle of the question "which are the corner cases the Collatz Sequences?". We pursue this question and to this end examine ascending continued fractions – the so called Engel expansions. We demonstrate that Engel expansions form worst case sequences  $v_1, v_2, \dots, v_n, v_{n+1}$  maximizing  $v_{n+1}$  and maximizing the product  $(1 + 1/3v_1)(1 + 1/3v_2) \cdots (1 + 1/3v_n)(1 + 1/3v_{n+1})$ .

## Fundamentals short and sweet

$[a]_n$	The residue class (also termed congruence class) of the integers for a modulus $n$ is the set $[a]_n = \{a + kn   k \in \mathbb{Z}\}$ and sometimes denoted by $\bar{a}_n$ or by $a + n\mathbb{Z}$ , see [1, p. 15], [2, p. 120], [3, p. 25].
$\mathbb{Z}/n\mathbb{Z}$	The set of all residue classes $[a]_n$ is called the ring of integers modulo $n$ and denoted by $\mathbb{Z}/n\mathbb{Z} = \{[a]_n   a \in \mathbb{Z}\}$ and trivially $\mathbb{Z}/0\mathbb{Z} = \mathbb{Z}$ and for all $n \neq 0$ we have $\mathbb{Z}/n\mathbb{Z} = \{[0], [1], \dots, [n-1]\}$ , see [1, p. 15], [3, p. 25].
unit	An element $a$ of a ring $R$ is called a "unit" (an invertible element) if there exist an element $b$ such that $ab = 1$ [9, p. 24]. Units are elements with inverses with respect to multiplication in the ring. Let $F$ be a field, then an element $a$ of $F$ is a non-unit iff $a = 0$ . The sum of any two non-units in $F$ is again a non-unit in $F$ .
unitary ring	A unitary ring is a ring with a multiplicative identity 1 (which differs from the additive identity $1 \neq 0$ ) such that $1a = a = a1$ for all elements $a$ of the ring.
Ideal	Let $(R, +, \cdot)$ be a commutative unitary ring. Then the subset $I \subseteq R$ is called an ideal of $R$ if $(I, +)$ is a commutative group and if $xI \subseteq I$ for all $x \in R$ , see [1, p. 66-67].

---

2010 *Mathematics Subject Classification.* 37P99.

*Key words and phrases.* Engel Expansions, Collatz Sequences.

direct prod.	If $R_1, R_2, \dots, R_n$ are rings, the cartesian product $R_1 \times R_2 \times \dots \times R_n$ forms the set of all ordered $n$ -tuples $(r_1, r_2, \dots, r_n)$ , where $r_i \in R_i$ . The addition and multiplication of these $n$ -tuples is defined "coordinatewise" by components. The resulting ring is called a "direct product" of the original rings $R_i$ [1, p. 51], [8, p. 169].
princip. ideal	A "principle ideal" is an ideal in a ring $R$ which is generated by a single element $a$ of $R$ through multiplication by every element of $R$ . There are some rings in which every ideal is a principle ideal, so-called "principle ideal rings" [1, p. 68].
max. ideal	A proper Ideal $M$ of a ring $R$ is called "maximal ideal" of $R$ if there is no other proper ideal $N$ of $R$ properly containing $M$ [8, p. 247], [9, p. 37]. A Note on "proper containment": If $R$ is any set, then $R$ is the improper subset of $R$ . Any other subset $N \neq R$ is a proper subset of $R$ and denoted by $N \subset R$ or $N \subsetneq R$ [8, p. 2].
prime ideal	Let $a$ and $b$ are two elements of $R$ and $P$ a proper ideal such that their product $ab$ is an element of $P$ . $P$ is called a prime ideal if at least one of $a$ and $b$ belongs to $P$ , in other words from $ab \in P$ and $a \notin P$ always follows $b \in P$ [9, p. 9].
max. prime ideal	A proper prime ideal $P$ is said to be a "maximal prime ideal" of the ring $R$ , if there is no other proper prime ideal containing $P$ [9, p. 23].
local ring	A commutative ring $R$ is called a local ring if it has a unique maximal ideal $M$ [10, p. 522].
Noeth. ring	A ring $R$ is called "Noetherian" when in $R$ the maximal condition for ideals is satisfied, in other words if every ideal $I$ of $R$ is finitely generated, that is, if we can find a finite set $a_1, a_2, \dots, a_n$ of elements, such that $I = Ra_1 + Ra_2 + \dots + Ra_n$ [9, p. 19, 101].
semi- local ring	A semi-local ring is a Noetherian ring which has only a finite number of maximal ideals [9, p. 107].

zero seq.	A zero sequence is a sequence, which converges towards 0 [5, p. 154]. Given the context of ideal theory, let $R$ be a ring and $I$ an ideal. In the ring $R^{\mathbb{N}} = \prod_{n \in \mathbb{N}} R$ , which is the repeated direct product of $R$ with itself, a sequence $(x_n)_{n \in \mathbb{N}}$ is called a zero sequence if for every $s \in \mathbb{N}$ there exist a $N \in \mathbb{N}$ (depending on $s$ ) such that $x_n \in I^s$ for all $n > N$ .
Cauchy seq. in $\mathbb{Q}, \mathbb{R}$	A sequence $(x_n)_{n \in \mathbb{N}}$ in $\mathbb{Q}$ or $\mathbb{R}$ is a Cauchy sequence if for any $\epsilon > 0$ there exists a positive integer $N$ such that $ x_n - x_m  < \epsilon$ for all $n, m \geq N$ , see [5, p. 153], [11, p. 24].
Cauchy seq. in a ring	Let $(x_n)$ be a sequence of elements in $R^{\mathbb{N}}$ , the repeated direct product of a ring with itself, and $I$ an ideal in $R$ . The sequence $(x_n)$ is a Cauchy sequence if for every $s \in \mathbb{N}$ there exist a $N \in \mathbb{N}$ such that $x_n - x_m \in I^s$ for all $n, m > N$ . Let $(x_n)$ be a sequence of elements in a local ring $R$ and $M$ is the maximal ideal of $R$ . The sequence $(x_n)$ is a Cauchy sequence if, given any $s \in \mathbb{N}$ , we can always find an integer $N$ such that $x_n - x_m \in M^s$ whenever $n > m > N$ , see [9, p. 63, 85]. The sequence $(x_n)$ is a Cauchy sequence iff $x_n - x_{n-1} \rightarrow 0$ as $n \rightarrow \infty$ [9, p. 85].
compl. of a ring	Let $R$ be a ring and $I$ an ideal.
concor. ext.	Let $R, S$ be local rings. If a sequence of elements of $S$ is a Cauchy sequence in $S$ iff it is a Cauchy sequence in $R$ , then we say that $R$ is a "concordant extension" of $S$ [9, p. 87]. When $R, S$ are semi-local rings $R \subseteq S$ , $R$ is said to be a "concordant extension" of $S$ if a sequence $(s_n)$ of elements in $S$ is regular in $S$ iff $(s_n)$ is regular in $R$ [12].
compl. of a local ring	Let $S$ be a local ring. A local ring $R$ will be called a completion of $S$ if $R$ is a concordant extension of $S$ and $R$ is complete and if every element of $R$ is the limit of a sequence of elements of $S$ . Each local ring has a completion [9, p. 92].
compl. local ring	A local ring $R$ is called "complete" if every Cauchy sequence composed of elements of $R$ has a limit in $R$ [9, p. 85], [13, p. 184].
$p$ -adic val. for $\mathbb{Z}$	Fix a prime number $p$ in $\mathbb{Z}$ . The $p$ -adic valuation of a nonzero integer $n = r \cdot p^{v_p(n)}$ is the highest exponent $v_p(n)$ such that $p^{v_p(n)}$ divides $n$ (we say $p^{v_p(n)}$ divides $n$ "exactly"). Hence $p$ and $r$ are coprime. If $n, p$ are coprime then $v_p(n) = 0$ , and by convention $v_p(0) = \infty$ , see [4].

$p$ -adic val. for $\mathbb{Q}$	The $p$ -adic valuation can be extended to the field of rational numbers. Let $x = n \cdot s^{-1}$ be a rational number, then $v_p(x) = v_p(n) - v_p(s)$ . Any nonzero rational number $x$ can be uniquely represented as $x = rp^{v_p(x)}s^{-1}$ , where $r, s \in \mathbb{Z}$ , $s > 0$ , and $\gcd(r, s) = \gcd(r, p) = \gcd(s, p) = 1$ , see [5, p. 154], [6].
$p$ -adic norm	Let $x$ be any number in $\mathbb{Q}$ , for which we already know that it can be written as $x = rp^{v_p(x)}s^{-1}$ , where $p$ is a prime number, $s > 0$ and $r$ are integers not divisible by $p$ . The $p$ -adic norm of $x$ is defined by $ x _p = p^{-v_p(x)}$ for $x \neq 0$ , and $ 0 _p = 0$ , see [4], [5, p. 154], [7].
$\mathbb{Z}_p$	The ring $\mathbb{Z}_p$ is the completion of $\mathbb{Z}$ with respect to the $p$ -adic norm. That is, $\mathbb{Z}_p$ is the set of all equivalence classes of Cauchy sequences $(a_n)$ where $(a_n)$ and $(b_n)$ are equivalent if $\lim_{n \rightarrow \infty}  a_n - b_n _p = 0$ , see [14].

## 1. INTRODUCTION

The Collatz conjecture is a well-known number theory problem and is the subject of numerous publications. An overview is provided by Lagarias [15]. Therefore, our description of the topic will be brief. The mathematician Lothar Collatz introduced a function  $g : \mathbb{N} \rightarrow \mathbb{N}$  as follows:

$$(1) \quad g(x) = \begin{cases} 3x + 1 & 2 \nmid x \\ x/2 & \text{otherwise} \end{cases}$$

In the following, we only consider compressed Collatz sequences that solely contain the odd members, such as described by Bruckman [16], who used the more convenient function that opts out all even integers:

$$(2) \quad f(x) = (3x + 1) \cdot 2^{-\alpha(x)}, \text{ where } 2^{\alpha(x)} \parallel (3x + 1)$$

Note that  $\alpha(x)$  is the largest possible exponent for which  $2^{\alpha(x)}$  exactly divides  $3x + 1$ . Especially for prime powers, one often says  $p^\alpha$  *divides* the integer  $x$  *exactly*, denoted as  $p^\alpha \parallel x$ , if  $p^\alpha$  is the greatest power of the prime  $p$  that divides  $x$ .

A (compressed) Collatz sequence  $v_1, v_2, \dots, v_n, v_{n+1}$  allowed at most one division by 2 between two successive members. Dividing only once between two successive members, maximizes  $v_{n+1}$ . Such a sequence forms the following ascending continued fraction (cf. also [17, p. 11]):

$$(3) \quad v_{n+1} = \frac{3 \frac{3 \frac{3 \frac{3v_1 + 1}{2} + 1}{2} + 1}{2}}{2} \dots = \frac{3^n v_1 + \sum_{i=0}^{n-1} 3^i 2^{n-1-i}}{2^n} = \frac{3^n(v_1 + 1) - 2^n}{2^n}$$

**Example 1.1.** A concrete example for such a sequence is  $v_1 = 31$ ,  $v_2 = 47$ ,  $v_3 = 71$ ,  $v_4 = 107$ ,  $v_5 = 161$ . And, to follow that example, we can calculate  $v_5$  in a straightforward way:

$$v_5 = v_{n+1} = \frac{3^4(31 + 1) - 2^4}{2^4} = 161$$

Besides, by choosing a starting number  $v_1 = 2^{n+1} - 1$ , we are able to infinitely generate sequences each forming an ascending continued fraction. As per equation 3 the last member in this sequence is the odd number  $v_{n+1} = 3^n \cdot 2 - 1$ .

**Remark 1.2.** Ascending variants of a continued fraction, such as used in equation 3, shall not be confused with continued fractions as treated in [18], [19], [20]. Ascending continued fractions used in our case correspond to the so-called "Engel Expansions" [21].

As illustrated below, we can formulate the ascending continued fractions in a generalized fashion, whereas the analogy to 3 is given by  $b_1 = b_2 = b_3 = b_4 = 2$  and  $a_1 = 3^0$ ,  $a_2 = 3^1$ ,  $a_3 = 3^2$  and  $a_4 = 3^3 + 3^4 v_1$ :

$$\frac{a_1 + \frac{a_2 + \frac{a_3 + \frac{a_4}{b_4}}{b_3}}{b_2}}{b_1} \dots = \frac{a_1}{b_1} + \frac{a_2}{b_1 b_2} + \frac{a_3}{b_1 b_2 b_3} + \frac{a_4}{b_1 b_2 b_3 b_4} + \dots$$

The generalized form of equation 3 may be used to compute any of the above-named ascending continued fraction that has  $a_i = k^{i-1}$ ,  $b_i = b$  for  $i \in \mathbb{N}$  and  $a_n = k^{n-1} + k^n v_1$ :

$$(4) \quad v_{n+1} = \frac{k^n(kv_1 - bv_1 + 1) - b^n}{b^n(k - b)}$$

## 2. INCLUDE MORE DIVISIONS BY TWO INTO AN ENGEL EXPANSION

For calculating the largest possible  $v_{n+1}$ , we considered so far Engel expansions which contain only  $n$  division by two within a Collatz sequence of  $n + 1$  members. In the following we include  $m$  additional divisions by two and thus a total of  $m + n$  divisions.

Let us take a look at two corner cases:

- the one where we do the additional  $m$  divisions by 2 at the end and
- the one where we do these additional divisions at the very beginning.

**The first case** is our starting point to examine how the swapping a division by two affects the node  $v_{n+1}$ . For this, let us compare the Engel expansion where we divide by  $2^m$  afterwards with one where we divide by 2 in the penultimate step and by  $2^{m-1}$  in last step. One can immediately recognize the following inequality with a mere look:

$$\frac{1 + \frac{3 + \frac{3^2 + \frac{3^3 + 3^4 v_1}{2}}{2}}{2}}{2 \cdot 2^m} < \frac{1 + \frac{3 + \frac{3^2 + \frac{3^3 + 3^4 v_1}{2}}{2}}{2 \cdot \mathbf{2}}}{2 \cdot 2^{m-1}}$$

To put it simply, in the expansion on the right side of the above-shown inequality we perform one division by two a little bit earlier as we do it in the expansion on the left side of the expansion. Almost all summands of both expansions cancel out each other:

$$\frac{1}{2 \cdot 2^m} + \frac{3}{\cancel{2^2 \cdot 2^m}} + \frac{3^2}{2^3 \cdot 2^m} + \frac{3^3 + 3^4 v_1}{2^4 \cdot 2^m} < \frac{1}{2 \cdot 2^{m-1}} + \frac{3}{\cancel{2^2 \cdot \mathbf{2} \cdot 2^{m-1}}} + \frac{3^2}{2^3 \cdot \mathbf{2} \cdot 2^{m-1}} + \frac{3^3 + 3^4 v_1}{2^4 \cdot \mathbf{2} \cdot 2^{m-1}}$$

**The second case** deals with Engel expansions where we perform that additional  $m$  divisions by two as early as possible. The resulting value decreases, when we make a division by two later:

$$\frac{1 + \frac{3 + \frac{3^2 + \frac{3^3 + 3^4 v_1}{2 \cdot 2^{m-1}}}{2 \cdot \mathbf{2}}}{2}}{2} < \frac{1 + \frac{3 + \frac{3^2 + \frac{3^3 + 3^4 v_1}{2 \cdot 2^m}}{2}}{2}}{2}$$

Also here almost all summands of both Engel expansions, they cancel each other out:

$$\frac{1}{2} \cancel{+ \frac{3}{2^2}} + \frac{3^2}{2^3 \cdot \mathbf{2}} + \frac{3^3 + 3^4 v_1}{\cancel{2^4 \cdot \mathbf{2} \cdot 2^{m-1}}} < \frac{1}{2} \cancel{+ \frac{3}{2^2}} + \frac{3^2}{2^3} + \frac{3^3 + 3^4 v_1}{\cancel{2^4 \cdot 2^m}}$$

While the first case minimizes the value of  $v_{n+1}$ , the second case maximizes it. The difference between the maximum and the minimum is given by the following equation:

$$\frac{3^{n-1} \left( \frac{3v_1+1}{2 \cdot 2^m} + 1 \right) - 2^{n-1}}{2^{n-1}} - \frac{3^n (v_1 + 1) - 2^n}{2^{n+m}} = \left( \frac{3^{n-1}}{2^{n-1}} - 1 \right) \left( 1 - \frac{1}{2^m} \right)$$

This has the consequence that for a given sequence consisting of  $n + 1$  members, between which a total of  $n + m$  divisions have taken place, the permutation of these divisions has a very limited effect on the node  $v_{n+1}$  as described by theorem 2.1.

**Theorem 2.1.** *Let  $v_1, v_2, \dots, v_n, v_{n+1}$  be a sequence in which a total of  $n + m$  divisions by two took place. No matter how these divisions are permuted, i.e. performed sooner or later, the last member  $v_{n+1}$  can differ at most by the following product:*

$$\left( \frac{3^{n-1}}{2^{n-1}} - 1 \right) \left( 1 - \frac{1}{2^m} \right)$$

The Engel expansion of the second case (that maximizes the value of  $v_{n+1}$ ) provides the following formula for calculating the sequence member  $v_{n+1}$ :

$$(5) \quad v_{n+1} = \left( \frac{k}{2} \right)^{n-1} \left( \frac{kv_1 + 1}{2^{\alpha_1}} + \frac{1}{k-2} \right) - \frac{1}{k-2} = \left( \frac{k}{2} \right)^{n-1} \left( v_2 + \frac{1}{k-2} \right) - \frac{1}{k-2}$$

Let us refer to example 1.1 using the sequence starting at  $v_1 = 31$ ,  $v_2 = 47$  and so forth. We calculate  $v_5$  directly as follows:

$$v_5 = \left( \frac{3}{2} \right)^{4-1} \left( 47 + \frac{1}{3-2} \right) - \frac{1}{3-2} = 161$$

**Example 2.2.** *As another example we choose the sequence for  $k = 5$  starting at  $v_1 = 67$  continuing with  $v_2 = 21$ ,  $v_3 = 53$ ,  $v_4 = 133$ ,  $v_5 = 333$ ,  $v_6 = 833$  and  $v_7 = 2083$ . The last sequence member can be calculated by equation 5 directly as follows:*

$$v_7 = \left( \frac{5}{2} \right)^{6-1} \left( 21 + \frac{1}{5-2} \right) - \frac{1}{5-2} = 2083$$

### 3. SUM OF RECIPROCATED COLLATZ MEMBERS

A product  $\prod(1+a_n)$  with positive terms  $a_n$  is convergent if the series  $\sum a_n$  converges, see Knopp [22, p. 220]. A similar statement provides Murphy [23], who write the factors in the form  $c_n = 1+a_n$  and explains that if  $\prod c_n$  is convergent then  $c_n \rightarrow 1$  and therefore if  $\prod(1+a_n)$  is convergent then  $a_n \rightarrow 0$ .

We write the sum of reciprocated Collatz members as  $1/kv_1 + 1/kv_2 + \dots + 1/kv_n + 1/kv_{n+1}$ . In order to formulate this sum independently from the successive members  $v_2, v_3, \dots$ , we substitute these as follows:

$$\begin{aligned}
 v_1 &= v_1 \\
 v_2 &= \frac{kv_1 + 1}{2^{\alpha_1}} \\
 v_3 &= \frac{k^2v_1 + k + 2^{\alpha_1}}{2^{\alpha_1+\alpha_2}} \\
 v_4 &= \frac{k^3v_1 + k^2 + k \cdot 2^{\alpha_1} + 2^{\alpha_1+\alpha_2}}{2^{\alpha_1+\alpha_2+\alpha_3}} \\
 &\vdots \\
 v_{n+1} &= \frac{k^n v_1 + \sum_{j=1}^n k^{j-1} 2^{\alpha_1+\dots+\alpha_n-\sum_{l>n-j} \alpha_l}}{2^{\alpha_1+\dots+\alpha_n}}
 \end{aligned}
 \tag{6}$$

The sum of the reciprocal Collatz sequence members can be expressed as a term that only depends from  $v_1$  and from the number of divisions by two  $\alpha_1, \alpha_2, \alpha_3, \dots$  between two successive members:

$$\sum_{i=1}^{n+1} \frac{1}{kv_i} = \frac{1}{k} \left( \frac{1}{v_1} + \sum_{i=1}^n \frac{1}{v_{i+1}} \right) = \frac{1}{k} \left( \frac{1}{v_1} + \sum_{i=1}^n \frac{2^{\alpha_1+\dots+\alpha_i}}{k^i v_1 + \sum_{j=1}^i k^{j-1} 2^{\alpha_1+\dots+\alpha_n-\sum_{l>i-j} \alpha_l}} \right)$$

### 4. THE PRODUCT OF RECIPROCATED COLLATZ MEMBERS INCREMENTED BY ONE

In a similar way to deduce the sum of reciprocal vertices depending only on  $v_1$  as performed in 3, we evolve the formula for the product of reciprocated Collatz members (incremented by one):



$$(8) \quad \prod_{i=1}^{n+1} \left( 1 + \frac{1}{kv_i} \right) = 1 + \frac{2^{\alpha_1+\dots+\alpha_n} + k \cdot 2^{\alpha_1+\dots+\alpha_{n-1}} + \dots + k^{n-1} \cdot 2^{\alpha_1} + k^n}{k^{n+1}v_1}$$

$$(9) \quad = 1 + \frac{2^{\alpha_1+\dots+\alpha_n} + k \cdot \sum_{j=1}^i k^{j-1} 2^{\alpha_1+\dots+\alpha_n - \sum_{l>i-j} \alpha_l}}{k^{n+1}v_1}$$

$$(10) \quad = \frac{2^{\alpha_1+\dots+\alpha_n} (1 + kv_{n+1})}{k^{n+1}v_1}$$

We inserted the sum used in equation 7 into the above-given equation 8 and then obtained equation 9. Now let us divide this product by the last factor in order to retrieve the product which iterates to  $n$  instead of  $n+1$ :

$$(11) \quad \prod_{i=1}^n \left( 1 + \frac{1}{kv_i} \right) = \frac{\prod_{i=1}^{n+1} \left( 1 + \frac{1}{kv_i} \right)}{\frac{kv_{n+1}+1}{kv_{n+1}}} = \frac{2^{\alpha_1+\dots+\alpha_n} v_{n+1}}{k^n v_1}$$

The above-shown equation 11 becomes simplified, when we replaced the numerator by equation 10. The question which sequence maximizes its last member  $v_{n+1}$  ties into the question: Which sequence maximizes the product? The product formula 11 does not depend from all vertices  $v_1, v_2, \dots, v_n$ , it depends only from  $2^\alpha = 2^{\alpha_1+\dots+\alpha_n}$ , from the first sequence member  $v_1$  and the final one  $v_{n+1}$ .

Consider a Collatz sequence containing  $n$  elements and starting at a given integer  $v_1$ . The corresponding product given by equation 11 becomes as large as possible if we maximize the last member  $v_{n+1}$ . This maximum occurs when the sequence is an Engel expansion, id est when we run the most divisions by two at the beginning. Consequently, the exponent alpha (the total number of divisions by two) is the sum of a large  $\alpha_1$  and the remaining alpha values which are all one:

$$\alpha = n + m = \alpha_1 + \alpha_2 + \dots + \alpha_n = \alpha_1 + 1 + \dots + 1 = \alpha_1 + n - 1$$

The product of reciprocated Collatz members (incremented by one) for an Engel expansion is given by the following equation:

$$(12) \quad \prod_{i=1}^n \left( 1 + \frac{1}{kv_i} \right) = 1 + \frac{1}{kv_1} + \frac{2^{\alpha_1}}{k(k-2)v_1} \left( 1 - \left( \frac{2}{k} \right)^{n-1} \right)$$

$$(13) \quad = 1 + \frac{1}{kv_1} + \frac{kv_1+1}{k(k-2)v_1v_2} \left( 1 - \left( \frac{2}{k} \right)^{n-1} \right)$$

**Example 4.1.** An example for  $k = 3$  provides the sequence  $v_1 = 661$ ,  $v_2 = 31$ ,  $v_3 = 47$ , and  $v_4 = 71$ . In this case  $\alpha_1 = 6 = m + 1$  and  $\alpha_2 = \alpha_3 = \alpha_4 = 1$ . We now calculate the product of reciprocated Collatz sequence members by inserting  $v_1 = 661$  and  $v_2 = 31$  together with  $k = 3$  and  $n = 4$  into equation 13:

$$\begin{aligned} \prod_{i=1}^4 \left(1 + \frac{1}{3v_i}\right) &= \left(1 + \frac{1}{3 \cdot 661}\right) \left(1 + \frac{1}{3 \cdot 31}\right) \left(1 + \frac{1}{3 \cdot 47}\right) \left(1 + \frac{1}{3 \cdot 71}\right) \\ &= 1 + \frac{1}{3 \cdot 661} + \frac{3 \cdot 661 + 1}{3 \cdot (3 - 2) \cdot 661 \cdot 31} \left(1 - \left(\frac{2}{3}\right)^{4-1}\right) \\ &= 1.0232158532713247 \end{aligned}$$

Maximizing the product of reciprocated Collatz sequence members  $\prod_{i=1}^n (1 + 1/kv_i)$  requires us to maximize the equation 13.

$k$	product formula	maximum case	resulting product
3	$1 + \frac{1}{3v_1} + \frac{3v_1+1}{3v_1v_2} \left(1 - \left(\frac{2}{3}\right)^{n-1}\right)$	$v_1 = 1, n = 1$	$\frac{4}{3}$
5	$1 + \frac{1}{5v_1} + \frac{5v_1+1}{15v_1v_2} \left(1 - \left(\frac{2}{5}\right)^{n-1}\right)$	$v_1 = 1, v_2 = 3, n = 2$	$\frac{32}{25}$
7	$1 + \frac{1}{7v_1} + \frac{7v_1+1}{35v_1v_2} \left(1 - \left(\frac{2}{7}\right)^{n-1}\right)$	$v_1 = 1, n = 1$	$\frac{8}{7}$

TABLE 1. Formulas that calculate the Engel expansion's product for  $k = 3, 5, 7$

## 5. CONDITION FOR A LIMITED GROWTH OF THE ENGEL EXPANSION

Let us look now into the question of what condition must be met to prevent a greater growth than a decline in Collatz sequences. Specifically we consider an Engel expansion comprising  $n + 1$  sequence members that include  $m$  additional divisions by two at the beginning. The last member  $v_{n+1}$  in such a sequence can be calculated by formula ???. In order to restrict the growth of this sequence, we require that the last member has to be smaller than the first one. For this we define the condition  $v_{n+1} < v_1$ :

$$\frac{3^n v_1 + 3^{n-1} + 3^{n-1} 2^{m+1}}{2^{m+n}} - 1 < v_1$$

Reshaping this inequality leads to the following condition:

$$(14) \quad \frac{3^{n-1} (2^{m+1} - 2)}{2^{m+n} - 3^n} - 1 < v_1$$

## REFERENCES

- [1] J. Wolfart. *Einführung in die Zahlentheorie und Algebra*. Vieweg+Teubner, Wiesbaden, Germany, 2 edition, 2011.
- [2] M. Schubert. *Mathematik für Informatiker*. Vieweg+Teubner, Wiesbaden, Germany, 2009.
- [3] S. Müller-Stach and J. Piontkowski. *Elementare und algebraische Zahlentheorie*. Vieweg+Teubner, Wiesbaden, Germany, 2 edition, 2011.
- [4] T. C. Herwig. The p-adic completion of  $\mathbb{Q}$  and hensel's lemma. Technical report, Department of Mathematics, University of Chicago, 2011.
- [5] A. Schmidt. *Einführung in die algebraische Zahlentheorie*. Springer, Berlin, Germany, 2007.
- [6] E. W. Weisstein. "p-adic number." from mathworld—a wolfram web resource.
- [7] E. W. Weisstein. "p-adic norm." from mathworld—a wolfram web resource.
- [8] J. B. Fraleigh. *A First Course in Abstract Algebra*. Pearson, Harlow, United Kingdom, 7 edition, 2014.
- [9] D. G. Northcott. *Ideal Theory*, volume 42 of *Cambridge Tracts in Mathematics and Mathematical Physics*. Cambridge University Press, Cambridge, United Kingdom, 1953.
- [10] J. J. Rotman. *A First Course in Abstract Algebra*. Prentice Hall, Upper Saddle River, NJ, 3 edition, 2005.
- [11] N. J. Higham. *The Princeton Companion to Applied Mathematics*. Princeton University Press, Princeton, NJ, 2015.
- [12] E. H. Batho. Some remarks on non-commutative extensions of local rings. *Nagoya Mathematical Journal*, 14.
- [13] G. Kemper. *A Course in Commutative Algebra*. Springer, Heidelberg, Germany, 2011.
- [14] A. Gupta. The p-adic integers, analytically and algebraically. Technical report, Department of Mathematics, University of Chicago, 2018.
- [15] J. C. Lagarias. *The Ultimate Challenge: The  $3x+1$  Problem*. American Mathematical Society, Providence, RI, 2010.
- [16] P. S. Bruckman. Retracted article: A proof of the collatz conjecture. *International Journal of Mathematical Education in Science and Technology*, 39(3):403–407, 2008.
- [17] T.M.M. Laarhoven. The  $3n + 1$  conjecture, 7 2009.
- [18] C. G. Moore. *Introduction to Continued Fractions*. National Council of Teachers of Mathematics, 1964.
- [19] D. Hensley. *Continued Fractions*. World Scientific Publishing, 2006.
- [20] J. Borwe, A. van der Poorten, J. Shallit, and W. Zudlin. *Neverending Fractions: An Introduction to Continued Fractions*. Cambridge University Press, 2014.
- [21] C. Kraaikamp and J. Wu. On a new continued fraction expansion with non-decreasing partial quotients. *Monatshefte für Mathematik*, 143:285–298, 2004.
- [22] K. Knopp. *Theorie und Anwendung der Unendlichen Reihen*. Springer, 2 edition, 1924.
- [23] T. Murphy. 2006 course 4281: Prime numbers, 2006.

FIRST LASTNAME, GRADUATE SCHOOL OF MATHEMATICS, XYZ UNIVERSITY, CITY, ADRESSZUSATZ,  
ZIP, GERMANY

*Email address:* `first.last@university.de`

FIRST LASTNAME, GRADUATE SCHOOL OF MATHEMATICS, XYZ UNIVERSITY, CITY, ADRESSZUSATZ,  
ZIP, GERMANY

*Email address:* `first.last@university.de`