

CLINICAL DATA ANALYSIS AND REPORTING

Simulated Interim Analysis Report for a Virtual Clinical Study

Virtual RCT of a Smartphone-Delivered CBT Application for Mild-to-Moderate Generalized Anxiety in Adults

Protocol Number: VCR-2026-001

Report Version: 1.0

Date: August 29, 2026

Study Type: Hypothetical / Training Exercise — Simulated Dataset

Prepared by:

[SUMEDH DIGAMBARRAO JOSHI]

Clinical Research Assistant Intern – YuvaIntern

Week 4 Task: Clinical Data Analysis and Reporting

Table of Contents

Sr. No	CONTENTS	PAGE NUMBER
1	Introduction	3
2	Data Analysis Methodology	3
3	Results	3
3.1	Participant Flow and Attrition	3
3.2	Primary Outcome: GAD-7 Anxiety Symptoms	4
3.3	Secondary Outcome: PHQ-9 Depressive Symptoms	5
3.4	Engagement and Usability	6
4	Discussion	7
5	Conclusions and Recommendations	8
6	References	8

1. Introduction

This report presents a simulated interim analysis of outcome data for study VCR-2026-001, the virtual randomized controlled trial evaluating an 8-week smartphone-delivered CBT application against a waitlist control for adults with mild-to-moderate generalized anxiety. Since real participant data does not yet exist for this hypothetical study, a synthetic dataset of 120 participants (60 per arm) was generated in Python to mimic the score distributions, attrition pattern, and treatment effect size specified in the Week 1 protocol and Week 1 statistical analysis plan, so that the analysis pipeline, reporting format, and interpretation approach could be developed and validated before real data collection begins.

The purpose of this exercise is to demonstrate the full analysis workflow the study team would execute at each planned interim look: computing descriptive statistics, testing the primary and secondary hypotheses, visualizing trends, and translating the statistical results into clinically meaningful conclusions.

2. Data Analysis Methodology

2.1 Dataset

The simulated dataset contains 120 enrolled participants (60 CBT App, 60 Waitlist), each with a baseline GAD-7 score drawn from a truncated normal distribution (mean 9.2, SD 2.1, bounded to the protocol's 5–14 inclusion range). A 20% cumulative dropout by Week 12 was built into the simulation, consistent with the attrition assumption in the Week 1 power calculation. A treatment effect was simulated for the CBT App arm consistent with the between-group difference (≈ 2.5 GAD-7 points) the Week 1 sample-size calculation was powered to detect.

2.2 Statistical Methods

- Descriptive statistics (n, mean, standard deviation) computed for GAD-7 and PHQ-9 scores at each timepoint, by arm.
- Primary and secondary endpoint analysis: within-participant change score from Baseline to Week 8, compared between arms using Welch's independent-samples t-test (which does not assume equal variances between arms).
- Effect size: Cohen's d computed for the primary endpoint using the pooled standard deviation of the change scores.
- Association between engagement (modules completed) and outcome: Pearson correlation coefficient.
- All analysis performed in Python (pandas 2.x, scipy.stats 1.17), with $\alpha = 0.05$ as the significance threshold, matching the Week 1 statistical analysis plan.

Justification: Welch's t-test was used instead of the standard (Student's) t-test because the two arms showed unequal variance in change scores in this simulated sample, and Welch's version is more robust to that violation without requiring an additional equal-variance test step.

3. Results

3.1 Participant Flow and Attrition

Enrolled: 120 participants (60 CBT App, 60 Waitlist)

Completed through Week 12: 96 participants (80.0%)

Overall attrition: 20.0%, matching the rate assumed in the Week 1 sample-size calculation

Dropout was modeled as somewhat more likely among participants with higher baseline anxiety severity, reflecting a realistic pattern in which more symptomatic participants are more prone to disengage from a self-guided digital intervention — a pattern documented in prior digital-mental-health trials and worth monitoring closely once real recruitment begins.

3.2 Primary Outcome: GAD-7 Anxiety Symptoms

Table 1 summarizes mean GAD-7 scores by arm and time point.

Time point	Arm	n	Mean	SD
Baseline	CBT App	60	8.87	1.94
Baseline	Waitlist	60	9.18	1.72
Week 4	CBT App	57	7.37	3.19
Week 4	Waitlist	57	8.47	4.01
Week 8	CBT App	55	5.67	3.89
Week 8	Waitlist	54	9.06	3.70
Week 12	CBT App	49	5.04	3.58
Week 12	Waitlist	47	7.96	3.37




Figure 1. Mean GAD-7 score declined steadily in the CBT App arm while remaining relatively flat in the Waitlist arm, with both arms trending toward the shaded “minimal anxiety” range (GAD-7 < 5) only in the CBT App arm by Week 12.




Figure 2. Primary endpoint comparison: mean GAD-7 reduction from Baseline to Week 8, by arm, with SEM error bars.

Primary endpoint analysis (Baseline → Week 8 change): the CBT App arm showed a mean reduction of 2.96 points (SD 3.42, n=55), compared with a mean change of -0.11 points (SD 3.28, n=54) in the Waitlist arm — i.e., the Waitlist arm's symptoms were essentially unchanged. The between-arm difference of 3.07 points was statistically significant (Welch's $t=4.80$, $p<0.0001$), with a large effect size (Cohen's $d=0.92$).

This result exceeds the 2.5-point between-group difference the Week 1 protocol's sample-size calculation was powered to detect, meaning the study — in this simulated dataset — was adequately powered to detect the primary treatment effect.

3.3 Secondary Outcome: PHQ-9 Depressive Symptoms

Timepoint	Arm	n	Mean	SD
Baseline	CBT App	60	6.88	2.21
Baseline	Waitlist	60	6.88	2.65
Week 4	CBT App	57	5.61	3.00
Week 4	Waitlist	57	6.33	4.43
Week 8	CBT App	55	4.78	3.46
Week 8	Waitlist	54	6.24	3.53
Week 12	CBT App	49	4.04	3.05
Week 12	Waitlist	47	5.53	3.68

Baseline PHQ-9 scores were nearly identical between arms (both means = 6.88), confirming successful randomization on this secondary measure. By Week 8, the CBT App arm improved by a mean of 1.96 points (SD 2.00) versus 0.22 points (SD 2.14) in the Waitlist arm — a statistically significant difference (Welch's $t=3.07$, $p=0.0027$), though smaller in magnitude than the primary GAD-7 effect, consistent with the app's design as an anxiety-focused (not depression-focused) intervention.

3.4 Engagement and Usability

Metric (CBT App arm, n=60)	Mean	SD	Min	Max
Modules completed (of 8)	5.8	1.2	3	8
Total session time (minutes)	144.6	49.5	20	260.9
System Usability Scale (SUS) score	77.2	9.8	52.7	98.6

The mean SUS score of 77.2 falls in the “good” usability range, and 85.0% of participants scored above 68 — the threshold Bangor et al. (2009) identify as above-average usability across benchmarked digital products — suggesting the application interface itself was not a major barrier to engagement.

Figure 3. Distribution of Module Completion — CBT App Arm (n=60)




Figure 3. Most CBT App participants completed 5–6 of the 8 available modules; very few completed fewer than 3.

Figure 4. Module Completion vs. Symptom Improvement (CBT App Arm)




Figure 4. Modules completed vs. GAD-7 improvement (CBT App arm), with a linear trend line.

Contrary to the intuitive expectation that more app usage would predict better outcomes, the correlation between modules completed and GAD-7 improvement was weak and not statistically significant (Pearson $r=-0.23$, $p=0.0945$, $n=55$). This is reported honestly as a null finding rather than a confirmed relationship — the direction and non-significance should not be over-interpreted given the modest sample size, but it does suggest that module count alone is an incomplete proxy for therapeutic engagement, and that future analyses should examine engagement quality (e.g., completion of specific high-yield modules like cognitive restructuring) rather than raw count.

4. Discussion

The simulated results support the primary hypothesis: participants using the CBT application showed a significantly greater reduction in anxiety symptoms than the waitlist control group by Week 8, with the effect largely maintained through the Week 12 follow-up (CBT App mean 5.04 vs. Waitlist mean 7.96). The magnitude of improvement (≈ 3 GAD-7 points) is clinically meaningful — GAD-7 reductions of this size are generally considered to reflect a noticeable improvement in day-to-day functioning, not just a statistical artifact.

The secondary PHQ-9 finding suggests the intervention's benefit may extend modestly to co-occurring depressive symptoms, which is consistent with the substantial symptom overlap between anxiety and depression and with CBT's transdiagnostic mechanism (cognitive restructuring and behavioral activation techniques apply to both conditions).

The usability data (mean SUS 77.2, 85% above the above-average threshold) indicate the application itself was generally easy to use, which is an important precondition for a digital intervention — a good intervention delivered through a confusing interface would show weaker real-world effects than what a controlled trial detects. However, the null engagement-outcome correlation is a meaningful caution: it suggests the study team should not assume that simply pushing participants toward higher module counts will improve outcomes, and should instead investigate whether specific modules or usage patterns (e.g., completing exposure exercises specifically) matter more than total volume.

Limitations of this analysis: (1) this is simulated, not real, data, so the specific effect sizes should not be treated as evidence of the actual application's efficacy; (2) attrition, while modeled at the assumed 20% rate, was concentrated somewhat more heavily among higher-severity participants, which could bias a real study's results toward the less-symptomatic survivors if not addressed with the multiple-imputation approach specified in the Week 1 statistical analysis plan; (3) the Week 12 sample sizes ($n=49$ and $n=47$) are noticeably smaller than the Week 8 sample, so the durability finding should be interpreted with more caution than the Week 8 primary result.

5. Conclusions and Recommendations

1. The primary analysis pipeline is validated: the simulated data confirms the planned statistical approach (Welch's t-test on Baseline-to-Week-8 change scores) can detect a treatment effect of the magnitude the study was powered for, and the reporting format (descriptive tables + trend and comparison figures) clearly communicates both statistical and clinical significance.
2. When real data collection begins, prioritize monitoring differential attrition by baseline severity in real time, since this simulation shows it can plausibly emerge and would need to be addressed analytically (multiple imputation) rather than ignored.
3. Add an engagement-quality metric (e.g., which specific modules were completed) in addition to raw module count, since raw count alone showed no significant relationship to outcome in this simulation and may be an incomplete engagement measure.
4. Continue tracking the Week 12 durability outcome separately from the Week 8 primary endpoint, given its smaller completer sample size, and consider a sensitivity analysis restricted to Week-12 completers versus an imputed intent-to-treat estimate.

5. Retain the current SUS-based usability check as a standard component of future interim reports, since a usability score below the above-average threshold would be an early warning sign of an implementation (rather than efficacy) problem.

6. References

Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine*, 166(10), 1092–1097.

Kroenke, K., Spitzer, R. L., Williams, J. B., & Löwe, B. (2010). The Patient Health Questionnaire somatic, anxiety, and depressive symptom scales. *General Hospital Psychiatry*, 32(4), 345–359.

Bangor, A., Kortum, P., & Miller, J. (2009). Determining what individual SUS scores mean: Adding an adjective rating scale. *Journal of Usability Studies*, 4(3), 114–123.

Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences* (2nd ed.). Lawrence Erlbaum Associates.

Virtual Study Protocol, VCR-2026-001, Week 1 deliverable (statistical analysis plan and sample-size calculation).

Data Collection and Management Strategy, VCR-2026-001, Week 2 deliverable.