


PH3101 Classical Mechanics  
@2025

Rajesh Kumble Nayak

Lecture 07

**Lagrange Equation of Motion**



PH3101 Classical Mechanics  
@2025

Rajesh Kumble Nayak

Lecture 07

**Lagrange Equation of Motion**

# Newton's Equation of motion

# Newton's Equation of motion

In the last class we see that, the generalised coordinate systems are best suited for applying laws of motion, because it is free from the constraints.

# Newton's Equation of motion

In the last class we see that, the **generalised coordinate systems** are best suited for applying laws of motion, because it is free from the constraints.

These are the detailed formal steps

First we need the relation between the coordinate system

# Newton's Equation of motion

In the last class we see that, the **generalised coordinate systems** are best suited for applying laws of motion, because it is free from the constraints.

These are the detailed formal steps

First we need the relation between the coordinate system

$$\vec{r}^i = \vec{r}^i(q_j) \text{ and } q_k = g_k(\vec{r}^i). \quad i = 1, 2, \dots, N \text{ and } k = 1, 2, \dots, n$$

In addition we have  $M$  holonomic constraint  $f_\alpha(\vec{r}^i) = 0$  for  $\alpha = 1, 2, \dots, M$

# Newton's Equation of motion

In the last class we see that, the **generalised coordinate systems** are best suited for applying laws of motion, because it is free from the constraints.

These are the detailed formal steps

First we need the relation between the coordinate system

$$\vec{r}^i = \vec{r}^i(q_j) \text{ and } q_k = g_k(\vec{r}^i). \quad i = 1, 2, \dots, N \text{ and } k = 1, 2, \dots, n$$

$$\text{In addition we have } M \text{ holonomic constraint } f_\alpha(\vec{r}^i) = 0 \text{ for } \alpha = 1, 2, \dots, M$$

We also need relation between the basis vectors  $\{\hat{e}_i\} \leftrightarrow \{\hat{q}_j\}$

# Newton's Equation of motion

In the last class we see that, the **generalised coordinate systems** are best suited for applying laws of motion, because it is free from the constraints.

These are the detailed formal steps

First we need the relation between the coordinate system

$$\vec{r}^i = \vec{r}^i(q_j) \text{ and } q_k = g_k(\vec{r}^i). \quad i = 1, 2, \dots, N \text{ and } k = 1, 2, \dots, n$$

$$\text{In addition we have } M \text{ holonomic constraint } f_\alpha(\vec{r}^i) = 0 \text{ for } \alpha = 1, 2, \dots, M$$

We also need relation between the basis vectors  $\{\hat{e}_i\} \leftrightarrow \{\hat{q}_j\}$

Using this, we can relate the components of forces and momentum in the generalised coordinate

# Newton's Equation of motion

In the last class we see that, the **generalised coordinate systems** are best suited for applying laws of motion, because it is free from the constraints.

These are the detailed formal steps

First we need the relation between the coordinate system

$$\vec{r}^i = \vec{r}^i(q_j) \text{ and } q_k = g_k(\vec{r}^i). \quad i = 1, 2, \dots, N \text{ and } k = 1, 2, \dots, n$$

$$\text{In addition we have } M \text{ holonomic constraint } f_\alpha(\vec{r}^i) = 0 \text{ for } \alpha = 1, 2, \dots, M$$

We also need relation between the basis vectors  $\{\hat{e}_i\} \leftrightarrow \{\hat{q}_j\}$

Using this, we can relate the components of forces and momentum in the generalised coordinate

$$\vec{F} = \sum_{i=1}^N F_i \vec{e}_i = \sum_{kj=1}^n Q_k \hat{q}_k$$

# Newton's Equation of motion

In the last class we see that, the **generalised coordinate systems** are best suited for applying laws of motion, because it is free from the constraints.

These are the detailed formal steps

First we need the relation between the coordinate system

$$\vec{r}^i = \vec{r}^i(q_j) \text{ and } q_k = g_k(\vec{r}^i). \quad i = 1, 2, \dots, N \text{ and } k = 1, 2, \dots, n$$

In addition we have  $M$  holonomic constraint  $f_\alpha(\vec{r}^i) = 0$  for  $\alpha = 1, 2, \dots, M$

We also need relation between the basis vectors  $\{\hat{e}_i\} \leftrightarrow \{\hat{q}_j\}$

Using this, we can relate the components of forces and momentum in the generalised coordinate

$$\vec{F} = \sum_{i=1}^N F_i \vec{e}_i = \sum_{kj=1}^n Q_k \hat{q}_k$$

$F_i$  are components of forces

$Q_k$  are components of forces in generalised coordinate

# Newton's Equation of motion

We also need relation between the basis vectors  $\{\hat{e}_i\} \leftrightarrow \{\hat{q}_j\}$

Using this, we can relate the components of forces and momentum in the generalised coordinate

$$\vec{F} = \sum_{i=1}^N F_i \vec{e}_i = \sum_{kj=1}^n Q_k \hat{q}_k$$

$F_i$  are components of forces  
 $Q_k$  are components of forces in generalised coordinate

# Newton's Equation of motion

We also need relation between the basis vectors  $\{\hat{e}_i\} \leftrightarrow \{\hat{q}_j\}$

Using this, we can relate the components of forces and momentum in the generalised coordinate

$$\vec{F} = \sum_{i=1}^N F_i \vec{e}_i = \sum_{kj=1}^n Q_k \hat{q}_k$$

$F_i$  are components of forces  
 $Q_k$  are components of forces in generalised coordinate

This step is not difficult, Next we need to transform momentum, and differentiate it substitute in NLOM.

# Newton's Equation of motion

We also need relation between the basis vectors  $\{\hat{e}_i\} \leftrightarrow \{\hat{q}_j\}$

Using this, we can relate the components of forces and momentum in the generalised coordinate

$$\vec{F} = \sum_{i=1}^N F_i \vec{e}_i = \sum_{kj=1}^n Q_k \hat{q}_k$$

$F_i$  are components of forces  
 $Q_k$  are components of forces in generalised coordinate

This step is not difficult, Next we need to transform momentum, and differentiate it substitute in NLOM.

$$\vec{p} = \sum_{kj=1}^n p_k \hat{q}_k \text{ for momentum, we need } \frac{d\vec{p}}{dt}$$

# Newton's Equation of motion

We also need relation between the basis vectors  $\{\hat{e}_i\} \leftrightarrow \{\hat{q}_j\}$

Using this, we can relate the components of forces and momentum in the generalised coordinate

$$\vec{F} = \sum_{i=1}^N F_i \vec{e}_i = \sum_{kj=1}^n Q_k \hat{q}_k$$

$F_i$  are components of forces  
 $Q_k$  are components of forces in generalised coordinate

This step is not difficult, Next we need to transform momentum, and differentiate it substitute in NLOM.

$$\vec{p} = \sum_{kj=1}^n p_k \hat{q}_k \text{ for momentum, we need } \frac{d\vec{p}}{dt}$$

if the basis vectors  $\{\hat{q}_j\}$  are not constant, i.e.  $\frac{d\hat{q}_j}{dt} \neq 0$ , we have seen this results in the many terms and often difficult to deal with.

# Work done

# Work done

We have seen earlier, the work done  $W = \vec{F} \cdot \vec{dr}$  and energy can carry information about the NLOM

# Work done

We have seen earlier, the work done  $W = \vec{F} \cdot \vec{dr}$  and energy can carry information about the NLOM

Work done is a scalar, so it can be easily transformed from one coordinate system to another while retaining information about the EOM, which can then be reconstructed in the new (generalized) coordinate system. We follow this approach.

# Work done

We have seen earlier, the work done  $W = \vec{F} \cdot \vec{dr}$  and energy can carry information about the NLOM

Work done is a scalar, so it can be easily transformed from one coordinate system to another while retaining information about the EOM, which can then be reconstructed in the new (generalized) coordinate system. We follow this approach.

When a force  $\vec{F}$  acting on a system with momentum  $\vec{p}$ , we have

# Work done

We have seen earlier, the work done  $W = \vec{F} \cdot \vec{dr}$  and energy can carry information about the NLOM

Work done is a scalar, so it can be easily transformed from one coordinate system to another while retaining information about the EOM, which can then be reconstructed in the new (generalized) coordinate system. We follow this approach.

When a force  $\vec{F}$  acting on a system with momentum  $\vec{p}$ , we have

$$\vec{F} = \frac{d\vec{p}}{dt}$$

External force

Reaction offered by the physical system against force.

# Work done

We have seen earlier, the work done  $W = \vec{F} \cdot \vec{dr}$  and energy can carry information about the NLOM

Work done is a scalar, so it can be easily transformed from one coordinate system to another while retaining information about the EOM, which can then be reconstructed in the new (generalized) coordinate system. We follow this approach.

When a force  $\vec{F}$  acting on a system with momenti

$$\vec{F} = \frac{d\vec{p}}{dt}$$

External force

Reaction offered by the physical system against force.



# Work done

We have seen earlier, the work done  $W = \vec{F} \cdot \vec{dr}$  and energy can carry information about the NLOM

Work done is a scalar, so it can be easily transformed from one coordinate system to another while retaining information about the EOM, which can then be reconstructed in the new (generalized) coordinate system. We follow this approach.

When a force  $\vec{F}$  acting on a system with momenti

$$\vec{F} = \frac{d\vec{p}}{dt}$$

External force

Reaction offered by the physical system against force.



As per Newton, these should balance—that is, the third law of motion. The total external force and the reaction/restoring force offered by the system are in equilibrium.

# Principle of Virtual Work

# Principle of Virtual Work

Let's start on an inertial frame

$$\vec{F} = \frac{d\vec{p}}{dt} \quad \text{let } \vec{dr} \text{ be a small displacement.}$$

# Principle of Virtual Work

Let's start on an inertial frame       $\vec{F} = \frac{d\vec{p}}{dt}$       let  $\vec{dr}$  be a small displacement.

taking dot product on both side we, get,       $\vec{F}_i \cdot \vec{dr}_i = \frac{d\vec{p}_i}{dt} \cdot \vec{dr}_i$       for  $N$  particles

# Principle of Virtual Work

Let's start on an inertial frame

$$\vec{F} = \frac{d\vec{p}}{dt} \quad \text{let } \vec{dr} \text{ be a small displacement.}$$

taking dot product on both side we, get,  $\vec{F}_i \cdot \vec{dr}_i = \frac{d\vec{p}_i}{dt} \cdot \vec{dr}_i$  for  $N$  particles

System is under dynamical equilibrium, between the applied force and reaction force offered by the system. Combined with the work done, we get the condition

# Principle of Virtual Work

Let's start on an inertial frame

$$\vec{F} = \frac{d\vec{p}}{dt} \quad \text{let } \vec{dr} \text{ be a small displacement.}$$

taking dot product on both side we, get,  $\vec{F}_i \cdot \vec{dr}_i = \frac{d\vec{p}_i}{dt} \cdot \vec{dr}_i$  for  $N$  particles

System is under dynamical equilibrium, between the applied force and reaction force offered by the system. Combined with the work done, we get the condition

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \vec{dr}_i = 0$$

# Principle of Virtual Work

Let's start on an inertial frame

$$\vec{F} = \frac{d\vec{p}}{dt} \quad \text{let } \vec{dr} \text{ be a small displacement.}$$

taking dot product on both side we, get,  $\vec{F}_i \cdot \vec{dr}_i = \frac{d\vec{p}_i}{dt} \cdot \vec{dr}_i$  for  $N$  particles

System is under dynamical equilibrium, between the applied force and reaction force offered by the system. Combined with the work done, we get the condition

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \vec{dr}_i = 0$$

**D'Alembert's Principle:** D'Alembert realised that, it is not need to have real displacement , it sufficient to use virtual displacement and virtual work results in the principle of virtual work

# D'Alembert's Principle

# D'Alembert's Principle

For a dynamical system subject to constraints, the difference between the applied forces and the inertial forces on each particle, projected onto any virtual displacement consistent with the constraints, is zero:

# D'Alembert's Principle

For a dynamical system subject to constraints, the difference between the applied forces and the inertial forces on each particle, projected onto any virtual displacement consistent with the constraints, is zero:

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

# D'Alembert's Principle

For a dynamical system subject to constraints, the difference between the applied forces and the inertial forces on each particle, projected onto any virtual displacement consistent with the constraints, is zero:

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

where  $\vec{F}_i$  is the total applied force on the  $i$ -th particle,  $\vec{p}_i$  its momentum , and  $\delta \vec{r}_i$  is any virtual displacement allowed by the constraints.

# D'Alembert's Principle

For a dynamical system subject to constraints, the difference between the applied forces and the inertial forces on each particle, projected onto any virtual displacement consistent with the constraints, is zero:

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

where  $\vec{F}_i$  is the total applied force on the  $i$ -th particle,  $\vec{p}_i$  its momentum , and  $\delta \vec{r}_i$  is any virtual displacement allowed by the constraints.

We use this principle for transforming EOM from inertial frame to generalised coordinate

# D'Alembert's Principle

For a dynamical system subject to constraints, the difference between the applied forces and the inertial forces on each particle, projected onto any virtual displacement consistent with the constraints, is zero:

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

where  $\vec{F}_i$  is the total applied force on the  $i$ -th particle,  $\vec{p}_i$  its momentum , and  $\delta \vec{r}_i$  is any virtual displacement allowed by the constraints.

We use this principle for transforming EOM from inertial frame to generalised coordinate

$$\left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \vec{\delta r}_i = 0 \text{ and}$$

# D'Alembert's Principle

For a dynamical system subject to constraints, the difference between the applied forces and the inertial forces on each particle, projected onto any virtual displacement consistent with the constraints, is zero:

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

where  $\vec{F}_i$  is the total applied force on the  $i$ -th particle,  $\vec{p}_i$  its momentum , and  $\delta \vec{r}_i$  is any virtual displacement allowed by the constraints.

We use this principle for transforming EOM from inertial frame to generalised coordinate

$$\left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \vec{\delta r}_i = 0 \text{ and}$$

$$\left( \vec{Q} - \frac{d\vec{p}}{dt} \right) \cdot \vec{\delta q} = 0$$

# D'Alembert's Principle

For a dynamical system subject to constraints, the difference between the applied forces and the inertial forces on each particle, projected onto any virtual displacement consistent with the constraints, is zero:

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

where  $\vec{F}_i$  is the total applied force on the  $i$ -th particle,  $\vec{p}_i$  its momentum , and  $\delta \vec{r}_i$  is any virtual displacement allowed by the constraints.

We use this principle for transforming EOM from inertial frame to generalised coordinate

$$\left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \vec{\delta r}_i = 0 \text{ and}$$

$$\left( \vec{Q} - \frac{d\vec{p}}{dt} \right) \cdot \vec{\delta q} = 0$$

Here, Coefficient of  $\delta q$   
represent the EOM

# Why Virtual Work ?

# Why Virtual Work?

Cartesian Coordinate



Generalised Coordinate

# Why Virtual Work?

Cartesian Coordinate

The relation with generalised coordinates

$$\vec{r}^i = \vec{r}^i(q_1, q_2, \dots, q_n, t)$$



Generalised Coordinate

# Why Virtual Work?

Cartesian Coordinate

The relation with generalised coordinates

$$\vec{r}^i = \vec{r}^i(q_1, q_2, \dots, q_n, t)$$



Generalised Coordinate

The relation with Cartesian coordinates

$$q_k = g_k(\vec{r}^1, \vec{r}^2, \dots, \vec{r}^N, t)$$

# Why Virtual Work?

Cartesian Coordinate

The relation with generalised coordinates

$$\vec{r}^i = \vec{r}^i(q_1, q_2, \dots, q_n, t)$$

holonomic constraint  $f_\alpha(\vec{r}^i, t) = 0$

$$\alpha = 1, 2, \dots, M$$



Generalised Coordinate

The relation with Cartesian coordinates

$$q_k = g_k(\vec{r}^1, \vec{r}^2, \dots, \vec{r}^N, t)$$

No constraints!

# Why Virtual Work?

Cartesian Coordinate

The relation with generalised coordinates

$$\vec{r}^i = \vec{r}^i(q_1, q_2, \dots, q_n, t)$$

holonomic constraint  $f_\alpha(\vec{r}^i, t) = 0$   
 $\alpha = 1, 2, \dots, M$

work done on total system  
while displacement  $\vec{dr}_i$

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

Generalised Coordinate

The relation with Cartesian coordinates

$$q_k = g_k(\vec{r}^1, \vec{r}^2, \dots, \vec{r}^N, t)$$

No constraints!

# Why Virtual Work?

Cartesian Coordinate

The relation with generalised coordinates

$$\vec{r}^i = \vec{r}^i(q_1, q_2, \dots, q_n, t)$$

holonomic constraint  $f_\alpha(\vec{r}^i, t) = 0$   
 $\alpha = 1, 2, \dots, M$

work done on total system  
while displacement  $\vec{dr}_i$

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

Generalised Coordinate

The relation with Cartesian coordinates

$$q_k = g_k(\vec{r}^1, \vec{r}^2, \dots, \vec{r}^N, t)$$

No constraints!

work done on total system in the  
generalised coordinate system can be  
expressed in displacement  $\vec{dq}_k$  as

$$W = \sum_{k=1}^n Q_k dq_k$$

# Why Virtual Work?

Cartesian Coordinate

The relation with generalised coordinates

$$\vec{r}^i = \vec{r}^i(q_1, q_2, \dots, q_n, t)$$

holonomic constraint  $f_\alpha(\vec{r}^i, t) = 0$   
 $\alpha = 1, 2, \dots, M$

work done on total system  
while displacement  $\vec{dr}_i$

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

Generalised Coordinate

The relation with Cartesian coordinates

$$q_k = g_k(\vec{r}^1, \vec{r}^2, \dots, \vec{r}^N, t)$$

No constraints!

work done on total system in the  
generalised coordinate system can be  
expressed in displacement  $\vec{dq}_k$  as

$$W = \sum_{k=1}^n Q_k dq_k$$

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

# Why Virtual Work?

Cartesian Coordinate

The relation with generalised coordinates

$$\vec{r}^i = \vec{r}^i(q_1, q_2, \dots, q_n, t)$$

holonomic constraint  $f_\alpha(\vec{r}^i, t) = 0$   
 $\alpha = 1, 2, \dots, M$

work done on total system  
while displacement  $\vec{dr}_i$

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

Generalised Coordinate

The relation with Cartesian coordinates

$$q_k = g_k(\vec{r}^1, \vec{r}^2, \dots, \vec{r}^N, t)$$

No constraints!

work done on total system in the  
generalised coordinate system can be  
expressed in displacement  $\vec{dq}_k$  as

$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$


work done on total system while displacement  $\vec{dr}_i$

work done on total system while displacement  $\vec{dq}_k$

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$



$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

work done on total system while displacement  $\vec{dr}_i$

work done on total system while displacement  $\vec{dq}_k$

# Why Virtual Work?

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

And the virtual displacement  $\vec{\delta r}_i$  is given by

$$\vec{\delta r}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} \delta q_k$$

$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

work done on total system while displacement  $\vec{dr}_i$

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

And the virtual displacement  $\vec{\delta r}_i$  is given by

$$\vec{\delta r}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} \delta q_k$$

work done on total system while displacement  $\vec{dq}_k$

$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

And the virtual displacement  $\delta q_i$  is given by

$$\delta q_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{\delta r}_i$$

work done on total system while displacement  $\vec{dr}_i$

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

And the virtual displacement  $\vec{\delta r}_i$  is given by

$$\vec{\delta r}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} \delta q_k$$

work done on total system while displacement  $\vec{dq}_k$

$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

And the virtual displacement  $\delta q_i$  is given by

$$\delta q_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{\delta r}_i$$

Time is frozen hence no differentiation with t

# Why Virtual Work?

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$



$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

# Why Virtual Work?

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

From this we get

$$\vec{F} \cdot \vec{dr}_i = \vec{F} \cdot \left\{ \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt \right\}$$



$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

# Why Virtual Work?

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

From this we get

$$\vec{F} \cdot \vec{dr}_i = \vec{F} \cdot \left\{ \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt \right\}$$



$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

$$Q_k dq_k = Q_k \left\{ \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt \right\}$$

# Why Virtual Work?

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

From this we get

$$\vec{F} \cdot \vec{dr}_i = \vec{F} \cdot \left\{ \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt \right\}$$



$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

$$Q_k dq_k = Q_k \left\{ \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt \right\}$$

there is now. real way to compare the last term and hence may be left all together

# Why Virtual Work?

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

From this we get

$$\vec{F} \cdot \vec{dr}_i = \vec{F} \cdot \left\{ \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt \right\}$$

$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

$$Q_k dq_k = Q_k \left\{ \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt \right\}$$

there is now. real way to compare the last term and hence may be left all together

?

# Why Virtual Work?

$$W = \sum_{i=1}^N \vec{F}_i \cdot \vec{dr}_i = \sum_{i=1}^{3N} F_i dx_i$$

$\vec{dr}_i$  in terms of  $dq_k$  as follows

$$\vec{dr}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt$$

From this we get

$$\vec{F} \cdot \vec{dr}_i = \vec{F} \cdot \left\{ \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} dq_k + \frac{\partial r^i}{\partial t} dt \right\}$$

$$W = \sum_{k=1}^n Q_k dq_k$$

$\vec{dq}_k$  in terms of  $dq_i$  as follows

$$dq_k = \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt$$

$$Q_k dq_k = Q_k \left\{ \sum_{i=1}^N \frac{\partial q^k}{\partial r_i} \vec{dr}_i + \frac{\partial q^k}{\partial t} dt \right\}$$

there is now. real way to compare the last term and hence may be left all together

Virtual work give's more freedom to work with only one of the virtual displacement is real

# Lagrange Equation of Motion

# Lagrange Equation of Motion

We start with

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \delta\vec{r}_i = 0$$

Ref: Goldstein, Herbert; Poole, Charles; Safko, John; published by Pearson Education, Inc.,

# Lagrange Equation of Motion

We start with

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \delta\vec{r}_i = 0$$

We have already seen,

$$\delta\vec{r}_i = \sum_{k=1}^n \frac{\partial r^i}{\partial q_k} \delta q_k$$

Ref: Goldstein, Herbert; Poole, Charles; Safko, John; published by Pearson Education, Inc.,

# Lagrange Equation of Motion

We start with

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

We have already seen,

$$\delta \vec{r}_i = \sum_{k=1}^n \frac{\partial \vec{r}^i}{\partial q_k} \delta q_k$$

$$\sum_i \vec{F}_i \cdot \delta \vec{r}_i = \sum_{i,j} \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left( \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) \delta q_j$$

Ref: Goldstein, Herbert; Poole, Charles; Safko, John; published by Pearson Education, Inc.,

# Lagrange Equation of Motion

We start with

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

We have already seen,

$$\delta \vec{r}_i = \sum_{k=1}^n \frac{\partial \vec{r}_i}{\partial q_k} \delta q_k$$

$$\sum_i \vec{F}_i \cdot \delta \vec{r}_i = \sum_{i,j} \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left( \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) \delta q_j$$

$$\boxed{\sum_i \vec{F}_i \cdot \delta \vec{r}_i = \sum_j Q_j \delta q_j}$$

Ref: Goldstein, Herbert; Poole, Charles; Safko, John; published by Pearson Education, Inc.,

# Lagrange Equation of Motion

We start with

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

We have already seen,

$$\delta \vec{r}_i = \sum_{k=1}^n \frac{\partial \vec{r}_i}{\partial q_k} \delta q_k$$

$$\sum_i \vec{F}_i \cdot \delta \vec{r}_i = \sum_{i,j} \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left( \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) \delta q_j$$

$$\boxed{\sum_i \vec{F}_i \cdot \delta \vec{r}_i = \sum_j Q_j \delta q_j}$$

Where  $Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j}$  is called component of generalised force.

Ref: Goldstein, Herbert; Poole, Charles; Safko, John; published by Pearson Education, Inc.,

# Lagrange Equation of Motion

We start with

$$\sum_{i=1}^N \left( \vec{F}_i - \frac{d\vec{p}_i}{dt} \right) \cdot \delta \vec{r}_i = 0$$

We have already seen,

$$\delta \vec{r}_i = \sum_{k=1}^n \frac{\partial \vec{r}^i}{\partial q_k} \delta q_k$$

$$\sum_i \vec{F}_i \cdot \delta \vec{r}_i = \sum_{i,j} \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left( \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) \delta q_j$$

$$\boxed{\sum_i \vec{F}_i \cdot \delta \vec{r}_i = \sum_j Q_j \delta q_j}$$

Where  $Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j}$  is called component of generalised force.

It is just transformation of components of vector under the coordinate transformation

Ref: Goldstein, Herbert; Poole, Charles; Safko, John; published by Pearson Education, Inc.,

# Lagrange Equation of Motion

# Lagrange Equation of Motion

The remaining term is

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i=1} m_i \ddot{\vec{r}}_i \cdot \delta\vec{r}_i$$

# Lagrange Equation of Motion

The remaining term is

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i=1} m_i \ddot{\vec{r}}_i \cdot \delta\vec{r}_i$$

For a particle  $\vec{p}_i = m_i \dot{\vec{r}}_i$

# Lagrange Equation of Motion

The remaining term is

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i=1} m_i \ddot{\vec{r}}_i \cdot \delta\vec{r}_i$$

For a particle  $\vec{p}_i = m_i \dot{\vec{r}}_i$

Substituting for  $\vec{\delta r}_i = \sum_{k=1}^n \frac{\partial \vec{r}^i}{\partial q_k} \delta q_k$  we get,

# Lagrange Equation of Motion

The remaining term is

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i=1} m_i \ddot{\vec{r}}_i \cdot \delta\vec{r}_i$$

For a particle  $\vec{p}_i = m_i \dot{\vec{r}}_i$

Substituting for  $\vec{\delta r}_i = \sum_{k=1}^n \frac{\partial \vec{r}^i}{\partial q_k} \delta q_k$  we get,

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i,j} m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left[ \sum_i m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] \delta q_j$$

# Lagrange Equation of Motion

The remaining term is

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i=1} m_i \ddot{\vec{r}}_i \cdot \delta\vec{r}_i$$

For a particle  $\vec{p}_i = m_i \dot{\vec{r}}_i$

Substituting for  $\vec{\delta r}_i = \sum_{k=1}^n \frac{\partial \vec{r}^i}{\partial q_k} \delta q_k$  we get,

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i,j} m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left[ \sum_i m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] \delta q_j$$

The rest is a series of algebraic steps to bring the equation into a simpler form.

# Lagrange Equation of Motion

The remaining term is

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i=1} m_i \ddot{\vec{r}}_i \cdot \delta\vec{r}_i$$

For a particle  $\vec{p}_i = m_i \dot{\vec{r}}_i$

Substituting for  $\vec{\delta r}_i = \sum_{k=1}^n \frac{\partial \vec{r}^i}{\partial q_k} \delta q_k$  we get,

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i,j} m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left[ \sum_i m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] \delta q_j$$

*Let's check this term*

The rest is a series of algebraic steps to bring the equation into a simpler form.

# Lagrange Equation of Motion

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i,j} m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] \delta q_j$$

The rest is a series of algebraic steps to bring the equation into a simpler form.

# Lagrange Equation of Motion

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i,j} m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] \delta q_j$$

The rest is a series of algebraic steps to bring the equation into a simpler form.

Let's start with the coefficients of  $\delta q_j$

# Lagrange Equation of Motion

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i,j} m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] \delta q_j$$

The rest is a series of algebraic steps to bring the equation into a simpler form.

Let's start with the coefficients of  $\delta q_j$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \left[ \sum_i m_i \frac{d\dot{\vec{r}}_i}{dt} \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right]$$

# Lagrange Equation of Motion

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i,j} m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] \delta q_j$$

The rest is a series of algebraic steps to bring the equation into a simpler form.

Let's start with the coefficients of  $\delta q_j$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \left[ \sum_i m_i \frac{d\dot{\vec{r}}_i}{dt} \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right]$$

expand  $\frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right)$  and adjust  
the terms we should get

# Lagrange Equation of Motion

$$\sum_{i=1} \frac{d\vec{p}_i}{dt} \cdot \delta\vec{r}_i = \sum_{i,j} m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \delta q_j = \sum_j \left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] \delta q_j$$

The rest is a series of algebraic steps to bring the equation into a simpler form.

Let's start with the coefficients of  $\delta q_j$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \left[ \sum_i m_i \frac{d\dot{\vec{r}}_i}{dt} \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right]$$

expand  $\frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right)$  and adjust  
the terms we should get

$$= \sum_i \left[ \frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i \dot{\vec{r}}_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

$$\left[\sum_im_i\ddot{\vec{r}}_i\cdot\frac{\partial\vec{r}_i}{\partial q_j}\right]=\sum_i\left[\frac{d}{dt}\left(m_i\dot{\vec{r}}_i\cdot\frac{\partial\vec{r}_i}{\partial q_j}\right)-m_i\dot{\vec{r}}_i\cdot\frac{d}{dt}\left(\frac{\partial\vec{r}_i}{\partial q_j}\right)\right]$$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i \dot{\vec{r}}_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

with  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$ , we get,

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i \dot{\vec{r}}_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$


with  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$ , we get,

$$= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i \dot{\vec{r}}_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

with  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$ , we get,

$$= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$



2

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i \dot{\vec{r}}_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

with  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$ , we get,

$$= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

Expanding second term, 2, we get



$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i \dot{\vec{r}}_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

with  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$ , we get,

$$= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

Expanding second term,  , we get

$$= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \left( \sum_k \frac{\partial^2 \vec{r}_i}{\partial q_j \partial q_k} \dot{q}_k + \frac{\partial^2 \vec{r}_i}{\partial q_j \partial t} \right) \right]$$



$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i \dot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i \dot{\vec{r}}_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

with  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$ , we get,

$$= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \cdot \frac{d}{dt} \left( \frac{\partial \vec{r}_i}{\partial q_j} \right) \right]$$

Expanding second term, 2, we get

$$= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \left( \sum_k \frac{\partial^2 \vec{r}_i}{\partial q_j \partial q_k} \dot{q}_k + \frac{\partial^2 \vec{r}_i}{\partial q_j \partial t} \right) \right]$$

Note this is  
 $\dot{r}_i = v_i$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial}{\partial q_j} \left( \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial^2 \vec{r}_i}{\partial t} \right) \right]$$



We need more simplification, let's look at  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$  under transformation to new coordinate

We need more simplification, let's look at  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$  under transformation to new coordinate

$$v_i = \frac{d\vec{r}_i}{dt} = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t}$$

We need more simplification, let's look at  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$  under transformation to new coordinate

$$v_i = \frac{d\vec{r}_i}{dt} = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t}$$

$$\frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial}{\partial \dot{q}_j} \left[ \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t} \right]$$

Do not deprnd  
on  $\dot{q}_j$

Do not deprnd  
on  $\dot{q}_j$

We need more simplification, let's look at  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$  under transformation to new coordinate

$$v_i = \frac{d\vec{r}_i}{dt} = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t}$$

$$\frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial}{\partial \dot{q}_j} \left[ \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t} \right] = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \delta_{jk}$$

Do not deprnd  
on  $\dot{q}_j$

Do not deprnd  
on  $\dot{q}_j$

We need more simplification, let's look at  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$  under transformation to new coordinate

$$v_i = \frac{d\vec{r}_i}{dt} = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t}$$

$$\frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial}{\partial \dot{q}_j} \left[ \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t} \right] = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \delta_{jk} \implies \frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial \vec{r}_i}{\partial q_j}$$

Do not deprnd  
on  $\dot{q}_j$

Do not deprnd  
on  $\dot{q}_j$

We need more simplification, let's look at  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$  under transformation to new coordinate

$$v_i = \frac{d\vec{r}_i}{dt} = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t}$$

$$\frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial}{\partial \dot{q}_j} \left[ \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t} \right] = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \delta_{jk} \implies \frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial \vec{r}_i}{\partial q_j}$$

$$\frac{\partial \vec{v}_i}{\partial \dot{q}_j} = \frac{\partial \vec{r}_i}{\partial q_j}$$

Do not deprnd  
on  $\dot{q}_j$

Do not deprnd  
on  $\dot{q}_j$

We need more simplification, let's look at  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$  under transformation to new coordinate

$$v_i = \frac{d\vec{r}_i}{dt} = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t}$$

$$\frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial}{\partial \dot{q}_j} \left[ \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t} \right] = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \delta_{jk} \implies \frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial \vec{r}_i}{\partial q_j}$$

Do not deprnd  
on  $\dot{q}_j$ 
Do not deprnd  
on  $\dot{q}_j$

$$\frac{\partial \vec{v}_i}{\partial \dot{q}_j} = \frac{\partial \vec{r}_i}{\partial q_j}$$


Going back to original equation,

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right]$$

We need more simplification, let's look at  $v_i = \frac{d\vec{r}_i}{dt} = \dot{\vec{r}}_i$  under transformation to new coordinate


$$v_i = \frac{d\vec{r}_i}{dt} = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t}$$

$$\frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial}{\partial \dot{q}_j} \left[ \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \dot{q}_k + \frac{\partial \vec{r}_i}{\partial t} \right] = \sum_k \frac{\partial \vec{r}_i}{\partial q_k} \delta_{jk} \implies \frac{\dot{\vec{r}}_i}{\partial \dot{q}_j} = \frac{\partial \vec{r}_i}{\partial q_j}$$



$$\frac{\partial \vec{v}_i}{\partial \dot{q}_j} = \frac{\partial \vec{r}_i}{\partial q_j}$$

Going back to original equation,

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right]$$


$$\begin{aligned}
\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] &= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{v}_i}{\partial \dot{q}_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right]
\end{aligned}$$

$$\begin{aligned}
\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] &= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{v}_i}{\partial \dot{q}_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \frac{d}{dt} \left[ \frac{\partial}{\partial \dot{q}_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right) \right] - \frac{\partial}{\partial q_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right)
\end{aligned}$$

$$\begin{aligned}
\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] &= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{v}_i}{\partial \dot{q}_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \frac{d}{dt} \left[ \frac{\partial}{\partial \dot{q}_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right) \right] - \frac{\partial}{\partial q_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right)
\end{aligned}$$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j}$$

$$\begin{aligned}
\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] &= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{v}_i}{\partial \dot{q}_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \frac{d}{dt} \left[ \frac{\partial}{\partial \dot{q}_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right) \right] - \frac{\partial}{\partial q_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right)
\end{aligned}$$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j}$$

$$= \sum_{j=1} \left( \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} \right) \delta q_j$$

$$\begin{aligned}
\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] &= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{v}_i}{\partial \dot{q}_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \frac{d}{dt} \left[ \frac{\partial}{\partial \dot{q}_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right) \right] - \frac{\partial}{\partial q_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right)
\end{aligned}$$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j}$$

$$\sum_{i=1}^N \frac{\vec{p}_i}{dt} \cdot \delta \vec{r}_i = \sum_{j=1} \left( \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} \right) \delta q_j$$

$$\begin{aligned}
\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] &= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \sum_i \left[ \frac{d}{dt} \left( m_i v_i \cdot \frac{\partial \vec{v}_i}{\partial \dot{q}_j} \right) - m_i v_i \frac{\partial v_i}{\partial q_j} \right] \\
&= \frac{d}{dt} \left[ \frac{\partial}{\partial \dot{q}_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right) \right] - \frac{\partial}{\partial q_j} \left( \sum_i \frac{1}{2} m_i v_i^2 \right)
\end{aligned}$$

$$\left[ \sum_i m_i \ddot{\vec{r}}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} \right] = \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j}$$

$$\sum_{i=1}^N \frac{\vec{p}_i}{dt} \cdot \delta \vec{r}_i = \sum_{j=1} \left( \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} \right) \delta q_j$$

Finally we have **Lagrange Equation of Motion**

Finally we have **Lagrange Equation of Motion**  
Cartesian Coordinate

$$\sum_{i=1}^N \left( F_i - \frac{dp_i}{dt} \right) \cdot \delta r_i = 0$$

Finally we have **Lagrange Equation of Motion**

Cartesian Coordinate

$$\sum_{i=1}^N \left( F_i - \frac{dp_i}{dt} \right) \cdot \delta r_i = 0$$

Generalised Coordinate

$$\sum_{i=j} \left( Q_j - \frac{dp_j}{dt} \right) \cdot \delta q_j = 0$$

Finally we have **Lagrange Equation of Motion**

Cartesian Coordinate

$$\sum_{i=1}^N \left( F_i - \frac{dp_i}{dt} \right) \cdot \delta r_i = 0$$

Generalised Coordinate

$$\sum_{i=j} \left( Q_j - \frac{dp_j}{dt} \right) \cdot \delta q_j = 0$$



Finally we have **Lagrange Equation of Motion**

Cartesian Coordinate

$$\sum_{i=1}^N \left( F_i - \frac{dp_i}{dt} \right) \cdot \delta r_i = 0$$

Generalised Coordinate

$$\sum_{i=j} \left( Q_j - \frac{dp_j}{dt} \right) \cdot \delta q_j = 0$$

$$\sum_j \left\{ Q_j - \left( \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} \right) \right\} \delta q_j$$

Finally we have **Lagrange Equation of Motion**

Cartesian Coordinate

$$\sum_{i=1}^N \left( F_i - \frac{dp_i}{dt} \right) \cdot \delta r_i = 0$$

Generalised Coordinate

$$\sum_{i=j} \left( Q_j - \frac{dp_j}{dt} \right) \cdot \delta q_j = 0$$

$$\sum_j \left\{ Q_j - \left( \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} \right) \right\} \delta q_j$$

Where  $Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j}$  is called component of generalised force.

Finally we have **Lagrange Equation of Motion**

Cartesian Coordinate

$$\sum_{i=1}^N \left( F_i - \frac{dp_i}{dt} \right) \cdot \delta r_i = 0$$

Generalised Coordinate

$$\sum_{i=j} \left( Q_j - \frac{dp_j}{dt} \right) \cdot \delta q_j = 0$$

$$\sum_j \left\{ Q_j - \left( \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} \right) \right\} \delta q_j$$

Where  $Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j}$  is called component of generalised force.

The equation of motion in the new coordinate system is

Finally we have **Lagrange Equation of Motion**

Cartesian Coordinate

$$\sum_{i=1}^N \left( F_i - \frac{dp_i}{dt} \right) \cdot \delta r_i = 0$$

Generalised Coordinate

$$\sum_{i=j} \left( Q_j - \frac{dp_j}{dt} \right) \cdot \delta q_j = 0$$

$$\sum_j \left\{ Q_j - \left( \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} \right) \right\} \delta q_j$$

Where  $Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j}$  is called component of generalised force.

The equation of motion in the new coordinate system is

$$\frac{d}{dt} \left( \frac{\partial T}{\partial \dot{q}_j} \right) - \frac{\partial T}{\partial q_j} = Q_j$$

## Finally we have **Lagrange Equation of Motion**

Cartesian Coordinate

$$\sum_{i=1}^N \left( F_i - \frac{dp_i}{dt} \right) \cdot \delta r_i = 0$$

Generalised Coordinate

$$\sum_{i=j} \left( Q_j - \frac{dp_j}{dt} \right) \cdot \delta q_j = 0$$

$$\sum_j \left\{ Q_j - \left( \frac{d}{dt} \left[ \frac{\partial T}{\partial \dot{q}_j} \right] - \frac{\partial T}{\partial q_j} \right) \right\} \delta q_j$$

Where  $Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j}$  is called component of generalised force.

The equation of motion in the new coordinate system is

$$\frac{d}{dt} \left( \frac{\partial T}{\partial \dot{q}_j} \right) - \frac{\partial T}{\partial q_j} = Q_j$$

This is general form of equation of motion for a given force  $Q_j$

# Lagrange Equation of Motion

When the forces are derivable from a scalar potential function  $V$ ;  $F_i = -\nabla_i V$

# Lagrange Equation of Motion

When the forces are derivable from a scalar potential function  $V$ ;  $F_i = -\nabla_i V$

Then the generalized forces can be written as

# Lagrange Equation of Motion

When the forces are derivable from a scalar potential function  $V$ ;  $F_i = -\nabla_i V$

Then the generalized forces can be written as

$$Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} = \sum_i -\nabla_i V \cdot \frac{\partial \vec{r}_i}{\partial q_j}$$

# Lagrange Equation of Motion

When the forces are derivable from a scalar potential function  $V$ ;  $F_i = -\nabla_i V$

Then the generalized forces can be written as

$$Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} = \sum_i -\nabla_i V \cdot \frac{\partial \vec{r}_i}{\partial q_j}$$

We can also directly transform  $V(r^i)$  to  $V(q_j)$  under coordinate transformation then we have

# Lagrange Equation of Motion

When the forces are derivable from a scalar potential function  $V$ ;  $F_i = -\nabla_i V$

Then the generalized forces can be written as

$$Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} = \sum_i -\nabla_i V \cdot \frac{\partial \vec{r}_i}{\partial q_j}$$

We can also directly transform  $V(r^i)$  to  $V(q_j)$  under coordinate transformation then we have

$$Q_j = -\frac{\partial V}{\partial q_j}$$

# Lagrange Equation of Motion

When the forces are derivable from a scalar potential function  $V$ ;  $F_i = -\nabla_i V$

Then the generalized forces can be written as

$$Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} = \sum_i -\nabla_i V \cdot \frac{\partial \vec{r}_i}{\partial q_j}$$

We can also directly transform  $V(r^i)$  to  $V(q_j)$  under coordinate transformation then we have

$$Q_j = -\frac{\partial V}{\partial q_j}$$

We define  $\mathcal{L} = T - V$ , which is called the Lagrangian. Then the EOM of motion takes the form

# Lagrange Equation of Motion

When the forces are derivable from a scalar potential function  $V$ ;  $F_i = -\nabla_i V$

Then the generalized forces can be written as

$$Q_j = \sum_i \vec{F}_i \cdot \frac{\partial \vec{r}_i}{\partial q_j} = \sum_i -\nabla_i V \cdot \frac{\partial \vec{r}_i}{\partial q_j}$$

We can also directly transform  $V(r^i)$  to  $V(q_j)$  under coordinate transformation then we have

$$Q_j = -\frac{\partial V}{\partial q_j}$$

We define  $\mathcal{L} = T - V$ , which is called the Lagrangian. Then the EOM of motion takes the form

$$\boxed{\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{q}_j} \right) - \frac{\partial \mathcal{L}}{\partial q_j} = 0}$$

