# **Two arrays**

You are given two integer arrays, A and B, each containing N integers. The size of the array is less than or equal to 1000. You are free to permute the order of the elements in the arrays.

Now here's the real question: Is there an permutation A', B' possible of A and B, such that,  $A'_i + B'_i \ge K$  for all i, where  $A'_i$  denotes the i<sup>th</sup> element in the array A' and  $B'_i$  denotes i<sup>th</sup> element in the array B'.

## **Input Format**

The first line contains an integer, T, the number of test-cases. T test cases follow. Each test case has the following format:

The first line contains two integers, N and K. The second line contains N space separated integers, denoting array A. The third line describes array B in a same format.

#### **Output Format**

For each test case, if such an arrangement exists, output "YES", otherwise "NO" (without quotes).

#### **Constraints**

```
1 <= T <= 10
```

$$1 <= N <= 1000$$

$$1 <= K <= 10^9$$

$$0 <= A_i, B_i \le 10^9$$

#### **Sample Input**

```
2
3 10
2 1 3
7 8 9
4 5
1 2 2 1
3 3 3 4
```

## **Sample Output**

```
YES
NO
```

#### **Explanation**

The first input has 3 elements in Array A and Array B, we see that the one of the arrangements, 3 2 1 and 7 8 9 has each pair of elements (3+7, 2 + 8 and 9 + 1) summing upto 10 and hence the answer is "YES".

The second input has array B with three 3s. So, we need at least three numbers in A that are greater than 1. As this is not the case, the answer is "NO".