Week 4 Reading: Physical Constraints and Projection

1. Derive the sum equality from steady balances

For first-order consecutive reactions A → B → C, steady state requires r_A = r_B = r_C = 0. Solve A first, then substitute into B and C:

$$C_A = \frac{C_{\mathrm{Af}}}{1 + k_1\tau}$$

$$C_B = \frac{k_1\tau C_A}{1 + k_2\tau}, C_C = k_2\tau C_B$$

The B balance gives $k_1\tau C_A = (1 + k_2\tau)C_B$, hence $C_B + C_C = k_1\tau C_A$. Combine with $C_{Af} = (1 + k_1\tau)C_A$ from the A balance to obtain the concentration-sum equality.

Check: 330 K, τ = 5 min, and C_{Af} = 1.2 mol/L give [0.6, 0.4, 0.2].

2. The stationary point of a soft penalty

For J(z) = ½(z − a)² + ½λ(z − b)² with λ ≥ 0, dJ/dz = (z − a) + λ(z − b). The second derivative is 1 + λ > 0, so the stationary point is the unique minimizer.

$$z_* = \frac{a + \lambda b}{1 + \lambda}$$

$$z_* - b = \frac{a - b}{1 + \lambda}$$

Exercise: set a = 1.4 and b = 1.2; compute the solution and residual for λ = 1 and 9.

Answer check: solutions are 1.30 and 1.22; residuals are 0.10 and 0.02. A finite penalty can leave a residual even at its exact minimizer.

3. Equality QP and its KKT derivation

Let $B \in \mathbb{R}^{mxn}$ have rank m and let $d \in \mathbb{R}^{m}$. Hold the input and raw prediction fixed and solve:

$$y_p = \text{argmin}_z \frac{1}{2}\left\| z - \hat{y} \right\|^2 \text{subject to} Bz = d\,(x)$$

$$z - \hat{y} + B^T\lambda = 0, Bz = d$$

Stationarity gives $z = \hat{y} - B^T\lambda$. Substitution into feasibility gives $BB^T\lambda = B\hat{y} - d$. Invertibility makes λ and z unique. The equality set is convex and the objective Hessian is I.

4. Equality guarantee and projector identities

$$G = BB^T, G\lambda = B\hat{y} - d$$
$$y_p = \hat{y} - B^T G^{-1}\left(B\hat{y} - d\right)$$

Multiplying by B shows that correction cancels the original residual.

$$By_p = B\hat{y} - GG^{-1}\left(B\hat{y} - d\right) = d$$

$P = I - B^T(BB^T)^{-1}B$ satisfies $P^T = P$, $P^2 = P$, and $BP = 0$. It projects onto the null space of B. Feasible raw outputs receive zero correction, and projecting an already projected output changes nothing.

5. Hand calculation: correction and error

$$B = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}, d = 1.2$$

$$y_p = \begin{bmatrix} .70 \\ .45 \\ .25 \end{bmatrix} - \frac{.20}{3}\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} .633333 \\ .383333 \\ .183333 \end{bmatrix}$$

Against reference [0.6, 0.4, 0.2], the raw sum of squared errors is 0.015 (mol/L)². Squared correction norm is 0.04/3 and projected squared error is 1/600 ≈ 0.001667.

The A error changes from +0.10 to +0.033333. B and C change from +0.05 to −0.016667. Calculate aggregate and individual errors separately.

6. Chain rule for parameters and inputs

Let J_f,θ be the raw-output parameter Jacobian. For fixed A, B, b:

$$\nabla_\theta L = {J_{f,\theta}}^T P^T \nabla_{y_p} L$$

$$\frac{\partial y_p}{\partial x} = P\frac{\partial f_\theta}{\partial x} - B^T G^{-1} A$$

Input gradients include the direct path through d(x) = b − Ax. If B depends on inputs, P does too and additional derivative terms are required; the fixed-projector formula is insufficient.

Check: upstream [1, 0, 0] and B = [1, 1, 1] yield [2/3, −1/3, −1/3].

7. Equality and distance in standardized coordinates

$$y = \mu + Dz, BDz = d - B\mu$$
$$D = \mathrm{diag}\left(s_1, \ldots, s_n\right)$$

Euclidean projection in z minimizes $\|D^{-1}(y - \hat{y})\|^2$ in physical coordinates. Its metric differs from physical Euclidean projection.

Example: raw [0.70, 0.45, 0.25], total 1.20, and s = [0.2, 0.5, 0.1] give restored standardized-projection output [0.673333, 0.283333, 0.243333]. Corrections are proportional to s_j^2.

Both methods satisfy the sum. For outputs with different units, even a physical Euclidean distance depends on unit choices; design the metric explicitly.

8. Redundant equalities and inconsistent right sides

$$B = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix}$$

The second row is twice the first, so rank is 1. With d = [1.2, 2.4], both equations define the same set; retaining one independent row is sufficient.

With d = [1.2, 2.5], twice the first equation contradicts the second. No feasible output exists. A pseudoinverse can return a least-squares result, but cannot make every residual zero.

When discarding small singular values, recheck residuals of removed equations in their original units.

9. Combining residuals with different units

The sum residual has units mol/L; species-balance residuals have units mol/(L·min). Adding their squares without scales makes the meaning of λ depend on the unit convention.

For example, choose reference concentration c_* and time t_*; divide the sum residual by c_* and species residuals by c_*/t_*. Still specify loss averaging and λ after nondimensionalization.

The lab penalizes only the sum. Adding known kinetics as residuals is possible, but this example already has an analytic solution, so explain the educational purpose of learning a surrogate.

10. Accuracy conditions and incorrect physics

$$\left\|\hat{y} - y^*\right\|^2 = \left\|\hat{y} - y_p\right\|^2 + \left\|y_p - y^*\right\|^2$$

This decomposition requires y^* in the same affine set. The correction is normal to the set while y_p − y^* is tangent, so their inner product vanishes.

If the true total is 1.2 but an incorrect equality enforces 1.1, the true output lies outside the set. The minimum Euclidean distance from truth is already 0.1/√3 ≈ 0.057735 mol/L.

Check sensor bias, omitted species, and stoichiometric errors first. Projection enforces an incorrectly specified equality too.

11. Requiring nonnegativity and the sum together

Clamping negative values after equality projection changes the sum. Requiring both the sum and nonnegativity gives:

$$\min_z \frac{1}{2}\left\| z - \hat{y} \right\|^2 \text{s.t.} 1^T z = C_{\text{Af}}, z \geq 0$$

For C_Af ≥ 0 the feasible set is a simplex. The solution is affine within a fixed active-set region; derivatives need care where a component reaches zero.

The required lab stops at linear equalities. Report results and gradient checks separately if implementing this inequality extension.

12. Derive the dynamic total-concentration transition

Adding dynamic species balances gives dS/dt = (C_Af − S)/τ. With constant C_Af and τ within an interval, the derivative of S − C_Af is −(S − C_Af)/τ, hence:

$$S_{k+1} = C_{\mathrm{Af}} + \left(S_k - C_{\mathrm{Af}}\right) e^{-\frac{\Delta t}{\tau}}$$

Exercise: use S_k = 1.0, C_Af = 1.2 mol/L, τ = 5 min, and Δt = 0.2 min. Compute the next total.

Answer check: 1.2 − 0.2 exp(−0.04) ≈ 1.007842 mol/L. Forcing the steady-state total immediately would change this transition. Project the raw next state onto this right side instead.

13. General-B implementation and gradient check

raw has shape (batch, n), rhs (batch, m), and B (m, n). B is fixed and must have independent rows.

```
def linear_project(raw, rhs, B):
    residual = raw @ B.T - rhs
    lam = torch.linalg.solve(B @ B.T, residual.T).T
    return raw - lam @ B

# Use float64 and independent inputs for finite differences.
assert torch.autograd.gradcheck(
    lambda a, d: linear_project(a, d, B), (raw, rhs)
)
```

Create checked raw and rhs tensors with requires_grad=True. The lab’s –check-only checks feasibility, idempotence, Jacobians, scaling, and counterexamples.

14. Define the comparison metrics

Component-j RMSE is the square root of mean squared prediction error over test rows. Pooled RMSE averages over rows and components. Sum MAE and maximum use the absolute sum residual.

Count equality violations when the absolute sum residual exceeds 10^{-6} mol/L and negative entries when a predicted concentration is below -10^{-8} mol/L. Also report the minimum concentration to expose the values behind the tolerance.

Species-balance RMS averages r_j^2 over all three species and test rows, then takes the square root; units are mol/(L·min). Do not select settings using test. Assess variation across seeds in a separate repeated experiment.

15. Exercises with short answer checks

1. How many output directions remain trainable for $B = I_3$? Answer: $P = 0$ fixes the output to d; there are no free directions.
2. How many remain for $B = [1, 1, 1]$? Answer: null-space dimension is $3 - 1 = 2$.
3. What happens to a raw output whose sum is already 1.2? Answer: zero residual means no change.
4. Does zero test sum-violation count make the hard model fully physically feasible? Answer: inspect negative concentrations and species balances separately.
5. Does post-projection reduce every component error? Answer: no. Nonincrease applies to total squared error in the same metric with a feasible reference.

16. Primary-source reading guide

Chen, H., Constante Flores, G. E., & Li, C. (2024). *Physics-informed neural networks with hard linear equality constraints*. C&CE 189, 108764. DOI · Manuscript · Author repository. Compare Section 3's fixed projection and assumptions; read one Section 4 process case.

Raissi, M., Perdikaris, P., & Karniadakis, G. E. (2019). *Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations*. JCP 378, 686-707. DOI · 2017 Part I. Read Part I's continuous-time residual and coordinate differentiation.

17. Sources for mathematics and implementation

Boyd, S., & Vandenberghe, L. (2004). *Convex Optimization*. Official book. Use Section 5.5.3 on KKT conditions and Section 10.1 on equality-constrained minimization to rederive companion pages 3-4.

PyTorch: torch.linalg.solve solves a linear system without explicitly forming an inverse. gradcheck compares automatic differentiation with finite differences; default tolerances assume double precision.

Return to the lecture. After running the lab, distinguish mathematical guarantees from observed numerical results.