

Entry Combo

Samuel de Araújo Brandão

October 5, 2025

This document collects my solutions to the OTIS problem sets from the **Entry Combo** unit, written during my preparation for mathematical olympiads.

The solutions reflect my understanding and problem-solving approach at the time of writing. Some arguments were informed by discussions, official notes, or published sources; when so, attribution is provided (see [section 3](#)).

If you find errors or have suggestions, please contact me at samuelbaraujo19@gmail.com.

Contents

1	Practice Problems	2
2	Solutions	4
2.1	Lecture Notes	4
2.1.1	USAJMO 2019 P1	4
2.1.2	Shortlist 2012 C1	5
2.1.3	USAMO 2015 P4	6
2.2	Mandatory	7
2.2.1	USAMO 2011 P2	7
2.2.2	Canadian MO 2019 P3	8
2.2.3	USAJMO 2021 P4	9
2.2.4	USAMO 2008 P4	10
2.3	Not mandatory	11
2.3.1	PAGMO 2021 P5	11
2.3.2	XOOK 2025 P1	12
2.3.3	Shortlist 2018 N1	13
2.3.4	IMO 1974 P4	14
2.3.5	EGMO 2018 P4	15
3	References	16

1 Practice Problems

- 19JM01 (USAJMO 2019 P1)** There are $a + b$ bowls arranged in a row, numbered 1 through $a + b$, where a and b are given positive integers. Initially, each of the first a bowls contains an apple, and each of the last b bowls contains a pear. A legal move consists of moving an apple from bowl i to bowl $i + 1$ and a pear from bowl j to bowl $j - 1$, provided that the difference $i - j$ is even. We permit multiple fruits in the same bowl at the same time. The goal is to end up with the first b bowls each containing a pear and the last a bowls each containing an apple. Show that this is possible if and only if the product ab is even.
- 12SLC1 (Shortlist 2012 C1)** Several positive integers are written in a row. Iteratively, Alice chooses two adjacent numbers x and y such that $x > y$ and x is to the left of y , and replaces the pair (x, y) by either $(y + 1, x)$ or $(x - 1, x)$. Prove that she can perform only finitely many such iterations.
- 12AM04 (USAMO 2015 P4)** Steve is piling $m \geq 1$ indistinguishable stones on the squares of an $n \times n$ grid. Each square can have an arbitrarily high pile of stones. After he finished piling his stones in some manner, he can then perform stone moves, defined as follows. Consider any four grid squares, which are corners of a rectangle, i.e. in positions $(i, k), (i, l), (j, k), (j, l)$ for some $1 \leq i, j, k, l \leq n$, such that $i < j$ and $k < l$. A stone move consists of either removing one stone from each of (i, k) and (j, l) and moving them to (i, l) and (j, k) respectively, or removing one stone from each of (i, l) and (j, k) and moving them to (i, k) and (j, l) respectively.
- Two ways of piling the stones are equivalent if they can be obtained from one another by a sequence of stone moves.
- How many different non-equivalent ways can Steve pile the stones on the grid?
- 11AM02 (USAMO 2011 P2)** An integer is assigned to each vertex of a regular pentagon so that the sum of the five integers is 2011. A turn of a solitaire game consists of subtracting an integer m from each of the integers at two neighboring vertices and adding $2m$ to the opposite vertex, which is not adjacent to either of the first two vertices. (The amount m and the vertices chosen can vary from turn to turn.) The game is won at a certain vertex if, after some number of turns, that vertex has the number 2011 and the other four vertices have the number 0. Prove that for any choice of the initial integers, there is exactly one vertex at which the game can be won.
- 19CAN3 (Canadian MO 2019 P3)** You have a $2m$ by $2n$ grid of squares coloured in the same way as a standard checkerboard. Find the total number of ways to place mn counters on white squares so that each square contains at most one counter and no two counters are in diagonally adjacent white squares.
- 21JM04 (USAJMO 2021 P4)** Carina has three pins, labeled A, B , and C , respectively, located at the origin of the coordinate plane. In a move, Carina may move a pin to an adjacent lattice point at distance 1 away. What is the least number of moves that Carina can make in order for triangle ABC to have area 2021? (A lattice point is a point (x, y) in the coordinate plane where x and y are both integers, not necessarily positive.)
- 08AM04 (USAMO 2008 P4)** Let \mathcal{P} be a convex polygon with n sides, $n \geq 3$. Any set of $n - 3$ diagonals of \mathcal{P} that do not intersect in the interior of the polygon determine a triangulation of \mathcal{P} into $n - 2$ triangles. If \mathcal{P} is regular and there is a triangulation of \mathcal{P} consisting of only isosceles triangles, find all the possible values of n .

21PAGM05 (PAGMO 2021 P5) Celeste has an unlimited amount of each type of n types of candy, numerated type 1, type 2, ... type n . Initially she takes $m > 0$ candy pieces and places them in a row on a table. Then, she chooses one of the following operations (if available) and executes it:

1. She eats a candy of type k , and in its position in the row she places one candy type $k - 1$ followed by one candy type $k + 1$ (we consider type $n + 1$ to be type 1, and type 0 to be type n).
2. She chooses two consecutive candies which are the same type, and eats them.

Find all positive integers n for which Celeste can leave the table empty for any value of m and any configuration of candies on the table.

25X00K1 (XOOK 2025 P1) For the Exeter disco party, Oron wants to tile a 2024×2024 square in the middle of the dance floor with 1×1 tiles which have arrows pointing either up, left, right, or down.

After placing down all the tiles in some configuration, he realizes that he wants to rotate some of the tiles so that no matter which tile a dancer starts on, after following the arrows they end up leaving the grid through some fixed cell. What's the minimum number of tiles he needs to rotate so that this is always possible?

18SLN1 (Shortlist 2018 N1) Determine all pairs (n, k) of distinct positive integers such that there exists a positive integer s for which the number of divisors of sn and of sk are equal.

74IM04 (IMO 1974 P4) Consider decompositions of an 8×8 chessboard into p non-overlapping rectangles subject to the following conditions: (i) Each rectangle has as many white squares as black squares. (ii) If a_i is the number of white squares in the i -th rectangle, then $a_1 < a_2 < \dots < a_p$. Find the maximum value of p for which such a decomposition is possible. For this value of p , determine all possible sequences a_1, a_2, \dots, a_p .

18EGM04 (EGMO 2018 P4) A domino is a 1×2 or 2×1 tile. Let $n \geq 3$ be an integer. Dominoes are placed on an $n \times n$ board in such a way that each domino covers exactly two cells of the board, and dominoes do not overlap. The value of a row or column is the number of dominoes that cover at least one cell of this row or column. The configuration is called balanced if there exists some $k \geq 1$ such that each row and each column has a value of k . Prove that a balanced configuration exists for every $n \geq 3$, and find the minimum number of dominoes needed in such a configuration.

2 Solutions

2.1 Lecture Notes

2.1.1 USAJMO 2019 P1

Problem Statement

There are $a + b$ bowls arranged in a row, numbered 1 through $a + b$, where a and b are given positive integers. Initially, each of the first a bowls contains an apple, and each of the last b bowls contains a pear. A legal move consists of moving an apple from bowl i to bowl $i + 1$ and a pear from bowl j to bowl $j - 1$, provided that the difference $i - j$ is even. We permit multiple fruits in the same bowl at the same time. The goal is to end up with the first b bowls each containing a pear and the last a bowls each containing an apple. Show that this is possible if and only if the product ab is even.

2.1.2 Shortlist 2012 C1

Problem Statement

Several positive integers are written in a row. Iteratively, Alice chooses two adjacent numbers x and y such that $x > y$ and x is to the left of y , and replaces the pair (x, y) by either $(y + 1, x)$ or $(x - 1, x)$. Prove that she can perform only finitely many such iterations.

2.1.3 USAMO 2015 P4

Problem Statement

Steve is piling $m \geq 1$ indistinguishable stones on the squares of an $n \times n$ grid. Each square can have an arbitrarily high pile of stones. After he finished piling his stones in some manner, he can then perform stone moves, defined as follows. Consider any four grid squares, which are corners of a rectangle, i.e. in positions $(i, k), (i, l), (j, k), (j, l)$ for some $1 \leq i, j, k, l \leq n$, such that $i < j$ and $k < l$. A stone move consists of either removing one stone from each of (i, k) and (j, l) and moving them to (i, l) and (j, k) respectively, or removing one stone from each of (i, l) and (j, k) and moving them to (i, k) and (j, l) respectively.

Two ways of piling the stones are equivalent if they can be obtained from one another by a sequence of stone moves.

How many different non-equivalent ways can Steve pile the stones on the grid?

2.2 Mandatory

2.2.1 USAMO 2011 P2

Problem Statement

An integer is assigned to each vertex of a regular pentagon so that the sum of the five integers is 2011. A turn of a solitaire game consists of subtracting an integer m from each of the integers at two neighboring vertices and adding $2m$ to the opposite vertex, which is not adjacent to either of the first two vertices. (The amount m and the vertices chosen can vary from turn to turn.) The game is won at a certain vertex if, after some number of turns, that vertex has the number 2011 and the other four vertices have the number 0. Prove that for any choice of the initial integers, there is exactly one vertex at which the game can be won.

2.2.2 Canadian MO 2019 P3

Problem Statement

You have a $2m$ by $2n$ grid of squares coloured in the same way as a standard checkerboard. Find the total number of ways to place mn counters on white squares so that each square contains at most one counter and no two counters are in diagonally adjacent white squares.

2.2.3 USAJMO 2021 P4

Problem Statement

Carina has three pins, labeled A , B , and C , respectively, located at the origin of the coordinate plane. In a move, Carina may move a pin to an adjacent lattice point at distance 1 away. What is the least number of moves that Carina can make in order for triangle ABC to have area 2021? (A lattice point is a point (x, y) in the coordinate plane where x and y are both integers, not necessarily positive.)

2.2.4 USAMO 2008 P4

Problem Statement

Let \mathcal{P} be a convex polygon with n sides, $n \geq 3$. Any set of $n - 3$ diagonals of \mathcal{P} that do not intersect in the interior of the polygon determine a triangulation of \mathcal{P} into $n - 2$ triangles. If \mathcal{P} is regular and there is a triangulation of \mathcal{P} consisting of only isosceles triangles, find all the possible values of n .

2.3 Not mandatory

2.3.1 PAGMO 2021 P5

Problem Statement

Celeste has an unlimited amount of each type of n types of candy, numerated type 1, type 2, ... type n . Initially she takes $m > 0$ candy pieces and places them in a row on a table. Then, she chooses one of the following operations (if available) and executes it:

1. She eats a candy of type k , and in its position in the row she places one candy type $k - 1$ followed by one candy type $k + 1$ (we consider type $n + 1$ to be type 1, and type 0 to be type n).
2. She chooses two consecutive candies which are the same type, and eats them.

Find all positive integers n for which Celeste can leave the table empty for any value of m and any configuration of candies on the table.

2.3.2 XOOK 2025 P1

Problem Statement

For the Exeter disco party, Oron wants to tile a 2024×2024 square in the middle of the dance floor with 1×1 tiles which have arrows pointing either up, left, right, or down.

After placing down all the tiles in some configuration, he realizes that he wants to rotate some of the tiles so that no matter which tile a dancer starts on, after following the arrows they end up leaving the grid through some fixed cell. What's the minimum number of tiles he needs to rotate so that this is always possible?

2.3.3 Shortlist 2018 N1

Problem Statement

Determine all pairs (n, k) of distinct positive integers such that there exists a positive integer s for which the number of divisors of sn and of sk are equal.

2.3.4 IMO 1974 P4

Problem Statement

Consider decompositions of an 8×8 chessboard into p non-overlapping rectangles subject to the following conditions: (i) Each rectangle has as many white squares as black squares. (ii) If a_i is the number of white squares in the i -th rectangle, then $a_1 < a_2 < \dots < a_p$.

2.3.5 EGMO 2018 P4

Problem Statement

A domino is a 1×2 or 2×1 tile. Let $n \geq 3$ be an integer. Dominoes are placed on an $n \times n$ board in such a way that each domino covers exactly two cells of the board, and dominoes do not overlap. The value of a row or column is the number of dominoes that cover at least one cell of this row or column. The configuration is called balanced if there exists some $k \geq 1$ such that each row and each column has a value of k . Prove that a balanced configuration exists for every $n \geq 3$, and find the minimum number of dominoes needed in such a configuration.

3 References