

Qwen3-8B [same_as_student]

1 0 0 0 .\n\n So the key is to compute the total number of subsets (i .e ., \$ 2 ^{ 1 0 \ cdot 1 0 } = 1 0 2 4 0 0 \$) and subtract the term \$ f (0) \$, which is the number of subsets that sum to 0 (i .e ., the empty set).\n\n \$\$ \n f (0) = 1 \ \Rightarrow P = 1 0 2 4 0 0 - 1 = \boxed{ 1 0 2 3 9 9 }\n \$\$\$ \n\n ---\n\n ### 🧐 Final Answer :\n\n \$\$ \n \boxed{ 1 0 2 3 9 9 }\n \$\$\$

Qwen3-1.7B [full]

1 0 0 0 .\n\n So the key is to compute the total number of subsets (i .e ., \$ 2 ^{ 1 0 \ cdot 1 0 } = 1 0 2 4 0 0 \$) and subtract the term \$ f (0) \$, which is the number of subsets that sum to 0 (i .e ., the empty set).\n\n \$\$ \n f (0) = 1 \ \Rightarrow P = 1 0 2 4 0 0 - 1 = \boxed{ 1 0 2 3 9 9 }\n \$\$\$ \n\n ---\n\n ### 🧐 Final Answer :\n\n \$\$ \n \boxed{ 1 0 2 3 9 9 }\n \$\$\$

Qwen3-1.7B [answer_only]

1 0 0 0 .\n\n So the key is to compute the total number of subsets (i .e ., \$ 2 ^{ 1 0 \ cdot 1 0 } = 1 0 2 4 0 0 \$) and subtract the term \$ f (0) \$, which is the number of subsets that sum to 0 (i .e ., the empty set).\n\n \$\$ \n f (0) = 1 \ \Rightarrow P = 1 0 2 4 0 0 - 1 = \boxed{ 1 0 2 3 9 9 }\n \$\$\$ \n\n ---\n\n ### 🧐 Final Answer :\n\n \$\$ \n \boxed{ 1 0 2 3 9 9 }\n \$\$\$