



Shah Md. Tanvir Siddiquee

**Assistant Professor, Department of CSE
Daffodil International University**

 +8801712560297  tanvir.cse@diu.edu.bd

 1st Floor, Room No. 111, Daffodil Tower5

 <https://sites.google.com/diu.edu.bd/tanvirsiddiquee/>

Canonical Table

Given That, $S \rightarrow A A$
 $A \rightarrow a A \mid b$

We will denote an expression as $S' \rightarrow .S$


$S \rightarrow .A A$
 $A \rightarrow .a A \mid .b$

We will denote an expression as $S' \rightarrow .S$

$S \rightarrow .AA$

$A \rightarrow .aA \mid .b$

So, the canonical view:




We will denote an expression as $S' \rightarrow .S$

$S \rightarrow .AA$

$A \rightarrow .aA \mid .b$

So, the canonical view:




Here we can say,

$$S \rightarrow . A A \text{ -----1}$$

$$A \rightarrow . a A \text{ -----2}$$

$$A \rightarrow . b \text{ -----3}$$

States	Action			Goto	
	a	b	\$	A	S
I ₀	S3	S4		2	1
I ₁			Accepted		
I ₂	S3	S4		5	
I ₃	S3	S4		6	
I ₄	r3	r3	r3		
I ₅	r1	r1	r1		
I ₆	r2	r2	r2		



Shift (S) = S_i (1,2,3)
 Reduce (r) = r_i (1,2,3)

THANK YOU