



Lecture 27

Produced by Dr. Worldwide

Welcome to the 305

Binomial Probability



- Consider tossing a coin with probability of heads equal to p a total of 6 times
- Q: What is the probability that we get exactly 3 heads?
- We could express all possible outcomes of tossing a coin 6 times using a tree diagram that goes on forever, but we all have lives
- Let's consider a few of the outcomes (sequences) where we get exactly 3 heads
- If $A =$ Event of Exactly 3 Heads, then $A = \{HHHTTT, TTTHHH, HTHTHT, \dots\}$
- For each outcome where A occurs, the probability is $p^3(1 - p)^3$ because each coin flip is independent
- Q: How many such sequences exist where A occurs?

Binomial Probability



- Number of ways in which we can choose k items out of n distinct things is called **n choose k**, denoted by $\binom{n}{k}$, and computed by

$$\binom{n}{k} = \frac{n!}{(n-k)! k!}$$

where $n! = n \times (n-1) \times (n-2) \cdots 3 \times 2 \times 1$ (**n factorial**)

- The numbers $\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n-1}, \binom{n}{n}$ are called **binomial coefficients**, since

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$$

Binomial Probability



- From coin example,
$$P(\text{Exactly 3 Heads}) = P(A) = \binom{6}{3} p^3 (1 - p)^3$$
- **Bernoulli process** is a repetition of **fixed number** of **independent** trials with a **binary** outcome where the probability of each outcome remains **constant**
- Each trial/experiment is called a **Bernoulli trial**
- For a Bernoulli process, the probability of k successes in n trials is
$$\binom{n}{k} p^k (1 - p)^{n-k}$$
- These probabilities build the **binomial distribution**
- Excel formula is `=BINOM.DIST(n, k, p, FALSE)`

Binomial Probability




- From coin example,
$$P(\text{Exactly 3 Heads}) = P(A) = \binom{6}{3} p^3 (1 - p)^3$$
- **Bernoulli process** is a repetition of **fixed number** of **independent** trials with a **binary** outcome where the probability of each outcome remains **constant**
- Each trial/experiment is called a **Bernoulli trial**
- For a Bernoulli process, the probability of k successes in n trials is
$$\binom{n}{k} p^k (1 - p)^{n-k}$$
- These probabilities build the **binomial distribution**
- Excel formula is `=BINOM.DIST(n, k, p, FALSE)`

Ex: Balls and Urns



- Suppose we have a fair coin and two urns with six balls each
 - Urn 1 has 4 white balls and 2 red balls
 - Urn 2 has 5 red balls and 1 blue ball
- Experiment
 - Flip a fair coin
 - If heads, draw a ball from urn 1
 - If tails, draw a ball from urn 2




- Q: What are the possible colors of the ball we draw?



Ex: Balls and Urns



- Probability tree to describe all possible outcomes



- List of all outcomes
 - Flip heads and grab red ball ($H \cap R$)
 - Flip heads and grab white ball ($H \cap W$)
 - Flip tails and grab red ball ($T \cap R$)
 - Flip tails and grab blue ball ($T \cap B$)

Ex: Balls and Urns



- Probability of all outcomes

- $P(H \cap R) = \frac{1}{2} \times \frac{2}{6} = \frac{1}{6}$
- $P(H \cap W) = \frac{1}{2} \times \frac{4}{6} = \frac{1}{3}$
- $P(T \cap R) = \frac{1}{2} \times \frac{5}{6} = \frac{5}{12}$
- $P(T \cap B) = \frac{1}{2} \times \frac{1}{6} = \frac{1}{12}$

- Sample space is $\Omega = \{H \cap R, H \cap W, T \cap R, T \cap B\}$ & $P(\Omega) = \frac{1}{6} + \frac{1}{3} + \frac{5}{12} + \frac{1}{12} = 1$


- Probabilities of each color

- $P(R) = \frac{1}{6} + \frac{5}{12} = \frac{7}{12}$
- $P(W) = \frac{1}{3}$
- $P(B) = \frac{1}{12}$

Ex: Balls and Urns



- Recall the probability tree



- Q: What is probability of a white ball given we flip heads?
 $P(W|H) = ??$
- Q: Given we have a white ball, what is the probability the coin flip was heads?
 $P(H|W) = ??$

Bayesian Analysis



- Recall the formula for conditional probability

$$P(X|Y) = \frac{P(X \cap Y)}{P(Y)}$$

- Based off this formula,

$$P(Y|X) = \frac{P(X \cap Y)}{P(X)} \quad \rightarrow \quad P(Y|X)P(X) = P(X \cap Y)$$

- With simple substitution,

$$P(X|Y) = \frac{P(Y|X)P(X)}{P(Y)}$$

- Since there are two events, $P(Y) = P(Y \cap X) + P(Y \cap X^c)$
- This last fact comes from the two branches of a probability tree

Bayesian Analysis



- Based off past statements
 - $P(Y \cap X) = P(Y|X)P(X)$
 - $P(Y \cap X^c) = P(Y|X^c)P(X^c)$

- Bayes' rule


$$P(X|Y) = \frac{P(Y|X)P(X)}{P(Y)} = \frac{P(Y|X)P(X)}{P(Y|X)P(X) + P(Y|X^c)P(X^c)}$$

- Bayesian analysis
 - Focus on finding posterior probability $P(X|Y)$
 - Prior probability is found in $P(X)$
 - The term $P(Y|X)$ and is typically the likelihood based off gathered data

Ex: Balls and Urns



- Back to the probability tree



- Q: Given we have a white ball, what is the probability the coin flip was heads?

$$P(H|W) = \frac{P(W|H)P(H)}{P(W|H)P(H) + P(W|T)P(T)} = \frac{\left(\frac{4}{6}\right) \left(\frac{1}{2}\right)}{\left(\frac{4}{6}\right) \left(\frac{1}{2}\right) + (0) \left(\frac{1}{2}\right)} = 1$$

- Q: Given we have a red ball, what is the probability the coin flip was heads?

$$P(H|R) = \frac{P(R|H)P(H)}{P(R|H)P(H) + P(R|T)P(T)} = \frac{\left(\frac{2}{6}\right) \left(\frac{1}{2}\right)}{\left(\frac{2}{6}\right) \left(\frac{1}{2}\right) + \left(\frac{5}{6}\right) \left(\frac{1}{2}\right)} = \frac{2}{7}$$

Random Variables



- A **random variable** X is a function that assigns a real number to each point in the sample space

$$X(s): S \rightarrow \mathbb{R}$$

- Examples
 - Tossing a 6-sided die, $X = \text{Result of Die}$, $X \in \{1, 2, 3, 4, 5, 6\}$
 - Observe weather tomorrow, $X = \begin{cases} 1, & \text{if it rains} \\ 0, & \text{if it doesn't rain} \end{cases}$ $X \in \{0, 1\}$
 - Wait for the bus, $X = \text{Time Spent Waiting}$, $X \in [0, \infty)$
 - Flip a coin 12 times, $X = \text{Number of Heads}$, $X \in \{0, 1, 2, 3, \dots, 12\}$
- Two types
 - A **discrete** random variable can take on at most a **countable** number of values
 - A **continuous** random variable has an **uncountable** number of values

Random Variables



- Examples of discrete random variables
 - Outcome of a toss of a die (Recoded to binary)
 - Response to a survey question (1 to 5)
 - Number of heads in first 10 tosses
 - Number of cars that pass in front of the Old Well between 8AM and 6PM
 - Grade in school
 - Spread between scores in a basketball game
- Examples of continuous random variables
 - Weight of a baby
 - Height of a giraffe
 - Time a person spends walking per day
 - Age of person
- Q: Can you count the set of integers $\mathbb{Z} = \{0, 1, 2, 3, \dots\}$?

Discrete Random Variable



- The **probability mass function** $p(x)$ (**pmf**) is a function that assigns a probability to every possible value of a discrete random variable X

$$p(x): \{x_1, x_2, \dots\} \rightarrow [0,1]$$

- Since $p(x)$ is a probability law, $\sum_{i=1}^{\infty} p(x_i) = 1$
- Although $p(x)$ is a function, we interpret it as $p(x) = P(X = x)$
- For any $A \subseteq \{x_1, x_2, \dots\}$, we have $P(X \in A) = \sum_{x_i \in A} p(x_i)$
- The **cumulative distribution function** $F(x) = P(X \leq x)$
- For a discrete random variable, $F(x) = P(X \leq x) = \sum_{x_i \leq x} p(x_i)$

Ex: Tossing Coin



- We toss a fair coin 10 times
 - Q: What is the probability there will be at most 2 heads?
 - The random variable $X = \text{number of heads}$
 - The variable X is a binomial random variable

$$p(x) = \binom{10}{x}(0.5)^x(0.5)^{10-x} \text{ where } x \in \{0,1,2, \dots, 10\}$$

- We want to compute $F(2) = P(X \leq 2)$

$$\begin{aligned} F(2) &= P(X \leq 2) = p(0) + p(1) + p(2) \\ &= (0.5)^{10} + \binom{10}{1}(0.5)^1(0.5)^9 + \binom{10}{2}(0.5)^2(0.5)^8 \\ &= \left(\frac{1}{2}\right)^{10} (1 + 10 + 45) = \frac{7}{2^7} \approx 0.0547 \end{aligned}$$

- In Excel, $F(2) = \text{BINOM.DIST}(2,10,0.5, \text{TRUE})$
 $= \text{BINOM.DIST}(0,10,0.5, \text{FALSE})$
 $+ \text{BINOM.DIST}(1,10,0.5, \text{FALSE})$
 $+ \text{BINOM.DIST}(2,10,0.5, \text{FALSE})$

Ex: Rolling Die



- Toss a fair die until the first 6 shows up
 - Q: What is the probability we will need to toss the die at least 4 times?
 - The random variable $X = \text{number of tosses to get the first 6}$
 - The variable X is a geometric random variable

$$p(x) = \left(\frac{5}{6}\right)^{x-1} \left(\frac{1}{6}\right) \text{ where } x \in \{1, 2, 3, \dots\}$$

- We want to compute $P(X \geq 4) = 1 - P(X < 4) = 1 - F(3)$

$$\begin{aligned} 1 - P(X < 4) &= 1 - [p(1) + p(2) + p(3)] = 1 - p(1) - p(2) - p(3) \\ &= 1 - \frac{1}{6} - \frac{5}{6} \times \frac{1}{6} - \left(\frac{5}{6}\right)^2 \times \frac{1}{6} = \frac{5^3}{6^3} \approx 0.5787 \end{aligned}$$

- In Excel, there is not a "GEOM.DIST" function that can be used
- Can be calculated using Excel's calculator functions



The End



Dale

