

Guided Tour of Machine Learning in Finance

Week 3-1-2-2: Unsupervised Learning

Principal Component Analysis (PCA)

Igor Halperin

NYU Tandon School of Engineering, 2017

PCA approach: eigen-portfolios

Try PCA as a way to **extract factors** directly from the data

Data: history of $N + 1$ daily prices of M stocks $S_n^{(i)} \equiv S_{t_n}^{(i)}$ for $i = 1, \dots, M$ measured at times $t = [t_0, t_1, \dots, t_N]$

1. Compute **daily returns**:

$$R_{ni} = \frac{S_n^{(i)} - S_{n-1}^{(i)}}{S_{n-1}^{(i)}} \simeq \log \frac{S_n^{(i)}}{S_{n-1}^{(i)}}, \quad n = 1, \dots, N, \quad i = 1, \dots, M$$

2. **Standardized returns** (data normalization):

$$X_{ni} = \frac{R_{ni} - \bar{R}_i}{\bar{\sigma}_i}, \quad \bar{R}_i = \frac{1}{N} \sum_{n=1}^N R_{ni}, \quad \bar{\sigma}_i^2 = \frac{1}{N-1} \sum_{n=1}^N (R_{ni} - \bar{R}_i)^2$$


3. The empirical **correlation matrix** is the covariance matrix of standardized returns:

$$C_{ij} = \frac{1}{N-1} \sum_{n=1}^N X_{ni} X_{nj} = \frac{1}{N-1} (\mathbf{X}^T \mathbf{X})_{ij}$$

This matrix is not diagonal!

Distribution of pairwise correlations in the DJI


Histogram of pairwise return correlations for stocks in the DJI index:



PCA as a coordinate transform

How to make correlation matrix \mathbf{C} diagonal?

Introduce a linear transform (linear encoder) of the data $\mathbf{Z} = \mathbf{XV}$ parametrized by a $p \times k$ orthogonal matrix \mathbf{V} with $\mathbf{V}\mathbf{V}^T = \mathbf{I}$



PCA as a coordinate transform

How to make correlation matrix \mathbf{C} diagonal?

Introduce a linear transform (linear encoder) of the data $\mathbf{Z} = \mathbf{XV}$ parametrized by a $p \times k$ orthogonal matrix \mathbf{V} with $\mathbf{V}\mathbf{V}^T = \mathbf{I}$

$$\begin{matrix} \text{width } k \\ \text{height } N \end{matrix} \mathbf{Z} = \begin{matrix} \text{width } p \\ \text{height } N \end{matrix} \mathbf{X} \times \begin{matrix} \text{width } k \\ \text{height } p \end{matrix} \mathbf{V}$$

A **decoded signal** (inverse transform) is obtained as $\hat{\mathbf{X}} = \mathbf{ZV}^T$

$$\begin{matrix} \text{width } p \\ \text{height } N \end{matrix} \hat{\mathbf{X}} = \begin{matrix} \text{width } k \\ \text{height } N \end{matrix} \mathbf{Z} \times \begin{matrix} \text{width } p \\ \text{height } k \end{matrix} \mathbf{V}^T$$

PCA: Eigenvalue decomposition


Eigenvalue decomposition of the correlation matrix (note that \mathbf{C} is non-negative definite!)

$$\mathbf{C} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$$

where (assuming that $N > p$):

$\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_p)$ ($\lambda_1 \geq \dots \geq \lambda_p$) is a diagonal matrix of ordered eigenvalues

\mathbf{U} is a $N \times N$ orthogonal matrix ($\mathbf{U}\mathbf{U}^T = \mathbf{I}$) that stores eigenvectors column-wise



PCA: Eigenvalue decomposition

Eigenvalue decomposition of the correlation matrix (note that \mathbf{C} is non-negative definite!)

$$\mathbf{C} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$$

where (assuming that $N > p$):

$\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_p)$ ($\lambda_1 \geq \dots \geq \lambda_p$) is a diagonal matrix of ordered eigenvalues

\mathbf{U} is a $N \times N$ orthogonal matrix ($\mathbf{U}\mathbf{U}^T = \mathbf{I}$) that stores eigenvectors column-wise

Use this to compute the covariance matrix of $\mathbf{Z} = \mathbf{X}\mathbf{V}$

$$\text{Cov}[\mathbf{Z}] = \frac{1}{N-1} \mathbf{Z}^T \mathbf{Z} = \frac{1}{N-1} \mathbf{V}^T \mathbf{X}^T \mathbf{X} \mathbf{V} = \mathbf{V}^T \mathbf{C} \mathbf{V} = \mathbf{V}^T \mathbf{U} \mathbf{\Lambda} (\mathbf{V}^T \mathbf{U})^T$$

PCA: Eigenvalue decomposition

Eigenvalue decomposition of the correlation matrix (note that \mathbf{C} is non-negative definite!)

$$\mathbf{C} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$$


where (assuming that $N > p$):

$\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_p)$ ($\lambda_1 \geq \dots \geq \lambda_p$) is a diagonal matrix of ordered eigenvalues

\mathbf{U} is a $N \times N$ orthogonal matrix ($\mathbf{U}\mathbf{U}^T = \mathbf{I}$) that stores eigenvectors column-wise

Use this to compute the covariance matrix of $\mathbf{Z} = \mathbf{X}\mathbf{V}$

$$\text{Cov}[\mathbf{Z}] = \frac{1}{N-1} \mathbf{Z}^T \mathbf{Z} = \frac{1}{N-1} \mathbf{V}^T \mathbf{X}^T \mathbf{X} \mathbf{V} = \mathbf{V}^T \mathbf{C} \mathbf{V} = \mathbf{V}^T \mathbf{U} \mathbf{\Lambda} (\mathbf{V}^T \mathbf{U})^T \xrightarrow{\mathbf{V}=\mathbf{U}} \mathbf{\Lambda}$$



PCA as a coordinate transform

Eigenvalue decomposition of the correlation matrix (note that \mathbf{C} is non-negative definite!)

$$\mathbf{C} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$$

where (assuming that $N > p$):

$\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_p)$ ($\lambda_1 \geq \dots \geq \lambda_p$) is a diagonal matrix of ordered eigenvalues

\mathbf{U} is a $N \times N$ orthogonal matrix ($\mathbf{U}\mathbf{U}^T = \mathbf{I}$) that stores eigenvectors column-wise

Use this to compute the covariance matrix of $\mathbf{Z} = \mathbf{X}\mathbf{V}$

$$\text{Cov}[\mathbf{Z}] = \frac{1}{N-1} \mathbf{Z}^T \mathbf{Z} = \frac{1}{N-1} \mathbf{V}^T \mathbf{X}^T \mathbf{X} \mathbf{V} = \mathbf{V}^T \mathbf{C} \mathbf{V} = \mathbf{V}^T \mathbf{U} \mathbf{\Lambda} (\mathbf{V}^T \mathbf{U})^T \xrightarrow{\mathbf{V}=\mathbf{U}} \mathbf{\Lambda}$$

Therefore, when $\mathbf{V} = \mathbf{U}$ (and $k = p$), encoding $\mathbf{Z} = \mathbf{X}\mathbf{V}$ **preserves the total variation of data**



$$\text{TotVar}[\mathbf{X}] = \frac{1}{N-1} \text{Tr}[\mathbf{X}^T \mathbf{X}] = \text{Tr}[\mathbf{C}] = \text{Tr}[\mathbf{U}\mathbf{\Lambda}\mathbf{U}^T] = \text{Tr}[\mathbf{\Lambda}\mathbf{U}^T \mathbf{U}] = \text{Tr}[\mathbf{\Lambda}]$$

$$\text{TotVar}[\mathbf{Z}] = \frac{1}{N-1} \text{Tr}[\mathbf{Z}^T \mathbf{Z}] = \text{Tr}[\mathbf{V}^T \mathbf{U} \mathbf{\Lambda} (\mathbf{V}^T \mathbf{U})^T] \xrightarrow{\mathbf{V}=\mathbf{U}} \text{Tr}[\mathbf{\Lambda}]$$







PCA for stocks in DJI: eigenvectors

Eigenvectors = weights of stocks in “eigen-portfolios”

Approximates
the market (DJI)



Orthogonal to
the market (DJI)



Control question

Select all correct answers

1. The PCA implements a Linear Encoder that converts correlated inputs \mathbf{X} into uncorrelated features \mathbf{Z} by a linear transform $\mathbf{Z} = \mathbf{XV}$
2. If the orthogonal matrix \mathbf{V} has dimension $p \times p$, i.e. it keeps all eigenvectors of correlation matrix of \mathbf{X} , then \mathbf{Z} preserves the total variation of \mathbf{X} .
3. When the data is noisy or non-stationary, \mathbf{Z} can have a higher variance than \mathbf{X} .
4. The PCA is a probabilistic method that enables simulating from data.

Correct answer: 1, 2