# Triangular grids

Shantanu Nene

Polymath Jr 2023


August 2023

## Half-rectangular grids

An  $m \times n$  half-rectangular-grid is the set of non-negative integer points (x,y) such that  $0 \le x \le n-1$  and  $0 \le y \le m-1$  and  $(m-1)x \le (n-1)y$ .

## Half-rectangular grids

An  $m \times n$  half-rectangular-grid is the set of non-negative integer points (x,y) such that  $0 \le x \le n-1$  and  $0 \le y \le m-1$  and  $(m-1)x \le (n-1)y$ .



## Covering while omitting a point

#### Theorem

Let G be an  $m \times n$  half-rectangular grid, and let  $P = (x_0, y_0) \in G$  be any point. The minimum number of lines required to cover  $G \setminus P$  without covering P is  $n - \lceil \frac{n-m}{m-1} y_0 \rceil - 1$ .

## Covering while omitting a point

#### **Theorem**

Let G be an  $m \times n$  half-rectangular grid, and let  $P = (x_0, y_0) \in G$  be any point. The minimum number of lines required to cover  $G \setminus P$  without covering P is  $n - \lceil \frac{n-m}{m-1} y_0 \rceil - 1$ .

As a special case, in case of m = n, the minimum number of lines to cover  $G \setminus P$  without covering P is n - 1, regardless of position of P.

# Covering with multiplicity


# Covering with multiplicity

### Theorem (Basit, Clifton, Horn (2023))


Number of lines required to cover an  $n \times n$  half-rectangular grid with multiplicity k is greater than  $\frac{2}{3}nk$ 

We define a general triangular grid T to be a subset of an  $n \times n$  grid such that the  $i^{th}$  row from the top has exactly i points.

We define a general triangular grid T to be a subset of an  $n \times n$  grid such that the  $i^{th}$  row from the top has exactly i points.




We define a general triangular grid T to be a subset of an  $n \times n$  grid such that the  $i^{th}$  row from the top has exactly i points.



Note that an  $n \times n$  half-rectangular grid is an example of a general triangular grid.

We define a general triangular grid T to be a subset of an  $n \times n$  grid such that the  $i^{th}$  row from the top has exactly i points.



Note that an  $n \times n$  half-rectangular grid is an example of a general triangular grid.

### Theorem

Number of lines required to cover any general triangular grid with multiplicity k is at least nk  $\left(1-e^{\frac{1}{2n}-1}-\frac{2}{n}\right)\geq nk\left(1-\frac{1}{e}-O\left(\frac{1}{n}\right)\right)$ .

# Some Open Problems

• Is the constant  $1 - \frac{1}{e}$  in the previous theorem optimal?

## Some Open Problems

- **1** Is the constant  $1 \frac{1}{e}$  in the previous theorem optimal?
- ② What is a good lower bound on covering  $m \times n$  half-rectangular grids with multiplicity?