

# Set Sizes and Cardinal Numbers

Shantanu Nene

July 2021

# Introduction

Natural numbers can be thought of as counting numbers.

# Introduction

Natural numbers can be thought of as counting numbers. They indicate the size of finite sets.

# Introduction

Natural numbers can be thought of as counting numbers. They indicate the size of finite sets. However, we want to compare infinite sets in a similar way as well.

Natural numbers can be thought of as counting numbers. They indicate the size of finite sets. However, we want to compare infinite sets in a similar way as well. Thus we need to formalize our definition of 'size'.

# Bijections and Injections

We say that two sets  $A, B$  have the same **size** or **cardinality** if there is a bijection from  $A$  to  $B$ , i.e., a function that is both one-one and onto.

# Bijections and Injections

We say that two sets  $A, B$  have the same **size** or **cardinality** if there is a bijection from  $A$  to  $B$ , i.e., a function that is both one-one and onto. We also say that size of  $A$  is  $\leq$  size of  $B$  if there is an injective or one-one function from  $A$  to  $B$ .

# Bijections and Injections

We say that two sets  $A, B$  have the same **size** or **cardinality** if there is a bijection from  $A$  to  $B$ , i.e., a function that is both one-one and onto. We also say that size of  $A$  is  $\leq$  size of  $B$  if there is an injective or one-one function from  $A$  to  $B$ . We denote the size of a set  $A$  by  $|A|$ .



# Bijections and Injections

We say that two sets  $A, B$  have the same **size** or **cardinality** if there is a bijection from  $A$  to  $B$ , i.e., a function that is both one-one and onto. We also say that size of  $A$  is  $\leq$  size of  $B$  if there is an injective or one-one function from  $A$  to  $B$ . We denote the size of a set  $A$  by  $|A|$ .

## Examples

- $f(x) = 2x$  is a bijection from the set of integers to the even integers, which proves that they both have the same size.

# Bijections and Injections

We say that two sets  $A, B$  have the same **size** or **cardinality** if there is a bijection from  $A$  to  $B$ , i.e., a function that is both one-one and onto. We also say that size of  $A$  is  $\leq$  size of  $B$  if there is an injective or one-one function from  $A$  to  $B$ . We denote the size of a set  $A$  by  $|A|$ .

## Examples

- $f(x) = 2x$  is a bijection from the set of integers to the even integers, which proves that they both have the same size.
- The function  $f(a, b) = 2^{a-1}(2b - 1)$  from  $\mathbb{N} \times \mathbb{N}$  to  $\mathbb{N}$  is a bijection.

# Bijections and Injections

We say that two sets  $A, B$  have the same **size** or **cardinality** if there is a bijection from  $A$  to  $B$ , i.e., a function that is both one-one and onto. We also say that size of  $A$  is  $\leq$  size of  $B$  if there is an injective or one-one function from  $A$  to  $B$ . We denote the size of a set  $A$  by  $|A|$ .

## Examples

- $f(x) = 2x$  is a bijection from the set of integers to the even integers, which proves that they both have the same size.
- The function  $f(a, b) = 2^{a-1}(2b - 1)$  from  $\mathbb{N} \times \mathbb{N}$  to  $\mathbb{N}$  is a bijection.
- The function  $f(x) = (x - 1) \left( \frac{1 - (-1)^x}{4} \right) - x \left( \frac{1 + (-1)^x}{4} \right)$  is a bijection from  $\mathbb{N}$  to  $\mathbb{Z}$ .

# Schroder-Bernstein Theorem

In some cases, it isn't easy to construct a direct bijection.

# Schroder-Bernstein Theorem

In some cases, it isn't easy to construct a direct bijection. Consider the set of positive rational numbers  $\mathbb{Q}^+$ .

# Schroder-Bernstein Theorem

In some cases, it isn't easy to construct a direct bijection. Consider the set of positive rational numbers  $\mathbb{Q}^+$ . It is easy to see that

$$|\mathbb{Q}^+| \leq |\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$

# Schroder-Bernstein Theorem

In some cases, it isn't easy to construct a direct bijection. Consider the set of positive rational numbers  $\mathbb{Q}^+$ . It is easy to see that

$$|\mathbb{Q}^+| \leq |\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$

Also,

$$|\mathbb{N}| \leq |\mathbb{Q}^+|$$

# Schroder-Bernstein Theorem

In some cases, it isn't easy to construct a direct bijection. Consider the set of positive rational numbers  $\mathbb{Q}^+$ . It is easy to see that

$$|\mathbb{Q}^+| \leq |\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$

Also,

$$|\mathbb{N}| \leq |\mathbb{Q}^+|$$

Does this necessarily mean that the two sets have the same size?



# Schroder-Bernstein Theorem

The answer is **yes**.

## Theorem 1:

For sets  $A, B$ ,  $|A| \leq |B|$  and  $|B| \leq |A|$  imply  $|A| = |B|$ .

# Schroder-Bernstein Theorem

The answer is **yes**.

## Theorem 1:

For sets  $A, B$ ,  $|A| \leq |B|$  and  $|B| \leq |A|$  imply  $|A| = |B|$ .

The proof is skipped because it is a bit involved.

# Cantor's Theorem

Another question to ask: for a given set, can we always find a set which has a strictly bigger size?

# Cantor's Theorem

Another question to ask: for a given set, can we always find a set which has a strictly bigger size? The answer is **yes**.

## Theorem 2:

$|A| < |P(A)|$  for all non-empty sets  $A$ .

# Cantor's Theorem

Another question to ask: for a given set, can we always find a set which has a strictly bigger size? The answer is **yes**.

## Theorem 2:

$|A| < |P(A)|$  for all non-empty sets  $A$ .

**Proof:** Clearly  $|A| \leq |P(A)|$  so assume there is some bijection  $f$  between them.

# Cantor's Theorem

Another question to ask: for a given set, can we always find a set which has a strictly bigger size? The answer is **yes**.

## Theorem 2:

$|A| < |P(A)|$  for all non-empty sets  $A$ .

**Proof:** Clearly  $|A| \leq |P(A)|$  so assume there is some bijection  $f$  between them. Define a new subset  $T$  of  $S$  as follows:  $x \in T$  iff there exists an  $A \subseteq S$  such that  $f(A) = x$  and  $x \notin A$  (note that  $A$  must be unique, if it exists).

# Cantor's Theorem

Another question to ask: for a given set, can we always find a set which has a strictly bigger size? The answer is **yes**.

## Theorem 2:

$|A| < |P(A)|$  for all non-empty sets  $A$ .

**Proof:** Clearly  $|A| \leq |P(A)|$  so assume there is some bijection  $f$  between them. Define a new subset  $T$  of  $S$  as follows:  $x \in T$  iff there exists an  $A \subseteq S$  such that  $f(A) = x$  and  $x \notin A$  (note that  $A$  must be unique, if it exists). Now it is easy to see that  $f(T)$  cannot exist, contradiction!

# Cantor's Diagonal Argument

The technique that we used in the last proof is called **Cantor's Diagonal Argument** and was first discovered by Georg Cantor.



# Cantor's Diagonal Argument

The technique that we used in the last proof is called **Cantor's Diagonal Argument** and was first discovered by Georg Cantor. He also used it to prove the following result:

**Theorem 3:**

$$|\mathbb{R}| > |\mathbb{N}|$$

# Cantor's Diagonal Argument

The technique that we used in the last proof is called **Cantor's Diagonal Argument** and was first discovered by Georg Cantor. He also used it to prove the following result:

**Theorem 3:**

$$|\mathbb{R}| > |\mathbb{N}|$$

**Proof:** We will in fact show that  $[0, 1]$  has a bigger size than  $\mathbb{N}$ .

# Cantor's Diagonal Argument

The technique that we used in the last proof is called **Cantor's Diagonal Argument** and was first discovered by Georg Cantor. He also used it to prove the following result:

## Theorem 3:

$$|\mathbb{R}| > |\mathbb{N}|$$

**Proof:** We will in fact show that  $[0, 1]$  has a bigger size than  $\mathbb{N}$ . Suppose we have a list of all numbers in  $[0, 1]$  in their decimal expansions:  
 $a_1, a_2, \dots$  (with the convention of repeating 9s at the end).

# Cantor's Diagonal Argument

The technique that we used in the last proof is called **Cantor's Diagonal Argument** and was first discovered by Georg Cantor. He also used it to prove the following result:

## Theorem 3:

$$|\mathbb{R}| > |\mathbb{N}|$$

**Proof:** We will in fact show that  $[0, 1]$  has a bigger size than  $\mathbb{N}$ . Suppose we have a list of all numbers in  $[0, 1]$  in their decimal expansions:  $a_1, a_2, \dots$  (with the convention of repeating 9s at the end). Now consider a new number  $x$ , such that its  $i^{\text{th}}$  decimal place is 9 if the  $i^{\text{th}}$  decimal place of  $a_i$  is 8 and is 8 otherwise.

# Cantor's Diagonal Argument

The technique that we used in the last proof is called **Cantor's Diagonal Argument** and was first discovered by Georg Cantor. He also used it to prove the following result:

## Theorem 3:

$$|\mathbb{R}| > |\mathbb{N}|$$

**Proof:** We will in fact show that  $[0, 1]$  has a bigger size than  $\mathbb{N}$ . Suppose we have a list of all numbers in  $[0, 1]$  in their decimal expansions:  $a_1, a_2, \dots$  (with the convention of repeating 9s at the end). Now consider a new number  $x$ , such that its  $i^{\text{th}}$  decimal place is 9 if the  $i^{\text{th}}$  decimal place of  $a_i$  is 8 and is 8 otherwise. It is easy to see that this number is not on the list, contradiction! In fact, it can be shown that  $|\mathbb{R}| = |P(\mathbb{N})|$

# Defining the Cardinals

Cardinals are just "numbers" associated with sets of the same size.

# Defining the Cardinals

Cardinals are just "numbers" associated with sets of the same size. Finite sets have natural numbers associated with size, while infinite sets have "infinite cardinals".

# Defining the Cardinals

Cardinals are just "numbers" associated with sets of the same size. Finite sets have natural numbers associated with size, while infinite sets have "infinite cardinals". For example, the size of  $\mathbb{N}$  is denoted by  $\aleph_0$ .



# Defining the Cardinals

Cardinals are just "numbers" associated with sets of the same size. Finite sets have natural numbers associated with size, while infinite sets have "infinite cardinals". For example, the size of  $\mathbb{N}$  is denoted by  $\aleph_0$ . This can be proved to be the smallest infinite cardinal.

# Defining the Cardinals

Cardinals are just "numbers" associated with sets of the same size. Finite sets have natural numbers associated with size, while infinite sets have "infinite cardinals". For example, the size of  $\mathbb{N}$  is denoted by  $\aleph_0$ . This can be proved to be the smallest infinite cardinal. In fact, it can be proved that the cardinals are well-ordered, i.e., every non-empty set of cardinals has a smallest element.

# Defining the Cardinals

Cardinals are just "numbers" associated with sets of the same size. Finite sets have natural numbers associated with size, while infinite sets have "infinite cardinals". For example, the size of  $\mathbb{N}$  is denoted by  $\aleph_0$ . This can be proved to be the smallest infinite cardinal. In fact, it can be proved that the cardinals are well-ordered, i.e., every non-empty set of cardinals has a smallest element. Thus, we can talk about the next smallest infinite cardinal,  $\aleph_1$ , followed by  $\aleph_2$  and so on.

# Defining the Cardinals

Cardinals are just "numbers" associated with sets of the same size. Finite sets have natural numbers associated with size, while infinite sets have "infinite cardinals". For example, the size of  $\mathbb{N}$  is denoted by  $\aleph_0$ . This can be proved to be the smallest infinite cardinal. In fact, it can be proved that the cardinals are well-ordered, i.e., every non-empty set of cardinals has a smallest element. Thus, we can talk about the next smallest infinite cardinal,  $\aleph_1$ , followed by  $\aleph_2$  and so on. Another sequence of infinite cardinals is the beth numbers:  $\beth_0 = \aleph_0$ , and  $\beth_{k+1}$  is the cardinality of the power set of a set with size  $\beth_k$ .

# Defining the Cardinals

Cardinals are just "numbers" associated with sets of the same size. Finite sets have natural numbers associated with size, while infinite sets have "infinite cardinals". For example, the size of  $\mathbb{N}$  is denoted by  $\aleph_0$ . This can be proved to be the smallest infinite cardinal. In fact, it can be proved that the cardinals are well-ordered, i.e., every non-empty set of cardinals has a smallest element. Thus, we can talk about the next smallest infinite cardinal,  $\aleph_1$ , followed by  $\aleph_2$  and so on. Another sequence of infinite cardinals is the beth numbers:  $\beth_0 = \aleph_0$ , and  $\beth_{k+1}$  is the cardinality of the power set of a set with size  $\beth_k$ . The result that  $\aleph_1 = \beth_1$  is called the Continuum Hypothesis, and it can be shown to be independent of the ZFC axioms.

Suppose we have two cardinals  $u, v$  corresponding to disjoint sets  $A, B$ . Then we can define  $u + v$ ,  $u \cdot v$  and  $u^v$  as the cardinality of the sets  $A \cup B$ ,  $A \times B$  and the set of functions from  $B$  to  $A$  respectively.

# Cardinal Arithmetic

Suppose we have two cardinals  $u, v$  corresponding to disjoint sets  $A, B$ . Then we can define  $u + v$ ,  $u \cdot v$  and  $u^v$  as the cardinality of the sets  $A \cup B$ ,  $A \times B$  and the set of functions from  $B$  to  $A$  respectively. Then cardinals also follow the same arithmetic rules as natural numbers!

# Cardinal Arithmetic

Suppose we have two cardinals  $u, v$  corresponding to disjoint sets  $A, B$ . Then we can define  $u + v$ ,  $u \cdot v$  and  $u^v$  as the cardinality of the sets  $A \cup B$ ,  $A \times B$  and the set of functions from  $B$  to  $A$  respectively. Then cardinals also follow the same arithmetic rules as natural numbers! Further, for infinite cardinals  $u, v$ , we will have

$$u + v = u \cdot v = \max\{u, v\}$$



*Thank You!*