

Cardiovascular autonomic dysfunction impact on cardiovascular complications across glucose metabolism

Jonas Rosborg Schaarup

2025

Acknowledgements

Thanks for all the fish.

Supervisors and assessment committee

PhD student:

Jonas Rosborg Schaarup, Msc in Public Health
Department of Public Health, Aarhus University, Denmark

Supervisors:

Professor Daniel R. Witte, PhD (main supervisor)
Department of Public Health, Aarhus University, Denmark
Steno Diabetes Center Aarhus, Denmark

Lasse Bjerg, MD PhD
Department of Public Health, Aarhus University, Denmark
Steno Diabetes Center Aarhus, Denmark

Signe Toft Andersen, MD PhD
Department of Public Health, Aarhus University, Denmark Steno Diabetes Center
Aarhus, Denmark

Christian Stevens Hansen, MD PhD
Steno Diabetes Center Copenhagen, Denmark

Assessment committee:

Professor Morten Schmidt (chair and moderator of the defence) Department of Clinical Medicine - Department of Clinical Epidemiology, Aarhus University

Associate Professor Ilonca Vaartjes University Medical Center Utrecht, Julius Center, Department of Epidemiology and Health Economics

Cardiologist Peter Godsk Jørgensen Herlev and Gentofte Hospital, Capital Region of Denmark

Papers in the dissertation

Study I

Jonas R. Schaarup, Lasse Bjerg, Christian S. Hansen, Signe T. Andersen, Marleen van Greevenbroek, Miranda T. Schram, Bastiaan E. de Galan, Coen Stehouwer, Daniel R. Witte (2025). Cardiovascular autonomic dysfunction is linked with arterial stiffness across glucose metabolism: The Maastricht Study. medRxiv 2024.12.03.24317865; doi: <https://doi.org/10.1101/2024.12.03.24317865> (under peer-review at BMJ Open Diabetes Research & Care)

Study II

Jonas R. Schaarup, Lasse Bjerg, Christian S. Hansen, Erik L. Grove, Signe T. Andersen, Dorte Vistisen, Søren Brage, Annelli Sandbæk, Daniel R. Witte (2025). Cardiovascular autonomic dysfunction precedes cardiovascular disease and all-cause mortality: 11-year follow-up of the ADDITION-PRO study. medRxiv 2024.12.18.24319131; doi: <https://doi.org/10.1101/2024.12.18.24319131> (accepted at Diabetes, Obesity and Metabolism)

Study III

Cardiovascular autonomic neuropathy and indices of heart failure in type 2 diabetes: The CANCAN Study - add pieces!!!!

Additional publications

The 2 following original research studies and 2 preprints have been published during the PhD period, but have not been included in the dissertation.

Peer-reviewed

Schaarup JR, Christensen MS, Hulman A, Hansen CS, Vistisen D, Tabák AG, Witte DR, Bjerg L. Autonomic dysfunction is associated with the development of arterial stiffness: The Whitehall II cohort. *GeroScience*, 2023. <https://doi.org/10.1007/s11357-023-00762-0>

Schaarup JR, Aggarwal R, Dalsgaard E-M, Norman K, Dollerup OL, Ashrafi H, Witte DR, Sandbæk A, Hulman A. Perception of artificial intelligence-based solutions in healthcare among people with and without diabetes: A cross-sectional survey from the health in Central Denmark cohort. *Diabetes Epidemiology and Management*, 2023. <https://doi.org/10.1016/j.deman.2022.100114>

Pre-prints

Jonas R. Schaarup, Anders Aasted Isaksen, Kasper Norman, Lasse Bjerg, Adam Hulman. (2025). Trust in large language model-based solutions in healthcare among people with and without diabetes: a cross-sectional survey from the Health in Central Denmark cohort. medRxiv 2025.02.24.25322734; doi: <https://doi.org/10.1101/2025.02.24.25322734> (under review at BMJ digital health and AI)

Anders Aasted Isaksen, **Jonas R. Schaarup**, Lasse Bjerg, Adam Hulman. (2025). Changes in public perception of AI in healthcare after exposure to ChatGPT. medRxiv 2025.01.23.25321048; doi: <https://doi.org/10.1101/2025.01.23.25321048> (under review at npj digital medicine)

Table of contents

Acknowledgements	1
Supervisors and assessment committee	2
Papers in the dissertation	3
Abbreviations	10
Resume	12
1. Summary	13
2. Introduction	14
3. Background	16
3.1. T2D and prediabetes	16
3.2. Cardiovascular disease	18
3.2.1. Arteriosclerosis	18
3.2.2. Atherosclerosis	19
3.2.3. Heart failure	20
3.3. Cardiovascular autonomic dysfunction	21
3.4. Risk-stratification	23
4. Aim and hypothesis	25
5. Materials and methods	26
5.1. Overview of the studies	26
5.1.1. Study population	27
5.2. Study variables	30
5.2.1. Measures for cardiovascular autonomic dysfunction/ neuropathy .	30
5.2.2. Confounders and variables for instrumental bias	34
5.3. Outcomes	35
5.3.1. Arterial stiffness	35
5.3.2. Indicators of heart failure	36

Table of contents

5.3.3. Cardiovascular events	37
5.4. Statistical Methods	37
5.4.1. Cross-sectional analysis	37
5.4.2. Time-to-event analysis	38
5.4.3. Effect modification [det kan evt. kortes ned]	38
5.4.4. Multiple imputed by chained equations	39
5.4.5. Instrumental bias	39
6. Results	41
6.1. Study I	41
6.1.1. Descriptive	41
6.1.2. 24-hour HRV and arterial stiffness	41
6.1.3. Effect modification of diabetes status	43
6.2. Study II	44
6.2.1. Descriptive	44
6.2.2. Multiday HRV and MACE, heart failure, and all-cause mortality . .	45
6.2.3. Hourly HRV and MACE, heart failure, and all-cause mortality . .	46
6.3. Study III	49
6.3.1. Descriptive	49
6.3.2. CAN and indicators of heart failure	49
7. Discussion	51
7.1. Summary of findings	52
7.2. Cardiovascular autonomic dysfunction and its impact on heart disease across glucose metabolism	53
7.3. Clinical implications	58
7.3.1. Public health	58
7.3.2. Primary care	60
7.3.3. Secondary care	61
7.4. Strengths and limitations	63
7.4.1. Study design	63
7.4.2. Internal validity	64
7.4.3. External validity	67
8. Conclusion	69
9. Perspective	76
9.1. Continuous monitoring of cardiovascular health	76
9.2. Risk-stratification	78
9.3. Effective causal modifiable marker	80

Table of contents

References	82
Resume	92
10. Summary	93
Appendices	95
A. More results	95
B. Another appendix	96

List of Figures

3.1.	Risk-stratification based on preclinical disease. (Source: Author)	24
5.1.	Study populations	27
5.2.	Left: Holter monitor Middle: Actiheart Right: Handheld Vagus™ device .	30
5.3.	Heart rate variability. (Source: Author)	33
5.4.	CART	34
5.5.	Measures of arterial stiffness, measured dynamically at the aortic and local carotid sites. (Source: Author)	36
6.1.	?(caption)	44
6.2.	Multiday SDNN levels (100 ms, 120 ms, 160 ms) by age-specific incidence rates for A) major adverse cardiovascular events, B) heart failure, and C) all-cause mortality	47
6.3.	Hourly SDNN levels (100 ms, 120 ms, 160 ms) by age-specific incidence rates for A) major adverse cardiovascular events, B) heart failure, and C) all-cause mortality. Figure adapted from [authors]. —. (Paper II, appendix)	48
6.4.	Relationship between CAN and indicators for heart failure. Figure from [authors]. Cardiovascular autonomic neuropathy and indices of heart failure in T2D: The CANCAN Study. (Paper III, appendix)	50
7.1.	Autonomic dysfunction and arterial stiffness. (Source: Author). —.	54
9.1.	HRV feedback in response to lifestyle and treatment interventions. (Source: Author)	77
9.2.	Risk-stratification by autonomic dysfunction. (Source: Author)	79
9.3.	Mediation of HRV by intervention in prevention of CVD. (Source: Author)	80

List of Tables

5.1.	Table 1: Overview of studies	26
5.2.	Box 1 Time-domain indices reflections of autonomic function	31
5.4.	Box 2 Frequency-domain indices reflections of autonomic function	32
5.6.	?(caption)	37
6.1.	Study charteristics by diabetes status	42
6.2.	Study participants charteristics	45

Abbreviations

- BMI:** Body mass index
CAN: Cardiovascular autonomic neuropathy
CARTs: Cardiovascular autonomic reflex tests
CD: Carotid artery distensibility coefficient
cf-PWV: Carotid-femoral pulse wave velocity
CI: Confidence interval
CVD: Cardiovascular disease
DKD: Diabetic kidney disease
eGFR: Estimated glomerular filtration rate
FPG: Fasting plasma glucose
GLP1RA: Glucagon-like peptide-1 receptor agonists
HDL: High-density lipoprotein cholesterol
HRV: Heart rate variability
IR: Incidence rate
IRR: Incidence rate ratio
HbA1c: Haemoglobin-A1c
LDL-C: Low-density lipoprotein cholesterol
MACE: Three-point major adverse cardiovascular events
NT-proBNP: OGTT: Oral glucose tolerance test
OR: Odds ratio
PAEE: Physical activity energy expenditure
RPAQ: Recent Physical Activity Questionnaire
SBP: Systolic blood pressure
DBP: Diastolic blood pressure
SGLT2i: Sodium glucose co-transporter type 2 inhibitors
SES: Socioeconomic status
SD: Standard deviation **SDNN:** Standard deviation of NN intervals
SDANN: Standard deviation of the averages of NN intervals in 5-minute segments
SDNN index: Mean of the SDs of all NN intervals for all 5-minute segments
pNN50: Proportion of NN intervals differing by more than 50 ms
RMSSD: Root mean square of successive differences between NN intervals
TP: Total power (variance of NN intervals 0.4 Hz)
ULF: Ultra low-frequency range (0.003 Hz)

Abbreviations

VLF: Very-low-frequency range (0.003–0.04 Hz)

LF: Low-frequency range (0.04–0.15 Hz)

HF: High-frequency range (0.15–0.4 Hz)

T2D: Type 2 diabetes mellitus

TC: Total cholesterol

TG: Triglycerides

UACR: Urine albumin-to-creatinine ratio

Resume

Denne afhandling undersøgte, om kardiovaskulær autonom dysfunktion er forbundet med kardiovaskulære komplikationer hos personer med normal glukosemetabolisme, prædiabetes og type 2-diabetes. Kardiovaskulær autonom dysfunktion blev målt igennme hjeretrymefvariabilitet (HRV) og standardiserede kardiovaskulære autonome refleksundersøgelser (CARTs). Blandt målingerne gav 24-timers HRV indsigt i den autonome balance og døgnrytmens skift mellem sympathisk og parasympatisk aktivitet. Lavere værdier var konsekvent forbundet med ugunstige kardiovaskulære udfald.

På tværs af tre studier blev autonom dysfunktion forbundet med højere niveauer af arteriel stivhed, en højere forekomst af iskæmiske hjertekarsygedom, hjertesvigt og dødelighed. Sammenhængen med arteriel stivhed blev observeret på tværs af hele spektret af glukosemetabolisme og var særligt forhøjet hos personer med prædiabetes, hvilket antyder, at autonom dysfunktion. På baggrund af dette fokuserede det efterfølgende studie på personer med høj risiko for at udvikle diabetes. I denne population var kardiovaskulær autonom dysfunktion forbundet med højere risiko for kardiovaskulær sygdom, hjertesvigt og dødelighed.

Kardiovaskulær autonom neuropati, defineret gennem abnormale CARTs, var forbundet med en højere risiko for hjertesvigt hos individer med type 2 diabetes, selv hos personer uden symptomer. Disse fund tyder på, at kardiovaskulær autonom dysfunktion kan fungere som en tidlig og uafhængig markør for risikoen for hjertesvigt, især i populationer med risiko for type 2-diabetes.

Afhandlingen konkluderer, at kardiovaskulær autonom dysfunktion er en klinisk relevant risikofaktor, som bør tiltrække større opmærksomhed i både forskning og klinisk praksis. Fremtidige studier bør undersøge, om forbedring af autonom funktion kan reducere kardiovaskulær risiko, og om målinger af autonom dysfunktion kan integreres i eksisterende risikomodeller. Derudover er potentialet i smarture til kontinuerlig overvågning og tidlig opsporing en mulighed, der kan vurderes i både kliniske og generelle populationer.

1. Summary

This dissertation investigated whether cardiovascular autonomic dysfunction is associated with cardiovascular complications in individuals with normal glucose metabolism, prediabetes, and type 2 diabetes. Cardiovascular autonomic dysfunction was assessed using heart rate variability and standardized cardiovascular autonomic reflex tests. Among the measures, 24-hour heart rate variability reflected autonomic balance and circadian shifts in sympathetic and parasympathetic activity. Lower long-term heart rate variability was consistently associated with adverse cardiovascular outcomes.

Across three studies, autonomic dysfunction was linked to higher levels of arterial stiffness, a higher incidence of ischemic cardiovascular events, heart failure, and all-cause mortality. The association with arterial stiffness was observed across the full spectrum of glucose metabolism and was particularly pronounced in individuals with prediabetes, suggesting that autonomic dysfunction may play a role early in the pathophysiological process. Building on this, the subsequent studies focused on individuals at high risk of developing diabetes. In this population, cardiovascular autonomic dysfunction was associated with a higher risk of cardiovascular disease, heart failure, and mortality.

Cardiovascular autonomic neuropathy, as identified through abnormal cardiovascular reflex testing, was associated with a higher risk of heart failure, even in individuals without symptoms. These findings suggest that cardiovascular autonomic dysfunction may serve as an early and independent marker of heart failure risk, particularly in populations at risk of type 2 diabetes.

The dissertation concludes that cardiovascular autonomic dysfunction is a clinically relevant risk factor that warrants further attention in both research and clinical practice. Future studies should explore whether improving autonomic function can reduce cardiovascular risk and whether measures of autonomic dysfunction can be integrated into existing risk models. Additionally, the potential of wearable technologies for continuous monitoring and early detection is a possibility to be evaluated in both clinical and general populations.

2. Introduction

Diabetes mellitus is a growing global health concern, posing pressing challenges for public health systems¹. As prevalence rises, more individuals are exposed to an increased risk of premature mortality and cardiovascular disease (CVD)¹. People live longer lives with diabetes they face longer periods under the burden of diabetes complications². Despite advancements in cardiovascular care, many complications are still detected at more advanced stages. These include coronary artery disease, often identified through ischemia or major CVD events and symptomatic heart failure. Early detection of CVD risk and asymptomatic heart failure, such as increased atheroma burden or subclinical heart failure, is desirable^{[3]4}.

Over the last decades, cardiovascular autonomic dysfunction has repeatedly gained attention as a risk factor for CVD⁵. Heart rate variability (HRV) is considered a reliable marker for measuring autonomic function, as it reflects the balance between sympathetic and parasympathetic modulation of heart rate intervals⁶. Despite its recognition as a CVD risk factor, cardiovascular autonomic dysfunction has not been implemented in healthcare practice. In diabetes, lower HRV is regarded as an early indicator of cardiovascular autonomic neuropathy (CAN), which is diagnosed using cardiovascular autonomic reflex tests (CARTs)⁷. Signs of autonomic dysfunction, may already be present in individuals with prediabetes⁸. Despite rising prevalence and increased CVD risk, people with prediabetes often remain outside structured treatment pathways [9]¹⁰. Although diabetes contributes to autonomic dysfunction, it is still unclear at what stage in the diabetes risk spectrum HRV and CARTs become clinically useful for assessing CVD risk.

In the past, measuring HRV needed special instruments like an electrocardiogram. Today, it's easy to track HRV with everyday devices like smartwatches^{[11]12}. This increased accessibility allows for continuous monitoring and a better understanding of HRV over extended periods and under various free-living conditions¹³. However, long-term HRV patterns and specific diurnal responses in relation to cardiovascular complications remain less well understood.

The overall aim of this dissertation is to understand how cardiovascular autonomic dysfunction/CAN affects cardiovascular disease risk (i.e. heart failure, stroke, myocardial infarction) and specific subclinical markers of CVD: carotid-femoral pulse wave velocity

and carotid artery distensibility in populations covering the whole glycemic continuum, from healthy glucose metabolism to type 2 diabetes.

3. Background

The concept of type 2 diabetes (T2D) and its associated risk with CVD. Following, I will provide an overview of various cardiovascular complications, including arteriosclerosis, atherosclerosis, and heart failure. Lastly, I describe cardiovascular autonomic function and its potential to enhance our understanding of CVD.

3.1. T2D and prediabetes

The progression from normal glucose metabolism to T2D is characterized by sustained elevations in blood glucose levels. T2D is characterized by a progressive decline in beta-cell function, most often as a consequence of chronic insulin resistance [14][15]. Insulin resistance occurs when certain tissues, such as muscle and liver tissues, lose their sensitivity to insulin. As a result, glucose is not effectively taken up by these tissues and remains in the blood circulation. Meanwhile, beta-cell function deteriorates, leading to a diminished insulin response to glucose intake. Years before diagnosis, these changes contribute to rising fasting and postprandial glucose levels¹⁴.

The body regulates glucose through various mechanisms to maintain glucose homeostasis. During fasting, pancreatic alpha cells secrete glucagon, which stimulates hepatic glucose production via glycogenolysis and gluconeogenesis. Meanwhile, glucose is endogenously produced by the liver and kidneys and utilized by body tissues. After a meal, rising blood glucose levels stimulate pancreatic beta cells to release insulin and trigger the secretion of incretins, such as glucagon-like peptide-1 (GLP-1) from the intestines. Insulin and incretins work together to suppress hepatic glucose production, while insulin promotes glucose uptake in muscle and adipose tissue. Excess glucose is primarily stored as glycogen in the liver and muscles, with some converted to triglycerides for long-term storage. Multiple organs, including the pancreas, liver, kidneys, intestines, muscle, and adipose tissue are involved in this coordinated process. The autonomic nervous system plays a supportive role in glucose homeostasis by modulating metabolic activity. Parasympathetic signals tend to reduce glucose production, while sympathetic signals enhance it, especially during hypoglycemia¹⁵.

3.1. T2D and prediabetes

Diabetes progression is a continuous process, with T2D defined based on glucose thresholds associated with an increased risk of diabetes-specific microvascular complications, particularly retinopathy. The World Health Organization (WHO)¹⁶ and American Diabetes Association (ADA)¹⁷ diagnostic criteria for T2D include fasting plasma glucose 7.0 mmol/L, 2-hour plasma glucose 11.1 mmol/L during an oral glucose tolerance test (OGTT), or hemoglobin A1c (HbA1c) 6.5% (48 mmol/mol). OGTT is defined by measuring glucose levels two hours after the ingestion of a standard 75-gram glucose load. Diabetes progression is a continuum, with T2D defined based on glucose thresholds associated with an increased risk of diabetes-specific microvascular complications, particularly retinopathy. However, many complications of diabetes, such as macrovascular disease, neuropathy, cancer, and cognitive impairment, may develop at earlier stages of dysglycemia^{[18]1920}. This stage is referred to as prediabetes or high risk of diabetes and is defined by fasting plasma glucose levels between 6.1–6.9 mmol/L, 2-hour plasma glucose levels between 7.8–11.0 mmol/L (WHO criteria), and HbA1c levels between 5.7–6.4% (39–47 mmol/mol) (ADA criteria)¹⁷. In parallel with the growing prevalence of T2D, the prevalence of prediabetes is also on the rise⁹.

Risk factors for progression to T2D and its complications range from genetic predisposition to lifestyle and socio-environmental factors. The most common precursor to diabetes is central obesity, characterized by excess body fat²¹. The accumulation of diabetes risk factors is linked with a combination of adverse changes in cardiometabolic markers, including increases in low-density lipoprotein (LDL) cholesterol, triglycerides, body mass index (BMI), and systolic blood pressure, along with decreases in high-density lipoprotein (HDL) cholesterol²².

Diabetes increases the risk of both microvascular and macrovascular complications, which are major contributors to the morbidity and mortality associated with the disease¹⁵. Diabetes and cardiovascular disease (CVD) share common risk factors, including obesity, hypertension, and hypercholesterolemia, as well as lifestyle-related factors. Beyond these, chronic hyperglycemia promotes the formation of harmful byproducts such as reactive oxygen species and advanced glycation end products, which drive oxidative stress and inflammation²³. These processes contribute to endothelial dysfunction and vascular damage²³. As individuals progress toward diabetes, their cardiovascular risk increases, making them more susceptible to developing CVD²². However, the identification of preclinical stages of CVD and how CVD risk differs among individuals at high risk of diabetes and individuals with T2D needs further definition.

3. Background

3.2. Cardiovascular disease

Globally, CVD remains the leading cause of death. At the population level, CVD risk is primarily attributable to modifiable lifestyle behaviors such as chronic stress, physical inactivity, unhealthy diet, excessive alcohol consumption, and smoking, as well as socio-environmental factors like socio-economic status and air pollution²⁴. At the individual level, these exposures often manifest through more proximal biological risk factors, including hypertension, hypercholesterolemia, diabetes, and obesity, which elevate the likelihood of developing CVD. Along the causal pathway, these intermediate conditions act as comorbidities that accelerate disease progression. These processes are underpinned by biomolecular mechanisms, including local and systemic inflammation, oxidative stress involving oxidized low-density lipoprotein (LDL), and dysregulated immune responses mediated by pro-inflammatory cytokines and signaling pathways. Risk factors across different types of CVD contribute to distinct pathophysiological mechanisms, involving structural, signaling, inflammatory, and hemodynamic changes within the cardiovascular system. Among these, cellular and molecular signaling pathways play a central role in regulating vascular tone, cardiac function, and inflammatory responses. These processes are closely modulated by the autonomic nervous system through sympathetic and parasympathetic nerve branches.

3.2.1. Arteriosclerosis

Hard CVD endpoints remain the primary focus of prevention strategies, emerging evidence emphasize the importance of vascular aging in early disease development²⁵. Arteriosclerosis, commonly referred to as arterial stiffness, is a hallmark of this process. Biologically, the medial layer of large arteries consists of a structured network of vascular smooth muscle cells together with elastic and collagen fibers, forming functional musculoelastic sheets²⁶. Arterial stiffness arises from progressive remodeling of the arterial wall [²⁵]²⁷. This remodeling is driven by changes in the structural interactions between elastin and collagen fibers, along with functional alterations in vascular smooth muscle cells and the accumulation of calcium and advanced glycation end products²⁶. Remodeling of the arterial wall increases systolic blood pressure and reduces coronary perfusion, thereby contributing to the development of hypertension and, eventually, cardiovascular disease²⁸. Additionally, they elevate the pulsatile load on the microcirculation, promoting the progression of chronic kidney disease, vascular dementia, and Alzheimer's disease²⁵.

3.2.2. Atherosclerosis

Atherosclerosis is characterized by the accumulation of cholesterol, lipids, and other substances within the arterial walls, forming plaques that narrow the arteries and reduce blood flow (ref.). This chronic process can lead to progressive occlusion of the vessel, contributing to reduced oxygen supply to the heart (ref.).

Atherosclerotic plaques can be classified into stable and unstable types, each with distinct structural characteristics and clinical implications. Stable plaques typically have a thick fibrous cap composed of collagen, a small lipid core, and low levels of inflammation. These plaques are less likely to rupture and tend to remain intact over time due to internal remodeling. In contrast, unstable plaques, also known as vulnerable plaques, often contain a large lipid-rich necrotic core, a thin fibrous cap, and infiltration by inflammatory cells such as macrophages. A well-recognized subtype of unstable plaque is the thin-cap fibroatheroma, which is particularly prone to rupture. When rupture occurs, the necrotic core becomes exposed to the bloodstream, initiating the formation of a thrombus or blood clot. This acute event can abruptly obstruct the artery, resulting in myocardial infarction. In some cases, fragments of the thrombus may dislodge and travel further along the arterial tree, leading to thromboembolic complications such as stroke or peripheral ischemia²⁹. Chronic ischemia due to reduced coronary perfusion can lead to myocardial remodeling, impaired contractility, and electrical instability, thereby increasing the risk of arrhythmias and heart failure^{[30][31]}.

Myocardial infarction

[Try to briefly mention the main advances that have powered this marked reduction in CVD events: much lower smoking prevalence, better management of hypertension and hyperlipidemia, but also thrombolysis, stents and CABGs becoming standard practice.]

Myocardial infarction (MI) occurs due to the rupture of an atherosclerotic plaque in the coronary arteries, triggering thrombus formation that blocks blood flow. This leads to oxygen deprivation (ischemia) and subsequent myocardial injury or necrosis. If untreated, this process can cause extensive cardiac damage and fatal arrhythmias. Over the past decades, the incidence of myocardial infarction (MI) has declined in high-income countries(ref.) with a marked reduction in MI-related mortality(ref.). These improvements are largely attributed to a combination of public health initiatives and medical advances. On the public health front, a substantial decrease in smoking prevalence has been the most important lifestyle-related factor contributing to the reduction in CVD [32]³³. Medically, the improved preventive management of hypertension and hyperlipidemia has reduced the burden of atherosclerotic disease. In acute care, the widespread adoption of evidence-based interventions such as thrombolytic therapy, percutaneous

3. Background

coronary interventions (including stenting), and coronary artery bypass grafting (CABG) has improved survival and outcomes following MI. In T2D, the risk of MI is elevated by 72%, with an approximately threefold risk among patients under 60 years compared to age under 60 without T2D³⁴. Similar to the general population, its incidence and fatality have declined in diabetes.

Stroke

The majority of strokes are ischemic, caused by an obstruction in a cerebral artery, often due to an atherosclerotic plaque or embolism. The second main cause is hemorrhagic stroke, which is characterized as a hypertensive small-vessel disease, leading to small lipohyalinotic aneurysms that subsequently rupture, causing intracerebral bleeding³⁵. Ischemic stroke remains one of the global leading contributor to mortality and disability³⁶. The incidence, prevalence, and cause-specific mortality of stroke remain high but have stagnated, although some declines have been observed in high-income countries³⁷. Stroke risk is already elevated at high levels of glucose (fasting plasma, OGTT, and HbA1c) among people in a pre-diabetic range where the the risk exceed of 26% higher risk compared to population without diabetes [³⁸]³⁹. In T2D, the ischemic stroke risk is elevated almost two-fold compared with individuals without diabetes³⁴.

3.2.3. Heart failure

Heart failure develops gradually with age and often accelerates with the progression of T2D. As prevention and treatment of CVD have improved survival in recent years, the prevalence of heart failure has increased, while the incidence remains stable, but may rise with aging populations⁴⁰.

Heart failure is commonly classified as either ischemic or non-ischemic in origin. It may arise as a consequence of atherosclerosis, arteriosclerosis, or both, contributing to myocardial ischemia, pressure overload, and structural cardiac changes. Heart failure is a clinical condition characterized by symptoms of breathlessness, fatigue, and fluid retention, often accompanied by clinical signs such as pulmonary crepitations, jugular venous elevation, and peripheral edema. Heart failure can be defined hemodynamically as the inability to maintain adequate cardiac output at rest or during exertion, or the ability to do so only with elevated cardiac filling pressures. It is a complex cardiovascular disease caused by structural and functional changes in the heart musculature, affecting systolic and/or diastolic pumping function. Heart failure is generally classified into two subtypes: heart failure with reduced ejection fraction (HFrEF) and heart failure with preserved ejection fraction (HFpEF). Both subtypes involve cardiac remodeling but are defined by left ventricular ejection fraction (LVEF). HFrEF is defined by an LVEF < 40%, while HFpEF is characterized by an LVEF 50% along with structural or functional

3.3. Cardiovascular autonomic dysfunction

cardiac abnormalities, as assessed by echocardiography. HFrEF is often a consequence of repeated, non-fatal myocardial infarctions. These events can leave behind scar tissue in the myocardium, impairing the heart's ability to contract effectively and leading to progressive systolic dysfunction.

The most common feature of HFpEF is left ventricular diastolic dysfunction, caused by impaired relaxation and increased stiffness, leading to elevated left atrial pressure and reduced diastolic reserve⁴¹. Over the past decades, the prevalence of HFpEF has increased with an aging population and more people living with conditions such as hypertension, diabetes, and obesity. It is diagnosed based on structural or functional abnormalities identified through echocardiographic measures, such as left ventricular hypertrophy, left atrial enlargement, or elevated filling pressures⁴². The diagnosis may seem straightforward, but it is often challenging in community settings, as patients frequently present without typical heart failure symptoms (e.g., shortness of breath) and are not routinely assessed with biomarkers like N-terminal pro-B-type natriuretic peptide (NT-proBNP) or brain-natriuretic-peptide (BNP). As a result, HFpEF is commonly underdiagnosed and consequently detected at more severe stages, leading to hospitalization⁴².

3.3. Cardiovascular autonomic dysfunction

The cardiovascular system is regulated by autonomic nervous system which influences heart rate and vasoconstriction through neurotransmitter release by the sympathetic and parasympathetic nerves. The primary neurotransmitter of the sympathetic nervous system is noradrenaline, while the parasympathetic nervous system primarily releases acetylcholine by stimulation through the Vagus nerve. Sympathetic activation increases heart rate and myocardial contractility by stimulating the sinoatrial (SA) node, atrioventricular (AV) node, and ventricular myocardium. In contrast, parasympathetic activation primarily reduces heart rate by directly modulating SA node activity through vagal stimulation. It also slows AV nodal conduction, predominantly via the left vagus nerve, thereby prolonging atrioventricular conduction time. Afferent nerves mainly carry sensory information (e.g., baroreceptor input from the carotid sinus and aortic arch) to the brain, which then adjusts efferent autonomic output to regulate arterial tone. Hence, the autonomic nervous system dynamically regulates heart rate and blood pressure to maintain homeostasis in response to physiological demands, such as rest and physical activity.

[insert figure of brain heart and sympathetic nerves]

In youth, the autonomic nervous system is highly adaptive and responsive to living conditions, maintaining autonomic balance. However, with aging, there is a gradual

3. Background

decline in parasympathetic function and an increase in sympathetic activity. Additionally, metabolic-related conditions such as obesity and diabetes have been shown to further contribute to autonomic dysfunction. Autonomic dysfunction reflects a stressed cardiometabolic environment, as both dysfunction in lipid and glucose metabolism are associated with increased sympathetic activity⁴³. This dysfunction may result from cumulative neural damage mediated by mechanisms such as hyperinsulinemia, insulin resistance, and elevated levels of adipokines. At the same time, autonomic dysfunction is known to disrupt lipid and glucose metabolism⁴³. Therefore, the relationship between autonomic dysfunction and cardiometabolic factors is likely a vicious cycle⁴⁴. The consequences can lead to cardiovascular autonomic dysfunction/neuropathy (CAN), resulting dysregulation in heart rate and vascular dynamics. In this dissertation, we will use ‘cardiovascular autonomic dysfunction’ as the broader term, while ‘CAN’ will refer specifically to autonomic dysfunction resulting from neuropathy in T2D.

Cardiovascular autonomic function can be assessed using heart rate variability (HRV) indices, which measure the variation in successive normal RR intervals in milliseconds. HRV provides time- and frequency-domain estimates of the balance between sympathetic and parasympathetic activity. High HRV reflects an autonomic nervous system with strong adaptability to the body’s demands, whereas low variation indicates poor adaptation to changing conditions. HRV changes in response to different physiological or environmental conditions (e.g., sleep, stress, posture, physical activity), and these changes can be observed in its natural 24-hour (diurnal) pattern¹³. Most studies have examined cardiovascular autonomic function using short-term ECG recordings at rest. However, extended HRV recordings across the circadian cycle may offer deeper insights into the influence of lower-frequency variability sources, such as very-low frequency (0.003–0.04 Hz) and ultra-low frequency (0.003 Hz)[reflecting what]. HRV has been applied across several research domains. For example, in psychology as a marker of mental stress, in exercise physiology as an indicator of recovery, in cardiovascular research as a marker of autonomic dysfunction due to cardiac complications, and in diabetes research as a marker of autonomic neuropathy(ref.,ref.ref.,ref.). T2D alters the expression of sympathetic bursts, as measured by resting muscle sympathetic nerve activity (MSNA). MSNA is elevated in individuals with both T2D and hypertension, compared to those who are normotensive, regardless of whether they have diabetes or not⁴⁵. Parasympathetic activity is also impaired in individuals with high cardiometabolic risk and T2D, as reflected by reduced baroreflex sensitivity⁴⁶ and lower HF and RMSSD short-term HRV. Before onset of diabetes and during progression of diabetes long-term (24-hour) HRV has shown to be lower compare to those with normal glucose metabolism [⁴⁴]⁸. Cardiovascular autonomic reflex tests (CARTs) and orthostatic hypotension are considered the gold standard for assessing CAN⁴⁷. The diagnosis includes assessing pulse rate ratio under test conditions, such as the deep breathing test, the lying-to-standing test, and the Valsalva maneuver⁴⁷. Both HRV and CARTs have shown to be associated with cardiovascular disease, heart

3.4. Risk-stratification

failure, and all-cause mortality, primarily in populations with T2D or established cardiovascular disease. However, it remains unclear at which stage in the progression of diabetes risk to pre-diabetes to diabetes these measures begin to influence the risk of cardiovascular complications.

3.4. Risk-stratification

Current cardiopreventive guidelines place strong emphasis on prevent and treat T2D. The 2022 ADA/EASD guidelines for the management of hyperglycemia in T2D recommend, cardioprotective medication (GLP-1 receptor analogues and SGLT2-inhibitors) as first-line options for individuals at high cardiovascular risk. Due to their benefits in heart failure, SGLT2 inhibitors are specifically recommended for patients with documented HFrEF or HFpEF. High cardiovascular risk is defined as the presence of at least two risk factors at age >55 years, such as obesity, hypertension, smoking, dyslipidemia, or albuminuria. However, no additional preclinical markers are recommended to identify individuals at higher CVD or HF risk. Despite their increased risk of cardiovascular complications, individuals at high risk of developing diabetes remain outside structured treatment options, even though diabetes risk and cardiometabolic markers can be successfully modified through lifestyle interventions and medication such as GLP-1 analogues [48] [49]. During the progression and following the onset of T2D, preclinical stages may be characterized by markers of elevated cardiovascular risk, highlighting the potential for early risk stratification. Risk stratification is the process of classifying or ranking individuals in increasing order of estimated risk, based on risk scores, biomarker levels, omic data (metabolomic, proteomics, and genomic) or preclinical conditions. This approach aids in identifying patients for prognostic or diagnostic purposes, identifying subgroups that require further evaluation, intensified treatment, or lifestyle modifications.

Cardiovascular autonomic dysfunction despite its relationship with cardiovascular complication has not been used in clinical practice. Larger epidemiological cohort studies encompassing various stages of diabetes risk, from normal glucose metabolism to prediabetes, onset of T2D, and longer term progression of T2D, serve as valuable resources for identifying risk-stratification opportunities. Epidemiological studies provide a broad representation of the target population, allowing understand the relationship between cardiovascular autonomic dysfunction and cardiovascular complications. They also have potential to determine when, along the trajectory of diabetes progression and duration, autonomic function are meaningful for cardiovascular risk-stratification.

3. Background




Figure 3.1.: Risk-stratification based on preclinical disease. (Source: Author)

4. Aim and hypothesis

The hypotheses of this dissertation are:

CAN and autonomic dysfunction is associated with CVD and acts as an early risk factor for heart failure and other cardiovascular complications, including stroke, and myocardial infarction in patients with prediabetes and/or T2D. In addition autonomic dysfunction is associated with higher levels of sub-clinical measures such as carotid-femoral pulse wave velocity and carotid artery distensibility.

This dissertation investigates the hypothesis by addressing the following three aims:

Study I: Quantify the cross-sectional association between 24-hour HRV and subclinical markers of cardiovascular complications: carotid-femoral pulse wave velocity and carotid artery distensibility, in participants with normal glucose metabolism, prediabetes or T2D.

Study II: Quantify the longitudinal association of week-long and hourly HRV with incidence ischemic-CVD, heart failure, and all-cause mortality in a population with high-risk of diabetes.

Study III: Quantify the cross-sectional association between CAN and heart failure. Heart failure will be defined by clinical measures i.e. N-terminal-pro-BNP (Pro-BNP), WATCH-DM risk, and New York Heart Association (NYHA) scores among individuals with T2D.

5. Materials and methods

5.1. Overview of the studies

Table 5.1.: Table 1: Overview of studies

	Study I	Study II	Study III
Title	Cardiovascular autonomic dysfunction is linked with arterial stiffness across glucose metabolism: The Maastricht Study	Cardiovascular autonomic dysfunction precedes cardiovascular disease and all-cause mortality: 11-year follow-up in the ADDITION-PRO study	Cardiovascular autonomic neuropathy and subclinical heart failure in T2D: The CANCAN study Descriptive
Design	Aetiological cross-sectional study	Aetiological prospective cohort study	cross-sectional study
Cohort	Maastricht study	ADDITION-PRO study	CANCAN study
Study	3673 people with normal glucose metabolism, prediabetes, and T2D 2082 people with high risk of diabetes 173 patients with T2D visiting outpatients clinics		
Data	Population-based cohort sources from The Maastricht Study in the Netherlands	Cohort study of selected people based on having high risk of diabetes	Clinical cohort study
Determination	24-hour HRV	multiday and hourly HRV	Cardiovascular autonomic reflex test
Primary outcome	Arterial stiffness	Major adverse cardiovascular events, heart failure, and all-cause mortality	NT-proBNP and NYHA classification?
Statistical analysis	Linear regression	Poisson regression	Logistic regression

5.1. Overview of the studies

Study I	Study II	Study III
Missing data Complete case analysis	Multiple imputation of chained equations for confounders	Complete case analysis and multiple imputation of chained equations for CART and confounders

5.1.1. Study population




Figure 5.1.: Study populations

5. Materials and methods

5.1.1.1. Study I - The Maastricht Study

The Maastricht Study is a prospective observational population-based study of the general population of the province of Limburg, in the southern part of the Netherlands. The study emphasized the recruitment of people with T2D, through the regional Diabetes Patient Registry, to extensively phenotype individuals with T2D and those in intermediate stages of the disease. The eligibility criteria included an age range of 40–70 years. Participants were recruited through mass media campaigns and mailings from municipal registries (Gemeentelijke Basis Administratie; GBA). In the analysis of Study I, the study among 7449 people included participants with measurements of 24-hour HRV and at least one measure of arterial stiffness (carotid-femoral pulse wave velocity or carotid artery distensibility), both of which were completed within a three-month period between November 2010 and December 2020. The study has been approved by the institutional medical ethics committee (NL31329.068.10) and the Minister of Health, Welfare and Sports of the Netherlands (Permit 131088-105234-PG). All participants gave written i

5.1.1.2. Study II - ADDITION-PRO

The ADDITION-PRO study is a prospective, population-based cohort nested within the Danish arm of the ADDITION-Europe study. ADDITION was originally designed as a stepwise screening program for T2D in general practice, aiming to identify individuals with screen-detected T2D for recruitment into the ADDITION trial. ADDITION-PRO aims to investigate early markers of CVD and metabolic dysfunction in individuals in different tiers of diabetes risk.

The ADDITION-Europe screening program identified a large number of individuals with impaired fasting glucose (IFG), impaired glucose tolerance (IGT), and normoglycemia despite having risk factors for diabetes and CVD. Participants for ADDITION-PRO were recruited from the original ADDITION-DK screening cohort, which included individuals from 190 general practices across Denmark. The recruitment strategy focused on individuals at high risk of diabetes without T2D, identified through a stepwise screening program that incorporated the Danish diabetes risk score from the Inter99. This assessment, conducted between 2001 and 2006, considered factors such as age, sex, history of gestational diabetes, family history of diabetes, known hypertension, BMI, and physical activity. High-risk individuals were further screened for T2D using blood measurements, including HbA1c, random blood glucose, FPG, and OGTT. Those with screen-detected diabetes, confirmed by a second OGTT, were invited to participate in the ADDITION trial. High risk individuals without T2D were further considered in as the sampling frame for ADDITION-PRO.

5.1. Overview of the studies

Between 2009 and 2011, a follow-up health examination was conducted at four ADDITION-DK study centers to establish a cohort baseline. Eligible participants were those still alive, residing near the research centers (Steno Diabetes Center Copenhagen, Aarhus University Hospital, Holstebro Hospital, and the Hospital of South West Jutland, Esbjerg), and who had not withdrawn consent. Eligibility criteria included individuals aged 40–70 years who had previously undergone diabetes screening in ADDITION-DK. Exclusion criteria included pregnancy, psychological or psychiatric disorder preventing informed consent, and life-limiting conditions. One key feature of the data collection was the precise measurement of physical activity and energy expenditure using a combined chest worn accelerometer/heart rate monitor (ActiHeart), which recorded acceleration and heart rate over a week. In study II, we included participants with a least 48-hour recording for our first analysis, and then include those participants with hourly measures of physical acceleration during the hourly HRV recording for the second analysis. We also excluded participant with prior CVD ten years before inclusion.

Disease history and follow-up data for the population were obtained from Denmark's unique national registry system, which allows linkage of health records using the personal Civil Registration Number assigned to all citizens. The following national registries were accessed to collect information on incident CVD and mortality, medication use, and healthcare utilization: the National Patient Registry (hospital admissions and outpatient contacts), the National Health Service Registry (general practice visits), the Medical Prescription Registry, the Diabetes Registry, and the Cause of Death Registry.

5.1.1.3. Study III - CANCAN

The CANCAN Study is an observational study conducted at two hospital outpatient clinics in Viborg Regional Hospital and Regional Hospital Gødstrup. It aims to implement a screening protocol for identifying high-risk individuals using CAN assessments, continuous glucose monitoring, and heart failure indicators. All measures were part of routine clinical care for T2D in Central Denmark. We included 200 adults (>18 years) with T2D with duration of over one year. Exclusion criteria were recent laser-treated eye disease (3 months), pregnancy, lactation, life-threatening illness, or cognitive impairment preventing consent. Participants were identified via electronic records and informed about the study by their doctor during a telephone call. Those interested attended a dedicated meeting before their annual diabetes exam, where study details were discussed. Recruitment took place from 2021 to 2024. In study III, participants without a valid NT-proBNP measurement were excluded.

5. Materials and methods

5.2. Study variables

5.2.1. Measures for cardiovascular autonomic dysfunction/ neuropathy



Figure 5.2.: Left: Holter monitor Middle: Actiheart Right: Handheld VagusTM device

Heart rate variability

In study I-III a device was used to capture the distance between each heartbeat defined as RR intervals from electrocardiogram traces either directly from heart-beat traces or indirectly from pulse traces. From this a sequence of successive heart beat intervals is extracted to calculate HRV. The pool of hearbeat data, we extrapolated time-domain and frequency-domain HRV indices.

Time-domain indices

Time-domain measures of HRV are based on the statistical distribution of normal-to-normal (NN) heartbeat intervals. Description of time-domain indices are summarized in ?@tbl-td.

Frequency-domain indices

Frequency-domain HRV indices are derived from sequences of NN intervals transformed into the spectral domain using Fourier transformation. These indices quantify heart rate oscillations over different timescales. Short-term variations, such as respiratory sinus arrhythmia, reflect rapid autonomic changes, while longer oscillations capture autonomic responses to posture changes, circadian rhythms, or other physiological processes. Description of frequency-domain indices are summarized in ?@tbl-fq.

Holter recordings in study I

All ECG recordings were obtained using a 12-lead Holter system (Fysiologic ECG Services, Amsterdam, the Netherlands) over 24 hours, as previously described. Participants were instructed to follow their regular daily activities but avoid showering during the

Table 5.2.: **Box 1** Time-domain indices reflections of autonomic function

Time-domain HRV	Description
Standard deviation of NN heart beat intervals (SDNN, in ms)	Measures the total variation in interbeat intervals and reflects both sympathetic and parasympathetic activity ⁶ .
SD of the averages of NN intervals in 5-minute segments throughout the recording (SDANN, in ms)	Measures variations in 5-minute mean interbeat intervals, primarily reflecting autonomic fluctuations associated with the circadian rhythm ⁶
Mean of the SDs of all NN intervals for all 5-minute segments (SDNN index, in ms)	Measures the average short-term variability in interbeat intervals across successive 5-minute periods, reflecting both sympathetic and parasympathetic modulation of heart rate ⁶
NN50 count divided by the total number of all NN intervals (pNN50, percentage)	Measures the proportion of successive interbeat intervals differing by more than 50 ms, primarily reflecting parasympathetic (vagal) activity ⁵⁰ .
Square root of the mean of the sum of squares of differences between adjacent NN intervals (RMSSD, in ms)	Measures variation in successive interbeat intervals during inhalation and exhalation, primarily reflecting parasympathetic (vagal) activity ⁵⁰

5. Materials and methods

Table 5.4.: **Box 2** Frequency-domain indices reflections of autonomic function

Frequency domain HRV	Description
Variance of all NN intervals 0.4 Hz, total power (TP, in ms²)	Measures the total variation in interbeat intervals, reflecting both short- and long-term autonomic regulation by the sympathetic and parasympathetic nervous system ⁶ .
Ultra low-frequency range (ULF, in ms² 0.003 Hz)	Measures very long-term oscillations in interbeat intervals, influenced by autonomic responses to circadian rhythms, physical activity, metabolic processes, and thermoregulation [51]52.
Very-low-frequency range (VLF, in ms²; 0.003–0.04 Hz)	Measures oscillations in interbeat intervals over 5-minute periods, reflecting the activity of the renin–angiotensin system and peaks in sympathetic nervous system activity, while also depending on parasympathetic modulation[53]54.
Low-frequency range (LF, in ms²; 0.04–0.15 Hz)	Measures intermediate oscillations in interbeat intervals, reflecting a combination of sympathetic and parasympathetic nervous system activity, particularly associated with baroreflex function and blood pressure regulation ⁵⁵ .
High-frequency range (HF, in ms²; 0.15–0.4 Hz)	Measures short-term oscillations during inspiration and expiration, reflecting parasympathetic modulation of heart rate via the vagus nerve, and closely associated with respiratory sinus arrhythmia ⁵⁶ .

5.2. Study variables




Figure 5.3.: Heart rate variability. (Source: Author)

recording. The ECG data were processed using proprietary Holter Analysis Software (Fysiologic ECG Services), where artefacts and ectopic beats were excluded through automated processing and manual validation. A minimum recording duration of 18 hours was required for further analysis. Inter-beat intervals between consecutive sinus beats were provided in milliseconds (ms). Time-domain HRV indices were calculated, including SDNN, SDANN, RMSSD, SDNN index, and pNN50. Frequency-domain measures were derived using Fast Fourier Transform, including TP, ULF, VLF, LF, and HF. Outliers were removed. HRV indices were standardised by their mean and SD, and composite Z-scores were computed for time and frequency-domain measures, respectively. This selection of indices covers the main sources of HRV variance.

ActiHeart heart rate and physical activity in study II

Heart rate was measured using a combined accelerometer and heart rate monitor (ActiHeart, CamNTech, Cambridge, UK), recording uniaxial acceleration and heart rate. The data collection and processing methods have been described previously. Mean heart rates were recorded in 30-second epochs, and HRV was derived as the variation between consecutive normal heartbeats on the ECG. HRV calculations were performed using the RHRV package (version 4.2.7) in R, including SDNN, SDANN, SDNN index, TINN, and mean HR (mHR). We tested our approach on a dataset with full access to all interbeat

5. Materials and methods

intervals to validate our algorithm⁵⁷. These indices have shown high validity for HRV indices based on global distribution (e.g. SDNN, SDANN, SDNNi) in 24-hour recordings. HRV indices were calculated by week, 24-hour cycle, and hour of the day, with hourly values averaged across recording days.

Vagus device for cardiovascular autonomic reflex test in study III

CAN was diagnosed using cardiovascular autonomic reflex tests (CARTs), the gold standard for CAN assessment. R-R intervals were derived from an ECG signal using the Vagus™ device (Medicus Engineering, Aarhus, Denmark). We used pulse rate ratios measured under different conditions. Three standardized cardiovascular autonomic reflex tests (CARTs) were performed—lying-to-standing, deep breathing, and the Valsalva manoeuvre, following a standardized protocol conducted between 8:00 a.m. and 2:00 p.m., after 10 minutes of supine rest. Smoking and caffeine intake were prohibited two hours before testing. Each test was conducted once by trained examiners.

Manifest CAN was defined as two or more abnormal CARTs using age-specific cut-off values (ref.). The Vagus™ device's accuracy has been validated against FDA standards and stationary devices, showing moderate to high reproducibility (ref.).

Cardiovascular autonomic reflex test



Figure 5.4.: CART

HRV was derived from all CARTs using autoregressive spectral analysis. Time domain measures included SDNN and RMSSD, while frequency domain measures included LF, HF, and total power. Orthostatic hypertension was defined as a sustained drop in systolic blood pressure of 20 mmHg or diastolic blood pressure of 10 mmHg within three minutes of standing (ref.).

5.2.2. Confounders and variables for instrumental bias

Across Studies I, II, and III, a comprehensive set of covariates and potential confounders were assessed, including lifestyle factors, clinical measurements, biochemical markers,

and socioeconomic indicators.

Smoking status was self-reported in all studies, categorized as never, former, or current (Study I), current/ex/never (Study II), and smoker/non-smoker (Study III). Alcohol consumption was recorded as average weekly units in all three studies. Physical activity was assessed via self-report in Studies I, II, III, with Study I capturing total and moderate-to-vigorous activity (hours/week), Study II used the Recent Physical Activity Questionnaire (RPAQ) to calculate physical activity energy expenditure (PAEE), and Study III classifying activity as sedentary or non-sedentary. In Study II also used combined accelerometry and heart rate monitoring (ActiHeart) to estimate PAEE. Study II included register-based data on socioeconomic status at baseline, including education length, income, and employment status. All studies included measurements of body mass index (BMI), waist circumference, and systolic and diastolic blood pressure, obtained during clinical examinations.

Blood samples were analyzed in all studies for HbA1c, fasting plasma glucose (FPG), triglycerides, total cholesterol, high-density lipoprotein (HDL), and low-density lipoprotein (LDL) cholesterol. Study I also included a 2-hour oral glucose tolerance test (OGTT) to classify glucose metabolism status based on FPG and OGTT (normal, prediabetes, T2D) using WHO 2006 criteria, excluding HbA1c as a diagnostic criterion. Study III additionally measured creatinine, estimated glomerular filtration rate (eGFR), and urine albumin-to-creatinine ratio.

Self-reported history of cardiovascular disease (CVD) and use of anti-hypertensive, glucose-lowering, and lipid-lowering medications were collected in all studies. In Study II, history of CVD events in the 10 years prior to baseline were retrieved from national registers. In Study III, history of CVD was collected electronic patient records.

5.3. Outcomes

5.3.1. Arterial stiffness

Arterial stiffness characterized arteriosclerosis and atherosclerosis properties of the arteries. The stiffness of different trees of the vascular musculature can assessed both locally and dynamically. Aortic and carotid stiffness were assessed as markers of arterial stiffness, following previously described procedures⁵⁸.

Pulse wave velocity

Aortic stiffness was measured by carotid-femoral pulse wave velocity (cf-PWV) using applanation tonometry (SphygmoCor, Atcor Medical, Sydney, Australia), with the median of at least three consecutive recordings included in the analysis. cf-PWV is calculated

5. Materials and methods

from the time between the ECG systole and the arrival of the pressure wave at the femoral and carotid measurement sites and the distance between these two measurement sites. It is measured with participants in a supine position following a 10-minute rest period. The aortic path length was determined using a tape measure by subtracting the carotid-to-sternal notch distance from the femoral-to-sternal notch distance⁵⁸.

Carotid artery distensibility

Carotid stiffness was assessed by the carotid artery distensibility coefficient (CD), based on ultrasound imaging of the left common carotid artery using a 7.5 MHz linear probe (MyLab 70, Esaote Europe, Maastricht, the Netherlands). CD was calculated as $\Delta D/\text{braPP}$, where ΔD represents carotid distension and braPP is brachial pulse pressure. Mean heart rate and mean arterial pressure (MAP) were recorded every five minutes using an oscillometer device (Accutorr Plus, Datascope, Montvale, NJ, USA)⁵⁸..




Figure 5.5.: Measures of arterial stiffness, measured dynamically at the aortic and local carotid sites. (Source: Author)

5.3.2. Indicators of heart failure

N-terminal prohormone of brain natriuretic peptide (NT-proBNP) is a neuretic peptide that can be used to detect patients with heart failure and the progression. It derives from B-type natriuretic peptid (BNP) which is a cardial neurohormon, that is syntezied and secreted as response to streched cardiomyocytes and cardiac volume overload. After secretion, proBNP is cleaved, releasing the active hormone BNP along with the remaining N-terminal fragment, known as NT-proBNP. In Study III, blood sample were taken at Study cite. Description of the NT-proBNP analysis of plasma samples is described in supplementary material [ref.].

We used a modified version validated The WATCH-DM heart failure risk score. The risk score is based on 9 variables: two binary (history of myocardial infarction and coronary artery bypass grafting) and seven continuous (age, BMI, systolic/diastolic BP, serum

Table 5.6.: ?(caption)

Outcome	Diagnosis codes
<i>Heart failure</i>	ICD: I50
<i>Three-point MACE</i>	
• Stroke	ICD: I61 - I64
• Myocardial infarction	ICD: I21-I24
• Cardiovascular death	ICD: I20-I28, I42, I46 SKA: KPAE10, KPAE25, KPAF10, KPAF20, KPAF21, KPAF22, Cardiovascular revasculariza- KPAH10, KPAH20, KPAH21, KPEE, KPEF, KPEH, KPEP, tion KPEQ, KPFE,, KPFH, KPFP, KPFQ

creatinine, HDL cholesterol, and HbA1c). Scores range from 0–39, categorized as very low (11), low (12–13), moderate (14–15), high (16–18), and very high (19) risk.

NYHA class stage I–IV was included. Heart failure symptoms were defined as NYHA class II–IV, assessed by a physician.

5.3.3. Cardiovascular events

Information on CVD events and mortality was obtained from the Danish National Patient Registers until 2021. ICD-10 codes for stroke, myocardial infarction, cardiovascular death, cardiovascular revascularization, and heart failure. We defined three-point major adverse cardiovascular events (MACE) as myocardial infarction, stroke, cardiovascular revascularization, and cardiovascular death.

5.4. Statistical Methods

5.4.1. Cross-sectional analysis

In Study I, we used multiple linear regression to investigate associations between multiday HRV and arterial stiffness. Model 1 adjusted for age, sex, education, glucose metabolism status, and mean arterial pressure (MAP) to account for the oversampling of individuals with T2D and potential instrumental bias of arterial pressure flow. Model

5. Materials and methods

2 included additional adjustments for smoking behavior, alcohol consumption, physical activity, body mass index, HbA1c, triglycerides, total-to-HDL cholesterol ratio, and medication use. Arterial stiffness measures were log-transformed to ensure normally distributed residuals and back-transformed into percentage change estimates. We add interaction sex to observe if the association differed between sex. We performed sensitivity analyses excluding individuals on antihypertensive treatment or glucose-lowering medication. In Study III, we applied logistic regression models to investigate the association between CAN and heart failure, using NT-proBNP as the primary outcome. We adjusted for age, sex, and diabetes duration, smoking behavior, alcohol consumption, body mass index, HbA1c, triglycerides, total cholesterol, and antihypertensive medication, eGFR and prior CVD. We performed sensitivity analyses excluded participants with beta-blocker treatment or prior CVD.

5.4.2. Time-to-event analysis

In Study II, we used Poisson regression models to quantify the associations between HRV and cardiovascular events, as follow-up data were undisturbed over time and to avoid assumptions of proportional hazards⁵⁹. Multiday HRV was modelled using splines with knots at predefined percentiles to assess non-linear associations. Hourly HRV was analysed separately for each hour to observe if the association of HRV had diurnal variation. Both HRV and mHR were standardized by their mean and standard deviation to ensure comparability. Based on assumptions about potential confounding pathways summarized in directed acyclic graphs (DAG), we fitted two models: Model 1 adjusted for age and sex, while Model 2 further adjusted for education, smoking, alcohol consumption, physical activity (physical activity energy expenditure (PAEE) calculated from Recent Physical Activity Questionnaire RPAQ), body mass index, total cholesterol, and HbA1c. Additional analyses were performed with HRV pre-adjusted for concurrent heart rate and physical acceleration to account the influence of these factors. Missing covariates were handled using multiple imputation. [Add age specific incidence rates]

5.4.3. Effect modification [det kan evt. kortes ned]

Effect modification is used to assess whether the association between an exposure and an outcome varies depending on the level of a third variable, known as the effect modifier. This means that the observed relationship between the exposure and the outcome is not uniform across all subgroups. Instead, it differs across strata defined by the effect modifier⁶⁰.

In Study I, we hypothesize that the association between 24-hour and arterial stiffness was stronger in strata of progression of diabetes (normal glucose metabolism, prediabetes,

T2D). We therefore first stratified by diabetes status to observe the size of the association across strata. We then combine all groups and include an interaction term between HRV and diabetes status. We did subsidiary analysis to check if the effect was modified by dysglycemia by stratifying HbA1c and fasting plasma glucose into deciles. In Study II, we quantified whether the association between multiday HRV and CVD endpoints varied by sex to explore potential biological dimorphism.

In Study III, we aimed to determine whether the association between CAN and elevated NT-proBNP is present in the subgroup without symptoms, defined as NYHA class < II. Hence, we hypothesized no significant effect modification between groups with and without symptoms. Similarly, we explored whether the association remains present in the group classified as low to moderate risk of heart failure, based on the WATCH-DM risk score.

A significant effect modification between the exposure and the effect modifier in all analyses was defined as an interaction term with a p-value < 0.05.

5.4.4. Multiple imputed by chained equations

Multiple Imputation by Chained Equations (MICE) is a method for handling missing data in datasets. This procedure imputes missing values through an iterative series of predictive models, generating plausible estimates while preserving the relationships within the data. To avoid one imputation for missing value could give the value the same confidence as the a non-missing value, we followed Rubins Rule. Rubin's rules in MICE combine results from multiple imputed datasets by pooling estimates of interest (e.g., means or regression coefficients) using their within- and between-imputation variances. Thus, we ensure valid statistical inferences by accounting for the uncertainty introduced by missing data.

In Study II, we imputed confounders to include as many participants and avoid excluding population with our without cardiovascular or mortality events. We imputed dataset 10 times. In Study III, we imputed missing CART, as a proportion of participants had non-valid test due to insufficient air in the valsalva manuevre, unstable heart beats or data error. These variables was used as auxiliary variables in imputation to reduce bias⁶¹. All available variables of biochemical measures, diagnosis, medication and cause of non-valid CART was used to impute each missing CART using predictive mean matching.

5.4.5. Instrumental bias

In Study I-III we are investigating the body properties by dynamic measures and biomarkers to quantify autonomic function, arterial stiffness, and cardiac function.

5. Materials and methods

Other conditions may affect the properties we are attempting to measure, and thus are causing instrumental bias.

Vascular Stiffness

In Study I, we used measurements of arterial stiffness using cf-PWV and carotid distensibility. Both measures are influenced by arterial pressure at the time of examination. Arterial pressure affects the propagation of the pressure wave through the aorta (cf-PWV) and the expansion and contraction of the carotid artery (carotid distensibility) [ref.]. To account for this, we adjusted for mean arterial pressure in our models.

Cardiovascular autonomic function

In Study II, we assessed cardiovascular autonomic function using multiday HRV recordings and hourly HRV measurements. Studies have highlighted that HRV is dependent on heart rate, and low HRV may simply reflect a higher resting heart rate (rHR). To adjust for this without overcorrecting for a collinear variable, we pre-adjusted HRV by regressing rHR on HRV, extracting the residuals, and using these as the pre-adjusted determinant. For hourly HRV, variability in heart rate may be influenced by changes in physical activity, creating a risk that HRV serves as a proxy for movement rather than autonomic function. To address this, we applied a similar pre-adjustment approach by regressing concurrent heart rate and physical acceleration to account for physical activity.

Biomarker of Heart Failure

In Study III, kidney function and overweight are known to influence NT-proBNP levels independently of heart failure. We adjusted the model to account for the blurred effect of eGFR on NT-proBNP levels in the analysis.

6. Results

In this section, I will summarize study population characteristics and findings from analysis.

6.1. Study I

6.1.1. Descriptive

In The Maastricht Study, [10,000 participated by Date], of those 1316 reported prior CVD. Participants who had valid 24-hour HRV measured was 4379 and of those 3673 had a valid measurement of either CD or cf-PWV. Study population included 3673 participants. Further characteristic are described in the study I manuscript [Table 1] [reffernce to study I].

6.1.2. 24-hour HRV and arterial stiffness

Time-domain HRV

In the fully adjusted model 2, cf-PWV was 2.8% (CI: 2.1; 3.4) lower, while CD was 3.3% (CI: 1.5; 5.1) higher per SD higher in HRV time-domain Z-score. Among the time-domain indices, SDNN, SDNNi, and SDANN showed the strongest associations, with cf-PWV being lower by 2.5% (CI: 2.0; 3.1), 2.5% (CI: 1.9; 3.4), and 2.2% (CI: 1.7; 2.7), respectively. Conversely, CD was higher by 3.2% (CI: 1.7; 4.7), 3.0 % (CI: 1.4; 4.6), and 2.8% (CI: 1.3; 4.3), respectively. RMSSD and pNN50 showed a weaker association with cf-PWV (-1.1% [CI: -1.4; -0.4], and -1.1 [-1.7; -0.6]), while no evidence for an association was found with CD.

Frequency-domain HRV

In the fully adjusted model 2, cf-PWV was 2.8% (CI: 2.1; 3.5) lower, while CD was 3.2% (CI: 1.3; 5.1) higher per SD higher in HRV frequency-domain Z-score. Among the frequency-domain indices, total power, VLF, and ULF showed the strongest associations, with cf-PWV being lower by 2.2% (CI: 1.7; 2.8), 2.4% (CI: 1.9; 4.0), and 2.1% (CI: 1.5;

6. Results

Table 6.1.: Study characteristics by diabetes status

Characteristic	**Normal glucose metabolism** N = 2,389	**Prediabetes** N = 538	**Type 2 Diabetes** N = 746
Sex			
Men	1,028 (43%)	280 (52%)	481 (64%)
Women	1,361 (57%)	258 (48%)	265 (36%)
Age (years)	58 (51, 64)	62 (57, 68)	63 (57, 68)
Total physical activity (hours/week)	13 (9, 19)	13 (9, 19)	12 (7, 17)
Moderate to vigorous physical activity (hours/week)	5.3 (3.0, 8.3)	4.5 (2.3, 7.5)	3.8 (1.5, 6.8)
BMI (kg/m^2)	25.0 (22.9, 27.4)	27.2 (24.9, 30.1)	28.8 (26.0, 31.7)
Waist (cm)	89 (81, 97)	98 (90, 105)	103 (96, 112)
HbA1c (%)	5.35 (5.17, 5.63)	5.63 (5.35, 5.90)	6.54 (6.08, 7.09)
Fasting plasma glucose (mmol/L)	5.10 (4.80, 5.40)	5.90 (5.40, 6.30)	7.40 (6.60, 8.50)
LDL (mmol/L)	3.20 (2.70, 3.90)	3.30 (2.60, 4.00)	2.40 (1.80, 3.10)
HDL (mmol/L)	1.60 (1.30, 2.00)	1.40 (1.20, 1.80)	1.30 (1.00, 1.50)
Total cholesterol (mmol/L)	5.50 (4.80, 6.20)	5.50 (4.80, 6.30)	4.50 (3.90, 5.20)
Triglycerides (mmol/L)	1.05 (0.80, 1.45)	1.39 (1.03, 1.90)	1.51 (1.08, 2.14)
Duration of type-2 diabetes (only for diagnosed participants)	NA (NA, NA)	NA (NA, NA)	3 (0, 9)
Mean IBI (ms)	838 (775, 907)	815 (760, 897)	806 (744, 889)
SDNN (ms)	138 (117, 164)	127 (106, 152)	116 (96, 139)
RMSSD (ms)	26 (21, 34)	24 (19, 33)	22 (17, 31)
SDANN (ms)	125 (103, 149)	113 (92, 139)	103 (84, 127)
SDNNi (ms)	55 (46, 65)	50 (41, 60)	44 (36, 54)
bNN50 (%)	7 (3, 13)	5 (2, 10)	4 (2, 9)
TP (ms^2)	12,596 (8,880, 17,498)	10,615 (7,134, 15,374)	8,880 (6,064, 12,722)
ULF (ms^2)	10,771 (7,392, 15,142)	8,948 (5,852, 13,374)	7,524 (5,036, 11,001)
VLF (ms^2)	1,198 (833, 1,692)	1,015 (685, 1,478)	816 (541, 1,267)
LF (ms^2)	421 (257, 651)	328 (200, 540)	261 (154, 422)
HF (ms^2)	94 (57, 158)	78 (47, 138)	63 (36, 117)
Systolic blood pressure (mmHg)	123 (114, 133)	129 (122, 140)	130 (122, 139)
Diastolic blood pressure (mmHg)	75 (71, 80)	78 (73, 83)	76 (72, 81)
Mean arterial pressure (mmHg)	95 (88, 102)	99 (93, 107)	98 (92, 105)
Carotid artery distensibility (10-3/kPa)	15.0 (11.8, 18.8)	13.5 (10.4, 16.9)	12.5 (9.9, 16.0)
Carotid-femoral pulse wave velocity (m/s)	8.08 (7.28, 9.16)	8.96 (7.84, 10.32)	9.36 (8.16, 10.80)
N_HT	833 (35%)	317 (59%)	590 (79%)
Antihypertensive medication	431 (18%)	199 (37%)	478 (64%)
med_HT_beta	149 (6.2%)	77 (14%)	195 (26%)
Lipid-lowering medication	280 (12%)	141 (26%)	484 (65%)

2.6), respectively. Conversely, CD was higher by 2.7% (CI: 1.2; 4.2), 2.4% (CI: 0.9; 4.1), and 2.6% (CI: 1.1; 4.1), respectively. HF showed a weaker association with cf-PWV (-0.9% [CI: -1.4; -0.4]), while no evidence for an association was found with CD. Mean interbeat interval was associated with 2.4 % (CI: 1.8; 2.9) lower cf-PWV and 4.5% (3.1; 6.1) higher CD.

6.1.3. Effect modification of diabetes status

The study population represented diabetes risk of normal glucose metabolism (65%), prediabetes (15%), and T2D (20%). The median (IQR) cf-PWV (aortic stiffness) became higher with diabetes status: NGM: 8.08 m/s (7.28, 9.16), prediabetes: 8.96 m/s (7.84, 10.32), and T2D: 9.36 m/s (8.16, 10.80). CD (carotid stiffness) decreased: NGM: 15.0 (11.8, 18.8), prediabetes: 13.5 (10.4, 16.9), and T2D: 12.5 (9.9, 16.0) $\times 10^3$ /kPa. SDNN (ms) was highest in NGM and decreased with worsening glucose metabolism: NGM: 138ms (117, 164), prediabetes: 127ms (106, 152), and T2D: 116ms (96, 139). Further description of charaterisitcs by diabetes are described in Table 6.1.

The association between HRV time-domain Z-scores and cf-PWV and CD was significantly modified by prediabetes (cf-PWV: -4.9 [CI: -6.523; -3.243] $interaction^{(*)p-value<0.01}$ CD: 8.0 [CI:3.8; 12.5] $^{*p-value<0.01}$) but not by T2D (cf-PWV: -3.5 % [CI: -4.8; -2.1] $^{*p-value<0.1}$ CD: 4.8 % [CI:1.3; 8.4] $^{*p-value<0.1}$). For the indices SDNN and SDANN, the association with both cf-PWV and CD was significantly modified by both prediabetes and T2D.

The association between HRV frequency-domain Z-score and cf-PWV was significantly modified from normal glucose metabolism by prediabetes (-5.7 %[CI:-7.4; -3.9] $^{*p-value<0.01}$) and T2D (-3.9 %[CI:-5.4; -2.3] $^{*p-value<0.05}$) while CD was only modified by prediabetes (8.3 %[CI:3.6; 13.2] $^{*p-value<0.01}$) but not by T2D (5.3 %[CI:1.4; 9.4] $^{*p-value<0.1}$). For the indices total power and ULF, the association with both cf-PWV and CD was significantly modified by both prediabetes and T2D. Mean inter beat interval association with cf-PWV or CD was not significantly modified by diabetes status.

As I did not observe a stepwise increase in the modification of glucose metabolism status from prediabetes to T2D, I excluded the subgroup with T2D to test whether the association was gradually modified by dysglycemia. In this subgroup, the association between HRV time and frequency domain Z-scores and measures of arterial stiffness was modified by HbA1c (range of interaction p-values: 0.1 to 0.005) (see Figure Figure 6.1). For example, per SD lower HRV frequency domain Z-score at HbA1c 6.4% was associated with a 5.4% higher (CI: 3.5; 7.2) cf-PWV, which was 2.0% to 4.0% higher compared to the magnitude of association at HbA1c levels of 5.6% and 4.8% (see see Figure 6.1 B). In CD, per SD lower HRV frequency domain Z-score at HbA1c 6.4% was associated

6. Results

with an 8.1% lower (CI: -13.5; -2.9) CD, which was 4.8% to 9.5% lower compared to the magnitude of association at HbA1c levels of 5.6% and 4.8% (see Figure 6.1 D). No association between HRV frequency domain Z-score and CD was observed at HbA1c levels between 4.8% and 5.2%.




Figure 6.1.: ?(caption)

6.2. Study II

6.2.1. Descriptive

The ADDITION-PRO population consisted of 1,627 participant with a least 48-hour HRV measures, while 1,432 had all hour represented with hourly HRV and physical ac-

Table 6.2.: Study participants characteristics

Characteristic	Overall, N = 1,625	<100, N = 148	100-120, N = 312	120-140, N = 457	140-160, N = 346	>160, N = 362
sex						
Men	866 (53%)	68 (46%)	148 (47%)	206 (45%)	203 (59%)	241 (67%)
Women	759 (47%)	80 (54%)	164 (53%)	251 (55%)	143 (41%)	121 (33%)
Age (years)	65.9 (6.8)	67.4 (6.9)	65.7 (6.9)	66.0 (6.7)	65.5 (6.6)	66.0 (7.0)
Physical activity energy expenditure (kJ / day)	53.1 (25.1)	46.8 (24.0)	49.4 (21.0)	50.7 (21.5)	57.6 (27.2)	57.5 (29.2)
Alcohol consumption (units per week)	9.2 (9.5)	11.3 (10.8)	10.2 (11.3)	8.9 (8.5)	8.5 (9.2)	8.7 (8.2)
Smoking status						
1	263 (16%)	40 (28%)	70 (23%)	65 (14%)	41 (12%)	47 (13%)
2	750 (47%)	58 (40%)	145 (47%)	214 (47%)	162 (47%)	171 (48%)
3	598 (37%)	47 (32%)	95 (31%)	174 (38%)	140 (41%)	142 (39%)
BMI (kg/m ²)	27.7 (4.7)	28.1 (5.4)	28.2 (4.6)	28.0 (4.7)	27.7 (4.9)	26.9 (4.2)
Waist circumference (cm)	96.7 (13.4)	98.0 (14.9)	98.2 (13.2)	96.7 (13.6)	96.7 (13.1)	94.8 (12.5)
Systolic blood pressure (mmHg)	133.7 (17.3)	134.2 (16.3)	133.7 (17.6)	133.5 (17.8)	133.4 (16.9)	133.8 (17.5)
Diastolic blood pressure (mmHg)	81.9 (10.4)	83.8 (10.1)	82.7 (10.2)	81.7 (10.6)	82.1 (10.2)	80.6 (10.3)
Pulse rate (bpm)	67.4 (10.9)	77.7 (11.2)	72.6 (9.3)	67.9 (9.3)	65.3 (9.3)	60.0 (9.8)
HbA1c (%)	5.8 (0.5)	5.9 (0.9)	5.9 (0.6)	5.8 (0.5)	5.7 (0.4)	5.7 (0.4)
Triglycerides (mmol/L)	1.3 (0.7)	1.5 (0.9)	1.4 (0.7)	1.3 (0.6)	1.2 (0.7)	1.1 (0.6)
Total cholesterol (mmol/L)	5.4 (1.1)	5.2 (1.0)	5.4 (1.2)	5.4 (1.1)	5.4 (1.0)	5.4 (1.0)
HDL cholesterol (mmol/L)	1.6 (0.4)	1.6 (0.4)	1.5 (0.5)	1.6 (0.4)	1.6 (0.4)	1.6 (0.4)
LDL cholesterol (mmol/L)	3.2 (1.0)	3.0 (1.0)	3.2 (1.1)	3.2 (1.0)	3.3 (0.9)	3.3 (0.9)
Urine albumin-creatinine ratio (mg/g)	25.9 (132.8)	36.4 (105.9)	47.9 (275.1)	19.6 (48.2)	19.4 (67.7)	16.4 (36.3)
vo2max	26.6 (7.8)	24.8 (7.5)	24.8 (7.5)	26.1 (6.8)	27.0 (8.0)	28.7 (8.7)
rest_hr	57.3 (7.3)	67.8 (5.7)	63.3 (5.0)	58.4 (4.5)	55.0 (4.2)	49.8 (4.9)
med_any_antihypertensive	753 (47%)	88 (61%)	149 (48%)	216 (47%)	147 (43%)	153 (43%)

celeration. The study population included different tiers of diabetes risk: 154 individuals at low risk (9%), 889 at high risk (51%), 314 with impaired fasting glucose (IFG) (18%), 226 with impaired glucose tolerance (IGT) (13%), and 161 with both IFG and IGT (9%). I splitted SDNN into categories by very-low (SDNN< 100 ms), low (SDNN 100-120 ms), middle (SDNN 121-140 ms), high (SDNN 141-160 ms) and very-high (SDNN >160 ms).

Charateristics are desribed in Table 6.2. Participants in the lowest SDNN group (<100 ms) were older (67.4 ± 6.9 years), had higher BMI (28.1 ± 5.4), HbA1c (5.9 ± 0.9), triglycerides (1.5 ± 0.9 mmol/L), and resting heart rate (67.8 ± 5.7 bpm), were more likely to use anti-hypertensive medication (61%), and had lower physical activity energy expenditure (46.8 ± 24.0 kJ/day) compared to those with higher SDNN levels.

6.2.2. Multiday HRV and MACE, heart failure, and all-cause mortality.

The mean multiday SDNN was 139.0 (32.3) ms, and the mean heart rate was 73.5 (9.1) bpm. In the fully adjusted model, SDNN per SD was associated with a lower incidence rate ratio (IRR) for MACE 0.82 (CI: 0.69; 0.97), heart failure 0.76 (CI: 0.58; 0.99), and mortality rate ratio of 0.79 (CI: 0.66; 0.94). When I pre-adjusted for resting heart rate, the proportion of the association explained between HRV and MACE, HF, and all-cause mortality was 14%, 25%, and 19%, respectively. I included knots in the model, which showed that the risk became higher when SDNN fell below 120 to 110 ms (approximately below the 20th percentile), suggesting a potential cut-off point for higher

6. Results

risk. I therefore calculated the incidence rate (IR) at SDNN levels of 100 ms, 120 ms, and 160 ms, respectively, and plotted these as a function of age.

At age 65, the IR per 1000 person-years for MACE was 44.2 (CI: 33.5; 58.3) at SDNN = 100 ms, which was higher than the rates observed at SDNN = 120 ms (IR: 37.6 [CI: 29.2; 48.3]) and SDNN = 160 ms (IR: 34.7 [CI: 27.0; 44.5]) (Figure 6.2 A). The IR became higher with age, reaching its peak at age 72. For heart failure at age 65, the IR was 20.5 (CI: 15.3; 27.5) at SDNN = 100 ms, slightly higher than at SDNN = 120 ms (IR: 19.6 [CI: 14.8; 25.9]) and SDNN = 160 ms (IR: 19.8 [CI: 15.0; 26.2]) (Figure 6.2 B). The IR remained stable until age 70, after which it became higher. For all-cause mortality at age 65, the IR was 11.6 (CI: 6.3; 21.4) at SDNN = 100 ms, higher than at SDNN = 120 ms (IR: 7.6 [CI: 4.3; 13.3]) and SDNN = 160 ms (IR: 6.0 [CI: 3.4; 10.4]) (Figure 6.2 C). The IR for all-cause mortality became higher with age.

6.2.3. Hourly HRV and MACE, heart failure, and all-cause mortality.

From the hourly recordings, I observed a clear periodicity in SDNN, heart rate, sleep patterns, and physical acceleration. Mean (SD) SDNN increased from 5–6 AM (70.2 [28.8] ms), peaking at 8–9 AM (92.1 [29.0] ms), followed by a gradual decline, reaching its lowest point around 2 AM the next day (64.1 [28.1] ms). A similar circadian pattern was observed in heart rate, although its peak occurred two hours later, starting at 9 AM (76.7 [10.9] bpm). After peaking, heart rate remained stable throughout the afternoon before gradually decreasing.

In Figure 6.3, I observe hourly SDNN (preadjusted for heart rate and physical acceleration), heart rate, and physical acceleration association. Models was adjusted for age, sex, education, alcohol consumption, smoking behavior, BMI, total cholesterol, and HbA1c. The morning response of SDNN was most indicative of MACE, with the strongest association observed from 6–7 AM (IRR: 0.84; 95% CI: 0.71 to 1.00 per SD higher SDNN) (see Figure 6.3 A). Heart rate between 12 AM and 6 AM showed a small trend toward higher risk of MACE (IRR range: 1.11 to 1.15 per SD higher heart rate), although none of the confidence intervals exceeded one (see Figure 6.3 D). Across all hours, there was a plausible association between SDNN and heart failure. However, this association disappeared after adjusting for physical acceleration and heart rate (see Figure 6.3 B). In contrast, heart rate between 10 PM and 9 AM was associated with heart failure (IRR range: 1.37 to 1.58 per SD higher heart rate) (see Figure 6.3 E). SDNN was consistently associated with all-cause mortality across all hours, with a stronger inverse association observed between 12 PM and 1 AM (IRR range: 0.79 to 0.88 per SD higher SDNN) (see Figure 6.3 C). No clear trends of association were observed between heart rate and all-cause mortality.




Figure 6.2.: Multiday SDNN levels (100 ms, 120 ms, 160 ms) by age-specific incidence rates for A) major adverse cardiovascular events, B) heart failure, and C) all-cause mortality

6. Results




Figure 6.3.: Hourly SDNN levels (100 ms, 120 ms, 160 ms) by age-specific incidence rates for A) major adverse cardiovascular events, B) heart failure, and C) all-cause mortality. Figure adapted from [authors]. —. (Paper II, appendix)

6.3. Study III

6.3.1. Descriptive

In study III, 179 participants with type 2 had measures of NT-proBNP and performed the CART test. CAN was present in 30% ($n = 54$) of participants (36% among those with valid CAN measurements (Figure 6.4 A)). Meanwhile, 24% ($n = 43$) were unable to complete the CART assessment adequately, primarily due to irregular heart rhythms ($n = 8$) or insufficient air pressure during the Valsalva manoeuvre ($n = 21$). Compare to those without CAN, the participants with CAN were more women (41 % vs 33 %), were more sedentary (45% vs 36%), had a higher proportion with prior major CVD (41% vs 20%) and declined eGFR (< 60) (36% vs 22%), higher levels of triglyceride (median 2.05 mmol/L vs 1.95 mmol/ L), were slightly older (median 62 years vs 61 years), had longer duration of T2D (median 19 years vs 15 years), and higher use SGTL2-inhibitors (65% vs 60%) but lower use of GLP-1 RA (63% vs 70%). No other difference in clinical characteristic was observed.

6.3.2. CAN and indicators of heart failure

A greater proportion of individuals with CAN exhibited elevated NT-proBNP levels (>125 pg/ml) (51.9%, $n=52/78$) compared to those without CAN (23.2%, $n=26/112$) (Figure 6.4 E). The fully adjusted odds ratio (OR) for elevated NT-proBNP in individuals with CAN was 5.69 (95% CI: 1.95; 18.49) relative to those without CAN. Among the cardiovascular autonomic reflex tests (CART), the Valsalva maneuver demonstrated the strongest association with NT-proBNP (OR 9.00, 95% CI: 2.88; 33.09; $n=51/75$), followed by deep breathing (OR 3.30, 95% CI: 1.17; 9.77; $n=33/133$) and orthostatic hypertension (OR 4.04, 95% CI: 1.27; 13.77; $n=24/146$). No significant association was identified for the lying-to-standing test (OR 0.80, 95% CI: 0.32; 1.97; $n=54/108$). After imputing missing CART data, the OR for CAN in relation to elevated NT-proBNP declined to 2.94 (95% CI: 1.37; 6.56). Sensitivity analyses, which excluded participants using beta-blockers or those with a history of CVD, resulted in a smaller sample size and wider confidence intervals, though the overall association remained unchanged. CAN was associated with elevated NT-proBNP in individuals both without (NYHA I; OR = 4.3, 95% CI: 1.1; 16.3) and with heart failure symptoms (NYHA II; OR = 16.4, 95% CI: 1.2; 222.0), though the interaction was not significant ($p = 0.4$). Similar associations were seen across WATCH-DM risk groups: very-low-to-moderate (OR = 6.1, 95% CI: 1.6; 23.5) and high-to-very-high (OR = 6.3, 95% CI: 0.83; 46.9). Participants with CAN had 1.7 (95% CI: 0.3 to 3.0) point higher WATCH-DM risk score compared to those without CAN. The OR of presenting with NYHA class II or higher was 5.51 (95% CI: 1.9 to 15.97) in the group with CAN.

6. Results




Figure 6.4.: Relationship between CAN and indicators for heart failure. Figure from [authors]. Cardiovascular autonomic neuropathy and indices of heart failure in T2D: The CANCAN Study. (Paper III, appendix)

7. Discussion

The aim of this dissertation is to understand how cardiovascular autonomic dysfunction and CAN affect the risk of CVD across stages of glucose metabolism. Given the rising prevalence of prediabetes and T2D, and their association with increased risks of CVD and heart failure, there is a pressing need for earlier indicators to help healthcare providers intervene in a timely manner and prevent progression to more advanced stages of cardiovascular complications. One promising approach involves leveraging data from wearable devices and standardized screening tools. Heart rate dynamics and variability across different circumstances may hold promise as accessible indicators for early cardiovascular risk stratification.

This chapter presents a summary of the main findings from this dissertation, interpreted in the context of existing evidence in the field, and discusses their clinical relevance across different levels of healthcare. Moreover, the strengths and limitations of the methods and results will be discussed.

7.1. Summary of findings

In this dissertation, autonomic dysfunction, defined by long-term HRV and standardized CARTs, and its relationship with cardiovascular complications were studied across three different cohorts representing populations at varying levels of prevention and care, including public health, primary care, and secondary care. In the Maastricht study (Study I), I investigated autonomic dysfunction, measured by 24-hour HRV, and arterial stiffness, assessed dynamically at the aortic site and locally at the carotid site among individuals with normal glucose metabolism, prediabetes, and T2D. Lower HRV was associated with higher aortic and carotid stiffness. This association was evident regardless of glucose metabolism status, and was more pronounced in individuals with prediabetes or T2D. While no significant difference was observed between prediabetes and T2D, the association was modified by HbA1c after excluding individuals with T2D, supporting the modifying effect of dysglycemia. Z-scores of time- and frequency-domain measures showed the strongest associations, primarily driven by HRV indices reflecting total variation in interbeat intervals (SDNN, SDANN, SDNN index, ULF, VLF, TP).

Following Study I, I focused on individuals at higher risk of developing diabetes, using data from the ADDITION-PRO (Study II) cohort of individuals representing high-risk individuals of diabetes. In study II, per lower SD in SDNN, measured over a multiple days, was associated with 18%, 24%, and 21% higher risk for ischemic-related CVD, hospitalization of heart failure, and all-cause mortality, respectively. The risk became higher at SDNN levels below 120 ms, supported by a greater difference in incidence rates between individuals with 100 ms and 120 ms than the difference observed between individuals with 120 ms and 160 ms. Hourly measures suggested a specific time point related to ischemic-related CVD, as lower SDNN recorded between 6:00 and 7:00 AM was associated with MACE. Residual adjustment of concurrent heart rate and physical movement did not explain the observed association. Hourly SDNN was associated with all-cause MRR, although no specific time point showed an exceptionally strong association. While no association between hourly SDNN and heart failure was observed, higher heart rate during the night hours from 02:00 to 06:00 AM was linked to an increased risk of heart failure hospitalization.

Our findings suggest that both long-term HRV measures and hourly HRV responses may serve as indicators of CVD risk. However, a key limitation is that long-term HRV was assessed under free-living conditions, which restricts the ability to perform standardized tests of autonomic function. In the CANCAN study (study III), I used standardized CARTs to define CAN and describe indicators of heart failure, defined by elevated NT-proBNP, WATCH-DM risk, and NYHA classification among those with and without CAN in a population with T2D. In CANCAN, two out five had CAN. Compared to individuals without CAN, these individual more often showed signs of heart failure, in-

cluding elevated NT-proBNP levels, higher WATCH-DM risk score, and higher classifications on the NYHA. CAN was association elevated NT-proBNP levels and the persisted even among individuals without heart failure symptoms based on NYHA classification, as well as those categorized as having low to moderate heart failure risk according to the WATCH-DM score.

In summary, various aspects of autonomic dysfunction and cardiovascular complications were investigated in populations with normal glucose metabolism, prediabetes, and T2D. The overall findings showed that autonomic function, assessed through heart rate dynamics of long-term HRV and diurnal HRV and heart rate responses, is associated with an increased risk of CVD and heart failure. This relationship appears to be stronger in more severe stages of dysglycemia. Moreover, among individuals with T2D, the presence of CAN may help identify those at higher risk of heart failure, even in the absence of heart failure symptoms.

7.2. Cardiovascular autonomic dysfunction and its impact on heart disease across glucose metabolism

This dissertation show that cardiovascular autonomic dysfunction, measured by HRV and CARTs, is associated with CVD risk across the spectrum of glucose metabolism dysregulation. This association is evident with measures of arteriosclerosis, atherosclerotic events, all-cause mortality, and heart failure in individuals at high risk of diabetes, as well as with indications of heart failure in patients with T2D.

7.2.0.1. Arteriosclerosis

In Study I, autonomic dysfunction, measured by 24-hour HRV, showed to be associated with arterial stiffness, assessed both dynamically (cf-PWV) and locally (CD). This suggests that autonomic responses under free-living conditions contribute to the development of arterial stiffness. Majority of studies have shown an association between autonomic dysfunction, as measured by short-term HRV during rest, and arterial stiffness in populations with either type 1 or T2D⁶². Study I extended this perspective by long-term HRV and including a population without diabetes or prediabetes.

Arterial stiffness is not only a structural marker of vascular ageing but is also dynamically modulated by local endothelial signals and autonomic nervous system activity. Several studies have demonstrated a link between elevated sympathetic tone and increased arterial stiffness^{[63][64]}. Two possible mechanisms may explain how autonomic dysfunction is related to arterial stiffness. First, autonomic dysfunction may increase the vascular

7. Discussion




Figure 7.1.: Autonomic dysfunction and arterial stiffness. (Source: Author). —.

tone of large arteries, thereby impairing arterial elasticity. This concept is supported by animal studies. In rats, proper autonomic regulation has been shown to be essential for maintaining aortic elasticity. Conversely, chronic overstimulation of sympathetic activity can lead to structural remodeling and increased arterial stiffness. Therefore, although the initial effects of autonomic dysfunction are dynamic and amenable to change by intervention, they may become progressively less reversible over time. While such findings cannot be directly extrapolated to humans, they suggest plausible biological pathways. Second, the autonomic nervous system regulates heart rate and cardiac contractility. Autonomic dysfunction typically manifests as both reduced HRV and elevated resting heart rate. Arterial shear stress increases as a result of heightened sympathetic activity and parasympathetic withdrawal. A higher resting heart rate may contribute to structurally stiffer arteries by altering blood flow dynamics and increasing shear stress. Our earlier study using data from the Whitehall II cohort showed that a steeper decrease in short-term (5-minute) HRV over a ten-year period was linked with higher levels of aortic stiffness in the subsequent five years⁶⁵.

The association between 24-hour HRV and arterial stiffness was modified by dysglycemia, suggesting dysglycemia introduce CAN that can affect arterial stiffness, even before the onset of T2D^[66]^[67]^[68]. Data from the Whitehall II study showed that aortic stiffness increased more steeply with higher HbA1c values among non-diabetic individuals⁶⁹. In the subpopulation in Study I without diabetes, a modification by HbA1c in both aortic and carotid stiffness was observed. The modifying effect of HbA1c suggests that hyperglycaemia amplifies the consequences of autonomic dysfunction.

7.2.0.2. Atherosclerosis

In Study II, I showed that individuals with a preclinical stage of autonomic dysfunction, measured by multiday HRV, face a higher risk of incident ischemic CVD, heart failure, and all-cause mortality.

We assessed week-long HRV to capture autonomic activity in real-life settings across several days. Our results are consistent with earlier research linking reduced HRV to cardiovascular outcomes and mortality⁵. Our findings build on existing evidence by (1) focusing on a population at elevated risk of diabetes, (2) utilizing multiday HRV recordings, and (3) identifying specific daily periods where HRV patterns were indicative of ischemic-related CVD risk. By using both week-long and hourly data, we identified specific periods of higher risk. A strength of using multiday HRV recordings, is to provide more robust insights into individual autonomic patterns by averaging cardiovascular responses across typical daily conditions. This reduces the influence of random fluctuations caused by factors such as physical activity, emotional states, or sleep on any single day⁷⁰.

Multiple mechanisms may explain how autonomic dysfunction contributes to the initiation and progression of ischemic events and stroke. First, as discussed in Study I, autonomic dysfunction may promote arteriosclerosis, leading to arterial stiffness through a dynamic and potentially modifiable process. Arterial stiffness impairs vasodilation, increasing hemodynamic stress and the risk of plaque rupture and thrombus formation [71]⁷². In this context, findings from Study I may not entirely distinguish between arterial stiffness and atherosclerosis, as shown by data from the Rotterdam Study⁷³. As plaques develop, the associated increase in sympathetic nerve density around the arteries could transiently reduce vascular tone which over time reduce arterial elasticity⁷⁴. In a smaller study of people with T2D, lower HRV was linked with increased carotid atherosclerosis⁷⁵.

Second, the autonomic nervous system innervates the adventitia layer of blood vessels, where it modulates vascular tone via sympathetic fibres. Although atherosclerotic plaques form in the intima layer, recent *in vivo* studies have demonstrated that increased plaque burden is associated with higher local sympathetic nerve density, likely mediated by neuroinflammatory processes. Notably, reducing sympathetic innervation has been shown to attenuate plaque formation in animal models⁷⁶. These findings suggest that autonomic dysfunction may not only reflect but also actively contribute to atherogenesis.

Third, autonomic nervous dysfunction has been shown to interfere with signalling pathways controlling heart rhythm, potentially leading to arrhythmias that disturb cardiac contraction. Data from the Atherosclerosis Risk in Communities study illustrated that

7. Discussion

lower short-term HRV was associated with incident atrial fibrillation over 20 years, with a higher risk among participants with T2D⁷⁷. This supports the role of autonomic dysfunction in arrhythmogenesis, which increases the risk of myocardial infarction and stroke. However, in Study II, I did not include incident atrial fibrillation as an outcome. We did not include atrial fibrillation as an outcome in Study II due to limitations in Danish registries, which often do not distinguish between short- and long-term AF, affecting diagnostic validity. Future research is needed to explore whether it could explain the higher risk of major adverse cardiovascular events.

A study of individuals with coronary artery disease showed that stress-induced HRV was associated with myocardial infarction, even more than resting HRV, suggesting that reduced parasympathetic modulation of heart rate under stress may play a role in ischemia⁷⁸. Our study focused on long-term HRV under free-living conditions, capturing stress-responsive periods such as morning awakening. These recordings likely reflect underlying autonomic dynamics relevant to cardiovascular risk. A Genome-Wide Association Study (GWAS) in the UK biobank of short-term HRV support this by identifying mechanisms involving G-protein signaling, pacemaker activity, and mitochondrial function. These pathways influence vagal control, cardiac excitability, and energy metabolism⁷⁹. Although derived from short-term recordings, the genetic associations may reflect autonomic traits that persist across different time scales, thereby reinforcing the biological basis of HRV. A Mendelian randomization study using data from the Rotterdam Study found that genetically predicted HRV was associated with an increased risk of atrial fibrillation⁸⁰. However, this association did not extend to all-cause mortality or cardiovascular death in the UK Biobank cohort, where only phenotypically measured HRV showed a significant relationship with these outcomes⁸¹. Interestingly, the genetic determinants of HRV exhibited pleiotropic relationships with several autonomic traits, including resting heart rate, heart rate response during exercise, and post-exercise recovery dynamics⁸¹. No GWAS has been conducted for long-term HRV. Therefore, it is unclear whether the genetic influences identified for short-term HRV are applicable to long-term HRV. Future GWAS efforts targeting long-term HRV could help establish causal relationships by leveraging methods such as Mendelian randomization, thereby advancing our understanding of the genetic architecture underlying autonomic regulation.

7.2.0.3. Heart failure

The relationship between cardiovascular autonomic dysfunction and heart failure is complex⁸². On one hand, autonomic dysfunction contributes to cardiac remodelling and eventual heart failure. On the other hand, it may reflect compensatory mechanisms of the progression of cardiac remodelling and declining cardiac output. Our find-

7.2. Cardiovascular autonomic dysfunction and its impact on heart disease across glucose metabolism

ings demonstrated a relationship between autonomic dysfunction and heart failure both cross-sectionally in a population with T2D and prospectively in individuals representing different tiers of diabetes risk. However, our data are limited in determining the extent to which this relationship supports one explanation over the other, as we lack baseline and follow-up measures of both heart failure and HRV. Earlier studies have demonstrated that both short-term and long-term HRV are associated with incident heart failure in populations with and without T2D.^{[83][84][85][86]} Beyond examining a population at elevated risk of diabetes using multiday HRV recordings, we expanded previous research by (1) unmasking the role of resting heart rate in the relationship between HRV and heart failure risk, and (2) identifying specific times of day when heart rate patterns signaled higher risk of heart failure.

Several mechanisms may underlie the role of autonomic dysfunction in advancing heart failure. Findings from Study I confirmed the relationship between autonomic dysfunction and arterial stiffness. It is well established that arterial stiffness is linked to cardiac remodelling, as increased cf-PWV leads to an earlier return of the reflected pulse wave to the aorta, which increases cardiac afterload and reduces coronary perfusion pressure^[87]. Therefore, autonomic dysfunction may have an indirect effect on heart failure, potentially mediated by arterial stiffness. However, structured analyses are needed to confirm these pathways. For example, through mediation analysis to assess the direct and indirect effects of arterial stiffness, with autonomic dysfunction as the main determinant. In Study II, I observed that multiday HRV was associated with incident heart failure, and approximately one-fourth of the risk was explained by resting heart rate. Data from the Rotterdam Study showed that short-term HRV was longitudinally associated with echocardiographic measures reflecting systolic function, suggesting that autonomic dysfunction contributes to cardiac remodelling^[88]. In contrast to MACE outcomes, findings from Study II showed no specific time point in hourly HRV that was associated with heart failure. Instead, it was the overall daily pattern captured by multiday HRV that was linked to heart failure risk. This suggests that the association is not driven by isolated shifts in autonomic activity, but rather by a consistently impaired autonomic balance under free-living conditions. The effect appears to be driven in part by a failure to show appropriate decreases in heart rate during rest, as individuals with higher hourly heart rates at night had an increased risk of heart failure. In Study III, I observed that individuals with CAN had higher risk of elevated levels of NT-proBNP, a biomarker of myocardial stress and early heart failure. Therefore, CAN is associated with hemodynamic consequences that contribute to both structural and functional cardiac remodeling, which in turn leads to elevated NT-proBNP levels.

We cannot exclude the possibility that autonomic dysfunction represents an elevated demand for compensatory mechanisms as heart failure progresses. Studies have shown that patients with heart failure and lower HRV tend to have a worse prognosis in terms of mortality. If low HRV or the presence of CAN were primarily driven by existing car-

7. Discussion

diac complications, it would suggest that individuals with these conditions exhibit more pronounced sympathetic overactivity as a consequence of heart failure progression, indicating potential reverse causation. Hence, elevated sympathetic activity during rest may reflect a greater reliance on compensatory mechanisms to maintain cardiac output. More precise measures are needed to assess whether sympathetic activity acts as a primary driver of heart failure or as a secondary compensatory response to cardiac dysfunction. In addition, it remains unclear to what extent the parasympathetic nervous system can act as a protective mechanism to counterbalance sympathetic dominance, and whether a decline in HRV reflects a breakdown of this balance. The two pathways (autonomic neuropathy and cardiac remodelling) are not mutually exclusive and may interact in a reinforcing cycle. Autonomic dysfunction can lead to increased sympathetic tone and reduced parasympathetic modulation, placing the heart under chronic stress and promoting structural and functional changes²³. In turn, cardiac remodelling may impair autonomic regulation, further exacerbating the imbalance. This interplay may create a self-perpetuating loop that accelerates the progression of heart failure. However, this remains beyond the scope of our current data and analysis.

7.3. Clinical implications

The dissertation investigates autonomic dysfunction in populations ranging from normal glucose metabolism to T2D and yields insights relevant for individuals engaged at different levels of the healthcare system. No specific role has yet been defined for autonomic dysfunction in clinical decision-making within healthcare, as current treatment and intervention options remain limited. Despite the fact that the results do not extend to the implementation of autonomic dysfunction in clinical practice, the findings from three included cohorts representing populations under public health, primary care, and secondary care. They mark a notable first step in addressing the impact of autonomic dysfunction on cardiovascular complications. In following section, the clinical implications of using autonomic dysfunction in the prevention of CVD will be discussed. If long-term HRV or CARTs are to be considered for improving risk stratification, it is important to determine at what stage in the progression of diabetes risk, and at which level of care, autonomic dysfunction becomes meaningful for early detection and intervention.

7.3.1. Public health

A central strategy in preventing CVD is the early identification and treatment of individuals at high risk⁸⁹. Public health initiatives support this by promoting healthy lifestyles, facilitating early screening for risk factors, and improving access to essential

7.3. Clinical implications

care and medications. Long-term HRV may enhance these efforts by identifying individuals with elevated cardiovascular risk and by tracking their physiological response to lifestyle changes.

Evidence from Study I showed that lower long-term HRV was associated with increased arterial stiffness, as measured by cf-PWV and CD, even in individuals without T2D. One standard deviation lower HRV corresponded to the effect of 2.7 additional years on cf-PWV and 1.6 years on CD⁹⁰. These cross-sectional findings suggest that HRV may serve as a marker of early vascular aging and cardiovascular risk. Supporting this, the Whitehall II study demonstrated a longitudinal relationship between short-term HRV and aortic stiffness. Together, these findings highlight the potential of HRV as a dynamic indicator of vascular health.

Within the public health setting, individuals with prediabetes represent a particularly vulnerable group at risk for comorbidities⁹¹. They often fall between structured care pathways, sometimes receiving regular follow-up and other times remaining outside formal healthcare systems. Notably, Studies I and II demonstrated that the associations between long-term HRV and CVD risk were especially pronounced in this population. In those at high risk of diabetes, a one standard deviation (33 ms) decrease in multiday SDNN was equivalent to 4.5 additional years of aging for major adverse cardiovascular events and 2.2 to 2.4 years for heart failure⁹². On a population level, lower HRV (SDNN: 100 ms) in individuals with prediabetes was associated with a higher incidence rate of CVD, heart failure, and all-cause mortality compared to individuals with normal-to-higher HRV (SDNN: 120–160 ms). These findings reinforce the role of HRV as an early and sensitive marker of cardiovascular health in populations at cardiometabolic risk.

While these findings highlight HRV's potential, practical implementation faces several challenges. Historically, long-term HRV monitoring has required specialized equipment such as Holter ECG recorders. However, the growing popularity of wearable devices offers a promising alternative. These devices provide a non-invasive, user-friendly way to collect heart rate and HRV data over time¹¹.

If HRV monitoring proves effective in helping individuals maintain a healthy, age-adjusted HRV range through lifestyle changes and prompts healthcare engagement when HRV deteriorates, it could become a meaningful tool for long-term health tracking. A cross-sectional study of 8 million individuals found that those who took more steps per day had higher HRV¹³, suggesting that HRV may also reflect behavioral adaptation.

A major public health challenge lies in ensuring equitable access to wearable technology. Individuals from lower socioeconomic backgrounds are significantly less likely to own such devices, raising concerns about health disparities. Despite this, there is encouraging evidence that the general population is receptive to digital health innovations. Many are

7. Discussion

willing to share health data with public institutions and support the use of AI in disease monitoring^{[11]93}.

Integrating wearable HRV monitoring into public health strategies could represent a transformative step in proactive cardiovascular care. It may support early detection, personalized prevention, and timely referral to primary care when risk levels increase.

7.3.2. Primary care

Cardiovascular risk in primary care is assessed using clinical evaluations and standardized risk prediction tools to identify individuals at elevated risk (ref.). Management focuses on lifestyle modification, pharmacological therapy, and regular monitoring to reduce cardiovascular events (ref.). In this context, long-term HRV may offer added value by improving the precision of cardiovascular risk stratification and by serving as a marker to monitor the effectiveness of preventive strategies.

Long-term HRV may enhance individual risk classification when added to established clinical risk scores. Tools such as SCORE2 and the Framingham Risk Score are widely used in primary care to guide cardiovascular risk assessment^{94,95}. In Study I, models adjusted for conventional CVD risk factors supported the potential added value of 24-hour HRV in relation to arterial stiffness, a surrogate marker of CVD risk. Study II extended this perspective by demonstrating associations between multiday HRV and incident CVD and heart failure. However, these findings are based on associations and do not include formal prediction modeling⁹⁶. A limitation of our studies is that we did not evaluate whether incorporating long-term HRV or CARTs into existing risk scores improves predictive performance beyond current guidelines. While most biomarkers have shown limited incremental value beyond established predictors (including age, sex, lipid profiles, diabetes status, and blood pressure), some studies suggest that 24-hour HRV may improve risk discrimination for CVD and all-cause mortality in individuals with T2D⁹⁷, and for stroke and heart failure in older adults^{[83]98}. However, these studies often lack calibration or validation in large-scale cohorts and have not been integrated with widely used risk scores such as SCORE2 or the Framingham Risk Score.

Long-term HRV may also help identify preclinical autonomic dysfunction, enabling targeted interventions in a subgroup of patients to prevent CVD. The increasing availability of wearable devices capable of capturing long-term HRV data presents a practical opportunity for continuous monitoring in primary care. These devices may facilitate earlier detection of autonomic dysfunction and support more personalized approaches to cardiovascular risk management. However, the clinical utility of stratifying patients based on preclinical autonomic dysfunction remains uncertain. These considerations are only actionable if interventions in this subgroup can be shown to reduce cardiovascular

7.3. Clinical implications

risk. Emerging evidence suggests that both pharmacological and lifestyle interventions can improve HRV in the short term^{99,100}. For example, high-intensity interval training has been shown to improve autonomic function in obese individuals with and without T2D¹⁰¹. Similarly, lifestyle changes in individuals with prediabetes have been associated with improvements in short-term HRV, which may partly explain a reduction in diabetes risk independently of weight loss¹⁰². Nevertheless, it remains unclear whether these effects on HRV are sustainable over time and whether they translate into long-term cardiovascular protection. In many cases, improvements in autonomic function may be mediated indirectly through changes in cardiometabolic markers such as glucose levels, lipid profiles, body weight, maximal oxygen uptake, and blood pressure.

Despite these uncertainties, monitoring autonomic function through long-term HRV may offer a valuable tool for assessing cardiovascular risk and tracking the impact of preventive strategies. In Denmark, prediabetes, defined by HbA1c, is present in 7.1% of adults¹⁰³. One in five of these individuals develops T2D within five years¹⁰³, while others either remain in the prediabetic stage or return to normoglycemia. Despite their increased risk of CVD and heart failure¹⁰⁴, individuals with prediabetes are not captured by existing preventive strategies. This underscores the need for early and precise risk assessment¹⁰. Given that the cardiovascular consequences of autonomic dysfunction appear to be more pronounced in individuals with prediabetes compared to those with normoglycemia, HRV has the potential to help identify those at elevated CVD risk within this group. However, evidence demonstrating improved risk prediction and sustained effects leading to better cardiovascular outcomes is needed to establish its relevance for integration into primary care.

7.3.3. Secondary care

In secondary care, endocrinologists assess cardiovascular and heart failure risk by integrating advanced diagnostics, biomarker analysis, and imaging to detect early dysfunction. The treatment of patients with T2D is guided by evidence-based therapies and multidisciplinary collaboration. The ADA/EASD 2022 consensus on Management of Hyperglycemia in T2D emphasizes that early detection of heart failure in individuals with T2D is crucial, as it enables timely initiation of therapies such as SGLT2 inhibitors, which have demonstrated significant benefits in reducing heart failure-related outcomes¹⁰⁵. A major challenge in diabetes care is detecting heart failure before symptoms appear, as patients with symptomatic heart failure face a higher risk of mortality and more frequent hospitalizations⁴. The AHA, ACC, and HFSA 2022 guidelines recommend identifying individuals at risk of heart failure based on factors such as diabetes, poor glycaemic control, uncontrolled hypertension, hyperlipidaemia, elevated BMI, albuminuria, renal dysfunction, and a history of CVD¹⁰⁶. Still, there is a need to identify

7. Discussion

optimal approaches for recognizing and diagnosing heart failure in clinical care, as broad echocardiographic screening in T2D is time-consuming and costly⁴.

Study III demonstrated that CAN may help identify individuals at increased risk of heart failure, beyond what is captured by symptoms or existing risk scores. Our findings support considering CAN as a relevant risk factor for heart failure and suggest it may have value in future risk stratification strategies in T2D. A clinical advantage of using CARTs is that they are standardized tests performed under controlled conditions. CARTs has proven to be reliable and reproducible, with reference values established in large population studies⁴⁷. Beyond these findings and the established evidence of increased heart failure risk, CAN in the T2D population also identifies individuals at high risk for overall CVD, kidney disease, and early mortality^{[107]108}. In Study III, I observed that two out of five participants had CAN, highlighting it as a prevalent complication. Therefore, detecting CAN may uncover an often-overlooked condition that is common in individuals with T2D.

Clinical stratification of care includes two key considerations: (1) CAN should be further evaluated for associated cardiovascular complications, such as heart failure; and (2) cardiopreventive strategies should be initiated earlier in this subgroup.

First, patients with CAN may benefit from further cardiovascular assessment, including the use of sensitive biomarkers or echocardiography. NT-proBNP is a strong predictor of heart failure and a validated biomarker for ruling out the condition (ref.). However, its specificity varies across heart failure phenotypes, being less specific for detecting HFpEF compared to HFrEF. Therefore, additional evaluation using echocardiography may be warranted. Beyond classify heart failure phenotypes, echocardiography identifies preclinical stages of heart failure through the detection of functional or structural cardiac abnormalities. Including CAN in structured assessments of heart failure could help clarify to which extent does CAN overlap with cardiac abnormalities. Determining the diagnostic and prognostic value of CAN, particularly its sensitivity and specificity in detecting HFrEF and HFpEF, requires further investigation.

Second, the presence of CAN may justify earlier initiation of protective therapies. SGLT2 inhibitors are recommended as second-line treatment in T2D and have demonstrated benefits in reducing the risk of heart failure, CVD, and kidney function decline, complications commonly associated with CAN. Current guidelines recommend initiating these therapies based on a history of CVD, heart failure, or the presence of conventional high-risk cardiovascular factors. However, the specific impact of SGLT2 inhibitors on the progression of cardiorenal outcomes in patients with CAN remains to be fully understood. Furthermore, while antihypertensive treatment is a cornerstone of cardiovascular risk management, whether specific classes of antihypertensive agents offer protective effects in patients with CAN remains to be explored.

The clinical implications of our findings in Study III are limited. The generalizability of our results is restricted, as our study population consisted of patients with T2D receiving secondary care. Two out of five patients with CAN showed to have a history of CVD—a group already at increased risk of heart failure due to their prior diagnosis. This overlap may influence the interpretation of CAN as an independent risk factor. Therefore, these findings need to be validated in a broader population with T2D, including individuals without a history of CVD. Doing so would allow for greater generalizability of our results to the broader T2D population, particularly those visiting primary care.

7.4. Strengths and limitations

7.4.1. Study design

Cross-sectional design

Studies I and III are based on cross-sectional data, with exposure and outcome measured within a three-month period. The main limitation of this design is that it does not allow us to determine whether the exposure led to the outcome or vice versa. As a result, we cannot establish temporality or confirm whether changes in the outcome were caused by the exposure. Based on prior evidence, the direction of the associations in Study I was inferred using physiological knowledge and findings from epidemiological and *in vivo* studies⁶⁵.

Study III focused on the clinical diagnosis of CAN and the presence of heart failure. The research question was oriented toward the clinical utility of CAN in identifying patients with T2D who may be progressing early toward heart failure. Whether cardiac function progressively worsens due to the underlying mechanisms of CAN remains to be fully established.

Longitudinal design

A major strength of Study II is its longitudinal design, where HRV was measured at baseline and outcomes were captured prospectively through national registries. This temporal structure ensures that the exposure (HRV) preceded the outcome, reducing the risk of reverse causation. The prospective design allows for stronger inference of directionality than cross-sectional studies. Furthermore, the use of high-quality registry data ensures complete outcome ascertainment and minimizes loss to follow-up bias.

Causality cannot be ascertained from the findings in Study I and Study II, and more causally focused methods are needed. Mendelian randomization, which uses genetic instruments for exposure, could help address this question. Additionally, structured mediation analysis involving modifications such as medication or lifestyle changes could

7. Discussion

clarify whether improving HRV or CART reduces cardiovascular risk, using data from intervention studies.

7.4.2. Internal validity

In this project, I aimed to assess cardiovascular autonomic function both in free-living conditions and in response to standardized test procedures during clinical visits. Additionally, I used dynamic measurements to evaluate arterial stiffness both locally and by velocity, and biomarker assessments to determine the presence of heart failure. In this section, I discuss the validity of 24-hour, multiday, and hourly HRV measurements, as well as the standardized tests of CAN. I also address the validity of the included outcomes and discuss the strengths and limitations of using MACE as a time-to-event outcome.

7.4.2.1. Long-term HRV (>24 hours) as measurement for autonomic function

[Further consideration is needed regarding the accuracy and relevance of different HRV indices. In long-term HRV, indices sensitive to respiration (such as RMSSD, HF, and pNN50) are more influenced by behavioral factors and have shown weaker associations with metabolic and cardiovascular outcomes compared to total variability measures (such as SDNN, SDANN, TP, ULF, VLF, and LF)^{[44]890}. However, their utility may improve when measured during rest in short segments, such as 5-minute recordings.] [Even if HRV proves to be causally linked with cardiovascular complications, its measurement is limited by the inability to distinguish between sympathetic and parasympathetic contributions.]

A main consideration in HRV analysis is the reliability of raw inter-beat interval data from ECG recordings. To ensure accurate various HRV measures, the intervals must be captured in a continuous and correctly sequenced manner. Frequency-domain analyses depend on the integrity of the inter-beat interval sequence, while some time-domain measures, such as RMSSD and pNN50, specifically quantify the variability in the differences between successive intervals.

In Study I, data from a 12-lead Holter system was used, which is considered the gold standard for long-term ECG recordings. With detailed and sequential inter-beat intervals, all HRV metrics were calculated.

In Study II, data from the Actiheart device was used for HRV. The device was configured to record continuously over an 8-day period. It captured 30-second epochs of mean heart rate intervals, along with upper and lower prediction intervals. From each epoch, I generated a distribution of inter-beat intervals. An algorithm was applied to

7.4. Strengths and limitations

estimate HRV from these distributions, and its validation showed strong agreement with established metrics, including SDNN, SDANN, and the SDNN index [6]⁵⁷. However, a limitation of this dataset is that it did not allow for the calculation of frequency-domain measures or specific time-domain metrics such as RMSSD or pNN50. HF power, RMSSD, and pNN50 typically show earlier diurnal peaks compared to low-frequency power and SDNN¹³. These measures are more reflective of parasympathetic activity during in- and expiration. It remains to be determined how these hourly measures relate to CVD, heart failure, and all-cause mortality.

In the context of this study, which focuses on long-term HRV in free-living conditions, it is important to acknowledge that the autonomic nervous function we aim to assess may also be influenced by behavioral factors such as physical activity, sleep, meal timing, emotions, smoking, caffeine intake, alcohol consumption, medication use, and prior cardiovascular complications. These factors can potentially mask or mimic underlying physiological dysfunction during recordings, but they can also elicit the HRV responses we are interested in. Therefore, reduced long-term HRV cannot be interpreted solely as a marker of autonomic function. HRV is also influenced by lifestyle patterns over time, making it sensitive not only to day-to-day behaviors but also to long-term habits that affect autonomic balance.

In Studies I and II, I accounted for habitual physical activity, and in Study II, I also adjusted hourly HRV for physical movement during recordings to test the influence of concurrent activity. However, further studies are needed to understand how lifestyle patterns affect long-term HRV recordings on subsequent days, in order to separate direct behavioural from physiological components. In both studies, I excluded participants with prior CVD to preserve the etiological order between autonomic dysfunction and cardiovascular outcomes.

Anti-hypertensive medications, especially beta-blockers, are known to increase HRV in randomized controlled trials¹⁰⁹. However, in cohort studies, participants using anti-hypertensives generally show lower HRV, likely reflecting a higher burden of cardiovascular complications [ref]. Because beta-blockers target the autonomic nervous system, they may introduce bias in HRV measurements by interfering with the function. In sensitivity analyses in Studies I and III, excluding participants on anti-hypertensive treatment did not materially change the estimates. Therefore, these participants were kept and adjusted for medication use in the full models.

HRV levels are influenced by heart rate, as lower resting heart rate allows for greater variability. In Study I, I chose not to adjust for heart rate in our models, as this could introduce multicollinearity. Additionally, elevated heart rate is driven by increased sympathetic activity and may act as a mediator in the pathway leading to arterial stiffness. Our use of full-day recordings captures HRV during both rest and activity, providing a robust representation of autonomic function over a typical day. In contrast, heart

7. Discussion

rate correction may be more relevant for short-term HRV recordings, where standardized conditions can be affected by random influences such as time of day, smoking, or caffeine intake. These factors would have been relevant in Study III had I included HRV measures. In Study II, I used the residuals method to pre-adjust HRV measures for resting heart rate, which accounted for part of the observed associations, particularly with heart failure and all-cause mortality, and to a lesser extent with ischemic-related CVD events. Similar trends were observed for hourly associations, where heart rate pre-adjustment had had comparable effects on the outcomes.

[When HRV is analyzed in shorter segments (e.g., hourly or in 5-minute intervals), measures like RMSSD, pNN50, and HF appear to offer new insights into autonomic function and its relevance in diabetes and CVD, e.g. during night-time [¹¹⁰]⁹⁹. SDANN and SDNNi aim to reduce the impact of short-term variability, such as that caused by physical activity, by calculating either the standard deviation of 5-minute segment mean IBI (SDANN) or the mean of standard deviations across 5-minute segments (SDNNi). [include lower-frequency points]. This helps smooth out transient fluctuations and better capture long-term autonomic modulation. Thus, behavioral patterns pose a limitation in physiological research aiming to disentangle the causal pathways between autonomic dysfunction and CVD when using long-term HRV measures. These patterns likely introduce high variability between observations that is not attributable to autonomic function itself.]

7.4.2.2. Cardiovascular autonomic reflex test

CART provides a practical approach for screening for autonomic dysfunction and has been shown to be a reliable method¹¹¹. Although certain indices from CARTs may be influenced by factors such as time of day or recent physical activity, these effects are generally minimal. Furthermore, no impact of caffeine intake has been observed on the reference age-based formula⁴⁷. A limitation of the CARTs in this study was the high prevalence of participants who were unable to complete the full battery of tests, primarily due to missing data from the Valsalva manoeuvre.

7.4.2.3. Measures of cardiovascular risk

[Arterial stiffness - NT-proBNP - MACE limitation in aetiological research - Non-specific heart failure and MI and stroke and death - HRV stronger link with MI or stroke - Perspective: decreasing number of events with prolonging time-to-event - Challenge for coming observational cohorts (need to increase sample size) [The Problem With Composite End Points in Cardiovascular Studies The Story of Major Adverse Cardiac Events and Percutaneous Coronary Intervention]]

7.4.3. External validity

7.4.3.1. Selection bias

The Maastricht Study

The target population in Study I was intended to represent individuals at different stages of glucose metabolism. However, our analysis may be affected by selection bias in the representation of individuals with T2D. The Maastricht Study recruited participants who were able and willing to attend multiple research visits and receive personal health feedback, which likely attracted health-conscious individuals with higher education levels. As a result, participants with T2D were relatively healthy, with a median disease duration of three years and a low prevalence of complications. Those who completed both long-term ECG and arterial stiffness assessments may represent an even healthier subgroup. This selection bias may limit the generalizability of the findings to the broader T2D population and could explain why the effect modification did not differ step-wise from that observed in individuals with prediabetes.

ADDITION-PRO

The target population in Study II was intended to represent individuals at high risk of developing T2D. Participants were recruited through a stepwise screening procedure. Initially, individuals were selected based on a risk score derived from a self-administered questionnaire sent by mail. Those with high scores were invited for further testing using HbA1c or random glucose measurements.

This recruitment strategy involved selection by design, as it defined the source population based on specific risk criteria. The questionnaire prioritized risk factors such as older age and hypertension, leading to overrepresentation of these groups¹¹². Prediabetes was identified only after biochemical testing, while the risk score was primarily designed to detect undiagnosed T2D. Although this selection process was intentional and aligned with the ADDITION-PRO objectives, it may limit the generalizability of the findings to the broader population at risk for T2D.

In addition, selection bias may have occurred due to differential participation in the ADDITION screening program. Healthier individuals were more likely to participate, both by completing the risk questionnaire and by attending follow-up testing¹¹³. As a result, the baseline risk for CVD in ADDITION-PRO participants may have been lower compared to the broader target population.

CANCAN

The target population in Study III was intended to represent individuals with type 2 diabetes treated in outpatient clinics. In Denmark, patients with type 2 diabetes are

7. Discussion

referred to diabetes specialists at outpatient clinics when their general practitioner is unable to stabilize their condition. A strength of the CANCAN sampling strategy is that patients were already attending endocrinology consultations, and the study examination required only additional time during their visit, without the need for extra transportation or appointments. Assessing selection bias in this study is challenging, as inclusion depended on referral practices by general practitioners¹¹⁴. These practices may vary individually, with differing thresholds for referring patients to specialized care based on clinical judgment and patient characteristics.

7.4.3.2. Generalisability

[As a result, CANCAN participants represent a higher-risk group among Danish diabetes patients, while more stable patients remain under general practitioner care. Consequently, the prevalence of heart failure indicators and CAN is likely higher in this selected group, than patients in primary care.]

The generalisability of our findings can be considered on two levels: how well the results apply to the general population within the country, and how they translate to populations with different ancestries in other countries.

Studies II and III include individuals at high risk of diabetes and those with T2D. Therefore, the associations between cardiovascular autonomic dysfunction and cardiovascular outcomes or surrogate biomarkers are relevant to individuals with some degree of diabetes risk. However, whether these associations hold in the general population remains uncertain. Study I suggests that the link between autonomic dysfunction and cardiovascular risk, as measured by arterial stiffness, is also present in individuals with normal glucose metabolism, though to a lesser extent. This finding was supported by replication in the Whitehall II cohort, strengthening the generalisability of the observed relationship⁶⁵.

The study populations in Studies II and III consist of individuals of Nordic descent, while Study I includes individuals of Western European descent. Since the constellation of diabetes risk factors may differ in populations of Asian, South American, African, and other ancestries, our findings may not be fully generalisable to these groups. This limitation affects the applicability and magnitude of the observed associations to an unknown degree. Further cohort studies including underrepresented populations are warranted. As our studies focus on diabetes risk, all participants were adults aged 40 years and older. Therefore, our findings are limited to this age group, and whether the results extend to younger adults or children remains to be confirmed. Overall, while our study benefits from including individuals across different levels of diabetes risk, limitations in generalisability remain, particularly for more diverse and younger populations.

8. Conclusion

This dissertation aimed to investigate whether cardiovascular autonomic dysfunction, assessed by heart rate variability and cardiovascular autonomic reflex tests, is associated with cardiovascular complications across different stages of glucose metabolism. The findings support the hypothesis that cardiovascular autonomic dysfunction is an early and independent marker of cardiovascular risk.

Across three studies, cardiovascular autonomic dysfunction was associated with increased arterial stiffness, a higher incidence of ischemic cardiovascular events, heart failure, and all-cause mortality. These associations were observed not only in individuals with T2D but also in those with prediabetes and normal glucose metabolism. This suggests that cardiovascular autonomic dysfunction may precede the development of overt metabolic disease. Furthermore, standardized cardiovascular autonomic reflex tests identified individuals with T2D who were at increased risk of heart failure, even when they were asymptomatic and not classified as high risk by conventional clinical tools.

In summary, cardiovascular autonomic dysfunction appears to be a clinically relevant marker of cardiovascular risk across the full spectrum of glucose metabolism. However, it remains unclear whether this dysfunction plays a causal role or reflects underlying pathophysiological processes. Further research is needed to determine its clinical utility in risk stratification and its potential as a target for intervention.

8. Conclusion

Autonomic dysfunction, assessed through 24-hour HRV, is associated with increased arterial stiffness. This relationship is already evident in individuals with normal glucose metabolism and becomes more pronounced in those with prediabetes and type 2 diabetes. In individuals at high risk of type 2 diabetes, lower long-term HRV measured over a week has been linked to ischemic events, heart failure, and all-cause mortality, highlighting HRV's potential as a marker of cardiovascular health. Both HRV and heart rate follow circadian patterns in relation to cardiovascular disease. Higher nighttime heart rate is associated with increased risk of heart failure, and specific morning patterns of HRV have been linked to ischemic events. These findings suggest that both long-term and hourly heart rate and HRV measures provide valuable prognostic information. Structured testing of cardiovascular autonomic function in individuals with type 2 diabetes can detect those with CAN and may help identify individuals at higher risk of heart failure.

8. Conclusion

We have established an association between HRV and cardiovascular complications. However, the underlying mechanisms remain unclear. It is not yet known whether autonomic dysfunction, as indicated by low HRV, is a marker of developing arteriosclerosis, atheroma, or cardiac remodeling, or whether it plays a causal role in their development. While the pathogenic pathways leading to cardiovascular risk appear similar across the spectrum of glucose metabolism, dysglycemia may amplify the impact of autonomic dysfunction. Whether lower long-term HRV in individuals with prediabetes or type 2 diabetes reflects a distinct physiological mechanism involving neuropathy, compared to those with normal glucose metabolism, remains an open question.

Structured studies assessing screening strategies may help determine the optimal timing, population, and methods for incorporating HRV and CART into clinical practice. Furthermore, trial designs that focus on either lifestyle interventions or targeted pharmacological modulation of HRV are needed to clarify the clinical role of HRV and CART in cardiovascular prevention. When using HRV and standardized CART, it is important to carefully consider how the data are applied in relation to specific research objectives, ranging from physiological mechanisms to clinical diagnosis. Long-term HRV and its hourly fluctuations provide insight into autonomic responses under free-living conditions. Further research is needed to determine whether modifying these measures can yield sustained preventive effects on cardiovascular disease and mortality.

8. Conclusion

CARTs offer a standardized approach for diagnosing CAN. Clarifying the clinical utility of CARTs in assessing cardiovascular and heart failure risk through the identification of CAN is essential for advancing precision care in individuals with type 2 diabetes.

Echocardiographic studies can help establish the link between CAN and the risk of specific heart failure subtypes. Future research should carefully select HRV measures aligned with specific clinical or research objectives. Long-term HRV and CART have demonstrated potential in cardiovascular risk assessment and should be integrated to evaluate whether autonomic function assessments can monitor treatment or lifestyle effectiveness or guide stratified cardiovascular risk decisions in individuals with prediabetes or type 2 diabetes.

[Early detection are important as CVD and heart failure leads to lower life-expectancy and quality of life.]

9. Perspective

We have investigated cardiovascular autonomic function impact on cardiovascular complications across different stages of glucose metabolism. Based on our findings and conclusions, we propose further perspectives to define its role in research and healthcare from three aspects: (1) continuous non-invasive health monitoring, (2) risk stratification, and (3) identification as a causal and modifiable marker.

9.1. Continuous monitoring of cardiovascular health

Understanding when and how physiological signals reflect elevated CVD risk is essential for developing early and effective prevention strategies. Incorporating HRV into digital health solutions could support personalized feedback mechanisms, enabling timely lifestyle or therapeutic interventions and contributing to more adaptive and preventive healthcare strategies. Wearable devices enable comprehensive data collection on behavioral (e.g., sleep and physical activity) and physiological (e.g., heart rate, ECG, temperature) parameters¹¹⁵. These devices offer a broader and more feasible approach to long-term heart rate monitoring. Despite growing interest in wearable-based monitoring, the integration of HRV into routine cardiometabolic risk assessment remains limited.

Two key aspects highlight the potential applications of monitoring: (1) identification of risk and (2) assessment of response to intervention.

Identification of risk

We ascertained that lower long-term HRV is a risk factor for CVD, as demonstrated by its association with arterial stiffness and hard CVD endpoints. In addition, our findings that specific time points of HRV and heart rate are associated with increased risk of CVD and heart failure suggest that distinct physiological responses captured under free-living conditions may enhance early risk detection. The majority of the observed associations were not explained by movement at the corresponding time points. For risk assessment, instead of merely adjusting for physical activity as a confounding factor, future predictive models could benefit from integrating multiple physiological signals such as HRV, sleep, and activity patterns to better reflect dynamic states of health. Machine learning techniques offer the ability to process complex raw time-series

9.1. Continuous monitoring of cardiovascular health

data, such as interbeat intervals and accelerometer signals, and to uncover patterns that may improve risk prediction beyond conventional summary measures of HRV. However, a key limitation of these models is their reduced explainability, which may limit their clinical applicability.

Assessment of response to intervention

HRV highlights a potential target for intervention, given that low HRV may be indicative of adverse lifestyle patterns. For instance, behavioral patterns such as disrupted sleep, inactivity, or irregular meal timing influences circadian fluctuations in HRV. Evidence from studies on night-shift workers suggests that meal timing affects HRV, with daytime meals leading to higher HRV during night hours⁹⁹. With short-term effects observed in both 24-hour and hourly HRV measures, medications such as beta-blockers increases HRV while GLP-1 receptor analogues showed to lower HRV [¹⁰⁹]¹¹⁶.




Figure 9.1.: HRV feedback in response to lifestyle and treatment interventions. (Source: Author)

9. Perspective

Future studies can leverage wearable devices to better understand the behavioral factors and treatment options that contribute to its improvement or deterioration of HRV. This approach may help identify effective lifestyle patterns or medications that improve success of cardiovascular health through modulation of HRV.

Standardization and transparency across different brands of wearable devices remain a challenge for both research and clinical implementation of heart rate and HRV monitoring. While smartwatches offer a convenient method for heart rate measurement, their accuracy can vary, as they rely on photoplethysmography to detect pulse rate at the wrist. This method can be imprecise under certain conditions, particularly during physical activity, due to motion artifacts and other external factors^{[117]118}. Despite these limitations, ongoing improvements in sensor technology and algorithm calibration are likely to enhance the reliability of wearable-derived heart rate and HRV data.

9.2. Risk-stratification

Based on the clinical implications, the distinct roles of long-term HRV and CART in risk stratification remain to be demonstrated and require further investigation.

From the perspective of wearable devices, it remains to be determined whether HRV can serve as an early indicator of CVD risk when used in a risk score alongside simple non-invasive markers such as age, sex, and BMI, and how it compares to current CVD risk scores that include blood-based biomarkers and blood pressure.

A limitation of long-term HRV measurement is the lack of standardization, as data are collected under free-living conditions and may be influenced by daily behaviors, potentially affecting risk classification. Therefore, standardized procedures may be necessary. CART has been shown to be a reliable, non-invasive method that typically takes approximately 10 minutes to complete. As such, a standardized and validated diagnosis of CAN can help identify patients with T2D who are at higher risk of CVD, nephropathy, and heart failure. However, the extent to which the diagnosis of CAN identifies heart failure risk and is generalizable to broader populations with T2D or even prediabetes, in a clinically relevant context, remains to be assessed.

Our findings suggest that long-term HRV and CAN may serve as potential markers for CVD and heart failure risk among individuals at elevated metabolic risk, helping to identify those who could benefit from preventive strategies. Future research should evaluate whether individuals classified as high-risk based on autonomic dysfunction or CAN should initiate cardiovascular-focused prevention or screening earlier, or receive specific interventions tailored to their condition.




Figure 9.2.: Risk-stratification by autonomic dysfunction. (Source: Author)

9.3. Effective causal modifiable marker

Our findings support an etiological link between long-term HRV and the risk of CVD, providing initial evidence of a potential causal relationship. However, the observed association does not confirm causation, and further research is necessary to determine whether the relationship between HRV and CVD risk is truly causal. Traditionally, epidemiological research has relied on randomized controlled trials to establish causality. However, conducting such trials to isolate the direct effect of HRV is particularly challenging. Interventions that influence HRV often do so indirectly, through changes in lifestyle factors such as weight loss, inflammation, and insulin sensitivity. Similarly, pharmacological treatments may improve HRV as a secondary effect, for example through blood pressure reduction from antihypertensive medications. As a result, isolating the direct modification of HRV is difficult. To address the limitations of clinical trials, modern epidemiological approaches such as Mendelian randomization¹¹⁹ and structured causal mediation analysis offer alternatives for inferring causality from observational genetic studies and for estimating indirect effects using trial data¹²⁰. To date, no GWAS has been conducted to investigate the genetic determinants of long-term HRV. Establishing such genetic associations is essential for understanding its genetic architecture and for providing unconfounded estimates by using genetic variants as proxies to assess the causal role of HRV in CVD.




Figure 9.3.: Mediation of HRV by intervention in prevention of CVD. (Source: Author)

9.3. Effective causal modifiable marker

Future cardiometabolic intervention trials, whether focused on lifestyle modification or pharmacological treatment, should, where feasible, include HRV measurements to enable structured mediation analyses and to better understand whether modifying autonomic function improves cardiovascular outcomes. This could help demonstrate whether modification of HRV through potential strategies such as medications e.g anti-hypertensives or lifestyle interventions including physical activity, diet, and sleep has a sustainable effect on CVD outcomes.

References

- 1 International Diabetes Federation. Diabetes atlas, 11th Edition. International Diabetes Federation, 2025.
- 2 Magliano DJ, Chen L, Carstensen B, *et al.* Trends in all-cause mortality among people with diagnosed diabetes in high-income settings: A multicountry analysis of aggregate data. *The Lancet Diabetes & Endocrinology* 2022; **10**: 112–9.
- 3 Zaman S, Wasfy JH, Kapil V, *et al.* The lancet commission on rethinking coronary artery disease: Moving from ischaemia to atheroma. *The Lancet* DOI:10.1016/S0140-6736(25)00055-8.
- 4 Pop-Busui R, Januzzi JL, Bruemmer D, *et al.* Heart failure: An underappreciated complication of diabetes. A consensus report of the american diabetes association. *Diabetes Care* 2022; **45**: 1670–90.
- 5 Hillebrand S, Gast KB, Mutsert R de, *et al.* Heart rate variability and first cardiovascular event in populations without known cardiovascular disease: Meta-analysis and dose-response meta-regression. *EP Europace* 2013; **15**: 742–9.
- 6 Task Force of the European Society of Cardiology the North American Society of Pacing Electrophysiology. Heart rate variability. *Circulation* 1996; **93**: 1043–65.
- 7 Eleftheriadou A, Spallone V, Tahrani AA, Alam U. Cardiovascular autonomic neuropathy in diabetes: An update with a focus on management. *Diabetologia* 2024; **67**: 2611–25.
- 8 Coopmans C, Zhou TL, Henry RMA, *et al.* Both prediabetes and type 2 diabetes are associated with lower heart rate variability: The maastricht study. *Diabetes Care* 2020; **43**: 1126–33.
- 9 Rooney MR, Fang M, Ogurtsova K, *et al.* Global prevalence of prediabetes. *Diabetes Care* 2023; **46**: 1388–94.
- 10 Birkenfeld AL, Franks PW, Mohan V. Precision medicine in people at risk for diabetes and atherosclerotic cardiovascular disease: A fresh perspective on prevention. *Circulation* 2024; **150**: 1910–2.
- 11 Dhingra LS, Aminorroaya A, Oikonomou EK, *et al.* Use of wearable devices in individuals with or at risk for cardiovascular disease in the US, 2019 to 2020. *JAMA Network Open* 2023; **6**: e2316634–4.

- 12 Bayoumy K, Gaber M, Elshafeey A, *et al.* Smart wearable devices in cardiovascular care: Where we are and how to move forward. *Nature Reviews Cardiology* 2021; **18**: 581–99.
- 13 Natarajan A, Pantelopoulos A, Emir-Farinás H, Natarajan P. Heart rate variability with photoplethysmography in 8 million individuals: A cross-sectional study. *The Lancet Digital Health* 2020; **2**: e650–7.
- 14 Tabák AG, Herder C, Rathmann W, Brunner EJ, Kivimäki M. Prediabetes: A high-risk state for diabetes development. *The Lancet* 2012; **379**: 2279–90.
- 15 Bonora E, DeFronzo R. Diabetes. Epidemiology, genetics, pathogenesis, diagnosis, prevention, and treatment. 2018 DOI:10.1007/978-3-319-27317-4.
- 16 World Health Organization. Diagnosis and management of type 2 diabetes (HEARTS-d). Geneva, 2020.
- 17 American Diabetes Association Professional Practice Committee. 2. Classification and diagnosis of diabetes: Standards of medical care in diabetes—2022. *Diabetes Care* 2021; **45**: S17–38.
- 18 Mokhtar SBA, Heide FCT van der, Oyaert KAM, *et al.* (Pre)diabetes and a higher level of glycaemic measures are continuously associated with corneal neurodegeneration assessed by corneal confocal microscopy: The maastricht study. *Diabetologia* 2023; **66**: 2030–41.
- 19 Huang Y, Cai X, Qiu M, *et al.* Prediabetes and the risk of cancer: A meta-analysis. *Diabetologia* 2014; **57**: 2261–9.
- 20 Stehouwer CDA. Microvascular dysfunction and hyperglycemia: A vicious cycle with widespread consequences. *Diabetes* 2018; **67**: 1729–41.
- 21 Taylor R. Type 2 diabetes: Etiology and reversibility. *Diabetes Care* 2013; **36**: 1047–55.
- 22 Færch K, Witte DR, Tabák AG, *et al.* Trajectories of cardiometabolic risk factors before diagnosis of three subtypes of type 2 diabetes: a post-hoc analysis of the longitudinal Whitehall II cohort study. *The lancet Diabetes & endocrinology* 2013; **1**: 43–51.
- 23 Ritchie RH, Abel ED. Basic mechanisms of diabetic heart disease. *Circulation Research* 2020; **126**: 1501–25.
- 24 Yusuf S, Joseph P, Rangarajan S, *et al.* Modifiable risk factors, cardiovascular disease, and mortality in 155 722 individuals from 21 high-income, middle-income, and low-income countries (PURE): A prospective cohort study. *The Lancet* 2020; **395**: 795–808.
- 25 Mitchell GF, Powell JT. Arteriosclerosis. *Arteriosclerosis, Thrombosis, and Vascular Biology* 2020; **40**: 1025–7.

References

- 26 Lee H-Y, Oh B-H. Aging and arterial stiffness. *Circulation Journal* 2010; **74**: 2257–62.
- 27 Lu Y, Kiechl SJ, Wang J, Xu Q, Kiechl S, Pechlaner R. Global distributions of age- and sex-related arterial stiffness: systematic review and meta-analysis of 167 studies with 509,743 participants. *EBioMedicine* 2023; **92**: 104619.
- 28 Zhong Q, Hu M-J, Cui Y-J, et al. Carotid–femoral pulse wave velocity in the prediction of cardiovascular events and mortality: An updated systematic review and meta-analysis. *Angiology* 2018; **69**: 617–29.
- 29 Kawai K, Kawakami R, Finn AV, Virmani R. Differences in stable and unstable atherosclerotic plaque. *Arteriosclerosis, Thrombosis, and Vascular Biology* 2024; **44**: 1474–84.
- 30 Alasad M, Shipp NJ, Brooks AG, et al. Myocardial infarction and atrial fibrillation. *Circulation: Arrhythmia and Electrophysiology* 2013; **6**: 738–45.
- 31 Elgendi IY, Mahtta D, Pepine CJ. Medical therapy for heart failure caused by ischemic heart disease. *Circulation Research* 2019; **124**: 1520–35.
- 32 Bilano V, Gilmour S, Moffiet T, et al. Global trends and projections for tobacco use, 1990–2025: An analysis of smoking indicators from the WHO comprehensive information systems for tobacco control. *The Lancet* 2015; **385**: 966–76.
- 33 Ockene IS, Miller NH. Cigarette smoking, cardiovascular disease, and stroke. *Circulation* 1997; **96**: 3243–7.
- 34 Shah AD, Langenberg C, Rapsomaniki E, et al. Type 2 diabetes and incidence of cardiovascular diseases: A cohort study in 1 · 9 million people. *The Lancet Diabetes & Endocrinology* 2015; **3**: 105–13.
- 35 Donnan GA, Fisher M, Macleod M, Davis SM. Stroke. *The Lancet* 2008; **371**: 1612–23.
- 36 Vos T, Lim SS, Abafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: A systematic analysis for the global burden of disease study 2019. *The Lancet* 2020; **396**: 1204–22.
- 37 Li X, Kong X, Yang C, et al. Global, regional, and national burden of ischemic stroke, 1990–2021: An analysis of data from the global burden of disease study 2021. *eClinicalMedicine* 2024; **75**. DOI:10.1016/j.eclinm.2024.102758.
- 38 Lee M, Saver JL, Hong K-S, Song S, Chang K-H, Ovbiagele B. Effect of pre-diabetes on future risk of stroke: Meta-analysis. *BMJ : British Medical Journal* 2012; **344**: e3564.
- 39 Barr ELM, Zimmet PZ, Welborn TA, et al. Risk of cardiovascular and all-cause mortality in individuals with diabetes mellitus, impaired fasting glucose, and impaired glucose tolerance. *Circulation* 2007; **116**: 151–7.

- 40 Khan MS, Shahid I, Bennis A, Rakisheva A, Metra M, Butler J. Global epidemiology of heart failure. *Nature Reviews Cardiology* 2024; **21**: 717–34.
- 41 Normand C, Kaye DM, Povsic TJ, Dickstein K. Beyond pharmacological treatment: An insight into therapies that target specific aspects of heart failure pathophysiology. *The Lancet* 2019; **393**: 1045–55.
- 42 Campbell P, Rutten FH, Lee MM, Hawkins NM, Petrie MC. Heart failure with preserved ejection fraction: everything the clinician needs to know. *Lancet (London, England)* 2024; **403**: 1083–92.
- 43 Schlaich M, Straznicky N, Lambert E, Lambert G. Metabolic syndrome: a sympathetic disease? *Lancet Diabetes Endocrinol* 2015; **3**: 148–57.
- 44 Rinaldi E, Heide FCT van der, Bonora E, et al. Lower heart rate variability, an index of worse autonomic function, is associated with worse beta cell response to a glycemic load in vivo—the maastricht study. *Cardiovascular Diabetology* 2023; **22**: 105.
- 45 Huggett RJ, Scott EM, Gilbey SG, Stoker JB, Mackintosh AF, Mary DASG. Impact of type 2 diabetes mellitus on sympathetic neural mechanisms in hypertension. *Circulation* 2003; **108**: 3097–101.
- 46 Cseh D, Climie RE, Offredo L, et al. Type 2 diabetes mellitus is independently associated with decreased neural baroreflex sensitivity. *Arteriosclerosis, Thrombosis, and Vascular Biology* 2020; **40**: 1420–8.
- 47 Hansen CS, Christensen MMB, Vistisen D, et al. Normative data on measures of cardiovascular autonomic neuropathy and the effect of pretest conditions in a large danish non-diabetic CVD-free population from the lolland-falster health study. *Clinical Autonomic Research* 2025; **35**: 101–13.
- 48 Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *New England Journal of Medicine* 2002; **346**: 393–403.
- 49 Kahn SE, Deanfield JE, Jeppesen OK, et al. Effect of semaglutide on regression and progression of glycemia in people with overweight or obesity but without diabetes in the SELECT trial. *Diabetes Care* 2024; **47**: 1350–9.
- 50 Umetani K, Singer DH, McCraty R, Atkinson M. Twenty-four hour time domain heart rate variability and heart rate: Relations to age and gender over nine decades. *Journal of the American College of Cardiology* 1998; **31**: 593–601.
- 51 Serrador JM, Finlayson HC, Hughson RL. Physical activity is a major contributor to the ultra low frequency components of heart rate variability. *Heart* 1999; **82**: e9.
- 52 Akselrod S, Gordon D, Ubel FA, Shannon DC, Berger AC, Cohen RJ. Power spectrum analysis of heart rate fluctuation: A quantitative probe of beat-to-beat cardiovascular control. *Science* 1981; **213**: 220–2.

References

- 53 Taylor JA, Carr DL, Myers CW, Eckberg DL. Mechanisms underlying very-low-frequency RR-interval oscillations in humans. *Circulation* 1998; **98**: 547–55.
- 54 Jeffrey J. Goldberger, Rishi Arora, Una Buckley, Kalyanam Shivkumar. Autonomic nervous system dysfunction. *Journal of the American College of Cardiology* 2019; **73**: 1189–206.
- 55 Mccraty R, Shaffer F. Heart rate variability: New perspectives on physiological mechanisms, assessment of self-regulatory capacity, and health risk. *Global Advances in Health and Medicine* 2015; **4**: 46–61.
- 56 Elghozi J-L, Laude D, Girard A. EFFECTS OF RESPIRATION ON BLOOD PRESSURE AND HEART RATE VARIABILITY IN HUMANS. *Clinical and Experimental Pharmacology and Physiology* 1991; **18**: 735–42.
- 57 Schaarup J. Actiheart validation of time-domain heart rate variability. 2024. https://figshare.com/articles/online_resource/Actiheart_validation_of_time-domain_heart_rate_variability/26182361.
- 58 Tan Lai Zhou, Ronald M. A. Henry, Coen D. A. Stehouwer, Thomas T. van Sloten, Koen D. Reesink, Abraham A. Kroon. Blood pressure variability, arterial stiffness, and arterial remodeling. *Hypertension* 2018; **72**: 1002–10.
- 59 Bendix Carstensen Steno Diabetes Center. Who needs the cox model anyway. *Stat Med* 2012; **31**: 10741088.
- 60 Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect modification and interaction. *International Journal of Epidemiology* 2012; **41**: 514–20.
- 61 Hughes RA, Heron J, Sterne JAC, Tilling K. Accounting for missing data in statistical analyses: Multiple imputation is not always the answer. *International Journal of Epidemiology* 2019; **48**: 1294–304.
- 62 Angela Beres, John Sluyter, Robert Keith Rhodes Scragg. Association of arterial stiffness and neuropathy in diabetes: A systematic review and meta-analysis. *BMJ Open Diabetes Research & Care* 2023; **11**: e003140.
- 63 Casey DP, Curry TB, Joyner MJ, Charkoudian N, Hart EC. Relationship between muscle sympathetic nerve activity and aortic wave reflection characteristics in young men and women. *Hypertension* 2011; **57**: 421–7.
- 64 Nardone M, Floras JS, Millar PJ. Sympathetic neural modulation of arterial stiffness in humans. *American Journal of Physiology-Heart and Circulatory Physiology* 2020; **319**: H1338–46.
- 65 Schaarup JR, Christensen MS, Hulman A, et al. Autonomic dysfunction is associated with the development of arterial stiffness: The whitehall II cohort. *Gerontology* 2023; **45**: 2443–55.

- 66 Kuehl M, Stevens MJ. Cardiovascular autonomic neuropathies as complications of diabetes mellitus. *Nature Reviews Endocrinology* 2012; **8**: 405–16.
- 67 Shah AS, El ghormli L, Vajravelu ME, et al. Heart rate variability and cardiac autonomic dysfunction: Prevalence, risk factors, and relationship to arterial stiffness in the treatment options for type 2 diabetes in adolescents and youth (TODAY) study. *Diabetes Care* 2019; **42**: 2143–50.
- 68 Coopmans C, Zhou TL, Henry RMA, et al. Both prediabetes and type 2 diabetes are associated with lower heart rate variability: The maastricht study. *Diabetes Care* 2020; **43**: 1126–33.
- 69 McEnery CM, Wilkinson IB, Johansen NB, et al. Nondiabetic glucometabolic status and progression of aortic stiffness: The whitehall II study. *Diabetes Care* 2017; **40**: 599–606.
- 70 Rietz M, Schmidt-Persson J, Gillies Banke Rasmussen M, et al. Facilitating ambulatory heart rate variability analysis using accelerometry-based classifications of body position and self-reported sleep. *Physiological Measurement* 2024; **45**: 055016.
- 71 Bentzon JF, Otsuka F, Virmani R, Falk E. Mechanisms of plaque formation and rupture. *Circulation Research* 2014; **114**: 1852–66.
- 72 Zieman SJ, Melenovsky V, Kass DA. Mechanisms, pathophysiology, and therapy of arterial stiffness. *Arteriosclerosis, Thrombosis, and Vascular Biology* 2005; **25**: 932–43.
- 73 Popele NM van, Grobbee DE, Bots ML, et al. Association between arterial stiffness and atherosclerosis. *Stroke* 2001; **32**: 454–60.
- 74 Holwerda SW, Luehrs RE, DuBose L, et al. Elevated muscle sympathetic nerve activity contributes to central artery stiffness in young and middle-age/older adults. *Hypertension* 2019; **73**: 1025–35.
- 75 Gottsäter A, Ahlgren ÅR, Taimour S, Sundkvist G. Decreased heart rate variability may predict the progression of carotid atherosclerosis in type 2 diabetes. *Clinical Autonomic Research* 2006; **16**: 228–34.
- 76 Mohanta SK, Peng L, Li Y, et al. Neuroimmune cardiovascular interfaces control atherosclerosis. *Nature* 2022; **605**: 152–9.
- 77 Agarwal Sunil K., Norby Faye L., Whitsel Eric A., et al. Cardiac autonomic dysfunction and incidence of atrial fibrillation. *JACC* 2017; **69**: 291–9.
- 78 Osei J, Vaccarino V, Wang M, et al. Stress-induced autonomic dysfunction is associated with mental stress–induced myocardial ischemia in patients with coronary artery disease. *Circulation: Cardiovascular Imaging* 2024; **17**: e016596.

References

- 79 Nolte IM, Munoz ML, Tragante V, *et al.* Genetic loci associated with heart rate variability and their effects on cardiac disease risk. *Nature Communications* 2017; **8**: 15805.
- 80 Geurts S, Tilly MJ, Arshi B, *et al.* Heart rate variability and atrial fibrillation in the general population: A longitudinal and mendelian randomization study. *Clinical Research in Cardiology* 2023; **112**: 747–58.
- 81 Tegegne BS, Said MA, Ani A, *et al.* Phenotypic but not genetically predicted heart rate variability associated with all-cause mortality. *Communications Biology* 2023; **6**: 1013.
- 82 Shen MJ, Zipes DP. Role of the autonomic nervous system in modulating cardiac arrhythmias. *Circulation Research* 2014; **114**: 1004–21.
- 83 Patel VN, Pierce BR, Bodapati RK, Brown DL, Ives DG, Stein PK. Association of holter-derived heart rate variability parameters with the development of congestive heart failure in the cardiovascular health study. *JACC: Heart Failure* 2017; **5**: 423–31.
- 84 Kaze AD, Yuyun MF, Erqou S, Fonarow GC, Echouffo-Tcheugui JB. Cardiac autonomic neuropathy and risk of incident heart failure among adults with type 2 diabetes. *European Journal of Heart Failure* 2022; **24**: 634–41.
- 85 Ostrowska B, Lind L, Blomström-Lundqvist C. An association between heart rate variability and incident heart failure in an elderly cohort. *Clinical Cardiology* 2024; **47**: e24241.
- 86 Shah SA, Kambur T, Chan C, Herrington DM, Liu K, Shah SJ. Relation of short-term heart rate variability to incident heart failure (from the multi-ethnic study of atherosclerosis). *The American Journal of Cardiology* 2013; **112**: 533–40.
- 87 Boutouyrie P, Chowienczyk P, Humphrey JD, Mitchell GF. Arterial stiffness and cardiovascular risk in hypertension. *Circulation Research* 2021; **128**: 864–86.
- 88 Arshi B, Geurts S, Tilly MJ, *et al.* Heart rate variability is associated with left ventricular systolic, diastolic function and incident heart failure in the general population. *BMC Medicine* 2022; **20**: 91.
- 89 Rose GA, Khaw K-T, Marmot M. Rose's strategy of preventive medicine: The complete original text. Oxford University Press, 2008.
- 90 Schaarup J, Bjerg L, Hansen C, *et al.* Cardiovascular autonomic dysfunction is linked with arterial stiffness across glucose metabolism: The maastricht study. 2024 DOI:10.1101/2024.12.03.24317865.
- 91 Yafei Wu, Xiude Fan, Yingzhou Shi, *et al.* Association of pre-diabetes with the risks of adverse health outcomes and complex multimorbidity: Evidence from population-based studies in the NIS and UK biobank. *BMJ Public Health* 2025; **3**: e001539.

- 92 Schaarup JR, Bjerg L, Hansen CS, *et al.* Cardiovascular autonomic dysfunction precedes cardiovascular disease and all-cause mortality: 11-year follow-up of the ADDITION-PRO study. *medRxiv* 2024; : 2024.12.18.24319131.
- 93 Schaarup JFR, Aggarwal R, Dalsgaard E-M, *et al.* Perception of artificial intelligence-based solutions in healthcare among people with and without diabetes: A cross-sectional survey from the health in central denmark cohort. *Diabetes Epidemiology and Management* 2023; **9**: 100114.
- 94 group S working, ESC Cardiovascular risk collaboration. SCORE2 risk prediction algorithms: New models to estimate 10-year risk of cardiovascular disease in europe. *European Heart Journal* 2021; **42**: 2439–54.
- 95 D'Agostino RB, Vasan RS, Pencina MJ, *et al.* General cardiovascular risk profile for use in primary care. *Circulation* 2008; **117**: 743–53.
- 96 Varga TV, Niss K, Estampador AC, Collin CB, Moseley PL. Association is not prediction: A landscape of confused reporting in diabetes – a systematic review. *Diabetes Research and Clinical Practice* 2020; **170**: 108497.
- 97 Cardoso CRL, Oliveira VAG de, Leite NC, Salles GF. Prognostic importance of cardiovascular autonomic neuropathy on cardiovascular and mortality outcomes in individuals with type 2 diabetes: The rio de janeiro type 2 diabetes cohort. *Diabetes Research and Clinical Practice* 2023; **196**: 110232.
- 98 Bodapati RK, Kizer JR, Kop WJ, Kamel H, Stein PK. Addition of 24-Hour Heart Rate Variability Parameters to the Cardiovascular Health Study Stroke Risk Score and Prediction of Incident Stroke: The Cardiovascular Health Study. *Journal of the American Heart Association* 2017; **6**. DOI:10.1161/JAHA.116.004305.
- 99 Chellappa SL, Gao L, Qian J, *et al.* Daytime eating during simulated night work mitigates changes in cardiovascular risk factors: Secondary analyses of a randomized controlled trial. *Nature Communications* 2025; **16**: 3186.
- 100 Picard M, Tauveron I, Magdasy S, *et al.* Effect of exercise training on heart rate variability in type 2 diabetes mellitus patients: A systematic review and meta-analysis. *PLOS ONE* 2021; **16**: e0251863.
- 101 Bönhof GJ, Strom A, Apostolopoulou M, *et al.* High-intensity interval training for 12 weeks improves cardiovascular autonomic function but not somatosensory nerve function and structure in overweight men with type 2 diabetes. *Diabetologia* 2022; **65**: 1048–57.
- 102 Carnethon MR, Prineas RJ, Temprosa M, *et al.* The association among autonomic nervous system function, incident diabetes, and intervention arm in the diabetes prevention program. *Diabetes Care* 2006; **29**: 914–9.

References

- 103 Nicolaisen SK, Pedersen L, Witte DR, Sørensen HT, Thomsen RW. HbA1c-defined prediabetes and progression to type 2 diabetes in denmark: A population-based study based on routine clinical care laboratory data. *Diabetes Research and Clinical Practice* 2023; **203**. DOI:10.1016/j.diabres.2023.110829.
- 104 Cai X, Liu X, Sun L, et al. Prediabetes and the risk of heart failure: A meta-analysis. *Diabetes, Obesity and Metabolism* 2021; **23**: 1746–53.
- 105 Davies MJ, Aroda VR, Collins BS, et al. Management of hyperglycemia in type 2 diabetes, 2022. A consensus report by the american diabetes association (ADA) and the european association for the study of diabetes (EASD). *Diabetes Care* 2022; **45**: 2753–86.
- 106 Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA guideline for the management of heart failure: A report of the american college of cardiology/american heart association joint committee on clinical practice guidelines. *Circulation* 2022; **145**: e895–1032.
- 107 Mahin Chowdhury, Sarah Nevitt, Aikaterini Eleftheriadou, et al. Cardiac autonomic neuropathy and risk of cardiovascular disease and mortality in type 1 and type 2 diabetes: A meta-analysis. *BMJ Open Diabetes Research & Care* 2021; **9**: e002480.
- 108 Tang Y, Ang L, Jaiswal M, et al. Cardiovascular autonomic neuropathy and risk of kidney function decline in type 1 and type 2 diabetes: Findings from the PERL and ACCORD cohorts. *Diabetes* 2023; **73**: 751–62.
- 109 Niemelä MJ, Airaksinen KE, Huikuri HV. Effect of beta-blockade on heart rate variability in patients with coronary artery disease. *Journal of the American College of Cardiology* 1994; **23**: 1370–7.
- 110 Hadad R, Larsen BS, Weber P, et al. Night-time heart rate variability identifies high-risk people among people with uncomplicated type 2 diabetes mellitus. *Diabetic Medicine* 2021; **38**: e14559.
- 111 Fleischer J, Nielsen R, Laugesen E, Nygaard H, Poulsen PL, Ejskjaer N. Self-monitoring of cardiac autonomic function at home is feasible. *Journal of diabetes science and technology* 2011; **5**: 107–12.
- 112 Glümer C, Carstensen B, Sandbæk A, Lauritzen T, Jørgensen T, Borch-Johnsen K. A danish diabetes risk score for targeted screening: The Inter99 study. *Diabetes Care* 2004; **27**: 727–33.
- 113 Christensen JO, Sandbaek A, Lauritzen T, Borch-Johnsen K. Population-based stepwise screening for unrecognised Type 2 diabetes is ineffective in general practice despite reliable algorithms. *Diabetologia* 2004; **47**: 1566–73.

- 114 Olivarius N de F, Lauritzen T, Beck-Nielsen H, Fog J, Mogensen CE, Working Party appointed by the DNB of H. Co-ordination of diabetes care in the primary and the secondary health care system in denmark. *Diabetic Medicine* 1994; **11**: 123–5.
- 115 Keshet A, Reicher L, Bar N, Segal E. Wearable and digital devices to monitor and treat metabolic diseases. *Nature Metabolism* 2023; **5**: 563–71.
- 116 Kumarathurai P, Anholm C, Larsen BS, *et al.* Effects of liraglutide on heart rate and heart rate variability: A randomized, double-blind, placebo-controlled crossover study. *Diabetes Care* 2016; **40**: 117–24.
- 117 Nelson BW, Allen NB. Accuracy of consumer wearable heart rate measurement during an ecologically valid 24-hour period: Intraindividual validation study. *JMIR Mhealth Uhealth* 2019; **7**: e10828.
- 118 Fuller D, Colwell E, Low J, *et al.* Reliability and validity of commercially available wearable devices for measuring steps, energy expenditure, and heart rate: Systematic review. *JMIR Mhealth Uhealth* 2020; **8**: e18694.
- 119 Davey Smith G, Hemani G. Mendelian randomization: Genetic anchors for causal inference in epidemiological studies. *Human Molecular Genetics* 2014; **23**: R89–98.
- 120 Lash T, VanderWeele TJ, Haneuse S, Rothman K. Modern epidemiology, 4th edition. Wolters Kluwer.

Resume

Denne afhandling undersøgte, om kardiovaskulær autonom dysfunktion er forbundet med kardiovaskulære komplikationer hos personer med normal glukosemetabolisme, prædiabetes og type 2-diabetes. Kardiovaskulær autonom dysfunktion blev målt igennme hjeretrymefvariabilitet (HRV) og standardiserede kardiovaskulære autonome refleksundersøgelser (CARTs). Blandt målingerne gav 24-timers HRV indsigt i den autonome balance og døgnrytmens skift mellem sympathisk og parasympatisk aktivitet. Lavere værdier var konsekvent forbundet med ugunstige kardiovaskulære udfald.

På tværs af tre studier blev autonom dysfunktion forbundet med højere niveauer af arteriel stivhed, en højere forekomst af iskæmiske hjertekarsygedom, hjertesvigt og dødelighed. Sammenhængen med arteriel stivhed blev observeret på tværs af hele spektret af glukosemetabolisme og var særligt forhøjet hos personer med prædiabetes, hvilket antyder, at autonom dysfunktion. På baggrund af dette fokuserede det efterfølgende studie på personer med høj risiko for at udvikle diabetes. I denne population var kardiovaskulær autonom dysfunktion forbundet med højere risiko for kardiovaskulær sygdom, hjertesvigt og dødelighed.

Kardiovaskulær autonom neuropati, defineret gennem abnormale CARTs, var forbundet med en højere risiko for hjertesvigt hos individer med type 2 diabetes, selv hos personer uden symptomer. Disse fund tyder på, at kardiovaskulær autonom dysfunktion kan fungere som en tidlig og uafhængig markør for risikoen for hjertesvigt, især i populationer med risiko for type 2-diabetes.

Afhandlingen konkluderer, at kardiovaskulær autonom dysfunktion er en klinisk relevant risikofaktor, som bør tiltrække større opmærksomhed i både forskning og klinisk praksis. Fremtidige studier bør undersøge, om forbedring af autonom funktion kan reducere kardiovaskulær risiko, og om målinger af autonom dysfunktion kan integreres i eksisterende risikomodeller. Derudover er potentialet i smarture til kontinuerlig overvågning og tidlig opsporing en mulighed, der kan vurderes i både kliniske og generelle populationer.

10. Summary

This dissertation investigated whether cardiovascular autonomic dysfunction is associated with cardiovascular complications in individuals with normal glucose metabolism, prediabetes, and type 2 diabetes. Cardiovascular autonomic dysfunction was assessed using heart rate variability and standardized cardiovascular autonomic reflex tests. Among the measures, 24-hour heart rate variability reflected autonomic balance and circadian shifts in sympathetic and parasympathetic activity. Lower long-term heart rate variability was consistently associated with adverse cardiovascular outcomes.

Across three studies, autonomic dysfunction was linked to higher levels of arterial stiffness, a higher incidence of ischemic cardiovascular events, heart failure, and all-cause mortality. The association with arterial stiffness was observed across the full spectrum of glucose metabolism and was particularly pronounced in individuals with prediabetes, suggesting that autonomic dysfunction may play a role early in the pathophysiological process. Building on this, the subsequent studies focused on individuals at high risk of developing diabetes. In this population, cardiovascular autonomic dysfunction was associated with a higher risk of cardiovascular disease, heart failure, and mortality.

Cardiovascular autonomic neuropathy, as identified through abnormal cardiovascular reflex testing, was associated with a higher risk of heart failure, even in individuals without symptoms. These findings suggest that cardiovascular autonomic dysfunction may serve as an early and independent marker of heart failure risk, particularly in populations at risk of type 2 diabetes.

The dissertation concludes that cardiovascular autonomic dysfunction is a clinically relevant risk factor that warrants further attention in both research and clinical practice. Future studies should explore whether improving autonomic function can reduce cardiovascular risk and whether measures of autonomic dysfunction can be integrated into existing risk models. Additionally, the potential of wearable technologies for continuous monitoring and early detection is a possibility to be evaluated in both clinical and general populations.

A. More results

Some results that wouldn't fit into the main thesis

B. Another appendix

Something else