



Suite 18.05,
Level 18 Centrepont North Mid Valley City,
Lingkaran Syed Putra,
59200 Kuala Lumpur W.P. Kuala Lumpur, Malaysia.

Certificate of Analysis

PRODUCT DETAILS

Name

Batch

Date

Cell Source and Passage

Human Umbilical Cord (Wharton's Jelly), Passage 3

Storage Temperature

4 – 8°C

Viable Cell Count

TEST	SPECIFICATION	TEST RESULTS	REMARKS
Infectious Diseases Screening	HBsAg; HCV Ab; HIV1/2 Ag/Ab; CMV Ab; RPR	Not Detected / Non-Reactive	QUALITY CONTROL APPROVED
Morphology of Cells	Spindle-like cells	Small spindle shaped cells	QUALITY CONTROL APPROVED
Cell Viability	$\geq 90\% \pm 0.1$ of requested cells	Within range	QUALITY CONTROL APPROVED
Sterility	No cloudiness of the test media	Negative	QUALITY CONTROL APPROVED
Bacterial Endotoxin Test (LAL)	Endotoxin sensitivity 0.125 EU/mL	Negative	QUALITY CONTROL APPROVED
Mycoplasma	Mean of absorbance <0.05	Negative	QUALITY CONTROL APPROVED
Immunophenotyping	Positive marker – CD73, CD90, CD105 ($\geq 95\%$) Negative marker – CD11b, CD19, CD34, CD45, HLA-DR ($\leq 2\%$)	Within range	QUALITY CONTROL APPROVED

The cell product delivered by Malaysian Genomics Resource Centre Berhad is manufactured and tested for research purpose only. Product is processed in a facility operating under current Good Manufacturing Practice (cGMP) guidelines in accordance with PIC/S standards.

Authorised Signature:

Quality Control Department
Date:

Powered By:



Malaysian
Genomics
Resource
Centre

Certificate of Analysis

Supporting Information

MORPHOLOGY OF CELLS

Figure 1: Mesenchymal stem cells as viewed under an inverted phase contrast microscope (10x).

STERILITY TEST ^{6, 8}

Aerobic microorganisms are incubated in Soybean Casein Digest Medium at 20°C - 25°C for 14 days.

Facultative microorganisms are incubated in Fluid Thioglycollate Medium at 30°C - 35°C for 14 days.

BACTERIAL ENDOTOXIN TEST (LIMULUS AMOEBOCYTE LYSATE, LAL) TEST ^{3,6}

Negative findings for Gram-negative bacteria in cell culture based on FDA Reference of Standard Endotoxin Guideline.

MYCOPLASMA TEST ^{4,6}

Negative findings for mycoplasma species *M. hyorhinis*, *M. arginini*, *M. fermentans*, *M. orale*, *M. Pirum*, *M. hominis*, *M. salivarium* and *A. laidlawii*.

IMMUNOPHENOTYPING

Marker	Positive			Negative (CD11b, CD19, CD34, CD45, HLA-DR)
	CD73+	CD90+	CD105+	
Result (%)				

Table 1: Flow cytometric analysis was performed to verify the characteristic surface marker expression profile of mesenchymal stem cells.

Powered By:



REFERENCES

1. Dominici, M., Blanc, K. Le, Mueller, I., Slaper-Cortenbach, I., Marini, F.C., Krause, D.S. Deans, R.J., Keating, A., Prockop, D.J., Horwitz, E.M. (2006) Minimal Criteria for Defining Multipotent Mesenchymal Stromal Cells. The International Society for Cellular Therapy Position Statement. Cytotherapy, 8, 4-8. 15-17.

2. Bacterial Endotoxins. European Pharmacopoeia 5.0; <https://gmpua.com/Validation/Method/LAL/EUPHARMACOPDEIA.pdf>

3. US Pharmacopeia <52> Mycoplasma Tests

4. Bobis, S., Jarocha, D., Majka, M. (2006). Mesenchymal stem Cells: Characteristics and Clinical Applications. Folia Histochemica et Cytobiologica. 44(4): 215-230.

5. USP Method. Sterilization of Medical Devices, Microbiological Methods.

6. ICH Harmonized Tripartite Guideline, Specifications: Test Procedures and Acceptance Criteria for Biotechnological / Biological Products Q6B.

7. Muntión, S., Sánchez-Guilló, F.M., Carrancio, S., Villardón, E., López, O., Díez-Campelo, M., San Miguel, J.F. and del Cañizo, M.C. (2012). Optimisation of mesenchymal stromal cells karyotyping analysis: implications for clinical use. Transfusion Medicine. 22, 122-127.

8. US Pharmacopeia <71> Sterility Test