Circle — Definition, Forms, Reality Check

NOTEBOOK FLOW: read straight down the page, then continue on the next sheet.

LESSON 01 / 08

Definition: fixed distance from a centre

A circle with centre $C = (h, k)$ and radius $r \geq 0$ is the locus

$$(x - h)^2 + (y - k)^2 = r^2.$$

For $r > 0$ this is a proper circle; $r = 0$ is the single point C.

SHORT DERIVATION

Complete squares in the general equation

For $x^2 + y^2 + 2gx + 2fy + c = 0$, group the coordinate terms:

$$(x + g)^2 + (y + f)^2 = g^2 + f^2 - c.$$

Thus the centre is $(-g, -f)$ and $r^2 = g^2 + f^2 - c$. The equal coefficients of x^2, y^2, with no xy term, are essential for a circle in orthogonal Cartesian axes.

Three cases—do not call every equation a circle

- $g^2 + f^2 - c > 0$: a real circle of positive radius.
- $g^2 + f^2 - c = 0$: a point-circle (one real point).
- $g^2 + f^2 - c < 0$: no real points.

WORKED EXAMPLE

Centre, radius, then check

$x^2 + y^2 - 2x - 4y - 4 = 0$ gives $g = -1, f = -2, c = -4$, centre $(1, 2)$ and $r^2 = 1 + 4 + 4 = 9$, so $r = 3$. Check $(1, 5)$: it is distance 3 from $(1, 2)$ and satisfies the equation.

COMMON TRAP

Radius is the nonnegative square root

If a formula yields $r^2 < 0$, it is not a real circle. If $r^2 = 0$, do not describe it as a circle with positive circumference. A negative value cannot be repaired by dropping a sign.

LESSON 02 / 08

Three independent conditions determine a circle

A general circle has equation $x^2+y^2+Dx+Ey+F=0$ with three unknown coefficients. Substituting three noncollinear points gives a linear system for D, E, F. If the points are collinear, no proper circle passes through all three.

WORKED EXAMPLE

Through the origin and two unit intercepts

Through $(0,0), (1,0), (0,1)$: the origin gives $F=0$, then $1+D=0$ and $1+E=0$. The circle is $x^2+y^2-x-y=0$, centre $(1/2, 1/2)$, radius $1/\sqrt{2}$.

Diameter endpoints give a dot-product equation

For diameter endpoints $A=(x_1,y_1)$ and $B=(x_2,y_2)$, a point $P=(x,y)$ lies on the circle exactly when $\angle APB = 90^\circ$ (Thales' theorem). Therefore

$$(P-A)\cdot(P-B)=0,$$

or explicitly

$$(x-x_1)(x-x_2)+(y-y_1)(y-y_2)=0.$$

The centre is the midpoint of AB and the radius is $AB/2$.

WORKED EXAMPLE

Diameter through two points

For $A=(1,2)$, $B=(5,8)$, $(x-1)(x-5)+(y-2)(y-8)=0$, *i.e.* $(x-3)^2+(y-5)^2=13$. Both endpoints satisfy the diameter equation.

Parametric equations

For centre (h,k) and radius r,

$$x=h+r\cos t, \qquad y=k+r\sin t, \qquad 0\le t<2\pi.$$

The angle parameter traces every point once except the equivalent endpoint at 2π.

A rational parameter is

$$x=h+r\frac{1-u^2}{1+u^2}, \qquad y=k+r\frac{2u}{1+u^2}, \qquad u\in\mathbb{R}.$$

It traces the circle except $(h-r,k)$; that missing point is the limit $|u|\to\infty$. Include it separately when a problem asks for every point.

Determine a Circle and Parametrize It (cont.)

NOTEBOOK FLOW: read straight down the page, then continue on the next sheet.

COMMON TRAP

Parameter is not arc length

In the trigonometric parametrization, t is the polar angle about the centre. In the rational form, $u = \tan(t/2)$ and the omitted point comes from the half-angle coordinate's pole.

LESSON 03 / 08

Centre-to-line distance classifies the intersection

Let a circle have centre $C=(h,k)$, radius r, and let the line be $ax+by+c=0$. The distance from C to the line is

$$d=\frac{|ah+bk+c|}{\sqrt{a^2+b^2}}.$$

- $d<r$: two intersection points (a secant).
- $d=r$: one repeated intersection (a tangent).
- $d>r$: no real intersection.

SHORT DERIVATION

Why the chord length is twice a right-triangle leg

The perpendicular from the centre bisects a chord. If its length is ℓ, the right triangle has hypotenuse r and one leg d, hence $(\ell/2)^2=r^2-d^2$ and

$$\ell=2\sqrt{r^2-d^2}.$$


This is real precisely in the secant/tangent cases $d\le r$.

WORKED EXAMPLE

A secant, not merely a line equation

The line $x+y=5$ is at distance $5/\sqrt{2}$ from the centre $(0,0)$ of $x^2+y^2=25$. Since this is less than 5, it cuts a chord of length $2\sqrt{25-25/2}=5\sqrt{2}$.

Centre-to-line distance sets the chord length



Solve the intersection only after classifying it

Parametrize the line by $P(t) = P_0 + t(-b, a)$, where P_0 is any point on it. Substitute in $(x-h)^2 + (y-k)^2 = r^2$: a quadratic in t results. Its discriminant is positive, zero or negative exactly when $d < r$, $d = r$ or $d > r$.

COMMON TRAP

Tangency is a repeated root

A tangent line meets the circle in one geometric point but gives a double root after substitution. Do not infer "no intersection" from a zero discriminant.

LESSON 04 / 08

Tangent: perpendicular to the radius

For $(x-h)^2+(y-k)^2=r^2$ and a point $T=(x_1,y_1)$ on the circle, the tangent is

$$(x_1-h)(x-h)+(y_1-k)(y-k)=r^2.$$

The normal is the line through T and the centre (h,k).

SHORT DERIVATION

The tangent equation is a dot-product condition

The radius vector $T-C$ is normal to the tangent. Thus X is on the tangent exactly when $(T-C)\cdot(X-T)=0$. Rearranging gives $(T-C)\cdot(X-C)=|T-C|^2=r^2$.

For $x^2+y^2+2gx+2fy+c=0$, the same tangent is

$$xx_1+yy_1+g(x+x_1)+f(y+y_1)+c=0.$$


This formula assumes T lies on the circle; the "$T=0$" replacement is a mnemonic, not a substitute for checking incidence.

WORKED EXAMPLE

Unit-free tangent check

For $x^2+y^2=25$ at $T=(3,4)$, the tangent is $3x+4y=25$. It contains T, and its normal vector $(3,4)$ is the radius direction.

The radius is normal to the tangent



Tangent and Normal at a Point (cont.)

NOTEBOOK FLOW: read straight down the page, then continue on the next sheet.

COMMON TRAP

A tangent is attached to a point on the circle

If a proposed point is not on the circle, the at-point tangent formula does not apply. From an external point there are generally two tangents; derive them as lines through that point with discriminant zero or use the polar/chord-of-contact equation.

LESSON 05 / 08

Power records inside, on, or outside

For circle centre $C=(h,k)$ and radius r, define

$$\text{Pow}(P)=|P-C|^2-r^2.$$

It is negative inside, zero on, and positive outside. In the general equation $x^2+y^2+2gx+2fy+c=0$, the power at (x_0,y_0) is the left-hand side evaluated there.

SHORT DERIVATION

Tangent length from a right triangle

If P is outside and T is a tangent point, $CT\perp PT$. Therefore

$$PT^2=PC^2-CT^2=PC^2-r^2=\text{Pow}(P).$$

So each tangent from P has length $\sqrt{\text{Pow}(P)}$.

WORKED EXAMPLE

Tangent length is not centre distance

From $P=(5,5)$ to $x^2+y^2=9$, power is $25+25-9=41$. Each tangent has length $\sqrt{41}$, whereas $PC=\sqrt{50}$.

Chord of contact / polar of an external point

For a circle centred at the origin, $x^2+y^2=r^2$, the two tangent points from $P=(p,q)$ lie on the line

$$px+qy=r^2.$$

This line is the **chord of contact**. Its intersection with the circle gives the two contact points. For a translated circle, first use coordinates relative to its centre: $(P-C)\cdot(X-C)=r^2$.

WORKED EXAMPLE

Find both contacts by one line and one circle

From $P=(5,0)$ to $x^2+y^2=9$, the chord of contact is $5x=9$, so $x=9/5$. Then $y^2=9-81/25=144/25$, giving contacts $(9/5,\pm 12/5)$. The tangent length is $\sqrt{25-9}=4$.

COMMON TRAP

The polar formula needs the same circle centre

For a circle not centred at the origin, replace P and X by their vectors from the centre. Forgetting this translation shifts the chord of contact.

LESSON 06 / 08

Subtract normalized circle equations

Write $S_i = x^2 + y^2 + D_i x + E_i y + F_i$ for $i = 1, 2$; the corresponding circles are $S_i = 0$. Their power functions are $S_1(x, y)$ and $S_2(x, y)$. Equal power means $S_1 - S_2 = 0$; the quadratic terms cancel, leaving a line (unless the circles are concentric or identical).

SHORT DERIVATION

Why the line is perpendicular to the centres

The centre difference is parallel to $(D_1 - D_2, E_1 - E_2)$, the normal vector of $S_1 - S_2 = 0$. Thus the radical axis is perpendicular to the line of centres. When the real circles intersect, its points are exactly their common chord.

WORKED EXAMPLE

Subtract, then test equal powers

For $x^2 + y^2 = 9$ and $x^2 + y^2 - 4x + 6y - 3 = 0$, subtract to get $2x - 3y - 3 = 0$. At $(0, -1)$ both power values equal -8, checking the line's equal-power meaning.

Three circles and the radical centre

The radical axes of three circles are the pairwise equal-power lines. If the circles are not identical and the axes are not parallel, solve two axes to get their common point; equality of the third pair follows by subtraction. Degenerate cases need care: concentric unequal circles have no finite equal-power locus, identical circles have equal power everywhere, and parallel axes need not meet in a finite radical centre.

COMMON TRAP

A radical axis need not be a real common chord

Two nonconcentric circles can be disjoint or nested. Their equal-power line still exists, but it contains real points only as a locus of equal powers—not points on both circles. Call it a common chord only when the circles actually intersect.

LESSON 07 / 08

Compare centre distance and radii

For centres distance d and radii $r_1, r_2 > 0$:

Condition	Real intersection
$d > r_1 + r_2$	none; externally separate
$d = r_1 + r_2$	one; externally tangent
$\lvert r_1 - r_2 \rvert < d < r_1 + r_2$	two points
$d = \lvert r_1 - r_2 \rvert > 0$	one; internally tangent
$0 < d < \lvert r_1 - r_2 \rvert$	none; one strictly inside the other
$d = 0$	concentric: compare radii; identical if radii agree

The zero-radius point-circle is handled separately by testing whether its point lies on the other circle.

Efficient order of attack

1. Complete squares and identify centres/radii.
2. For a line, use centre-to-line distance before solving intersections.
3. For a tangent, check the point lies on the circle and use the radius as normal.
4. For a common chord, subtract normalized equations first.
5. For two circles, compare d with $r_1 + r_2$ and $\lvert r_1 - r_2 \rvert$.

WORKED EXAMPLE

Two equal circles

$(x-0)^2 + y^2 = 4$ and $(x-3)^2 + y^2 = 4$ have $d = 3$, $r_1 + r_2 = 4$, and $\lvert r_1 - r_2 \rvert = 0$. Hence they meet twice. Subtracting gives radical axis $x = 3/2$; substituting yields $y = \pm\sqrt{7}/2$.

ASSUMPTIONS & CONVENTIONS

JEE boundary


The official circle bullet includes circle forms, tangent, normal, chord, parametric equations and intersection with a straight line. Radical axes, power and two-circle position are useful direct applications; the official scope is not a promise to include every theorem about circles.

LESSON 08 / 08

A perpendicular from the centre bisects a chord

On $x^2 + y^2 = 25$, the vertical line $x = 3$ meets the circle at $(3, 4)$ and $(3, -4)$. The chord length is 8. More generally, a chord at distance d from the centre of a radius-r circle has length $2\sqrt{r^2 - d^2}$, provided $|d| \le r$.

Chord length follows from a right triangle



READ THE FIGURE
The right triangle has hypotenuse 5 and one leg 3, so half the chord is 4. The full chord is 2*4 = 8.

The same right triangle classifies a line: $d < r$ gives two intersections, $d = r$ gives tangency, and $d > r$ gives no real intersection. This is often shorter than forming a quadratic discriminant.


The tangent is perpendicular to the radius

At $P=(3,4)$ on $x^2+y^2=25$, the radius slope is $4/3$, so the tangent slope is $-3/4$. Its equation is

$$3(x-3)+4(y-4)=0 \quad\Longrightarrow\quad 3x+4y=25.$$

The normal at P is the line through O and P, namely $4x-3y=0$.

Tangent and normal at P(3,4) on x^2 + y^2 = 25



READ THE FIGURE


The tangent direction is (4,-3), whose dot product with the radius direction (3,4) is zero. The tangent touches the circle at exactly one point.

For $x^2+y^2+2gx+2fy+c=0$, the tangent at an on-circle point (x_1,y_1) uses the replacement rule $x^2\mapsto xx_1$, $y^2\mapsto yy_1$, $2x\mapsto x+x_1$, and $2y\mapsto y+y_1$. Verify first that the point lies on the circle.

Power of a point unifies secants and tangents

For external point P and a circle with centre O and radius r, the power is $PO^2 - r^2$. If a secant through P meets the circle at A and B, and a tangent touches at T, then $PA \cdot PB = PT^2 = PO^2 - r^2$ with directed lengths in the general formulation.

The same power appears on a secant and a tangent



READ THE FIGURE


PA = 3 and PB = 13 give PA*PB = 39. Also PO^2 - r^2 = 64 - 25 = 39, so PT = sqrt(39).

The power is positive outside, zero on, and negative inside the circle. Tangent length is real only for an external or on-circle point.

Radical axis: equal power, common chord

For $C_1 : x^2 + y^2 = 9$ and $C_2 : (x-4)^2 + y^2 = 16$, subtracting the equations gives $x = 9/8$. This is the radical axis; when the circles intersect, it is their common chord.

The radical axis x = 9/8 is the common chord



READ THE FIGURE

Subtracting the equations gives the chord line immediately. Substitution is needed only when the actual intersection coordinates are requested.

Position and number of common points


Centre distance condition	Geometric position	Common points
d > r + R	separate externally	0
d = r + R	touch externally	1
|r-R| < d < r+R	intersect	2
d = |r-R|, d > 0	touch internally	1
0 <= d < |r-R|	one inside the other	0

READ THE FIGURE

Use d = distance between centres and r, R > 0. Identical coincident circles are a separate degenerate case with infinitely many common points.

Circle strategy

Choose the shortest circle tool that preserves the geometry



READ THE FIGURE

Many circle questions collapse to one right triangle or one subtraction. Solving a large quadratic first often hides the simple geometric relationship.

WORKED EXAMPLE

A rapid chord check

For $x^2 + y^2 = 25$ and $x = 3$, the centre-line distance is 3, so the chord is $2\sqrt{25 - 9} = 8$. Direct substitution gives $y = \pm 4$, whose separation is also 8. The geometric and algebraic methods agree.