

# Assignment 3

Suraj - CS20BTECH11050

Download all python codes from

<https://github.com/Suraj11050/Assignments-AI1103/blob/main/Assignment%203/Assignment3.py>

Download Latex-tikz codes from

<https://github.com/Suraj11050/Assignments-AI1103/blob/main/Assignment%203/Assignment3.tex>

## 1 GATE 2009 (MA) PROBLEM 16

Let F, G and H be pair wise independent events such that  $\Pr(F) = \Pr(G) = \Pr(H) = \frac{1}{3}$  and  $\Pr(F \cap G \cap H) = \frac{1}{4}$  Then the probability that at least one event among F, G and H occurs is

- (A)  $\frac{11}{12}$     (B)  $\frac{7}{12}$     (C)  $\frac{5}{12}$     (D)  $\frac{3}{4}$

## 2 SOLUTION

If two Events  $X_1$  and  $X_2$  are independent then

$$\Pr(X_1 X_2) = \Pr(X_1) \times \Pr(X_2) \quad (2.0.1)$$

Using equation (2.0.1) we get the following results

$$\Pr(FG) = \frac{1}{3} \times \frac{1}{3} = \frac{1}{9} \quad (2.0.2)$$

$$\Pr(GH) = \frac{1}{3} \times \frac{1}{3} = \frac{1}{9} \quad (2.0.3)$$

$$\Pr(HF) = \frac{1}{3} \times \frac{1}{3} = \frac{1}{9} \quad (2.0.4)$$

At least one event among F, G, H should occur is  $\Pr(F + G + H)$  from Principal of inclusion and exclusion it is calculated using random variable as

$$\begin{aligned} \Pr(F + G + H) &= \Pr(F) + \Pr(G) + \Pr(H) \\ &\quad - (\Pr(FG) + \Pr(GH) + \Pr(HF)) + \Pr(FGH) \end{aligned} \quad (2.0.5)$$

Substituting above results from equation (2.0.2), (2.0.3), (2.0.4) in equation (2.0.5)

$$\Pr(F + G + H) = 3\left(\frac{1}{3}\right) - 3\left(\frac{1}{9}\right) + \frac{1}{4}$$

$$\therefore \Pr(F + G + H) = \frac{11}{12}$$

Hence Probability that at least one event among F, G, H occurs is  $\Pr(F + G + H) = \frac{11}{12}$  and correct answer is **Option (A)**

But we know that

$$(FG) = (FGH) + (FGH') \quad (2.0.6)$$

$$\Pr(FG) = \Pr(FGH) + \Pr(FGH') \quad (2.0.7)$$

$$\therefore \Pr(FG) \geq \Pr(FGH) \quad (2.0.8)$$

In the given question

$$\Pr(FGH) = \frac{1}{4} \quad (2.0.9)$$

$$\Pr(FG) = \frac{1}{9} \quad (2.0.10)$$

$$\Pr(FG) < \Pr(FGH) \quad (2.0.11)$$

Which is not possible according to equation (2.0.8). Similar case with  $\Pr(GH)$  and  $\Pr(HF)$

Some of the probabilities turnout to be negative like

$$\Pr(F'GH) = \frac{1}{9} - \frac{1}{4} = -\frac{5}{36} \quad (2.0.12)$$

$$\Pr(FG'H) = \frac{1}{9} - \frac{1}{4} = -\frac{5}{36} \quad (2.0.13)$$

$$\Pr(FGH') = \frac{1}{9} - \frac{1}{4} = -\frac{5}{36} \quad (2.0.14)$$

Probability  $P \in [0, 1]$  but because of the data in the question some of the probabilities turn out to be negative. Therefore **Question is incorrect**