

# USAMO 2024 Solution Notes

Evan Chen 《陳誼廷》

23 July 2025

This is a compilation of solutions for the 2024 USAMO. The ideas of the solution are a mix of my own work, the solutions provided by the competition organizers, and solutions found by the community. However, all the writing is maintained by me.

These notes will tend to be a bit more advanced and terse than the “official” solutions from the organizers. In particular, if a theorem or technique is not known to beginners but is still considered “standard”, then I often prefer to use this theory anyways, rather than try to work around or conceal it. For example, in geometry problems I typically use directed angles without further comment, rather than awkwardly work around configuration issues. Similarly, sentences like “let  $\mathbb{R}$  denote the set of real numbers” are typically omitted entirely.

Corrections and comments are welcome!

## Contents

<b>0 Problems</b>	<b>2</b>
<b>1 Solutions to Day 1</b>	<b>3</b>
1.1 USAMO 2024/1, proposed by Luke Robitaille . . . . .	3
1.2 USAMO 2024/2, proposed by Rishabh Das . . . . .	4
1.3 USAMO 2024/3, proposed by Krit Boonsiriseth . . . . .	7
<b>2 Solutions to Day 2</b>	<b>10</b>
2.1 USAMO 2024/4, proposed by Rishabh Das . . . . .	10
2.2 USAMO 2024/5, proposed by Anton Trygub . . . . .	12
2.3 USAMO 2024/6, proposed by Titu Andreescu, Gabriel Dospinescu . . . .	17

## §0 Problems

- Find all integers  $n \geq 3$  such that the following property holds: if we list the divisors of  $n!$  in increasing order as  $1 = d_1 < d_2 < \cdots < d_k = n!$ , then we have

$$d_2 - d_1 \leq d_3 - d_2 \leq \cdots \leq d_k - d_{k-1}.$$

- Let  $S_1, S_2, \dots, S_{100}$  be finite sets of integers whose intersection is not empty. For each non-empty  $T \subseteq \{S_1, S_2, \dots, S_{100}\}$ , the size of the intersection of the sets in  $T$  is a multiple of  $|T|$ . What is the smallest possible number of elements which are in at least 50 sets?
- Let  $(m, n)$  be positive integers with  $n \geq 3$  and draw a regular  $n$ -gon. We wish to triangulate this  $n$ -gon into  $n - 2$  triangles, each colored one of  $m$  colors, so that each color has an equal sum of areas. For which  $(m, n)$  is such a triangulation and coloring possible?
- Let  $m$  and  $n$  be positive integers. A circular necklace contains  $mn$  beads, each either red or blue. It turned out that no matter how the necklace was cut into  $m$  blocks of  $n$  consecutive beads, each block had a distinct number of red beads. Determine, with proof, all possible values of the ordered pair  $(m, n)$ .
- Point  $D$  is selected inside acute triangle  $ABC$  so that  $\angle DAC = \angle ACB$  and  $\angle BDC = 90^\circ + \angle BAC$ . Point  $E$  is chosen on ray  $BD$  so that  $AE = EC$ . Let  $M$  be the midpoint of  $BC$ . Show that line  $AB$  is tangent to the circumcircle of triangle  $BEM$ .
- Let  $n > 2$  be an integer and let  $\ell \in \{1, 2, \dots, n\}$ . A collection  $A_1, \dots, A_k$  of (not necessarily distinct) subsets of  $\{1, 2, \dots, n\}$  is called  $\ell$ -large if  $|A_i| \geq \ell$  for all  $1 \leq i \leq k$ . Find, in terms of  $n$  and  $\ell$ , the largest real number  $c$  such that the inequality

$$\sum_{i=1}^k \sum_{j=1}^k x_i x_j \frac{|A_i \cap A_j|^2}{|A_i| \cdot |A_j|} \geq c \left( \sum_{i=1}^k x_i \right)^2$$

holds for all positive integer  $k$ , all nonnegative real numbers  $x_1, x_2, \dots, x_k$ , and all  $\ell$ -large collections  $A_1, A_2, \dots, A_k$  of subsets of  $\{1, 2, \dots, n\}$ .

## §1 Solutions to Day 1

### §1.1 USAMO 2024/1, proposed by Luke Robitaille

Available online at <https://aops.com/community/p30216459>.

#### Problem statement

Find all integers  $n \geq 3$  such that the following property holds: if we list the divisors of  $n!$  in increasing order as  $1 = d_1 < d_2 < \cdots < d_k = n!$ , then we have

$$d_2 - d_1 \leq d_3 - d_2 \leq \cdots \leq d_k - d_{k-1}.$$

The answer is  $n \in \{3, 4\}$ . These can be checked by listing all the divisors:

- For  $n = 3$  we have  $(1, 2, 3, 6)$ .
- For  $n = 4$  we have  $(1, 2, 3, 4, 6, 8, 12, 24)$ .

We prove these are the only ones.

The numbers  $5 \leq n \leq 12$  all fail because:

- For  $n = 5$  we have  $20 - 15 > 24 - 20$ .
- For  $n = 6$  we have  $120 - 90 > 144 - 20$ .
- For  $7 \leq n \leq 12$  we have because  $14 - 12 > 15 - 14$  (and  $13 \nmid n!$ ).

Now assume  $n \geq 13$ . In that case, we have

$$\left\lfloor \frac{n}{2} \right\rfloor^2 - 1 \geq 2n.$$

So by Bertrand postulate, we can find a prime  $p$  such that

$$n < p < \left\lfloor \frac{n}{2} \right\rfloor^2 - 1.$$

However, note that

$$\left\lfloor \frac{n}{2} \right\rfloor^2 - 1 = \left( \left\lfloor \frac{n}{2} \right\rfloor - 1 \right) \left( \left\lfloor \frac{n}{2} \right\rfloor + 1 \right), \quad \left\lfloor \frac{n}{2} \right\rfloor^2$$

are consecutive integers both dividing  $n!$  (the latter number divides  $\left\lfloor \frac{n}{2} \right\rfloor \cdot (2 \left\lfloor \frac{n}{2} \right\rfloor)$ ). So the prime  $p$  causes the desired contradiction.

**§1.2 USAMO 2024/2, proposed by Rishabh Das**

Available online at <https://aops.com/community/p30216494>.

**Problem statement**

Let  $S_1, S_2, \dots, S_{100}$  be finite sets of integers whose intersection is not empty. For each non-empty  $T \subseteq \{S_1, S_2, \dots, S_{100}\}$ , the size of the intersection of the sets in  $T$  is a multiple of  $|T|$ . What is the smallest possible number of elements which are in at least 50 sets?

The answer is  $50 \binom{100}{50}$ .

¶ **Rephrasing (cosmetic translation only, nothing happens yet).** We encode with binary strings  $v \in \mathbb{F}_2^{100}$  of length 100. Write  $v \subseteq w$  if  $w$  has 1's in every component  $v$  does, and let  $|v|$  denote the number of 1's in  $v$ .

Then for each  $v$ , we let  $f(v)$  denote the number of elements  $x \in \bigcup S_i$  such that  $x \in S_i \iff v_i = 1$ . For example,

- $f(1 \dots 1)$  denotes  $|\bigcap_1^{100} S_i|$ , so we know  $f(1 \dots 1) \equiv 0 \pmod{100}$ .
- $f(1 \dots 10)$  denotes the number of elements in  $S_1$  through  $S_{99}$  but not  $S_{100}$  so we know that  $f(1 \dots 1) + f(1 \dots 10) \equiv 0 \pmod{99}$ .
- ... And so on.

So the problem condition means that  $f(v)$  translates to the statement

$$P(u) : \quad |u| \text{ divides } \sum_{v \supseteq u} f(v)$$

for any  $u \neq 0 \dots 0$ , plus one extra condition  $f(1 \dots 1) > 0$ . And the objective function is to minimize the quantity

$$A := \sum_{|v| \geq 50} f(v).$$

So the problem is transformed into an system of equations over  $\mathbb{Z}_{\geq 0}$  (it's clear any assignment of values of  $f(v)$  can be translated to a sequence  $(S_1, \dots, S_{100})$  in the original notation).

Note already that:

**Claim —** It suffices to assign  $f(v)$  for  $|v| \geq 50$ .

*Proof.* If we have found a valid assignment of values to  $f(v)$  for  $|v| \geq 50$ , then we can always arbitrarily assign values of  $f(v)$  for  $|v| < 50$  by downwards induction on  $|v|$  to satisfy the divisibility condition (without changing  $M$ ).  $\square$

Thus, for the rest of the solution, we altogether ignore  $f(v)$  for  $|v| < 50$  and only consider  $P(u)$  for  $|u| \geq 50$ .

¶ **Construction.** Consider the construction

$$f_0(v) = 2|v| - 100.$$

This construction is valid since if  $|u| = 100 - k$  for  $k \leq 50$  then

$$\begin{aligned} \sum_{v \supseteq u} f_0(v) &= \binom{k}{0} \cdot 100 + \binom{k}{1} \cdot 98 + \binom{k}{2} \cdot 96 + \cdots + \binom{k}{k} \cdot (100 - 2k) \\ &= (100 - k) \cdot 2^k = |u| \cdot 2^k \end{aligned}$$

is indeed a multiple of  $|u|$ , hence  $P(u)$  is true. In that case, the objective function is

$$A = \sum_{i=50}^{100} \binom{100}{i} (2i - 100) = 50 \binom{100}{50}$$

as needed.

**Remark.** This construction is the “easy” half of the problem because it coincides with what you get from a greedy algorithm by downwards induction on  $|u|$  (equivalently, induction on  $k = 100 - |u| \geq 0$ ). To spell out the first three steps,

- We know  $f(1 \dots 1)$  is a nonzero multiple of 100, so it makes sense to guess  $f(1 \dots 1) = 100$ .
- Then we have  $f(1 \dots 10) + 100 \equiv 0 \pmod{99}$ , and the smallest multiple of 99 which is at least 100 is 198. So it makes sense to guess  $f(1 \dots 10) = 98$ , and similarly guess  $f(v) = 98$  whenever  $|v| = 99$ .
- Now when we consider, say  $v = 1 \dots 100$  with  $|v| = 98$ , we get

$$f(1 \dots 100) + \underbrace{f(1 \dots 101)}_{=98} + \underbrace{f(1 \dots 110)}_{=98} + \underbrace{f(1 \dots 111)}_{=100} \equiv 0 \pmod{98}.$$

we obtain  $f(1 \dots 100) \equiv 96 \pmod{98}$ . That makes  $f(1 \dots 100) = 96$  a reasonable guess.

Continuing in this way gives the construction above.

¶ **Proof of bound.** We are going to use a smoothing argument: if we have a general working assignment  $f$ , we will mold it into  $f_0$ .

We define a **push-down on**  $v$  as the following operation:

- Pick any  $v$  such that  $|v| \geq 50$  and  $f(v) \geq |v|$ .
- Decrease  $f(v)$  by  $|v|$ .
- For every  $w$  such that  $w \subseteq v$  and  $|w| = |v| - 1$ , increase  $f(w)$  by 1.

**Claim —** Apply a push-down preserves the main divisibility condition. Moreover, it doesn't change  $A$  unless  $|v| = 50$ , where it decreases  $A$  by 50 instead.

*Proof.* The statement  $P(u)$  is only affected when  $u \subseteq v$ : to be precise, one term on the right-hand side of  $P(u)$  decreases by  $|v|$ , while  $|v| - |u|$  terms increase by 1, for a net change of  $-|u|$ . So  $P(u)$  still holds.

To see  $A$  doesn't change for  $|v| > 50$ , note  $|v|$  terms increase by 1 while one term decreases by  $-|v|$ . When  $|v| = 50$ , only  $f(v)$  decreases by 50.  $\square$

Now, given a valid assignment, we can modify it as follows:

- First apply pushdowns on  $1 \dots 1$  until  $f(1 \dots 1) = 100$ ;
- Then we may apply pushdowns on each  $v$  with  $|v| = 99$  until  $f(v) < 99$ ;
- Then we may apply pushdowns on each  $v$  with  $|v| = 98$  until  $f(v) < 98$ ;
- ...and so on, until we have  $f(v) < 50$  for  $|v| = 50$ .

Hence we get  $f(1 \dots 1) = 100$  and  $0 \leq f(v) < |v|$  for all  $50 \leq |v| \leq 100$ . However, by downwards induction on  $|v| = 99, 98, \dots, 50$ , we also have

$$f(v) \equiv f_0(v) \pmod{|v|} \implies f(v) = f_0(v)$$

since  $f_0(v)$  and  $f(v)$  are both strictly less than  $|v|$ . So in fact  $f = f_0$ , and we're done.

**Remark.** The fact that push-downs actually don't change  $A$  shows that the equality case we described is far from unique: in fact, we could have made nearly arbitrary sub-optimal decisions during the greedy algorithm and still ended up with an equality case. For a concrete example, the construction

$$f(v) = \begin{cases} 500 & |v| = 100 \\ 94 & |v| = 99 \\ 100 - 2|v| & 50 \leq |v| \leq 98 \end{cases}$$

works fine as well (where we arbitrarily chose 500 at the start, then used the greedy algorithm thereafter).

**§1.3 USAMO 2024/3, proposed by Krit Boonsiriseth**

Available online at <https://aops.com/community/p30216513>.

**Problem statement**

Let  $(m, n)$  be positive integers with  $n \geq 3$  and draw a regular  $n$ -gon. We wish to triangulate this  $n$ -gon into  $n - 2$  triangles, each colored one of  $m$  colors, so that each color has an equal sum of areas. For which  $(m, n)$  is such a triangulation and coloring possible?

The answer is if and only if  $m$  is a proper divisor of  $n$ .

Throughout this solution, we let  $\omega = \exp(2\pi i/n)$  and let the regular  $n$ -gon have vertices  $1, \omega, \dots, \omega^{n-1}$ . We cache the following frequent calculation:

**Lemma**

The triangle with vertices  $\omega^k, \omega^{k+a}, \omega^{k+b}$  has signed area


$$T(a, b) := \frac{(\omega^a - 1)(\omega^b - 1)(\omega^{-b} - \omega^{-a})}{4i}.$$

*Proof.* Rotate by  $\omega^{-k}$  to assume WLOG that  $k = 0$ . Apply complex shoelace to the triangles with vertices  $1, \omega^a, \omega^b$  to get

$$\frac{i}{4} \det \begin{bmatrix} 1 & 1 & 1 \\ \omega^a & \omega^{-a} & 1 \\ \omega^b & \omega^{-b} & 1 \end{bmatrix} = \frac{i}{4} \det \begin{bmatrix} 0 & 0 & 1 \\ \omega^a - 1 & \omega^{-a} - 1 & 1 \\ \omega^b - 1 & \omega^{-b} - 1 & 1 \end{bmatrix}$$

which equals the above.  $\square$

¶ **Construction.** It suffices to actually just take all the diagonals from the vertex 1, and then color the triangles with the  $m$  colors in cyclic order. For example, when  $n = 9$  and  $m = 3$ , a coloring with red, green, blue would be:



To see this works one can just do the shoelace calculation: fix a residue  $r \bmod m$  corresponding to one of the colors. Then

$$\begin{aligned}
\sum_{\substack{0 \leq j < n \\ j \equiv r \pmod m}} \text{Area}(\omega^0, \omega^j, \omega^{j+1}) &= \sum_{\substack{0 \leq j < n \\ j \equiv r \pmod m}} T(j, j+1) \\
&= \sum_{\substack{0 \leq j < n \\ j \equiv r \pmod m}} \frac{(\omega^j - 1)(\omega^{j+1} - 1)(\omega^{-(j+1)} - \omega^{-j})}{4i} \\
&= \frac{\omega - 1}{4i} \sum_{\substack{0 \leq j < n \\ j \equiv r \pmod m}} (\omega^{-j} - 1)(\omega^j - \omega^{-1}) \\
&= \frac{\omega - 1}{4i} \left( \frac{n}{m} (1 + \omega^{-1}) + \sum_{\substack{0 \leq j < n \\ j \equiv r \pmod m}} (\omega^{-1-j} - \omega^j) \right).
\end{aligned}$$

(We allow degenerate triangles where  $j \in \{0, m-1\}$  with area zero to simplify the notation above.)

However, if  $m$  is a proper divisor of  $n$ , then  $\sum_j \omega^j = \omega^r (1 + \omega^m + \omega^{2m} + \dots + \omega^{n-m}) = 0$ . For the same reason,  $\sum_j \omega^{-1-j} = 0$ . So the inner sum vanishes, and the total area of this color equals

$$\sum_{\substack{0 \leq j < n \\ j \equiv r \pmod m}} \text{Area}(\omega^0, \omega^j, \omega^{j+1}) = \frac{n}{m} \frac{(\omega - 1)(\omega^{-1} + 1)}{4i} = \frac{n}{m} \cdot \frac{\omega - \omega^{-1}}{4i}.$$

Because the right-hand side does not depend on the residue  $r$ , this shows all colors have equal area.

¶ **Proof of necessity.** It's obvious that  $m < n$  (in fact  $m \leq n-2$ ). So we focus on just showing  $m \mid n$ .

Repeating the same calculation as above, we find that if there was a valid triangulation and coloring, the total area of each color would equal

$$S := \frac{n}{m} \cdot \frac{\omega - \omega^{-1}}{4i}.$$

However:

**Claim —** The number  $4i \cdot S$  is not an algebraic integer when  $m \nmid n$ .

*Proof.* This is easiest to see if one knows the advanced result that  $K := \mathbb{Q}(\omega)$  is a number field whose ring of integers is known to be  $\mathcal{O}_K = \mathbb{Z}[\omega]$ , in which case it follows right away.  $\square$

**Remark.** We spell out the details in the proof a bit more explicitly here. It's enough to show that  $\omega \cdot 4i \cdot S = \frac{n}{m} \omega^2 - \frac{n}{m}$  is not an algebraic integer for completeness.

Take  $K = \mathbb{Q}(\omega)$  of degree  $d := \varphi(n) \geq 2$ ; as a  $\mathbb{Q}$ -module, it obeys  $K = \mathbb{Q} \oplus \omega \mathbb{Q} \oplus \dots \oplus \omega^{d-1} \mathbb{Q}$ . The theorem we are quoting is that, as  $\mathbb{Z}$ -modules, we have  $\mathcal{O}_K = \mathbb{Z}[\omega] = \mathbb{Z} \oplus \omega \mathbb{Z} \oplus \dots \oplus \omega^{d-1} \mathbb{Z}$  i.e.  $\mathcal{O}_K$  contains exactly those numbers in  $K$  for which the canonical  $\mathbb{Q}$ -coefficients happen



to be integers. (This is quite famous, but for a print reference, see Neukirch's *Algebraic Number Theory*, Section I.10.) And  $\omega \cdot 4i \cdot S$  fails this criteria, since  $\frac{n}{m} \notin \mathbb{Z}$ .

However, for any  $a$  and  $b$ , the number

$$4i \cdot T(a, b) = (\omega^a - 1)(\omega^b - 1)(\omega^{-b} - \omega^{-a})$$

is an algebraic integer. Since a finite sum of algebraic integers is also an algebraic integer, the sum of expressions of the form  $4i \cdot T(a, b)$  will never equal  $4i \cdot S$ .

**Remark.** If one wants to avoid citing the fact that  $\mathcal{O}_K = \mathbb{Z}[\omega]$ , then one can instead note that  $T(a, b)$  is actually always divisible by  $(\omega - 1)(\omega^{-1} + 1) = \omega - \omega^{-1}$  over the algebraic integers (at least one of  $\{\omega^a - 1, \omega^b - 1, \omega^{-a} - \omega^{-b}\}$  is a multiple of  $\omega + 1$ , by casework on  $a, b \pmod{2}$ ). Then one using  $\frac{4i}{(\omega - 1)(\omega^{-1} + 1)}$  as the scaling factor instead of  $4i$ , one sees that we actually need  $\frac{n}{m}$  to be an algebraic integer. But a rational number is an algebraic integer if and only if it's a rational integer; so  $\frac{n}{m}$  is an algebraic integer exactly when  $m$  divides  $n$ .

## §2 Solutions to Day 2

### §2.1 USAMO 2024/4, proposed by Rishabh Das

Available online at <https://aops.com/community/p30227198>.

#### Problem statement

Let  $m$  and  $n$  be positive integers. A circular necklace contains  $mn$  beads, each either red or blue. It turned out that no matter how the necklace was cut into  $m$  blocks of  $n$  consecutive beads, each block had a distinct number of red beads. Determine, with proof, all possible values of the ordered pair  $(m, n)$ .

The answer is  $m \leq n + 1$  only.

¶ **Proof the task requires  $m \leq n + 1$ .** Each of the  $m$  blocks has a red bead count between 0 and  $n$ , each of which appears at most once, so  $m \leq n + 1$  is needed.

¶ **Construction when  $m = n + 1$ .** For concreteness, here is the construction for  $n = 4$ , which obviously generalizes. The beads are listed in reading order as an array with  $n + 1$  rows and  $n$  columns. Four of the blue beads have been labeled  $B_1, \dots, B_n$  to make them easier to track as they move.

$$T_0 = \begin{bmatrix} R & R & R & R \\ R & R & R & B_1 \\ R & R & B & B_2 \\ R & B & B & B_3 \\ B & B & B & B_4 \end{bmatrix}.$$

To prove this construction works, it suffices to consider the  $n$  cuts  $T_0, T_1, T_2, \dots, T_{n-1}$  made where  $T_i$  differs from  $T_{i-1}$  by having the cuts one bead later also have the property each row has a distinct red count:

$$T_1 = \begin{bmatrix} R & R & R & R \\ R & R & B_1 & R \\ R & B & B_2 & R \\ B & B & B_3 & B \\ B & B & B_4 & R \end{bmatrix} \quad T_2 = \begin{bmatrix} R & R & R & R \\ R & B_1 & R & R \\ B & B_2 & R & B \\ B & B_3 & B & B \\ B & B_4 & R & R \end{bmatrix} \quad T_3 = \begin{bmatrix} R & R & R & R \\ B_1 & R & R & B \\ B_2 & R & B & B \\ B_3 & B & B & B \\ B_4 & R & R & R \end{bmatrix}.$$

We can construct a table showing for each  $1 \leq k \leq n + 1$  the number of red beads which are in the  $(k + 1)$ st row of  $T_i$  from the bottom:

$k$	$T_0$	$T_1$	$T_2$	$T_3$
$k = 4$	4	4	4	4
$k = 3$	3	3	3	2
$k = 2$	2	2	1	1
$k = 1$	1	0	0	0
$k = 0$	0	1	2	3

This suggests following explicit formula for the entry of the  $(i, k)$ th cell which can then be checked straightforwardly:

$$\#(\text{red cells in } k\text{th row of } T_i) = \begin{cases} k & k > i \\ k - 1 & i \geq k > 0 \\ i & k = 0. \end{cases}$$

And one can see for each  $i$ , the counts are all distinct (they are  $(i, 0, 1, \dots, k-1, k+1, \dots, k)$  from bottom to top). This completes the construction.

¶ **Construction when  $m < n + 1$ .** Fix  $m$ . Take the construction for  $(m, m - 1)$  and add  $n + 1 - m$  cyan beads to the start of each row; for example, if  $n = 7$  and  $m = 5$  then the new construction is

$$T = \begin{bmatrix} C & C & C & R & R & R & R \\ C & C & C & R & R & R & B_1 \\ C & C & C & R & R & B & B_2 \\ C & C & C & R & B & B & B_3 \\ C & C & C & B & B & B & B_4 \end{bmatrix}.$$

This construction still works for the same reason (the cyan beads do nothing for the first  $n + 1 - m$  shifts, then one reduces to the previous case). If we treat cyan as a shade of blue, this finishes the problem.

## §2.2 USAMO 2024/5, proposed by Anton Trygub


Available online at <https://aops.com/community/p30227196>.

### Problem statement

Point  $D$  is selected inside acute triangle  $ABC$  so that  $\angle DAC = \angle ACB$  and  $\angle BDC = 90^\circ + \angle BAC$ . Point  $E$  is chosen on ray  $BD$  so that  $AE = EC$ . Let  $M$  be the midpoint of  $BC$ . Show that line  $AB$  is tangent to the circumcircle of triangle  $BEM$ .

This problem has several approaches and we showcase a collection of them.

¶ **The author's original solution.** Complete isosceles trapezoid  $ABQC$  (so  $D \in \overline{AQ}$ ). Reflect  $B$  across  $E$  to point  $F$ .



**Claim** — We have  $DQCF$  is cyclic.


*Proof.* Since  $EA = EC$ , we have  $\overline{QF} \perp \overline{AC}$  as line  $QF$  is the image of the perpendicular bisector of  $\overline{AC}$  under a homothety from  $B$  with scale factor 2. Then

$$\begin{aligned} \angle FDC &= -\angle CDB = 180^\circ - (90^\circ + \angle CAB) = 90^\circ - \angle CAB \\ &= 90^\circ - \angle QCA = \angle FQC. \end{aligned} \quad \square$$

To conclude, note that

$$\angle BEM = \angle BFC = \angle DFC = \angle DQC = \angle AQC = \angle ABC = \angle ABM.$$

**Remark** (Motivation). Here is one possible way to come up with the construction of point  $F$  (at least this is what led Evan to find it). If one directs all the angles in the obvious way, there are really two points  $D$  and  $D'$  that are possible, although one is outside the triangle; they give corresponding points  $E$  and  $E'$ . The circles  $BEM$  and  $BE'M$  must then actually coincide since they are both alleged to be tangent to line  $AB$ . See the figure below.



One can already prove using angle chasing that  $\overline{AB}$  is tangent to  $(BEE')$ . So the point of the problem is to show that  $M$  lies on this circle too. However, from looking at the diagram, one may realize that in fact it seems


$$\triangle MEE' \simeq \triangle CDD'$$

is going to be true from just those marked in the figure (and this would certainly imply the desired concyclic conclusion). Since  $M$  is a midpoint, it makes sense to dilate  $\triangle EME'$  from  $B$  by a factor of 2 to get  $\triangle FCF'$  so that the desired similarity is actually a spiral similarity at  $C$ . Then the spiral similarity lemma says that the desired similarity is equivalent to requiring  $\overline{DD'} \cap \overline{FF'} = Q$  to lie on both  $(CDF)$  and  $(CD'F')$ . Hence the key construction and claim from the solution are both discovered naturally, and we find the solution above. (The points  $D'$ ,  $E'$ ,  $F'$  can then be deleted to hide the motivation.)

¶ **Another short solution.** Let  $Z$  be on line  $BDE$  such that  $\angle BAZ = 90^\circ$ . This lets us interpret the angle condition as follows:

**Claim** — Points  $A, D, Z, C$  are cyclic.


*Proof.* Because  $\angle ZAC = 90^\circ - \angle A = 180^\circ - \angle CDB = \angle ZDC$ . □



**Claim** — Points  $M, E, O, W$  are cyclic.

$$\begin{aligned}\angle MOE &= \angle(\overline{OM}, \overline{BC}) + \angle(\overline{BC}, \overline{AC}) + \angle(\overline{AC}, \overline{OE}) \\ &= 90^\circ + \angle BCA + 90^\circ \\ &= \angle BCA = \angle CAD = \angle CZD = \angle MWD = \angle MWE. \quad \square\end{aligned}$$
$$\begin{aligned}\angle MEB &= \angle MEW = \angle MOW \\ &= \angle(\overline{MO}, \overline{BC}) + \angle(\overline{BC}, \overline{AB}) + \angle(\overline{AB}, \overline{OW}) \\ &= 90^\circ + \angle CBA + 90^\circ = \angle CBA = \angle MBA.\end{aligned}$$

¶ **A Menelaus-based approach (Kevin Ren).** Let  $P$  be on  $\overline{BC}$  with  $AP = PC$ . Let  $Y$  be the point on line  $AB$  such that  $\angle ACY = 90^\circ$ ; as  $\angle AYC = 90^\circ - A$  it follows  $BDYC$  is cyclic. Let  $K = \overline{AP} \cap \overline{CY}$ , so  $\triangle ACK$  is a right triangle with  $P$  the midpoint of its hypotenuse.



**Claim** — Triangles  $BPE$  and  $DYK$  are similar.

*Proof.* We have  $\angle MPE = \angle CPE = \angle KCP = \angle PKC$  and  $\angle EBP = \angle DBC = \angle DYC = \angle DYK$ .  $\square$

**Claim** — Triangles  $BEM$  and  $YDC$  are similar.

*Proof.* By Menelaus  $\triangle PCK$  with respect to collinear points  $A, B, Y$  that

$$\frac{BP}{BC} \frac{YC}{YK} \frac{AK}{AP} = 1.$$

Since  $AK/AP = 2$  (note that  $P$  is the midpoint of the hypotenuse of right triangle  $ACK$ ) and  $BC = 2BM$ , this simplifies to


$$\frac{BP}{BM} = \frac{YK}{YC}.$$

To finish, note that

$$\angle DBA = \angle DBY = \angle DCY = \angle BME$$

implying the desired tangency.

¶ **A spiral similarity approach (Hans Yu).** As in the previous solution, let  $Y$  be the point on line  $AB$  such that  $\angle ACY = 90^\circ$ ; so  $BDYC$  is cyclic. Let  $\Gamma$  be the circle through  $B$  and  $M$  tangent to  $\overline{AB}$ , and let  $\Omega := (BCYD)$ . We need to show  $E \in \Gamma$ .



Denote by  $S$  the second intersection of  $\Gamma$  and  $\Omega$ . The main idea behind is to consider the spiral similarity

$$\Psi : \Omega \rightarrow \Gamma \quad C \mapsto M \text{ and } Y \mapsto B$$

centered at  $S$  (due to the spiral similarity lemma), and show that  $\Psi(D) = E$ . The spiral similarity lemma already promises  $\Psi(D)$  lies on line  $BD$ .

**Claim —** We have  $\Psi(A) = O$ , the circumcenter of  $ABC$ .

*Proof.* Note  $\triangle OBM \cong \triangle AYC$ ; both are right triangles with  $\angle BAC = \angle BOM$ .  $\square$

**Claim —**  $\Psi$  maps line  $AD$  to line  $OP$ .

*Proof.* If we let  $P$  be on  $\overline{BC}$  with  $AP = PC$  as before,

$$\angle(\overline{AD}, \overline{OP}) = \angle APO = \angle OPC = \angle YCP = \angle(\overline{YC}, \overline{BM}).$$

As  $\Psi$  maps line  $YC$  to line  $BM$  and  $\Psi(A) = O$ , we're done.  $\square$

Hence  $\Psi(D)$  should not only lie on  $BD$  but also line  $OP$ . This proves  $\Psi(D) = E$ , so  $E \in \Gamma$  as needed.



**§2.3 USAMO 2024/6, proposed by Titu Andreescu, Gabriel Dospinescu**

Available online at <https://aops.com/community/p30227192>.

**Problem statement**

Let  $n > 2$  be an integer and let  $\ell \in \{1, 2, \dots, n\}$ . A collection  $A_1, \dots, A_k$  of (not necessarily distinct) subsets of  $\{1, 2, \dots, n\}$  is called  $\ell$ -large if  $|A_i| \geq \ell$  for all  $1 \leq i \leq k$ . Find, in terms of  $n$  and  $\ell$ , the largest real number  $c$  such that the inequality

$$\sum_{i=1}^k \sum_{j=1}^k x_i x_j \frac{|A_i \cap A_j|^2}{|A_i| \cdot |A_j|} \geq c \left( \sum_{i=1}^k x_i \right)^2$$

holds for all positive integer  $k$ , all nonnegative real numbers  $x_1, x_2, \dots, x_k$ , and all  $\ell$ -large collections  $A_1, A_2, \dots, A_k$  of subsets of  $\{1, 2, \dots, n\}$ .

The answer turns out to be

$$c = \frac{n + \ell^2 - 2\ell}{n(n-1)}.$$

The following solution using indicator functions was suggested by the language reasoning system developed by Thang Luong and the team at Google DeepMind (<https://dpmid.ai/imo-silver>).

¶ **Rewriting the sum.** Abbreviate  $[n] := \{1, \dots, n\}$ , and WLOG assume  $\sum x_i = 1$ . To prove the bound, we first rewrite the left-hand side using indicator functions and switch the order of summation as

$$\begin{aligned} \text{LHS} &= \sum_{i=1}^k \sum_{j=1}^k \frac{x_i}{|A_i|} \frac{x_j}{|A_j|} |A_i \cap A_j|^2 \\ &= \sum_{i=1}^k \sum_{j=1}^k \frac{x_i}{|A_i|} \frac{x_j}{|A_j|} \left( \sum_{p \in [n]} \sum_{q \in [n]} \mathbf{1}_{p \in A_i} \mathbf{1}_{q \in A_j} \right) \\ &= \sum_{p \in [n]} \sum_{q \in [n]} \sum_{i=1}^k \sum_{j=1}^k \frac{x_i \mathbf{1}_{p \in A_i}}{|A_i|} \cdot \frac{x_j \mathbf{1}_{q \in A_j}}{|A_j|}. \end{aligned}$$

Hence, if we define the  $n^2$  variables

$$v_{p,q} := \sum_{i: p, q \in A_i} \frac{x_i}{|A_i|}$$

for each  $(p, q) \in [n]^2$  then the previous expression simplifies to just

$$\text{LHS} = \sum_{p \in [n]} \sum_{q \in [n]} v_{p,q}^2.$$

¶ **Proof of bound.** We split the sum into  $p = q$  versus  $p \neq q$  and bound by QM-AM separately in the obvious way. For the terms  $p = q$ ,

$$\sum_{p \in [n]} v_{p,p} = \sum_{i=1}^k |A_i| \cdot \frac{x_i}{|A_i|} = \sum_{i=1}^k x_i = 1$$

$$\implies \sum_{p \in [n]} v_{p,p}^2 \geq \frac{\left( \sum_{p \in [n]} v_{p,p} \right)^2}{n} = \frac{1}{n}.$$

When  $p \neq q$  we instead have

$$\begin{aligned} \sum_{p \neq q \in [n]} v_{p,q} &= \sum_{i=1}^k 2 \binom{|A_i|}{2} \cdot \frac{x_i}{|A_i|} = \sum_{i=1}^k (|A_i| - 1) x_i \geq \sum_{i=1}^k (\ell - 1) x_i = \ell - 1 \\ \implies \sum_{p \neq q \in [n]} v_{p,q}^2 &\geq \frac{\left( \sum_{p \neq q \in [n]} v_{p,q} \right)^2}{n(n-1)} \geq \frac{(\ell - 1)^2}{n(n-1)}. \end{aligned}$$

Combining the two bounds gives

$$\text{LHS} \geq \frac{1}{n} + \frac{(\ell - 1)^2}{n(n-1)} = \frac{n + \ell^2 - 2\ell}{n(n-1)}$$

which matches the answer claimed above.

¶ **Construction.** Equality occurs if  $v_{p,p}$  is the same for all  $p$  and  $v_{p,q}$  is the same for all  $p \neq q$ , and  $|A_i| = \ell$  for all  $i$ . This occurs if one takes, say,  $k = \binom{n}{\ell}$  and  $A_i$  to be all the subsets of size  $\ell$  each exactly once.

**Remark.** Originally, the write-up I used considered vectors in  $\mathbb{R}^{n^2}$  where entries will be indexed by ordered pairs  $(p, q) \in \{1, \dots, n\}^2$ . It was equivalent to the solution above, but looking back, I think this makes the problem look harder than it actually is.

For comparison, here is how the notation with vectors works, using  $\langle \bullet, \bullet \rangle$  for dot product, and  $\|\bullet\|$  for the vector norm. For  $i = 1, \dots, n$  define  $\mathbf{v}_i$  by

$$\mathbf{v}_i[p, q] := \begin{cases} \frac{1}{|A_i|} & p \in A_i \text{ and } q \in A_i \\ 0 & \text{otherwise;} \end{cases} \quad \mathbf{v} := \sum_i x_i \mathbf{v}_i.$$

Then

$$\begin{aligned} \sum_i \sum_j x_i x_j \frac{|A_i \cap A_j|^2}{|A_i| |A_j|} &= \sum_i \sum_j x_i x_j \langle \mathbf{v}_i, \mathbf{v}_j \rangle \\ &= \langle \mathbf{v}, \mathbf{v} \rangle = \|\mathbf{v}\|^2. \end{aligned}$$

Define two more vectors  $\mathbf{e}$  and  $\mathbf{1}$ ; the vector  $\mathbf{e}$  has 1 in the  $(p, q)^{\text{th}}$  component if  $p = q$ , and 0 otherwise, while  $\mathbf{1}$  has all-ones. In that case, note that

$$\begin{aligned} \langle \mathbf{e}, \mathbf{v} \rangle &= \sum_i x_i \langle \mathbf{e}, \mathbf{v}_i \rangle = \sum_i x_i \\ \langle \mathbf{1}, \mathbf{v} \rangle &= \sum_i x_i \langle \mathbf{1}, \mathbf{v}_i \rangle = \sum_i x_i |A_i|. \end{aligned}$$

That means for any positive real constants  $\alpha$  and  $\beta$ , by Cauchy-Schwarz for vectors, we should have

$$\begin{aligned} \|\alpha \mathbf{e} + \beta \mathbf{1}\| \|\mathbf{v}\| &\geq \langle \alpha \mathbf{e} + \beta \mathbf{1}, \mathbf{v} \rangle = \alpha \langle \mathbf{e}, \mathbf{v} \rangle + \beta \langle \mathbf{1}, \mathbf{v} \rangle \\ &= \alpha \cdot \sum_i x_i + \beta \cdot \sum_i x_i |A_i| \geq (\alpha + \ell \beta) \sum_i x_i. \end{aligned}$$

Set  $\mathbf{w} := \alpha \mathbf{e} + \beta \mathbf{1}$  for brevity. Then

$$\mathbf{w}[p, q] = \begin{cases} \alpha + \beta & \text{if } p = q \\ \beta & \text{if } p \neq q \end{cases} \implies \|\mathbf{w}\| = \sqrt{n \cdot (\alpha + \beta)^2 + (n^2 - n) \cdot \beta^2}.$$

Therefore, we get an lower bound

$$\frac{\|\mathbf{v}\|}{\sum x_i} \geq \frac{\alpha + \ell\beta}{\sqrt{n \cdot (\alpha + \beta)^2 + (n^2 - n) \cdot \beta^2}}$$

Letting  $\alpha = n - \ell$  and  $\beta = \ell - 1$  gives a proof that the constant

$$c = \frac{((n - \ell) + \ell(\ell - 1))^2}{n \cdot (n - 1)^2 + (n^2 - n) \cdot (\ell - 1)^2} = \frac{(n + \ell^2 - 2\ell)^2}{n(n - 1)(n + \ell^2 - 2\ell)} = \frac{n + \ell^2 - 2\ell}{n(n - 1)}$$

makes the original inequality always true. (The choice of  $\alpha : \beta$  is suggested by the example below.)

Meanwhile, for the construction (where  $k = \binom{n}{\ell}$ , etc.) it would be sufficient to show that  $\mathbf{w}$  and  $\mathbf{v}$  are scalar multiples, so that the above Cauchy-Schwarz is equality. However, we can compute

$$\mathbf{w}[p, q] = \begin{cases} n - 1 & \text{if } p = q \\ \ell - 1 & \text{if } p \neq q \end{cases}, \quad \mathbf{v}[p, q] = \begin{cases} \binom{n-1}{\ell-1} \cdot \frac{1}{\ell} & \text{if } p = q \\ \binom{n-2}{\ell-2} \cdot \frac{1}{\ell} & \text{if } p \neq q \end{cases}$$

which are indeed scalar multiples.