Homework 1

Qijun Han 12212635

September 15, 2024

1 **NE1.1**

1.1 Problem a

$$G(s) = \frac{1}{s^2 + 2s + 6}$$

ODE:

$$y'' + 2y' + 6y = u(t)$$

let $x_1 = y, x_2 = y'$, then we have

$$\begin{cases} x_1' = x_2 \\ x_2' = -2x_2 - 6x_1 + u(t) \end{cases}$$
 so the state space representation is

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -6 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u(t)$$

$$y = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$
(1)

thus
$$A = \begin{bmatrix} 0 & 1 \\ -6 & -2 \end{bmatrix}, B = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, C = \begin{bmatrix} 1 & 0 \end{bmatrix}, D = \begin{bmatrix} 0 \end{bmatrix}$$

1.2 Problem b

let
$$G_1(s) = \frac{1}{s^2 + 2s + 6} = \frac{W(s)}{U(s)}$$
, $G_2(s) = s + 3 = \frac{Y(s)}{W(s)}$, then

$$G(s) = G_1(s)G_2(s)$$

ODE:

$$w'' + w' + w = u(t)$$

$$y = w' + 3w$$
(2)

let $x_1 = w, x_2 = w'$, then we have

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -6 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u(t)$$

$$y = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$
(3)

and

$$y = \begin{bmatrix} 3 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$
 thus $A = \begin{bmatrix} 0 & 1 \\ -6 & -2 \end{bmatrix}$, $B = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$, $C = \begin{bmatrix} 3 & 1 \end{bmatrix}$, $D = \begin{bmatrix} 0 \end{bmatrix}$

1.3 Problem c

similar to (a), we have A =

- 2 Problem 2
- 3 Problem 3
- 4 Problem 4