

final_exam_prog

April 29, 2021

```
[2]: import numpy as np
import cvxpy as cp
```

1 Final exam

1.1 Question 03

1.1.1 Koidala Surya Prakash

1.1.2 EE18btech11026

1.1.3 3. a Convexity ?

We can reformulate the problem in the form :

objective := $x.T(Q)x + (c.T)x$ constraints := $Ax \leq b$

where : $x = [x_1, x_2]$

$$Q = \begin{bmatrix} 1 & -0.5 \\ -0.5 & 2 \end{bmatrix}$$
$$c = [-1, 0]$$
$$A = \begin{bmatrix} 1 & -2 \\ 1 & 4 \\ 5 & -76 \end{bmatrix}$$
$$b = \begin{bmatrix} u_1 \\ u_2 \\ 1 \end{bmatrix}$$

Since the obj is in quadratic form and constraints are in linear form

This is a QP problem

We know that all QP problems are convex optimisation problems . Hence this problem is convex.

1.1.4 3. b : Finding optimal variables

```
[3]: ### Constants
```

```
u1 = -2
u2 = -3

b = np.array([[u1,u2,1]]).T

Q = np.array([[1,-0.5],[-0.5, 2]])
c = np.array([[1,0]]).T

A = np.array([[1,-2],[1,4],[5,-76]])
```

```
[4]:
```

```
'''
Objective : A QP problem of the form  $(x.T)Q(x) + c.T*x$ 
with affine constraints :  $Ax \leq b$ 

Therefore this is a convex problem
'''

x = cp.Variable((2,1))
obj = cp.Minimize( cp.quad_form(x, Q) + c.T*x )
constraints = [A*x <= b]

prob = cp.Problem(obj, constraints)
prob.solve()

lam = constraints[0].dual_value

print("The following are the optimal primal variables : \n")

print('x1* = %.4f'%x.value[0][0])
print('x2* = %.4f'%x.value[1][0])
print('\n')

print("The following are the optimal dual variables : \n")
print('lamda_1* = %.4f' %lam[0])
print('lamda_2* = %.4f' %lam[1])
print('lamda_3* = %.4f' %lam[2])
```

The following are the optimal primal variables :

```
x1* = -2.3333
x2* = -0.1667
```

The following are the optimal dual variables :

```
lamda_1* = 2.8645
lamda_2* = 2.2980
lamda_3* = 0.0675
```

1.2 3.c Verifying KKT conditions

The primal variables and dual optimal variables satisfy the below mentioned kkt conditions.

1. $f_i(x^*) \leq 0 \implies Ax-b \leq 0$
2. $\lambda \geq 0$
3. $\lambda f_i(x) = 0$
4. $\frac{dfo(x)}{dx} + \sum(\lambda_i \frac{df_i(x)}{dx}) = 0$

```
[ ]: '''
Verifying KKT conditions..
'''

'''
1.  $f_i(x^*) \leq 0 \implies Ax-b \leq 0$ 
'''

print('Condition 1 :: obeys all constraints')
print('_____')
print('Ax* - b = \n', np.round(A@x.value - b, 3))

# 2.  $\lambda \geq 0$ 
print("\nCondition 2 :: all lamda values are positive")
print('_____')
print('lam_i = \n', lam)

# 3.  $\lambda * f_i(x^*) = 0$ 

print("\nCondition 3 :: since constraint func are zero")
print('_____')
print("lam_i * x* = \n", np.round(lam*(A@x.value - b), 3) )

# 4.  $\frac{dfo(x)}{dx} + \sum(\lambda_i \frac{df_i(x)}{dx}) = 0$ 
'''
 $\frac{dfo(x)}{dx} = Qx + c$ 
 $\frac{df_i(x)}{dx} = a \implies \sum(\lambda_i \frac{df_i(x)}{dx}) = A.T*\lambda$ 
'''

print("\nCondition 4 ::  $\frac{dfo(x)}{dx} + \sum(\lambda_i \frac{df_i(x)}{dx}) = 0$ ")
print('_____')
print(np.round(2*Q@x.value + c + A.T@lam, 3))
```

Condition 1 :: obeys all constraints

```
Ax* - b =
[[0.]
 [0.]
 [0.]]
```

Condition 2 :: all lamda values are positive

```
-----
lam_i =
[[2.86447804]
 [2.29803246]
 [0.0674979 ]]
```

Condition 3 :: since constraint func are zero

```
-----
lam_i * x* =
[[0.]
 [0.]
 [0.]]
```

Condition 4 :: $dfo(x)/dx + \sum(lam_i * dfi(x)/dx) == 0$

```
-----
[[ 0.]
 [-0.]]
```

1.3 3.d Plotting level curves

```
[ ]: ### Function finding p*(u1,u2)
def p(u1,u2):
    b = np.array([[u1,u2,1]]).T
    Q = np.array([[1,-0.5],[-0.5, 2]])
    c = np.array([[-1,0]]).T
    A = np.array([[1,-2],[1,4],[5,-76]])

    x = cp.Variable((2,1))
    obj = cp.Minimize( cp.quad_form(x, Q) + c.T@x )
    constraints = [A@x <= b]

    prob = cp.Problem(obj, constraints).solve()
    return obj.value
```

```
[ ]: import numpy as np
import matplotlib.pyplot as plt
import pylab

### Here u1 == X , u2 == Y , p*(u1,u2) = Z
```

```

# List of points in x axis
XPoints = []

# List of points in y axis
YPoints = []

# X and Y points are from -20 to +20 varying in steps of .5
lis = np.arange(-20,20,1)
for val in lis:
    XPoints.append(val)
    YPoints.append(val)

# Z values as a matrix
ZPoints = np.ndarray((len(XPoints),len(YPoints)))

for x in range(0, len(XPoints)):
    for y in range(0, len(YPoints)):
        ZPoints[x][y] = p(XPoints[x],YPoints[y])

# Set the x axis and y axis limits
pylab.xlim([-25,25])
pylab.ylim([-25,25])

# Provide a title for the contour plot
plt.title('Level curves of  $p(u_1,u_2)$ ')

# Set x axis label for the contour plot
plt.xlabel('u1')

# Set y axis label for the contour plot
plt.ylabel('u2')

# Create contour lines or level curves using matplotlib.pyplot module
contours = plt.contour(XPoints, YPoints, ZPoints)

# Display z values on contour lines
plt.clabel(contours, inline=1, fontsize=8)


# Display the contour plot

plt.scatter([-2], [-3])

plt.annotate(" $p(-2,-3) = 7.44$  ", (0,-5))

plt.show()

```



1.3.1 3.e Convexity of p^* from level curves

We can see that the gradient of the slopes are increasing as we move towards right part of the curve. This is seen in both the directions

====> Thus p^* is convex

since $d^2(p)/du_1^2 \geq 0$ $d^2(p)/du_2^2 \geq 0$

1.3.2 3.f Numerical verification