

# AI1110: Probability and Random Variables

## Assignment 11: Papoulis-Pillai Ex 8-28

Suruaansh Jain (cs21btech11057)

June 16, 2022

# Outline

## 1 Problem

## 2 Solution

- Definitions
- Transformation
- Hypothesis
- Test Statistic
- Decision

# Problem

Brand A batteries cost more than brand B batteries. Their life lengths are two normal and independent random variables  $x$  and  $y$ . We test 16 batteries of brand A and 26 batteries of brand B and find these values in hours:  $\bar{x} = 4.6$   $s_x = 1.1$   $\bar{y} = 4.2$   $s_y = 0.9$  Test the hypothesis  $\eta_x = \eta_y$ , against  $\eta_x > \eta_y$ , with  $\alpha = 0.05$

# Percentile

Given a distribution on random variable  $x$  as  $F_x(x)$ , we define the  $k^{th}$  percentile of this distribution as

$$Percentile_k = F_x^{-1}(k) \quad (1)$$

In other words, the  $k^{th}$  percentile returns the value of random variable  $x_0$  for which  $F_x(x_0) = k$ .

# Hypothesis

Let us assume, we are given a random variable  $x$  whose distribution is  $F(x, \theta)$  depending on some parameter  $\theta$  (The parameter might be mean, variance etc.). We are required to use evidence that either supports or rejects a given prediction of the actual value of  $\theta$ , which we will call  $\theta_0$ .

## Null Hypothesis

In the null hypothesis, we make the prediction that  $\theta = \theta_0$ . This is represented by  $H_0 : \theta = \theta_0$ .

## Alternate Hypothesis

In the alternate hypothesis, we make the prediction that  $\theta \neq \theta_0$ . This is represented by  $H_1 : \theta \neq \theta_0$ . Note that the null hypothesis may be defined differently based on utility.

# Testing Hypothesis

To test whether a given hypothesis is feasible based on evidence, we first define a random variable  $q$  whose density is convenient to plot and is a function of sample vector  $X$  as follows.

$$q = g(X) \quad (2)$$

We will call  $q$  as the test statistic.

The density of random variable  $q$  is given by  $p_q(q, \theta)$  where  $\theta$  is the parameter. Now consider the density  $p_q(q, \theta_0)$  (based on  $H_0$ ) and a region (Critical Region)  $R_c$  where  $p_q(q, \theta_0)$  is negligible. If we find that the value of  $q$  lies in  $R_c$ , then we reject  $H_0$ .

One can decide the region  $R_c$  using the significance level  $\alpha$ .  $\alpha$  represents the probability that  $q \in R_c$  when  $H_0$  is true. Hence, when given a value of  $\alpha$  one can determine  $R_c$  and thereby check the validity of the null hypothesis.

## Mean as Parameter: Unknown Variance

Consider a random variable  $x$ , from which we have obtained a sample vector  $X$ . We are required to reject or support the hypothesis  $H_0 : \eta = \eta_0$  against  $H_1 : \eta \neq \eta_0$ , where we check if the mean  $\eta$  equals a constant  $\eta_0$ . In the case that the variance is unknown but the sample mean  $\bar{x}$  and sample variance  $s^2$  are given, we must use a Student t distribution. Note that the sample vector  $X$  has  $n - 1$  degrees of freedom as we are constrained to ensure that the sum of the values of the vector  $X - \bar{x}$  must be 0.

Assuming random variable  $\bar{x}$  is represented by a normal distribution, we define test statistic  $q$  as follows:

$$q = \frac{\bar{x} - \eta_0}{s/\sqrt{n}} \quad (3)$$

## Mean as Parameter: Unknown Variance

For an alternate hypothesis  $H_1 : \eta \neq \eta_0$  and given significance value  $\alpha$ , we note that the critical region  $R_c$  is given by:

$$R_c = (t_{1-\alpha}(n-1), \infty) \quad (4)$$

where  $t_k(n-1)$  represents the  $k^{th}$  percentile (As explained in (1))

We consider the  $\alpha^{th}$  and its complementary percentile as the given hypothesis is single ended, i.e., it allows us accept values less than the hypothesised mean value.



# Stating the Transformation

Let,

$$w = \bar{x} - \bar{y} = 0.4 \quad (5)$$

$$\Rightarrow \sigma_w^2 = \frac{\sigma_x^2}{16} + \frac{\sigma_y^2}{26} = 0.32 \quad (6)$$

# Stating the Hypothesis

We state the null Hypothesis as

$$H_0 : \eta = 0 \quad (7)$$

and the alternate hypothesis as

$$H_1 : \eta > 0 \quad (8)$$

We are required to test the above hypotheses for significance value  
 $\alpha = 0.05$

## Calculate Test Statistic $q$

Given sample mean  $\bar{x} = 4.35$  and sample variance  $s = 0.32$ , we get our test statistic  $q$  from (3) as

$$q = \frac{0.4 - 0}{0.32} = 1.223 \quad (9)$$

# Making Decision

We shall determine the critical regions for given significance values  $\alpha_1$  and  $\alpha_2$  using (4)

For  $\alpha = 0.005$ , we find that  $t_{0.995} = 1.64$ . As  $q < 1.64$  we can accept this hypothesis.