

# Interpolation & Polynomial Approximation

## Lagrange Interpolating Polynomials I

Numerical Analysis (9th Edition)

R L Burden & J D Faires

Beamer Presentation Slides

prepared by

John Carroll

Dublin City University

© 2011 Brooks/Cole, Cengage Learning

# Outline

## 1 Weierstrass Approximation Theorem

# Outline

- 1 Weierstrass Approximation Theorem
- 2 Inaccuracy of Taylor Polynomials

# Outline

- 1 Weierstrass Approximation Theorem
- 2 Inaccuracy of Taylor Polynomials
- 3 Constructing the Lagrange Polynomial

# Outline

- 1 Weierstrass Approximation Theorem
- 2 Inaccuracy of Taylor Polynomials
- 3 Constructing the Lagrange Polynomial
- 4 Example: Second-Degree Lagrange Interpolating Polynomial

# Outline

- 1 Weierstrass Approximation Theorem
- 2 Inaccuracy of Taylor Polynomials
- 3 Constructing the Lagrange Polynomial
- 4 Example: Second-Degree Lagrange Interpolating Polynomial

# Weierstrass Approximation Theorem

## Algebraic Polynomials

# Weierstrass Approximation Theorem

## Algebraic Polynomials

One of the most useful and well-known classes of functions mapping the set of real numbers into itself is the **algebraic polynomials**, the set of functions of the form

$$P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$

where  $n$  is a nonnegative integer and  $a_0, \dots, a_n$  are real constants.



# Weierstrass Approximation Theorem

$$P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

## Algebraic Polynomials (Cont'd)

# Weierstrass Approximation Theorem

$$P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

## Algebraic Polynomials (Cont'd)

- One reason for their importance is that they uniformly approximate continuous functions.

# Weierstrass Approximation Theorem

$$P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

## Algebraic Polynomials (Cont'd)


- One reason for their importance is that they uniformly approximate continuous functions.
- By this we mean that given any function, defined and continuous on a closed and bounded interval, there exists a polynomial that is as “close” to the given function as desired.

# Weierstrass Approximation Theorem

$$P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

## Algebraic Polynomials (Cont'd)

- One reason for their importance is that they uniformly approximate continuous functions.
- By this we mean that given any function, defined and continuous on a closed and bounded interval, there exists a polynomial that is as “close” to the given function as desired.
- This result is expressed precisely in the **Weierstrass Approximation Theorem**.



## Weierstrass Approximation Theorem

Suppose that  $f$  is defined and continuous on  $[a, b]$ . For each  $\epsilon > 0$ , there exists a polynomial  $P(x)$ , with the property that

$$|f(x) - P(x)| < \epsilon, \quad \text{for all } x \text{ in } [a, b].$$

## Benefits of Algebraic Polynomials

## Benefits of Algebraic Polynomials

- Another important reason for considering the class of polynomials in the approximation of functions is that the **derivative** and **indefinite integral** of a polynomial are easy to determine and are also polynomials.

## Benefits of Algebraic Polynomials

- Another important reason for considering the class of polynomials in the approximation of functions is that the **derivative** and **indefinite integral** of a polynomial are easy to determine and are also polynomials.
- For these reasons, polynomials are often used for approximating continuous functions.



# Outline

- 1 Weierstrass Approximation Theorem
- 2 Inaccuracy of Taylor Polynomials**
- 3 Constructing the Lagrange Polynomial
- 4 Example: Second-Degree Lagrange Interpolating Polynomial

# The Lagrange Polynomial: Taylor Polynomials

## Interpolating with Taylor Polynomials

# The Lagrange Polynomial: Taylor Polynomials

## Interpolating with Taylor Polynomials

- The Taylor polynomials are described as one of the fundamental building blocks of numerical analysis.

# The Lagrange Polynomial: Taylor Polynomials

## Interpolating with Taylor Polynomials

- The Taylor polynomials are described as one of the fundamental building blocks of numerical analysis.
- Given this prominence, you might expect that polynomial interpolation would make heavy use of these functions.

# The Lagrange Polynomial: Taylor Polynomials

## Interpolating with Taylor Polynomials

- The Taylor polynomials are described as one of the fundamental building blocks of numerical analysis.
- Given this prominence, you might expect that polynomial interpolation would make heavy use of these functions.
- However this is not the case.

# The Lagrange Polynomial: Taylor Polynomials

## Interpolating with Taylor Polynomials

- The Taylor polynomials are described as one of the fundamental building blocks of numerical analysis.
- Given this prominence, you might expect that polynomial interpolation would make heavy use of these functions.
- However this is not the case.
- The Taylor polynomials agree as closely as possible with a given function at a specific point, but they concentrate their accuracy near that point.

# The Lagrange Polynomial: Taylor Polynomials

## Interpolating with Taylor Polynomials

- The Taylor polynomials are described as one of the fundamental building blocks of numerical analysis.
- Given this prominence, you might expect that polynomial interpolation would make heavy use of these functions.
- However this is not the case.
- The Taylor polynomials agree as closely as possible with a given function at a specific point, but they concentrate their accuracy near that point.
- A good interpolation polynomial needs to provide a relatively accurate approximation over an entire interval, and Taylor polynomials do not generally do this.

# The Lagrange Polynomial: Taylor Polynomials

Example:  $f(x) = e^x$

We will calculate the first six Taylor polynomials about  $x_0 = 0$  for  $f(x) = e^x$ .



# The Lagrange Polynomial: Taylor Polynomials

Example:  $f(x) = e^x$

We will calculate the first six Taylor polynomials about  $x_0 = 0$  for  $f(x) = e^x$ .

## Note

Since the derivatives of  $f(x)$  are all  $e^x$ , which evaluated at  $x_0 = 0$  gives 1.

# The Lagrange Polynomial: Taylor Polynomials

Example:  $f(x) = e^x$

We will calculate the first six Taylor polynomials about  $x_0 = 0$  for  $f(x) = e^x$ .

## Note

Since the derivatives of  $f(x)$  are all  $e^x$ , which evaluated at  $x_0 = 0$  gives 1.

The Taylor polynomials are as follows:

# Taylor Polynomials for $f(x) = e^x$ about $x_0 = 0$

$$P_0(x) = 1$$

$$P_1(x) = 1 + x$$


$$P_2(x) = 1 + x + \frac{x^2}{2}$$

$$P_3(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$$

$$P_4(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24}$$

$$P_5(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120}$$

# Taylor Polynomials for $f(x) = e^x$ about $x_0 = 0$



Notice that even for the higher-degree polynomials, the error becomes progressively worse as we move away from zero.

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

Example: A more extreme case

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

## Example: A more extreme case

- Although better approximations are obtained for  $f(x) = e^x$  if higher-degree Taylor polynomials are used, this is not true for all functions.

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

## Example: A more extreme case

- Although better approximations are obtained for  $f(x) = e^x$  if higher-degree Taylor polynomials are used, this is not true for all functions.
- Consider, as an extreme example, using Taylor polynomials of various degrees for  $f(x) = \frac{1}{x}$  expanded about  $x_0 = 1$  to approximate  $f(3) = \frac{1}{3}$ .

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

## Calculations



# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

## Calculations

Since

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = (-1)^2 2 \cdot x^{-3},$$

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

## Calculations

Since

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = (-1)^2 2 \cdot x^{-3},$$

and, in general,

$$f^{(k)}(x) = (-1)^k k! x^{-k-1},$$

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

## Calculations

Since

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = (-1)^2 2 \cdot x^{-3},$$

and, in general,

$$f^{(k)}(x) = (-1)^k k! x^{-k-1},$$

the Taylor polynomials are

$$P_n(x) = \sum_{k=0}^n \frac{f^{(k)}(1)}{k!} (x-1)^k = \sum_{k=0}^n (-1)^k (x-1)^k.$$

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

To Approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

To Approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$

- To approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$  for increasing values of  $n$ , we obtain the values shown below — rather a dramatic failure!

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

To Approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$

- To approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$  for increasing values of  $n$ , we obtain the values shown below — rather a dramatic failure!
- When we approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$  for larger values of  $n$ , the approximations become increasingly inaccurate.

# Taylor Polynomials for $f(x) = \frac{1}{x}$ about $x_0 = 1$

To Approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$

- To approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$  for increasing values of  $n$ , we obtain the values shown below — rather a dramatic failure!
- When we approximate  $f(3) = \frac{1}{3}$  by  $P_n(3)$  for larger values of  $n$ , the approximations become increasingly inaccurate.

$n$	0	1	2	3	4	5	6	7
$P_n(3)$	1	-1	3	-5	11	-21	43	-85

# The Lagrange Polynomial: Taylor Polynomials

## Footnotes



# The Lagrange Polynomial: Taylor Polynomials

## Footnotes

- For the Taylor polynomials, all the information used in the approximation is concentrated at the single number  $x_0$ , so these polynomials will generally give inaccurate approximations as we move away from  $x_0$ .

# The Lagrange Polynomial: Taylor Polynomials

## Footnotes

- For the Taylor polynomials, all the information used in the approximation is concentrated at the single number  $x_0$ , so these polynomials will generally give inaccurate approximations as we move away from  $x_0$ .
- This limits Taylor polynomial approximation to the situation in which approximations are needed only at numbers close to  $x_0$ .

# The Lagrange Polynomial: Taylor Polynomials

## Footnotes

- For the Taylor polynomials, all the information used in the approximation is concentrated at the single number  $x_0$ , so these polynomials will generally give inaccurate approximations as we move away from  $x_0$ .
- This limits Taylor polynomial approximation to the situation in which approximations are needed only at numbers close to  $x_0$ .
- For ordinary computational purposes, it is more efficient to use methods that include information at various points.

# The Lagrange Polynomial: Taylor Polynomials

## Footnotes

- For the Taylor polynomials, all the information used in the approximation is concentrated at the single number  $x_0$ , so these polynomials will generally give inaccurate approximations as we move away from  $x_0$ .
- This limits Taylor polynomial approximation to the situation in which approximations are needed only at numbers close to  $x_0$ .
- For ordinary computational purposes, it is more efficient to use methods that include information at various points.
- The **primary use** of Taylor polynomials in numerical analysis is not for approximation purposes, but for the derivation of numerical techniques and error estimation.

# Outline

- 1 Weierstrass Approximation Theorem
- 2 Inaccuracy of Taylor Polynomials
- 3 Constructing the Lagrange Polynomial**
- 4 Example: Second-Degree Lagrange Interpolating Polynomial

# The Lagrange Polynomial: The Linear Case

## Polynomial Interpolation

# The Lagrange Polynomial: The Linear Case

## Polynomial Interpolation

- The problem of determining a polynomial of degree one that passes through the distinct points

$$(x_0, y_0) \quad \text{and} \quad (x_1, y_1)$$

is the same as approximating a function  $f$  for which

$$f(x_0) = y_0 \quad \text{and} \quad f(x_1) = y_1$$

by means of a first-degree polynomial **interpolating**, or agreeing with, the values of  $f$  at the given points.

# The Lagrange Polynomial: The Linear Case

## Polynomial Interpolation

- The problem of determining a polynomial of degree one that passes through the distinct points

$$(x_0, y_0) \quad \text{and} \quad (x_1, y_1)$$

is the same as approximating a function  $f$  for which

$$f(x_0) = y_0 \quad \text{and} \quad f(x_1) = y_1$$

by means of a first-degree polynomial **interpolating**, or agreeing with, the values of  $f$  at the given points.

- Using this polynomial for approximation within the interval given by the endpoints is called polynomial **interpolation**.



# The Lagrange Polynomial: The Linear Case

Define the functions

$$L_0(x) = \frac{x - x_1}{x_0 - x_1} \quad \text{and} \quad L_1(x) = \frac{x - x_0}{x_1 - x_0}.$$

# The Lagrange Polynomial: The Linear Case

Define the functions

$$L_0(x) = \frac{x - x_1}{x_0 - x_1} \quad \text{and} \quad L_1(x) = \frac{x - x_0}{x_1 - x_0}.$$

## Definition

The linear **Lagrange interpolating polynomial** through  $(x_0, y_0)$  and  $(x_1, y_1)$  is

$$P(x) = L_0(x)f(x_0) + L_1(x)f(x_1) = \frac{x - x_1}{x_0 - x_1}f(x_0) + \frac{x - x_0}{x_1 - x_0}f(x_1).$$

# The Lagrange Polynomial: The Linear Case

$$P(x) = L_0(x)f(x_0) + L_1(x)f(x_1) = \frac{x - x_1}{x_0 - x_1}f(x_0) + \frac{x - x_0}{x_1 - x_0}f(x_1).$$

Note that

$$L_0(x_0) = 1, \quad L_0(x_1) = 0, \quad L_1(x_0) = 0, \quad \text{and} \quad L_1(x_1) = 1,$$

# The Lagrange Polynomial: The Linear Case

$$P(x) = L_0(x)f(x_0) + L_1(x)f(x_1) = \frac{x - x_1}{x_0 - x_1}f(x_0) + \frac{x - x_0}{x_1 - x_0}f(x_1).$$

Note that

$$L_0(x_0) = 1, \quad L_0(x_1) = 0, \quad L_1(x_0) = 0, \quad \text{and} \quad L_1(x_1) = 1,$$

which implies that

$$P(x_0) = 1 \cdot f(x_0) + 0 \cdot f(x_1) = f(x_0) = y_0$$

and

$$P(x_1) = 0 \cdot f(x_0) + 1 \cdot f(x_1) = f(x_1) = y_1.$$

# The Lagrange Polynomial: The Linear Case

$$P(x) = L_0(x)f(x_0) + L_1(x)f(x_1) = \frac{x - x_1}{x_0 - x_1}f(x_0) + \frac{x - x_0}{x_1 - x_0}f(x_1).$$

Note that

$$L_0(x_0) = 1, \quad L_0(x_1) = 0, \quad L_1(x_0) = 0, \quad \text{and} \quad L_1(x_1) = 1,$$

which implies that

$$P(x_0) = 1 \cdot f(x_0) + 0 \cdot f(x_1) = f(x_0) = y_0$$

and

$$P(x_1) = 0 \cdot f(x_0) + 1 \cdot f(x_1) = f(x_1) = y_1.$$

So  $P$  is the unique polynomial of degree at most 1 that passes through  $(x_0, y_0)$  and  $(x_1, y_1)$ .

# The Lagrange Polynomial: The Linear Case

## Example: Linear Interpolation

Determine the linear Lagrange interpolating polynomial that passes through the points  $(2, 4)$  and  $(5, 1)$ .

# The Lagrange Polynomial: The Linear Case

## Example: Linear Interpolation

Determine the linear Lagrange interpolating polynomial that passes through the points  $(2, 4)$  and  $(5, 1)$ .

## Solution

In this case we have

$$L_0(x) = \frac{x - 5}{2 - 5} = -\frac{1}{3}(x - 5) \quad \text{and} \quad L_1(x) = \frac{x - 2}{5 - 2} = \frac{1}{3}(x - 2),$$

# The Lagrange Polynomial: The Linear Case

## Example: Linear Interpolation

Determine the linear Lagrange interpolating polynomial that passes through the points  $(2, 4)$  and  $(5, 1)$ .

## Solution

In this case we have


$$L_0(x) = \frac{x-5}{2-5} = -\frac{1}{3}(x-5) \quad \text{and} \quad L_1(x) = \frac{x-2}{5-2} = \frac{1}{3}(x-2),$$

so

$$P(x) = -\frac{1}{3}(x-5) \cdot 4 + \frac{1}{3}(x-2) \cdot 1 = -\frac{4}{3}x + \frac{20}{3} + \frac{1}{3}x - \frac{2}{3} = -x + 6.$$




# The Lagrange Polynomial: The Linear Case



The linear Lagrange interpolating polynomial that passes through the points  $(2, 4)$  and  $(5, 1)$ .

# The Lagrange Polynomial: Degree $n$ Construction



To generalize the concept of linear interpolation, consider the construction of a polynomial of degree at most  $n$  that passes through the  $n + 1$  points

$$(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n)).$$

# The Lagrange Polynomial: The General Case

## Constructing the Degree $n$ Polynomial

# The Lagrange Polynomial: The General Case

## Constructing the Degree $n$ Polynomial

- We first construct, for each  $k = 0, 1, \dots, n$ , a function  $L_{n,k}(x)$  with the property that  $L_{n,k}(x_i) = 0$  when  $i \neq k$  and  $L_{n,k}(x_k) = 1$ .

# The Lagrange Polynomial: The General Case

## Constructing the Degree $n$ Polynomial

- We first construct, for each  $k = 0, 1, \dots, n$ , a function  $L_{n,k}(x)$  with the property that  $L_{n,k}(x_i) = 0$  when  $i \neq k$  and  $L_{n,k}(x_k) = 1$ .
- To satisfy  $L_{n,k}(x_i) = 0$  for each  $i \neq k$  requires that the numerator of  $L_{n,k}(x)$  contain the term

$$(x - x_0)(x - x_1) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n).$$

# The Lagrange Polynomial: The General Case

## Constructing the Degree $n$ Polynomial

- We first construct, for each  $k = 0, 1, \dots, n$ , a function  $L_{n,k}(x)$  with the property that  $L_{n,k}(x_i) = 0$  when  $i \neq k$  and  $L_{n,k}(x_k) = 1$ .
- To satisfy  $L_{n,k}(x_i) = 0$  for each  $i \neq k$  requires that the numerator of  $L_{n,k}(x)$  contain the term

$$(x - x_0)(x - x_1) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n).$$

- To satisfy  $L_{n,k}(x_k) = 1$ , the denominator of  $L_{n,k}(x)$  must be this same term but evaluated at  $x = x_k$ .

# The Lagrange Polynomial: The General Case

## Constructing the Degree $n$ Polynomial

- We first construct, for each  $k = 0, 1, \dots, n$ , a function  $L_{n,k}(x)$  with the property that  $L_{n,k}(x_i) = 0$  when  $i \neq k$  and  $L_{n,k}(x_k) = 1$ .
- To satisfy  $L_{n,k}(x_i) = 0$  for each  $i \neq k$  requires that the numerator of  $L_{n,k}(x)$  contain the term


$$(x - x_0)(x - x_1) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n).$$

- To satisfy  $L_{n,k}(x_k) = 1$ , the denominator of  $L_{n,k}(x)$  must be this same term but evaluated at  $x = x_k$ .
- Thus

$$L_{n,k}(x) = \frac{(x - x_0) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n)}{(x_k - x_0) \cdots (x_k - x_{k-1})(x_k - x_{k+1}) \cdots (x_k - x_n)}.$$

# The Lagrange Polynomial: The General Case

$$L_{n,k}(x) = \frac{(x - x_0) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n)}{(x_k - x_0) \cdots (x_k - x_{k-1})(x_k - x_{k+1}) \cdots (x_k - x_n)}.$$





# The Lagrange Polynomial: The General Case

Theorem:  $n$ -th Lagrange interpolating polynomial

# The Lagrange Polynomial: The General Case

## Theorem: $n$ -th Lagrange interpolating polynomial

If  $x_0, x_1, \dots, x_n$  are  $n + 1$  distinct numbers and  $f$  is a function whose values are given at these numbers,

# The Lagrange Polynomial: The General Case

## Theorem: $n$ -th Lagrange interpolating polynomial

If  $x_0, x_1, \dots, x_n$  are  $n + 1$  distinct numbers and  $f$  is a function whose values are given at these numbers, then a unique polynomial  $P(x)$  of degree at most  $n$  exists with

$$f(x_k) = P(x_k), \quad \text{for each } k = 0, 1, \dots, n.$$

# The Lagrange Polynomial: The General Case

## Theorem: $n$ -th Lagrange interpolating polynomial

If  $x_0, x_1, \dots, x_n$  are  $n + 1$  distinct numbers and  $f$  is a function whose values are given at these numbers, then a unique polynomial  $P(x)$  of degree at most  $n$  exists with

$$f(x_k) = P(x_k), \quad \text{for each } k = 0, 1, \dots, n.$$

This polynomial is given by

$$P(x) = f(x_0)L_{n,0}(x) + \cdots + f(x_n)L_{n,n}(x) = \sum_{k=0}^n f(x_k)L_{n,k}(x)$$

where, for each  $k = 0, 1, \dots, n$ ,  $L_{n,k}(x)$  is defined as follows:

# The Lagrange Polynomial: The General Case

$$P(x) = f(x_0)L_{n,0}(x) + \cdots + f(x_n)L_{n,n}(x) = \sum_{k=0}^n f(x_k)L_{n,k}(x)$$

## Definition of $L_{n,k}(x)$

$$\begin{aligned} L_{n,k}(x) &= \frac{(x - x_0)(x - x_1) \cdots (x - x_{k-1})(x - x_{k+1}) \cdots (x - x_n)}{(x_k - x_0)(x_k - x_1) \cdots (x_k - x_{k-1})(x_k - x_{k+1}) \cdots (x_k - x_n)} \\ &= \prod_{\substack{i=0 \\ i \neq k}}^n \frac{(x - x_i)}{(x_k - x_i)} \end{aligned}$$

We will write  $L_{n,k}(x)$  simply as  $L_k(x)$  when there is no confusion as to its degree.

# Outline

- 1 Weierstrass Approximation Theorem
- 2 Inaccuracy of Taylor Polynomials
- 3 Constructing the Lagrange Polynomial
- 4 Example: Second-Degree Lagrange Interpolating Polynomial

# The Lagrange Polynomial: 2nd Degree Polynomial

Example:  $f(x) = \frac{1}{x}$

# The Lagrange Polynomial: 2nd Degree Polynomial

Example:  $f(x) = \frac{1}{x}$

- (a) Use the numbers (called **nodes**)  $x_0 = 2$ ,  $x_1 = 2.75$  and  $x_2 = 4$  to find the second Lagrange interpolating polynomial for  $f(x) = \frac{1}{x}$ .



# The Lagrange Polynomial: 2nd Degree Polynomial

Example:  $f(x) = \frac{1}{x}$

- (a) Use the numbers (called **nodes**)  $x_0 = 2$ ,  $x_1 = 2.75$  and  $x_2 = 4$  to find the second Lagrange interpolating polynomial for  $f(x) = \frac{1}{x}$ .
- (b) Use this polynomial to approximate  $f(3) = \frac{1}{3}$ .

# The Lagrange Polynomial: 2nd Degree Polynomial

## Part (a): Solution

# The Lagrange Polynomial: 2nd Degree Polynomial

## Part (a): Solution

We first determine the coefficient polynomials  $L_0(x)$ ,  $L_1(x)$ , and  $L_2(x)$ :

$$L_0(x) = \frac{(x - 2.75)(x - 4)}{(2 - 2.5)(2 - 4)} = \frac{2}{3}(x - 2.75)(x - 4)$$

$$L_1(x) = \frac{(x - 2)(x - 4)}{(2.75 - 2)(2.75 - 4)} = -\frac{16}{15}(x - 2)(x - 4)$$

$$L_2(x) = \frac{(x - 2)(x - 2.75)}{(4 - 2)(4 - 2.5)} = \frac{2}{5}(x - 2)(x - 2.75)$$

# The Lagrange Polynomial: 2nd Degree Polynomial

## Part (a): Solution

We first determine the coefficient polynomials  $L_0(x)$ ,  $L_1(x)$ , and  $L_2(x)$ :

$$L_0(x) = \frac{(x - 2.75)(x - 4)}{(2 - 2.5)(2 - 4)} = \frac{2}{3}(x - 2.75)(x - 4)$$

$$L_1(x) = \frac{(x - 2)(x - 4)}{(2.75 - 2)(2.75 - 4)} = -\frac{16}{15}(x - 2)(x - 4)$$

$$L_2(x) = \frac{(x - 2)(x - 2.75)}{(4 - 2)(4 - 2.5)} = \frac{2}{5}(x - 2)(x - 2.75)$$

Also, since  $f(x) = \frac{1}{x}$ :

$$f(x_0) = f(2) = 1/2, \quad f(x_1) = f(2.75) = 4/11, \quad f(x_2) = f(4) = 1/4$$

# The Lagrange Polynomial: 2nd Degree Polynomial

## Part (a): Solution (Cont'd)

Therefore, we obtain

$$\begin{aligned}P(x) &= \sum_{k=0}^2 f(x_k)L_k(x) \\&= \frac{1}{3}(x - 2.75)(x - 4) - \frac{64}{165}(x - 2)(x - 4) + \frac{1}{10}(x - 2)(x - 2.75) \\&= \frac{1}{22}x^2 - \frac{35}{88}x + \frac{49}{44}.\end{aligned}$$

# The Lagrange Polynomial: 2nd Degree Polynomial

$$P(x) = \frac{1}{22}x^2 - \frac{35}{88}x + \frac{49}{44}$$

(b) Use this polynomial to approximate  $f(3) = \frac{1}{3}$ .

Part (b): Solution

# The Lagrange Polynomial: 2nd Degree Polynomial

$$P(x) = \frac{1}{22}x^2 - \frac{35}{88}x + \frac{49}{44}$$

(b) Use this polynomial to approximate  $f(3) = \frac{1}{3}$ .

## Part (b): Solution

An approximation to  $f(3) = \frac{1}{3}$  is

$$f(3) \approx P(3) = \frac{9}{22} - \frac{105}{88} + \frac{49}{44} = \frac{29}{88} \approx 0.32955.$$

# The Lagrange Polynomial: 2nd Degree Polynomial

$$P(x) = \frac{1}{22}x^2 - \frac{35}{88}x + \frac{49}{44}$$

(b) Use this polynomial to approximate  $f(3) = \frac{1}{3}$ .

## Part (b): Solution

An approximation to  $f(3) = \frac{1}{3}$  is

$$f(3) \approx P(3) = \frac{9}{22} - \frac{105}{88} + \frac{49}{44} = \frac{29}{88} \approx 0.32955.$$

Earlier, we we found that no Taylor polynomial expanded about  $x_0 = 1$  could be used to reasonably approximate  $f(x) = 1/x$  at  $x = 3$ .



# Second Lagrange interpolating polynomial for $f(x) = \frac{1}{x}$

