Complete Conic Section Theory Explained Clearly

Table of Contents

- Grade 12 Basic Mathematics (Chapter 8)
- Part 1: Understanding What is a Cone
- Part 2: What is a Conic Section?
- Part 3: Four Ways a Plane Can Cut a Cone
- Summary: Eccentricity Tells You Everything
- Part 4: Circle Theory (Section 8.2)
- Part 5: Special Cases of Circle Equations
- Part 6: Line and Circle Intersection
- Part 7: Tangent and Normal to Any Curve
- Part 8: Condition of Tangency (Section 8.3)
- Part 9: Equation of Tangent to Circle (Section 8.4)
- Part 10: Equation of Normal to Circle (Section 8.5)
- Part 11: Length of Tangent from External Point
- Quick Reference: All Important Formulas
- Final Tips for Understanding
- Key Takeaways

Grade 12 Basic Mathematics (Chapter 8)

Part 1: Understanding What is a Cone

What is a Cone?

Imagine a pointed hat (a cone). Here are the basic parts:

Key Parts of a Cone:

- **Vertex (O)**: The sharp point at the top of the cone
- **Axis (OC)**: The straight line going down the middle from the tip
- **Generator (OA)**: Any straight line on the surface of the cone from the tip to the bottom
- **Semi-vertical angle**: The angle between the axis and any generator

What is a Double Cone?

If you have two cones placed point-to-point (one facing up, one facing down), that is a double right cone. They are mirror images of each other.

Part 2: What is a Conic Section?

The Simple Definition

A conic section is the curve you get when a plane (flat surface) cuts through a cone.

The shape of the curve depends entirely on the angle at which the plane cuts the cone. Think of it like slicing through a cone at different angles—you get different shapes.

The Mathematical Definition

The locus of a point which moves in a plane such that the ratio of its distance from a fixed point to its distance from a fixed straight line is constant is called a conic section.

That sounds complicated, but here's what it means:

- Imagine a special point called the **focus**
- Imagine a straight line called the **directrix**
- Take any point on the curve
- Measure the distance from that point to the focus
- Measure the distance from that point to the directrix
- Divide the first distance by the second distance
- **You always get the same number!**

This constant number is called **eccentricity** (written as **e**).

Important Terms

- **Focus**: The special fixed point
- **Directrix**: The special fixed straight line
- **Eccentricity (e)**: The constant ratio—how stretched or squashed the curve is
- **Axis**: The line passing through the focus and perpendicular to the directrix
- **Vertex**: The point where the curve meets the axis (the closest point to the directrix)

Part 3: Four Ways a Plane Can Cut a Cone

When you cut a cone with a plane at different angles, you get four different conic sections:

1. Circle

When does this happen?
When the plane cuts the cone perpendicular to the axis (straight across, like cutting a cake horizontally).

What shape do you get?
A perfect circle.

Formula for Eccentricity:
$e = 0$ (exactly zero)

Visual: Imagine taking a knife and cutting straight down across the cone—you get a perfect circle.

2. Ellipse

When does this happen?
When the plane cuts the cone at an angle with the axis, but the angle is GREATER than the semi-vertical angle.

What shape do you get?
An oval shape (stretched circle).

Formula for Eccentricity:
(between 0 and 1, but not including 0 or 1)

Visual: Imagine tilting the knife slightly and cutting at an angle—you get an oval.

3. Parabola

When does this happen?
When the plane cuts the cone parallel to the generator (parallel to one of the slanted lines on the cone's surface), and the plane does NOT pass through the vertex.

What shape do you get?
A U-shaped curve that goes on forever.

Formula for Eccentricity:
$e = 1$ (exactly 1)

Visual: Imagine the knife tilted so it's parallel to the cone's slanted surface—you get a U shape that never closes.

4. Hyperbola

When does this happen?
When the plane cuts through BOTH parts of a double cone. The angle between the axis and the plane is LESS than the semi-vertical angle.

What shape do you get?
Two separate U-shaped curves, opening away from each other.

Formula for Eccentricity:
 (greater than 1)

Visual: Imagine cutting through both the top and bottom cone at a steep angle—you get two separate curves.

Summary: Eccentricity Tells You Everything

The eccentricity value instantly tells you which conic section you have:

Eccentricity Value	amp; **Shape**	amp; **How It Forms**	
$e = 0$	amp; Circle	amp; Cut perpendicular to axis	
	amp; Ellipse	amp; Cut at angle	gt; semi-vertical angle
$e = 1$	amp; Parabola	amp; Cut parallel to generator	
	amp; Hyperbola	amp; Cut through double cone	

Table 1: Eccentricity and Conic Sections

Part 4: Circle Theory (Section 8.2)

What is a Circle?

Definition: The locus of a moving point such that its distance from a fixed point is always constant.

Simple meaning: If you keep a point always the same distance from a center point, you draw a circle.

- **Fixed point** = Center
- **Constant distance** = Radius
- **Moving point** = Any point on the circle

Circle Equations

Type 1: Circle with Center at the Origin (0, 0)

Equation:
$x^2 + y^2 = r^2$

Where:

- r is the radius

Example: If $r = 5$, the equation is $x^2 + y^2 = 25$

All points (x, y) satisfying this equation are on the circle.

Type 2: Circle with Center at Any Point (h, k)

Equation:
$(x - h)^2 + (y - k)^2 = r^2$

Where:

- (h, k) is the center
- r is the radius

Example: If center is at $(3, 4)$ and radius is 5:
$(x - 3)^2 + (y - 4)^2 = 25$

Type 3: General Equation of a Circle

Equation:
$x^2 + y^2 + 2gx + 2fy + c = 0$

From this, we can find:

- **Center:** $(-g, -f)$
- **Radius:** $\sqrt{g^2 + f^2 - c}$

This is the most general form used in problems.

Part 5: Special Cases of Circle Equations

Case 1: Circle Touching the X-axis

When a circle touches the x-axis:

- The radius equals the y-coordinate of the center
- That is: $r = k$ (where k is the y-coordinate of center)

Equation:
$(x - h)^2 + (y - k)^2 = k^2$

Why? Because the lowest point of the circle just touches the x-axis, so the distance from center to x-axis equals the radius.

Case 2: Circle Touching the Y-axis

When a circle touches the y-axis:

- The radius equals the x-coordinate of the center
- That is: $r = h$ (where h is the x-coordinate of center)

Equation:
$(x - h)^2 + (y - k)^2 = h^2$

Why? Because the leftmost (or rightmost) point of the circle just touches the y-axis.

Case 3: Circle Touching Both Axes

When a circle touches both the x-axis AND y-axis:

- The radius must be equal to both coordinates
- That is: $r = h = k$

Equation:
$(x - h)^2 + (y - h)^2 = h^2$

Why? The circle must be equidistant from both axes.

Case 4: Circle with Diameter Endpoints Given

If the ends of a diameter are (x_1, y_1) and (x_2, y_2):

Equation:
$(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$

Why? Any point on the circle creates a right angle with the diameter ends (Thales' theorem).

Part 6: Line and Circle Intersection

How Does a Line Intersect a Circle?

Given:

- Line equation: $y = mx + c$
- Circle equation: $x^2 + y^2 = a^2$

To find intersection points, substitute the line equation into the circle:
$x^2 + (mx + c)^2 = a^2$

This expands to:
$(1 + m^2)x^2 + 2mcx + (c^2 - a^2) = 0$

This is a quadratic equation in x. The solutions tell us about intersection.

Three Possible Cases

Case 1: Line Cuts Circle at Two Different Points

Condition: The quadratic has two real and distinct roots

What it means: The line crosses the circle at two separate points

Case 2: Line Touches Circle at One Point

Condition: The quadratic has two real and equal roots (discriminant = 0)

What it means: The line is a tangent—just touching the circle at one point

Case 3: Line Doesn't Touch Circle

Condition: The quadratic has imaginary roots (discriminant < 0)

What it means: The line misses the circle completely

Part 7: Tangent and Normal to Any Curve

Understanding Tangent and Normal

What is a Tangent?

Definition: A tangent is a line that touches a curve at exactly one point.

How to imagine it:

1. Start with a point P on the curve
2. Take another point Q very close to P on the curve
3. Draw a line through P and Q (this is a secant)
4. Now move Q closer and closer to P
5. When Q reaches P, the secant becomes a tangent

Key property: A tangent meets the curve at two coincident (same) points.

What is a Normal?

Definition: A normal is a line perpendicular to the tangent at the point of contact.

For circles specifically:

- The normal always passes through the center of the circle
- The normal is perpendicular to the radius at the point of contact

Part 8: Condition of Tangency (Section 8.3)

When is a Line Tangent to a Circle?

Question: How do we know if a line touches a circle?

Answer: The perpendicular distance from the center to the line must equal the radius.

Finding the Tangency Condition

Given:

- Line: $y = mx + c$
- Circle: $x^2 + y^2 = a^2$ (center at origin, radius a)

Distance from center (0,0) to line $y = mx + c$:
$\text{Distance} = \frac{|c|}{\sqrt{1+m^2}}$

For tangency:
$\frac{|c|}{\sqrt{1+m^2}} = a$

Solving for c:
$c = \pm a\sqrt{1+m^2}$

Conclusion

The line $y = mx + c$ is tangent to the circle $x^2 + y^2 = a^2$ if:
$c = a\sqrt{1+m^2}$

Therefore, the tangent lines are:
$y = mx \pm a\sqrt{1+m^2}$

These two lines are always tangent to the circle $x^2 + y^2 = a^2$.

Part 9: Equation of Tangent to Circle (Section 8.4)

Method 1: General Method (Works for Any Curve)

Given: Point (x_1, y_1) on circle $x^2 + y^2 = a^2$

Steps:

1. Let $P(x_1, y_1)$ be on the circle
2. Take another point $Q(x_2, y_2)$ very close to P on the circle
3. Find the slope of line PQ:
   $\text{Slope} = \frac{y_2 - y_1}{x_2 - x_1}$
4. As Q approaches P, this becomes the tangent slope
5. Using the limiting process, we get:

Tangent equation at (x_1, y_1) on $x^2 + y^2 = a^2$:
$x_1x + y_1y = a^2$

How to remember: Replace x^2 with x_1x, and y^2 with y_1y.

Example: Find tangent to $x^2 + y^2 = 25$ at point $(3, 4)$

Using the formula: $3x + 4y = 25$

Method 2: Special Method for Circles

Why this works: In a circle, the tangent is always perpendicular to the radius at the point of contact.

Given: Point $P(x_1, y_1)$ on circle $x^2 + y^2 = a^2$

Step 1: The radius OP has slope $\frac{y_1}{x_1}$

Step 2: The tangent is perpendicular to the radius, so its slope is $-\frac{x_1}{y_1}$

Step 3: Equation of tangent passing through (x_1, y_1) with slope $-\frac{x_1}{y_1}$:
$y - y_1 = -\frac{x_1}{y_1}(x - x_1)$

Simplifying:
$x_1x + y_1y = x_1^2 + y_1^2 = a^2$

Same answer!

Tangent to General Circle

Given: Circle $x^2 + y^2 + 2gx + 2fy + c = 0$ and point (x_1, y_1) on it

Tangent equation:
$xx_1 + yy_1 + g(x + x_1) + f(y + y_1) + c = 0$

How to remember: Replace:

- x^2 with xx_1
- y^2 with yy_1
- $2x$ with $(x + x_1)$
- $2y$ with $(y + y_1)$

Part 10: Equation of Normal to Circle (Section 8.5)

Normal to Circle $x^2 + y^2 = a^2$

Given: Point $P(x_1, y_1)$ on the circle

Key fact: The normal passes through the center and the point of contact.

Step 1: Tangent equation at P is: $x_1x + y_1y = a^2$

Step 2: Slope of tangent $= -\frac{x_1}{y_1}$

Step 3: Slope of normal (perpendicular to tangent) $= \frac{y_1}{x_1}$

Step 4: Normal passes through (x_1, y_1) with slope $\frac{y_1}{x_1}$:
$y - y_1 = \frac{y_1}{x_1}(x - x_1)$

Simplifying:
$x_1y - x_1y_1 = y_1x - y_1x_1$

$x_1y = y_1x$

Or:
$\frac{x}{x_1} = \frac{y}{y_1}$

Key insight: The normal always passes through the center (origin in this case).

Part 11: Length of Tangent from External Point

What Does This Mean?

If you have a point outside a circle and draw a tangent line to the circle from that point, the tangent touches the circle at one point. The length from your external point to the touching point is the "length of tangent."

Finding the Length

Given:

- Circle: $x^2 + y^2 + 2gx + 2fy + c = 0$
- External point: $P(x_1, y_1)$

Center of circle: $C(-g, -f)$

Radius: $r = \sqrt{g^2 + f^2 - c}$

Using Pythagorean theorem:

The tangent, radius, and line from center to external point form a right triangle:

$$(\text{Length of tangent})^2 = (\text{Distance PC})^2 - (\text{Radius})^2$$

$$(\text{Length of tangent})^2 = [(x_1 + g)^2 + (y_1 + f)^2] - (g^2 + f^2 - c)$$

Simplifying:

$$(\text{Length of tangent})^2 = x_1^2 + y_1^2 + 2gx_1 + 2fy_1 + c$$

Therefore:

$$\text{Length of tangent} = \sqrt{x_1^2 + y_1^2 + 2gx_1 + 2fy_1 + c}$$

For Circle at Origin

If the circle is $x^2 + y^2 = a^2$:

$$\text{Length of tangent} = \sqrt{x_1^2 + y_1^2 - a^2}$$

Quick Reference: All Important Formulas

Topic	amp; **Formula**
Circle, center at $(0, 0)$	amp; $x^2 + y^2 = r^2$
Circle, center at (h, k)	amp; $(x - h)^2 + (y - k)^2 = r^2$
General circle	amp; $x^2 + y^2 + 2gx + 2fy + c = 0$
Center from general form	amp; $(-g, -f)$
Radius from general form	amp; $\sqrt{g^2 + f^2 - c}$
Tangent at (x_1, y_1) on $x^2 + y^2 = a^2$	amp; $x_1x + y_1y = a^2$
Normal at (x_1, y_1) on $x^2 + y^2 = a^2$	amp; $xy_1 = yx_1$
Length of tangent from (x_1, y_1)	amp; $\sqrt{x_1^2 + y_1^2 + 2gx_1 + 2fy_1 + c}$
Tangency condition for $y = mx + c$	amp; $c^2 = a^2(1 + m^2)$

Table 2: All Important Circle Formulas

Final Tips for Understanding

1. **Draw pictures** for everything. Visualization helps you understand.
2. **Learn the eccentricity concept first** before memorizing equations.
3. **Understand the logic** behind formulas, not just memorize them.
4. **Practice with real examples** and work through problems step by step.
5. **Remember that conic sections are everywhere** in real life—use that connection.
6. **Master circles first** because they are the foundation for understanding other conic sections.
7. **Take time to understand** tangents and normals—they appear in many problems.
8. **Don't rush.** Mathematics makes sense when you take time to learn properly.

Key Takeaways

- A conic section is formed by cutting a cone with a plane
- The eccentricity (e) value determines the type: $e = 0$ (circle), (ellipse), $e = 1$ (parabola), (hyperbola)
- Circles are defined by a center and radius
- A tangent touches a curve at exactly one point and is perpendicular to the radius at that point
- A normal is perpendicular to the tangent and passes through the center for circles
- All these concepts can be expressed as equations that help solve problems

Now you understand each part of the conic section theory! Keep practicing, and you'll master it completely.

[1]

⁂

1. Conic-sectin-initial-pdf.pdf