Detecting and quantifying change in extremes: a synthetic-data test of the Müller–Watson methods

Produced by Ken Caldeira with the help of Claude Code
https://github.com/Sustainable-Solutions-Lab/extreme-trend-detection

Draft, 4 November 2026

Abstract

Questions and approach. We asked four questions about the methods of Müller and Watson (2024) for extremes that change over time:

1. How reliably do they detect a change?
2. How accurately do they estimate the current 1-in-100-year level?
3. What happens when the model of the change is wrong?
4. How close to the true risk is their Bayesian model's posterior-predictive 1-in-100-year level, which builds in parameter uncertainty, and why?

To answer them against a known truth, we generated synthetic daily records of 30 to 150 years from four distributions (temperature-like, precipitation-like, heavy-tailed and bounded), changed their mean, spread or tail in known ways, and applied the tests and estimators to the annual maxima: thousands of records per case for the tests and 100 per case for the estimates (Table 1 lists the dimensions varied).

Detection. The parameter-constancy (Nyblom) test of Müller and Watson (2024) is a reliable detector of change in annual maxima: with one value per year its false-alarm rate stays at 4–6% for every parent distribution, record length and level of day-to-day persistence we tried. Its power depends strongly on the record length and on the shape of the daily distribution. A mean shift of half a daily standard deviation over the record is detected in 89% of 85-year records from a normal parent but in 24% from a skewed, precipitation-like (gamma) parent, and in 41% and 9% of 30-year records. With one value per year, however, it is no more powerful than the simpler Mann–Kendall rank test, which detects change about as often or more often in almost every case we simulated; the Nyblom test pulls ahead when it uses the three largest values per year, which must then be declustered.

Quantification. Saying how large the change is, or what the current 1-in-100-year level is, is much harder. Even with no change and the correct model, the 99% annual-maximum level estimated from 85 years is badly off in 44% of records, in the sense that its true chance of being exceeded is more than twice or less than half the nominal 1%: it understates the risk in 27% of records and overstates it in 18%. When extremes are changing, the methods we tested, including the authors' Bayesian random-walk model, are badly off in 51–59% of 85-year records and in about three-quarters of 30-year records. The Bayesian model's posterior median is no more accurate than a simple linear trend.

Using the wrong model. Change can be modelled as a function of time or of the mean state; which is right is usually unknown, the record rarely shows it, and the wrong choice matters most when the mean changes nonlinearly or the spread changes without the mean. Averaging the candidate models, weighted by how well each fits, comes close to the best single model whatever the truth.

The posterior-predictive level. Taking the 99th percentile of the Bayesian model's posterior predictive distribution, rather than the median of the draws' levels, gives a level that is right or conservative on average (true exceedance 0.8% at 85 years, 0.4–0.6% at 30–45 years) and rarely understates the risk. In a single record, however, it usually overstates the risk, often by a lot: its typical true exceedance is 0.4%, 0.05% and 0.01% at 85, 45 and 30 years, and it misses twofold in 68%, 84% and 92% of records. Its uncertainty comes from the short record and the prior, not from the computation, and it is as large in records with no change at all.

Contents

1 Questions **3**

2 Methods **3**

3 Detecting change **7**

4 Quantifying change **10**

5 Using the wrong model **30**

6 The Bayesian random-walk model **38**

7 The posterior-predictive 1-in-100-year level **42**

8 Limitations **49**

9 Summary **50**

A Cases and supplementary results **52**

B The posterior-predictive level at 30 years by kind of change **54**

1 Questions

Müller and Watson (2024) develop small-sample inference for extremes whose parameters may change over time: likelihood-ratio tests about the GEV parameters (the tail index, a high quantile, a Pareto tail) and the Nyblom test of parameter constancy, both with critical values that depend on the estimated tail index, and a Bayesian time-varying GEV model with random walks in its parameters, estimated by Hamiltonian Monte Carlo. They apply the methods to weather-disaster damages, stock returns, and city and firm sizes. Their simulations calibrate the size of the tests for data drawn exactly from the GEV model. They do not report power, do not test the methods on data whose extremes are only approximately GEV (the usual situation), treat serial dependence only briefly (an appeal to weak-dependence results and one simulation for the city-size data), and do not check whether the Bayesian model recovers known parameter paths. For climate applications these are the questions that matter:

1. **Detection.** How often is a change of realistic size detected with record lengths typical of reanalyses and stations (30, 45, 85 and 150 years), and how often is a change "detected" when there is none?

2. **Quantification.** How well do the methods estimate the current return level, for example the level the annual maximum exceeds with probability 1% this year ($X_{0.99}$), and its change over the record? Each method gives a best estimate of this fixed but unknown level; for the Bayesian model it is the posterior median.

3. **Using the wrong model.** A change can be modelled as a function of time (trends, random walks) or of the mean state (a covariate). Either could be physically right, and usually we do not know which. What is lost when the change follows one and we model it as a function of the other, and what can be done when we do not know?

4. **The posterior-predictive level.** A Bayesian user who must price risk reports the 99th percentile of the posterior predictive distribution rather than a statistic of the posterior draws. That level is not an estimate of $X_{0.99}$: it is the level whose chance of being exceeded is 1% once parameter uncertainty is averaged in. How close is it to the true risk, and where does its uncertainty come from?

2 Methods

2.1 Study design at a glance

Table 1 lists the dimensions we varied. It is not a full factorial: the persistence and values-per-year variations were run mainly at 85 years, and the quantification study uses one size of change per case.

	Detection (is there a change?)	**Quantification** (what is the current 99% level?)
Daily distribution (parent)	normal (temperature-like); gamma, shape 2 (precipitation-like); Student-t, 6 d.f. (heavy-tailed); beta$(4, 4)$ (bounded)	
What changes	mean; width; tail (gamma, t, beta); position within the bounds (beta)	the same, plus covariate-model cases (width and tail follow the mean; precipitation scaling) and their time-driven counterparts
Shape of the change	none; linear; step at mid-record; step at 80% of the record; accelerating	none; linear; step at mid-record; accelerating
Size of the change	four sizes per feature (e.g. mean shift 0.1–1.0 daily sd)	one size per feature (e.g. mean shift 0.5 daily sd)
Record length	30, 45, 85, 150 years	30, 45, 85 years
Values per year	1, 3 (all lengths); 10 (85 years)	1; 3 (45 and 85 years)
Declustering	none; values at least 3 days apart (10 per year)	values at least 3 days apart (3 per year)
Day-to-day persistence (AR(1) ϕ)	0.5; 0 and 0.8 at 85 years	0.5; 0 at 85 years
Methods	Nyblom test (Müller–Watson), location-trend LR test, two-halves test, Mann–Kendall	14 estimators: constant, last 30 years, two halves, location trend, location + width trend, all-parameter trend, mean-state covariate (two forms, and oracles with the true mean), AIC choice, AIC-weighted average, Bayesian random walks (posterior median and mean)
Bayesian prior	—	authors'; tighter (45 and 85 years)
Records per case	10,000 with no change; 2,000 per change	100
Scores	false-alarm rate (size); detection rate (power)	true exceedance probability of the last year's estimated level: missed $> 2\times$ either way, understated, overstated; error in the estimated change; Bayesian interval coverage and width

Table 1: Study design: the dimensions varied in the detection and quantification studies.

2.2 Extremes model and tests

For each year $t = 1, \ldots, T$ we keep the k largest daily values $X_t^{(1)} \geq \cdots \geq X_t^{(k)}$ (with $k = 1$, the annual maximum). By extreme-value theory these follow approximately the GEV_k distribution with location μ, scale σ and shape (tail index) ξ, with density

$$f(x_1, \ldots, x_k) = \sigma^{-k} \exp\Big[-(1 + \xi z_k)^{-1/\xi} \Big] \prod_{j=1}^{k} (1 + \xi z_j)^{-1/\xi - 1}, \qquad z_j = \frac{x_j - \mu}{\sigma}.$$

The tests compared are:

- **Nyblom** (Müller and Watson, eq. 12; Nyblom, 1989): with scores S_t of the constant-

parameter fit and average information V, $L_T = T^{-2}\sum_t \big(\sum_{l\ge t} S_l\big)' V^{-1} \big(\sum_{l\ge t} S_l\big)$. The test rejects when $L_T \exp(a_0 + a_1\hat{\xi} + a_2\hat{\xi}^2) > 1$, with coefficients calibrated by simulation so that the rejection rate is at most 5% for every true ξ on a grid and otherwise as close to 5% as possible (their Appendix B: ten values of ξ in $[-0.5, 1.5]$). We recalibrated the coefficients in the same way for each record length and k used here, with 40,000 simulated null records per grid value (they used 10,000).

- **Location-trend LR**: GEV with $\mu_t = \mu_0 + \mu_1 t$, likelihood-ratio test of $\mu_1 = 0$ against χ^2_1. This is the standard tool in climate science (e.g. Coles, 2001).
- **Two halves**: separate GEV fits to the first and second halves of the record, LR test of equal parameters, critical values calibrated like the Nyblom test.
- **Mann–Kendall** rank test for a monotone trend in the annual maxima.

2.3 Estimators of the 99% level

Each estimator gives a path $\hat{X}_{0.99}(t)$, the level the annual maximum of year t exceeds with probability 1% (the 100-year return level in that year's climate). All are fitted by maximum likelihood to the same records, except the Bayesian model.

Constant GEV with constant parameters (ignores change).

Last 30 years constant GEV fitted to the most recent 30 years only, a common practical shortcut (equal to the constant fit for 30-year records).

Two halves constant GEV fitted to the second half of the record.

Location trend μ_t linear in time, σ and ξ constant.

Location + width trend μ_t and $\log\sigma_t$ linear in time, ξ constant.

All-parameter trend μ_t, $\log\sigma_t$ and ξ_t linear in time.

Mean-state covariate μ_t and $\log\sigma_t$ linear in c_t, ξ constant, where c_t is a smoothed annual mean of *all* daily values (local-linear smoother, Gaussian kernel, bandwidth 10 years, fixed in advance). This is the usual approach in event attribution, where the covariate is a smoothed (global or local) mean temperature (Philip et al., 2020). A variant with all three parameters linear in the raw annual mean is unstable (the tail index follows the year-to-year noise of the covariate) and is shown only for reference.

Covariate (true mean) as above but with the true mean path as covariate: an oracle that shows the cost of estimating the covariate.

Data-chosen (AIC) for when the form of the change is unknown: of the constant, location-trend, location + width trend and mean-state covariate models, the one with the lowest AIC.

AIC-weighted average the average of the same four models' $X_{0.99}$ paths, with Akaike weights $w_i \propto \exp(-\Delta\mathrm{AIC}_i/2)$.

Bayesian the authors' four-random-walk model (their `GEV_4RW_Model.stan`, unchanged likelihood and priors), posterior median of $X_{0.99}(t)$. The parameters follow random walks in a logistic transform of $\xi_t \in (-0.9, 2.1)$, in $\ln \alpha_t$, $\ln s_t$ and m_t, with $\sigma_t = s_t \alpha_t^{\xi_t}$ and $\mu_t = m_t + (s_t/\xi_t)(\alpha_t^{\xi_t} - 1)$. The innovation standard deviations are $\gamma = \mathrm{sg} \cdot g/\sqrt{T}$ with $g \sim \mathrm{Exp}(1)$ and scale factors $\mathrm{sg} = 1, 0.5, 0.5$ for ξ, α, s (and the scale level for m), the prior means the paper calls g (their eq. 15). We use 4 chains of 1,000 warm-up and 1,000 sampling iterations (`adapt_delta` 0.95, against Stan's default of 0.8 in the authors' runs); the authors used 5,000 and 10,000 for single published fits. Like their replication scripts, we use the draws without the importance-sampling correction for the extended-support likelihood described in their Section 3.2.3, whose weights they report to be essentially 1. We also report the posterior mean, the 99th percentile of the posterior predictive distribution (Section 7), and a run with a tighter prior ($\mathrm{sg} = 0.25$ for all three).

2.4 Synthetic data

A daily standard Gaussian series Z_d with AR(1) correlation ϕ (0.5 unless stated; 0 and 0.8 also run) is mapped through the parent's quantile function, $X_d = m_t + s_t\, \varepsilon(Z_d; \mathrm{tail}_t)$, with m_t, s_t and the tail parameter constant within a year. The parents are standardized so that the baseline daily standard deviation is 1; all levels are in these units. For a daily temperature with a standard deviation of 3–4 K, a change of 0.5 is 1.5–2 K.

Parent	ε	Tail parameter (baseline)	Tail of the extremes
Normal	Z	—	light, $\xi \to 0$ slowly
Gamma	standardized gamma(a)	shape $a = 2$	exponential-type, skewed
Student-t	standardized t_ν	$\nu = 6$	heavy, $\xi = 1/\nu$
Beta	$3(2B-1)$, $B \sim \mathrm{beta}(\alpha, \beta)$	$\alpha = \beta = 4$	bounded, $\xi = -1/\beta$

Because the map from Z to X is increasing within a year, the annual maximum of X is the map applied to the annual maximum of Z, so the true $X_{0.99}(t)$ is known exactly: in closed form for independent days, and from 500,000 simulated years of Z for correlated days (error about 0.005 at $p = 0.99$).

The changes (Appendix, Table 9) move one feature at a time over the record, along a linear, mid-record step or accelerating path (exponential in time, so the feature changes at an increasing rate): the mean, the width ($\log s$), the tail parameter (log of a, ν or β), or, for the beta, the position of the mass within fixed bounds. A second set, the *covariate-model cases*, is built so that a mean-state model is correct:

- *Temperature-like* (normal, t, beta): the mean rises by 0.5 (beta 0.25) along an accelerating path, the width follows it ($\log s_t = 0.25\, m_t$), and for the t and beta so does the tail ($\log \nu_t$ or $\log \beta_t = -0.2\, m_t$, a heavier-tailed t and a more tightly bounded beta as it warms).
- *Precipitation-like* (gamma): the mean is scaled by $\lambda_t = e^{g_t}$ with a 22% rise over the record, so every quantile scales with the mean (a Clausius–Clapeyron-type change); in a variant the gamma shape also falls as it gets wetter (heavier tail).

A third set, the *time-driven* counterparts of the accelerating temperature-like cases, has the same mean path and the same first- and last-year distributions, but the width and tail change linearly in time between those states instead of following the mean (Section 5).

2.5 Scoring

For detection we report the rejection rate at the nominal 5% level with no change (size) and with change (power). For quantification, the main score is the *true exceedance probability* of each record's estimated level in the last year,

$$\pi = \Pr\left(M_T > \hat{X}_{0.99}(T)\right),$$

computed from the exact distribution of the last year's annual maximum M_T. An ideal estimator has $\pi = 1\%$. We report its median over records and count an estimate as a *miss* when $\pi > 2\%$ (risk understated at least twofold) or $\pi < 0.5\%$ (overstated at least twofold). The main score is the share of records missed in either direction, always given with its two parts. The two errors have different consequences: understating the risk leaves people unprepared, overstating it leads to over-design and false alarms. Estimators trade one for the other, so neither alone is a fair summary, and when the true change has a known direction a one-sided score rewards models that err in that direction. This score is comparable across parents and penalizes errors by their consequence for risk rather than by their size in data units. We also report the error of the estimated change $\hat{X}_{0.99}(T) - \hat{X}_{0.99}(1)$ and, for the Bayesian model, the coverage and width of its 90% credible intervals.

2.6 Monte Carlo design

Detection results ("Stage B" in the repository) use 10,000 simulated records with no change and 2,000 per change. Quantification results ("Stage C") use 100 records per case, fitted by all estimators. The same daily noise is used across cases, record lengths and estimators (common random numbers), so comparisons between estimators are paired. A bootstrap over records gives standard errors of 3–4 percentage points on the pooled shares and about 3 points on differences between estimators.

3 Detecting change

3.1 False alarms

With one value per year, all four tests hold their nominal size within Monte Carlo error at every record length and persistence level (Figure 1), except the location-trend LR test for short records (7–10% at $T = 30$), where the χ^2 approximation is poor. Using several values per year from a persistent daily series breaks the independence the GEV_k model assumes: with the three largest values per year, the Nyblom test rejects 7–9% at $\phi = 0.5$ and 21–23% at $\phi = 0.8$, and with ten values 46–51%. Requiring the values to be at least three days apart (declustering) brings the last case down to 9–12%. Mann–Kendall uses only the annual maxima and is unaffected.




Figure 1: False-rejection rate of the four tests with no change, by design (k values per year, day-to-day correlation ϕ, record length T). Dots: mean over the four parents; lines: range. Vertical line: nominal 5%.

3.2 Power

Power rises steeply with record length (Figure 2, Table 2). For a normal parent, a mean shift of 0.5 daily standard deviations over the record is detected by the Nyblom test in 41%, 60%, 89% and 99% of records of 30, 45, 85 and 150 years. The same shift is much harder to detect for skewed or heavy-tailed parents (gamma: 9–41%; t: 11–52%), whose annual maxima are more variable relative to the daily standard deviation. The location-trend LR test is more powerful against the change it is designed for, a linear shift in location (63% at 30 years and 96% at 85 years for the normal mean shift, though part of the edge at 30 years comes from its over-sizing), but much less powerful against a change in tail shape (t parent: 36% against 69% for the Nyblom test at 85 years).

With one value per year the Nyblom test is not the most powerful of the four. The rank-based Mann–Kendall test, which uses only the annual maxima, does well against both kinds of change (93% and 73% in the two examples). Over all 768 simulated changes with one value per year, with rejection rates corrected for size, it detects change at least as often as the Nyblom test, to within a point, in 95% of cases; the average power is 47% for Mann–Kendall, 45% for the Nyblom and LR tests and 37% for the two-halves test. The Nyblom test is ahead mainly for steps late in the record. Its advantage comes with more values per year, which Mann–Kendall cannot use: with the three largest values per year (size-corrected) it detects change at least as often as Mann–Kendall in 94% of cases (average 56% against 47%), and more often than the LR test for tail changes (49% against 39%). In a persistent daily series these values must be declustered to hold the false-alarm rate (Section 3.1).




Figure 2: Rejection rate (nominal 5% level) against the size of the true change in $X_{0.99}$ (computed for independent days; the correlated-day value differs by under 1%), for linear changes in mean, width and tail, one value per year, $\phi = 0.5$. Colour: record length. Solid: Nyblom; dashed: location-trend LR.

			$T = 30$				$T = 45$				$T = 85$				$T = 150$			
Parent	Change	$\Delta X_{0.99}$	Nyb.	LR	MK	Hlv.	Nyb.	LR	MK	Hlv.	Nyb.	LR	MK	Hlv.	Nyb.	LR	MK	Hlv.
Normal	mean 0.5	+0.50	41	**63**	52	24	60	**77**	69	40	89	**96**	93	71	99	**100**	99	95
Normal	width 0.2	+0.81	51	**70**	62	30	71	**84**	80	48	95	**98**	**98**	81	**100**	**100**	**100**	98
Gamma	mean 0.5	+0.50	9	**17**	11	8	15	**23**	18	10	24	**34**	27	17	42	**55**	45	30
Gamma	width 0.2	+1.58	21	**34**	28	14	34	**47**	43	21	58	**69**	68	42	84	**92**	91	69
Gamma	tail 1	-1.98	20	**27**	25	13	28	**35**	**35**	18	47	52	**59**	32	73	73	**83**	55
Student-t	mean 0.5	+0.50	11	**22**	14	8	19	**29**	19	13	30	**46**	32	22	52	**69**	50	39
Student-t	tail 1	-8.59	28	21	**34**	17	42	26	**47**	28	69	36	**73**	53	93	50	**94**	82
Beta	mean 0.25	+0.25	45	**67**	57	28	67	**82**	76	44	92	**97**	96	78	**100**	**100**	**100**	98
Beta	width 0.1	+0.28	44	**66**	55	28	65	**81**	74	44	91	**96**	95	77	**100**	**100**	**100**	98
Beta	tail 0.2	-0.11	38	**56**	49	22	55	**70**	66	34	85	**90**	**90**	68	98	**99**	**99**	92
Beta	alpha 0.5	+0.07	31	**45**	40	19	49	**59**	58	31	80	82	**83**	60	**98**	97	**98**	90
All (mean)	no change	0	4.5	8.0	4.7	4.9	5.0	6.8	5.1	4.9	5.1	5.8	4.9	5.1	5.6	5.5	5.0	5.5

Table 2: Power (%, nominal 5% level) of the Nyblom (Nyb.), location-trend LR, Mann–Kendall (MK) and two-halves (Hlv.) tests for selected linear changes, one value per year, $\phi = 0.5$; bold: the highest in each group. $\Delta X_{0.99}$: true change in the 99% level over the record (baseline daily sd). Last row: size (%) with no change, mean over parents; the LR test's edge at 30 years partly reflects its over-sizing.

4 Quantifying change

4.1 What the estimates look like

Figure 3 shows single 85-year records with the estimated and true paths of $X_{0.99}$. The records were chosen by a fixed rule, as the record of median difficulty (ranking records by the average over the estimators of $|\log(\pi/1\%)|$), so they are typical rather than best or worst cases. Figure 4 shows the spread of the estimated paths over all 100 records for three cases, and Figure 5 the effect of record length.

Figure 6 shows how much depends on the particular record. For a mean shift and for accelerating warming it shows an easy, a typical and a hard record (the 10th, 50th and 90th percentiles of the same difficulty ranking), each with the location trend, the AIC-weighted average and the Bayesian model. In the easy records the estimates end close to the truth (true exceedance probabilities 0.8–2.9%). In the hard ones they miss badly, understating the risk for the mean shift (4–9% against the nominal 1%) and overstating it for the warming (below 0.1% for the AIC average and the Bayesian model). Figure 7 shows typical records at 30, 45 and 85 years: at 30 years even the typical record understates the risk by a factor of 2 to 9.

Three features stand out. First, the estimates scatter widely about the truth: for a normal parent with 85 years and a mean shift of 0.5, the 5–95% range of the error of the last year's $X_{0.99}$ spans 0.7 daily standard deviations for the location trend and about 1.0 for the more flexible estimators, larger than the change being estimated. Second, a model with the wrong form for the change is biased in a predictable way: the constant fit is flat, and the location trend, a straight line, misses the late acceleration in the covariate-model cases. Third, the Bayesian posterior median follows the truth on average but with paths that are too flat toward the end of the record, and with 30 years its credible band is very wide upward.




Figure 3: Four typical 85-year records (points: annual maxima) with the true 99% level (black) and five estimates. Shaded: Bayesian 90% credible interval. Units: baseline daily standard deviation.




Figure 4: Spread of the estimated 99% level over 100 records of 85 years (median, 25–75% and 5–95% of the estimates in each year) for a linear mean shift (top) and two covariate-model cases (middle, bottom), with the truth in black.




Figure 5: Effect of record length: the location-trend (blue) and Bayesian (orange) estimates for 30, 45 and 85 years. The total change over the record is the same in each column, so the change per year is larger for shorter records.




Figure 6: The luck of the draw: for the same case, an easy, a typical and a hard 85-year record (10th, 50th and 90th percentiles of the average error of the three estimates in the last year). Points: annual maxima; black: true 99% level; lines: estimates; shaded: Bayesian 90% interval. Above each panel: the true probability that each estimate of the last year's level is exceeded (ideal 1%; Loc.: location trend, AIC: AIC-weighted average, Bayes: Bayesian median). Units: baseline daily sd.




Figure 7: Typical single records of 30, 45 and 85 years for the two cases of Figure 5. Same layout as Figure 6.

4.2 Single records, case by case

Figures 8–11 show, for every case of each parent, the typical 85-year record (chosen by the same rule as in Figure 3) with three estimates: the location trend, the AIC-weighted average and the Bayesian posterior median with its 90% interval. The line above each panel gives the true probability that each estimate of the last year's 99% level is exceeded (ideal 1%). Single records show things that pooled scores hide.

- **Even a typical record is usually off by a factor of 2 somewhere.** In 23 of the 24 typical records at least one of the three estimates misses the true risk by more than a factor of 2. Over all 85-year standard records, at least one of the three misses in 75% and all three in 36%.
- **Estimates from one record err together.** They are fitted to the same few extremes, so a record with unusually large or small maxima pulls them all the same way: the correlation between their log exceedance probabilities over records is 0.56–0.76. Agreement between methods applied to one record is therefore weak evidence that they are right.
- **Flexible estimates can invent a change.** In the typical no-change records for the gamma and t parents, a few large values in the second half make the AIC-weighted average and the

Bayesian model rise when nothing changes, so they overstate the current risk (exceedance 0.3–0.7%), while the location trend stays near the truth (0.9–1.2%).

- **A small error in level can be a large error in risk.** For the bounded beta parent the estimates look close to the truth, yet their exceedance probabilities range from below 0.1% to nearly 4%, because its annual maxima vary little from year to year, so the probability of exceeding a level changes quickly with the level.




Figure 8: Normal parent: the typical 85-year record for every case. Points: annual maxima; black: true 99% level; location trend (blue), AIC-weighted average (red, dashed), Bayesian median (orange) with its 90% interval (shaded). Above each panel: true probability that each estimate of the last year's level is exceeded (ideal 1%; Loc.: location trend, AIC: AIC-weighted average, Bayes: Bayesian median). Units: baseline daily sd.




Figure 9: Gamma (precipitation-like) parent: the typical 85-year record for every case. Points: annual maxima; black: true 99% level; location trend (blue), AIC-weighted average (red, dashed), Bayesian median (orange) with its 90% interval (shaded). Above each panel: true probability that each estimate of the last year's level is exceeded (ideal 1%; Loc.: location trend, AIC: AIC-weighted average, Bayes: Bayesian median). Units: baseline daily sd.




Figure 10: Student-t (heavy-tailed) parent: the typical 85-year record for every case. Points: annual maxima; black: true 99% level; location trend (blue), AIC-weighted average (red, dashed), Bayesian median (orange) with its 90% interval (shaded). Above each panel: true probability that each estimate of the last year's level is exceeded (ideal 1%; Loc.: location trend, AIC: AIC-weighted average, Bayes: Bayesian median). Units: baseline daily sd.




Figure 11: Beta (bounded) parent: the typical 85-year record for every case. Points: annual maxima; black: true 99% level; location trend (blue), AIC-weighted average (red, dashed), Bayesian median (orange) with its 90% interval (shaded). Above each panel: true probability that each estimate of the last year's level is exceeded (ideal 1%; Loc.: location trend, AIC: AIC-weighted average, Bayes: Bayesian median). Units: baseline daily sd.

4.3 Spread over records, case by case

Figures 12–15 repeat Figure 5 for every standard case, one figure per parent, comparing the location trend (blue) with the Bayesian posterior median (orange). Four patterns recur.

- **Mean and width changes are tracked on average by both estimators**, in all four parents. What changes with record length is the spread: the 5–95% band at 85 years is 37–50% as wide as at 30 years.
- **A step change is smoothed into a ramp.** Neither estimator locates the step; the location trend replaces it by a straight line, and the Bayesian median by a gentler rise.
- **Short records from skewed or heavy-tailed parents give the Bayesian model a long upper tail.** At 30 years the upper edge of its 5–95% band for the gamma and t parents lies far above the truth (up to about 2.5 times the true level for the t), because the posterior of the tail index is wide and right-skewed when there are few maxima.

- **Tail changes are where the two estimators differ most.** The location trend cannot represent a change in tail: it captures about a third of the fall in the 99% level for the gamma parent and almost none (5–7%) for the t. The Bayesian median captures more, 51–64% for the gamma and 37–50% for the t, increasing with record length. In the bounded beta parent the location trend does the opposite and overshoots the tail and position changes (by a factor of about 1.9 for the tail change), presumably because a straight-line location with fixed scale and shape is the only way it can absorb a change in the shape of the maxima near the bound.




Figure 12: Normal parent, all standard changes. Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Columns: record length; the total change over the record is the same in each column. Units: baseline daily sd.




Figure 13: Gamma (precipitation-like) parent, all standard changes. Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Columns: record length; the total change over the record is the same in each column. Units: baseline daily sd.




Figure 14: Student-t (heavy-tailed) parent, all standard changes. Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Columns: record length; the total change over the record is the same in each column. Units: baseline daily sd.




Figure 15: Beta (bounded) parent, all standard changes. Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Columns: record length; the total change over the record is the same in each column. Units: baseline daily sd.

4.4 The noise floor

Much of the error has nothing to do with change. With no change and the correctly specified constant model, the estimated $X_{0.99}$ misses the risk by more than a factor of 2 in 44% of 85-year records: it understates it at least twofold in 27% and overstates it at least twofold in 18%. For 45-year records 68% miss (38% and 30%), and for 30-year records 71% (44% and 27%). The median exceedance probability is 1.3% rather than 1%: the sampling distribution of an estimated high quantile is skewed, so the typical estimate is a little low. This floor is the yardstick for the results below. An estimator that allows for change should not be expected to beat it, and it pays for its flexibility with extra variance when the change is small.

4.5 Accuracy of the current 99% level

Figures 16 and 17 and Table 3 pool the results over the standard cases (20 cases including the four with no change) and the covariate-model cases (7 cases).

- **Estimators differ as much in the direction of their errors as in how often they miss.** At 85 years the location trend understates the risk twofold in 30% of records and overstates it in 26%; the Bayesian median understates in 39% and overstates in 20%. The Bayesian posterior mean is the extreme case: it rarely understates (20%) but overstates in 51%. Below we rank by the total and give the split; Table 3 has both for every estimator.
- **At 85 years an average of simple models does best** over the standard cases: the AIC-weighted average of the constant, location-trend, location + width and covariate models misses 51% of records (30% understated, 20% overstated), 5.5 ± 1.0 points fewer than the location trend or the location + width trend (both 56%). Ignoring change (constant, 65%) or using only the last 30 years (65%) is much worse, and trends in all three parameters (70%) pay for their flexibility with extra variance.
- **The Bayesian model is about as accurate as the simple trends**: it misses 75%, 70% and 59% of records at 30, 45 and 85 years, against 73%, 69% and 56% for the location trend (paired differences of 1–3 points, ± 1.2). Its errors lean toward overstatement at 30 years (39% overstated against 33%) and toward understatement at 85 years (39% understated against 30%). Its posterior mean is a poor point estimate: the posterior of a high quantile is right-skewed, so the mean overstates the level (median exceedance 0.5% at 85 years, and below 0.01% at 30 years, where the posterior has a very long right tail).
- **With 30 or 45 years the method hardly matters**: the location trend, location + width trend, covariate model, AIC choices and Bayesian model miss 73–76% of 30-year and 68–72% of 45-year records, against no-change floors of 71% and 68%.
- **Three values per year help** when declustered: at 85 years the AIC-weighted average misses 44% and the location + width trend 45% (from 51% and 56%), the location trend 51% and the Bayesian model 53%. For the trend models and the average both kinds of miss fall; the Bayesian model gains on understatement only. Figure 18 shows the bands narrowing a little, and the Bayesian median following the t tail change more closely; it also shows the unconverged Bayesian fits for the beta parent (Section 6) as a wide upper band early in the 85-year record.

- **Record length matters more than the method.** Every estimator is worse at 30 years than at 85, and the spread between these methods (3–4 points at 30 and 45 years, 8 at 85) is much smaller than the floor itself.




Figure 16: Distribution over records of the true probability that the estimated 99% level is exceeded in the last year (circle: median; bar: 25–75%; line: 5–95%). Shaded band: within a factor 2 of the ideal 1%.




Figure 17: Share of records in which the estimated 99% level misses the true risk by more than a factor of 2 in either direction (left), and its two parts: risk understated (middle) or overstated (right) at least twofold, pooled over the standard and the covariate-model cases, by record length.




Figure 18: One against three values per year (declustered, at least three days apart), at 45 and 85 years, for four cases. Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Units: baseline daily sd.

	30 yr				45 yr				85 yr				85 yr, 3/yr			
Estimator	med.	miss	>2%	<0.5%	med.	miss	>2%	<0.5%	med.	miss	>2%	<0.5%	med.	miss	>2%	<0.5%
Standard cases																
Constant	1.88	**78**	49	30	1.69	**73**	45	28	1.77	**65**	46	19	1.91	**64**	48	16
Last 30 years	1.88	**78**	49	30	1.62	**71**	44	27	1.94	**65**	49	15	1.96	**63**	49	14
Two halves	2.07	**82**	51	31	1.67	**70**	44	26	1.75	**61**	45	16	1.68	**57**	42	15
Location trend	1.39	**73**	40	33	1.13	**69**	33	35	1.14	**56**	30	26	1.24	**51**	28	23
Loc. + width trend	1.17	**75**	38	37	1.12	**68**	35	33	1.30	**56**	33	23	1.31	**45**	28	17
All-param. trend	1.81	**86**	48	39	1.21	**81**	41	40	1.47	**70**	42	28	1.68	**67**	43	24
Mean-state cov.	1.26	**76**	40	36	1.20	**72**	38	34	1.37	**59**	36	23	1.41	**51**	33	18
Cov., all param.	2.57	**85**	54	31	1.75	**78**	47	31	1.64	**70**	44	26	1.51	**64**	41	24
Data-chosen (AIC)	1.48	**76**	42	33	1.27	**71**	38	33	1.31	**55**	34	21	1.38	**45**	30	16
AIC-weighted avg.	1.47	**75**	41	34	1.17	**69**	36	33	1.30	**51**	30	20	1.40	**44**	28	15
Bayes (median)	1.13	**75**	36	39	0.94	**70**	34	36	1.50	**59**	39	20	1.30	**53**	31	22
Bayes (mean)	0.00	**92**	5	87	0.01	**88**	10	78	0.48	**71**	20	51	0.79	**60**	22	38
Cov., true mean	1.56	**77**	43	34	1.31	**71**	39	32	1.43	**59**	36	23	1.48	**55**	36	20
Cov., all, true mean	1.63	**80**	45	35	1.38	**74**	41	33	1.48	**64**	40	25	1.57	**61**	39	22
Covariate-model cases																
Constant	5.49	**86**	72	14	5.34	**80**	75	5	5.56	**87**	86	1				
Last 30 years	5.49	**86**	72	14	4.14	**77**	67	10	4.32	**78**	72	6				
Two halves	4.17	**87**	63	24	4.15	**80**	67	13	4.42	**80**	76	4				
Location trend	3.58	**83**	65	18	3.19	**76**	65	11	3.42	**74**	71	3				
Loc. + width trend	2.20	**79**	51	28	2.32	**75**	54	21	2.53	**69**	60	9				
All-param. trend	2.94	**90**	55	35	2.25	**81**	52	29	3.28	**77**	62	15				
Mean-state cov.	1.85	**78**	48	30	1.65	**71**	45	26	1.73	**62**	45	16				
Cov., all param.	3.72	**93**	55	39	2.65	**85**	54	31	2.83	**80**	57	23				
Data-chosen (AIC)	2.68	**82**	57	25	2.18	**74**	53	21	2.04	**63**	50	12				
AIC-weighted avg.	2.45	**82**	56	26	2.20	**73**	52	21	2.09	**64**	52	12				
Bayes (median)	1.67	**80**	47	33	1.67	**72**	44	28	2.46	**69**	57	12				
Bayes (mean)	0.00	**94**	5	89	0.01	**89**	12	77	0.69	**72**	28	45				
Cov., true mean	1.40	**77**	43	34	1.39	**70**	41	29	1.61	**63**	43	20				
Cov., all, true mean	2.23	**88**	52	36	1.46	**85**	46	39	2.01	**79**	50	29				

Table 3: Accuracy of the estimated 99% level in the last year: median true exceedance probability (%, ideal 1); share of records (%) in which the level misses the true risk by more than a factor of 2 either way (bold); and its two parts, risk understated (> 2%) or overstated (< 0.5%) at least twofold.

4.6 Estimating the size of the change

Figure 19 shows the estimated change in $X_{0.99}$ from the first year to the last for every case. The location trend is precise but estimates only the part of the change that is a shift in location: it captures between a third and four-fifths of a width change (depending on the parent) and a third or less of a tail change in the gamma and t parents. The location + width trend and the covariate model are roughly unbiased where their form is right, but their 10–90% ranges are as wide as the change itself. The Bayesian model systematically shrinks the change toward zero, most visibly for the large tail change in the t parent and for the precipitation-like cases. For a normal parent with 85 years and a true change of 0.5, the location trend's 10–90% range is 0.33–0.65. The location + width trend and the covariate model span about 0 to 0.9, so even the sign is wrong in about one record in ten. The Bayesian median change is 0.32, against the true 0.5.




Figure 19: Estimated change in the 99% level over 85-year records (median and 10–90% over records) against the truth (black bar), for every case.

5 Using the wrong model

A change in extremes can be modelled as a function of time (a constant, a linear trend in location, linear trends in location and width, or a random walk) or as a function of the mean state, here a smoothed annual mean of all daily values. Physically, either could be right. The width of the temperature distribution may change because the climate has warmed, a function of the mean state, or for reasons with their own time course, such as changes in circulation, land use or aerosols, a function of time. Usually we do not know which. There is therefore no single right model. The useful questions are what is lost when the model's driving variable is the wrong one, and what to do when it is unknown.

When the two kinds of model differ. They coincide when the mean state moves linearly in time, and they differ when it does not (a step, an acceleration) or when the distribution changes without any change in the mean. We classified the cases by their true process (Table 4) and compare two models with the same number of parameters that differ only in their driving variable: the location + width trend (μ and $\log\sigma$ linear in time) and the mean-state covariate model (μ and $\log\sigma$ linear in the smoothed annual mean). To complete the comparison we added time-driven counterparts of the accelerating warming cases: the same mean path and the same first- and last-year distributions, but with the width and tail changing linearly in time instead of following the mean. In each pair the two processes differ only in how the distribution gets from its first state to its last (Figure 21). At the coupling strength we use, the width in mid-record differs by about 4% between the two.

Score. We use the two-sided score of Section 4: the share of records in which the estimated level misses the true risk by more than a factor of 2 *in either direction* (true exceedance probability above 2% or below 0.5%). A one-sided score would mislead here, because the tail changes all lower the 99% level and the mean changes all raise it: a constant fit, for instance, never understates the risk after a tail change. With no change and the correct model, 44% of 85-year records and 71% of 30-year records miss.




Figure 20: Share of records in which the estimated 99% level in the last year misses the true risk by more than a factor of 2 in either direction, by true process (rows) and fitted model (columns). Bold: the model's form contains the true process (approximately); ×: it does not.

	Missed by > 2× either way, % of records							AIC picks, % of records			
True process	Const.	Loc.	Loc.+wid.	Bayes	Cov.	AIC	AIC avg.	Const.	Loc.	Loc.+wid.	Cov.
30 yr											
No change	**71**	**73**	**73**	**73**	**76**	74	74	64	16	5	16
Mean shift, linear in time	82	**72**	**73**	**76**	**74**	74	75	40	44	8	9
Mean shift, step	78	66	76	**74**	**74**	72	70	27	53	5	15
Width change, mean constant	83	76	**74**	**77**	78	79	79	29	40	15	16
Tail change, mean constant	75	79	74	**73**	77	78	74	29	36	15	20
Follows mean state, mean linear in time	83	78	**70**	**77**	**75**	80	74	12	60	11	17
Follows mean state, mean accelerating	87	85	82	**81**	**79**	83	85	18	38	10	34
Follows time, mean accelerating	86	83	78	**78**	79	81	81	13	48	12	27
85 yr											
No change	**44**	**49**	**55**	**53**	**60**	54	47	69	11	4	17
Mean shift, linear in time	67	**49**	**55**	**58**	**56**	52	46	22	59	2	17
Mean shift, step	59	54	55	**56**	**44**	46	44	6	20	2	72
Width change, mean constant	76	54	**55**	**63**	69	55	54	6	71	17	6
Tail change, mean constant	83	75	59	**60**	67	69	62	6	38	38	18
Follows mean state, mean linear in time	82	52	**54**	**64**	**52**	52	52	2	66	5	26
Follows mean state, mean accelerating	90	83	75	**71**	**65**	67	69	3	27	4	66
Follows time, mean accelerating	85	71	68	**68**	64	67	65	2	48	8	42

Table 4: Using the wrong model. Left: share of records (%) in which the estimated 99% level in the last year misses the true risk by more than a factor of 2 either way; bold where the model's form contains the true process. Right: how often AIC selects each candidate. Loc.: location trend; Loc.+wid.: location and width trend; Cov.: mean-state covariate. Appendix Table 10 splits the misses into under- and overstatement.

Findings.

- **At 85 years the wrong driving variable makes the estimate clearly worse.** When the distribution follows an accelerating mean state, the location + width trend misses 75% of records and the covariate model 65%. When the width changes while the mean stays constant, the covariate model, whose covariate then carries no signal, misses 69% and the location + width trend 55%.
- **Unneeded flexibility costs about as much.** With no change, the covariate model misses 60% of records against 44% for the constant fit, because it fits a relation to the year-to-year noise of the annual mean; the location + width trend misses 55%.
- **Where the two kinds of model coincide, the choice hardly matters**: with the mean linear in time, with or without the distribution following it, the location trend, location + width trend and covariate model miss 49–56% of records.
- **A step favours the covariate model** (44% against 54–55% for the time trends), because the smoothed annual mean follows the step and a straight line cannot.
- **For 30-year records the choice matters little**: every model misses 66–87% of records, and within each true process the models differ by 5–13 points, while sampling noise alone causes 71%.
- **When the mean rises at an increasing rate, the mean-state model loses little even if the width follows time.** In the time-driven counterparts of the accelerating warming cases the covariate model's assumption about the width is wrong, yet it misses 64% of 85-year records, against 68% for the location + width trend and the Bayesian model and 71% for the

location trend. The time-trend models do better than when the width follows the mean (68% against 75% for the location + width trend), and the covariate model about the same (64% against 65%). Following the accelerating mean, which a straight line in time cannot, matters more than the driver of the width, which differs little between the two (Figure 21). The exception is the heavy-tailed t parent, whose change is mostly in the tail: there the location + width trend does better (48% against 54%). Our parametric time models are linear; a time model with a curved mean might do better, but we have not tested one.

Can the data tell? AIC selects among the constant, location-trend, location + width trend and covariate models (Table 4, right). With 85 years it picks the covariate model in 66% of the accelerating mean-state cases and 72% of the step cases, but also in 17% of records with no change. It almost never detects a width change: when only the width changes it picks the location trend in 71% of records and the location + width trend in 17%. When the width follows time rather than the mean, it picks the covariate model less often (42% against 66%), so it leans the right way, but it cannot reliably tell the two apart. With 30 years it picks the constant fit or the location trend in most records whatever the truth.

Picking the lowest-AIC model is never far off, but it is sometimes 10 points worse than the best single model: with no change it misses 54% of 85-year records against 44% for the constant fit, and after a tail change 69% against 59%. Averaging the four models with Akaike weights does better (Table 4). At 85 years it is within 4 points of the best single model for every true process, and when the mean moves linearly it beats all of them (46% against 49% for the best). By contrast, every single model, including the Bayesian one, is 11–30 points worse than the best for at least one process. Over the standard cases it is also the best estimator at 85 years (51% missed against 56% for the location trend, Section 4), and in the covariate-model cases it is close to the covariate model (64% against 62%, and 69% for the location + width trend). With 30 and 45 years the differences between all the choices are small, and the average is within 6 points of the best model throughout.

What to do when we do not know.

- **Do not expect the record to reveal the driver.** The time and mean-state models coincide whenever the mean moves linearly in time, and AIC rarely detects a width change. Whether the change follows time or the mean state has to come from physical reasoning or from evidence outside the record.
- **Average rather than choose.** An AIC-weighted average of the constant, time-trend and mean-state models gives up a few points against the best model for the true process, whatever that is, and avoids the large losses of committing to a wrong one.
- **When the mean clearly moves nonlinearly, include a mean-state model.** With an accelerating mean or a step, the covariate model did best whether or not the width followed the mean.
- **Report the spread between plausible models** as part of the uncertainty of the current return level, alongside the sampling uncertainty of each.
- **With 30 years, the model matters less than the data.** Two-thirds or more of 30-year records miss the current 99% level by a factor of 2 whatever is fitted; only longer records or

additional information (other sites, or more declustered values per year) are likely to change that.




Figure 21: The same accelerating warming with the width (and for the t the tail) following the mean state ("width follows mean") or changing linearly in time ("width follows time"); the first- and last-year distributions are the same in each pair. Location + width trend (aqua) against the mean-state covariate model (yellow), which have the same number of parameters. Median (line), 25–75% and 5–95% of the estimated 99% level over 100 records, truth in black.




Figure 22: Cases in which the mean state does not track the change. For a width or tail change at constant mean the covariate model (yellow) has no signal and stays nearly flat; for a step it follows the smoothed step. Location + width trend in aqua. Median, 25–75% and 5–95% over 100 records, truth in black.

Single records. Figure 23 shows typical single records for six of the processes, with the location + width trend, the covariate model and the AIC-weighted average. When the width changes at constant mean, the covariate model stays almost flat and understates the current risk (4.9%), while the location + width trend follows the widening (1.8%). After a step the covariate model follows the smoothed step. With accelerating warming the covariate model ends closer to the truth than the location + width trend whether the width follows the mean (2.7% against 5.4%) or time (2.5% against 3.3%). The AIC-weighted average usually lies between the two.




Figure 23: Typical 85-year records for six true processes: annual maxima (points), true 99% level (black), location + width trend (aqua), mean-state covariate model (yellow) and AIC-weighted average (red, dashed). Above each panel: true probability that each estimate of the last year's level is exceeded (ideal 1%; L+W: location + width trend, Cov.: covariate model, AIC: AIC-weighted average). Units: baseline daily sd.

The covariate-model cases in detail. In these cases the width and tail of the daily distribution change with the mean, and the mean itself rises along an accelerating path. At 85 years the location trend misses the current risk by more than a factor of 2 in 74% of records, almost all by understating it (71%); the location + width trend and the Bayesian model miss 69% (60% and 57% understated). The covariate model does clearly better (62%: 45% understated, 16% overstated), as well as its oracle version with the true mean path (63%). Two things contribute: the covariate captures the acceleration, which a straight line in time cannot, and it is estimated from all daily values, not only the extremes, so it is much less noisy than the extremes themselves. Letting the tail index also depend on the covariate makes things worse (80%), because the tail index then follows the noise in the covariate.

Figures 24 and 25 show the paths. The covariate model (yellow) follows the accelerating rise; the location trend (blue), a straight line, cuts across it and ends well below the truth (by 0.4 daily sd for the normal parent, 40% of the true change), and this shortfall does not shrink with record length, because it comes from the form of the model, not from noise. The covariate model's median also ends somewhat low (by 0.2), but so does its oracle version with the true mean as covariate (by

0.14): this is mostly the downward tendency of a typical high-quantile estimate seen already in the no-change case. Its bands widen at the end of short records, where the smoothed covariate is least certain. For the t parent, whose tail gets heavier as it warms, the shortfall is larger (0.8 against 1.5 for the location trend), because the model holds the tail index constant.




Figure 24: Temperature-like covariate-model cases: the mean rises, and the width (and for the t and beta the tail) follow it. Location trend (blue) against the mean-state covariate model (yellow). Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Columns: record length; the total change over the record is the same in each column. Units: baseline daily sd.




Figure 25: Precipitation-like covariate-model cases (gamma): every quantile scales with the mean, and in the last row the tail also gets heavier. Location trend (blue) against the mean-state covariate model (yellow). Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Columns: record length; the total change over the record is the same in each column. Units: baseline daily sd.

6 The Bayesian random-walk model

Credible intervals. The 90% intervals for the last year's $X_{0.99}$ contain the truth in 94–100% of records, so they are conservative (Figure 26). Their median width is 1.3 to 3 times the 5–95% range of the actual errors of the posterior median. The intervals are honest but too wide to be informative for short records.




Figure 26: Bayesian model. Left: coverage of the nominal 90% interval for the last year's 99% level. Right: median interval width relative to the 5–95% range of the actual errors (1 = right width). One dot per parent; standard cases.

Convergence. Under the authors' prior most fits have some divergent transitions (median 9 of 4,000 at 85 years and 81 at 30 years). With one value per year, R-hat exceeds 1.05 in fewer than 1% of fits. With three values per year it exceeds 1.05 in 10–12% of fits, almost always far above (R-hat > 1.5, chains in different modes), and these are concentrated in the bounded beta parent (25–32% of its fits). Our chains are shorter than the authors', but disagreement between chains of this size is not a matter of chain length. The unconverged fits overstate the risk more often than the converged ones (37–49% against 19–32% overstated twofold), but they are a minority, and excluding them changes the pooled shares by at most 2 points.

Prior sensitivity. Shrinking the random-walk scales for ξ, α and s to 0.25 nearly eliminates the convergence problems (R-hat ≤ 1.05 in every fit, median 1–6 divergences). It improves accuracy slightly at 85 years (56% of records missed against 59%, mostly because fewer are understated: 35% against 39%), leaves 45 years unchanged (70%), and brings interval coverage closer to nominal (93–98%). See Appendix Figure 33. Figure 27 shows the cost: the tighter prior narrows the bands, but it also makes the paths stiffer, so for the t tail change the median follows less of the early, high values. Figure 28 compares both priors with the covariate model in the covariate-model cases. There the tighter prior misses as many records as the authors' (69% at 85 years, against 62% for the covariate model), more of them by understating the risk (60% against 57%), and its intervals cover the truth in only 88% of records (80–92% by case), because its stiffer paths lag the acceleration at the end of the record.




Figure 27: Bayesian posterior median with the authors' prior and a tighter prior on the random-walk scales, at 45 and 85 years. Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Units: baseline daily sd.




Figure 28: Covariate-model cases at 85 years: Bayesian posterior median (orange) with the authors' and the tighter prior, and the mean-state covariate model (yellow). Median (line), 25–75% (dark band) and 5–95% (light band) of the estimated 99% level in each year over 100 records, with the truth in black. Units: baseline daily sd.

Time reversal. The model's prior is symmetric in time, so fitting a record in reverse order should give the reversed posterior paths. For four test cases the largest difference between the forward and reversed posterior medians of $X_{0.99}$ was 0.003–0.11, at most 1% of the width of the 90% interval, consistent with MCMC noise.

7 The posterior-predictive 1-in-100-year level

A Bayesian posterior offers a second way to state the current 1-in-100-year level. Instead of a statistic (median or mean) of the levels implied by the individual posterior draws, one can average the draws' distributions into a single posterior predictive distribution and take its 99th percentile. We call this *the predictive level*. It is a single, well-defined number, it is what a learner who holds the posterior should use to price a contract that pays when the level is exceeded, and it is the natural choice in risk applications. This section asks how close it is to the truth and where its uncertainty comes from. We refitted every Bayesian run of Section 4 with its original seeds and settings (19,700 fits), keeping the draws' distributions; the posterior medians and means reproduce the earlier fits to within 3×10^{-6}, so all comparisons below are on the same records and draws.

The predictive level answers a different question from the estimators of Section 4. They try to estimate a fixed but unknown number, the true $X_{0.99}$, and the posterior median is the Bayesian model's best estimate of it. The predictive level instead states a risk: under everything the posterior still believes, including draws with heavy tails, the chance of exceeding it is 1%. Because it folds parameter uncertainty into the level, it lies above the posterior median by design, and lying above the true level is not in itself a fault. We score it the same way as the estimators, by the true chance of exceeding it, because that is what a user relying on it is exposed to. The question is whether the margin it builds in is the right size, and how much it varies from one record to the next.

7.1 Definition

The model gives, for each posterior draw $d = 1, \ldots, D$, GEV parameters $\theta_d(t) = (\mu_d(t), \sigma_d(t), \xi_d(t))$ for the annual maximum in year t. The draw's own 99th percentile is $q_d(t)$, with $P(Y_t > q_d(t) \mid \theta_d) = 1\%$. The *posterior median* estimate is the median of the $q_d(t)$ over draws. The *predictive level* $x^*(t)$ instead solves

$$\frac{1}{D} \sum_{d=1}^{D} P\big(Y_t > x^*(t) \mid \theta_d(t)\big) = 1\%, \tag{1}$$

the 99th percentile of the posterior predictive distribution of the year-t maximum. It averages probabilities, not levels. Since a probability cannot go above 1 but can fall to zero, and the upper tail of a GEV distribution is very sensitive to ξ, the average is pulled up by the draws with the heaviest tails: in our records x^* is above the posterior median in every record, and above the posterior mean of q_d in two-thirds of records at 85 years (a tenth at 30 years; Figure 31).

Throughout, the quantity studied is this predictive 99th percentile of the annual-maximum distribution, i.e. the 1-in-100-year level, in the final year of the record. A predictive percentile of another quantity (daily rainfall, say) works by the same mechanism, but its numbers will differ.

7.2 How close is the predictive level to the truth?

Table 5 and Figure 29 give the main result, pooled over the standard cases (the four distributions with and without change).

Table 5: True chance of exceeding the reported 1-in-100-year level in the final year, standard cases (2,000 records per design). For the predictive level: median and average over records, and the shares (%) understated ($> 2\%$) and overstated ($< 0.5\%$) by more than a factor of 2. Right: the share missed by the other estimators on the same records, and the difference predictive minus posterior median ($\pm$ one standard error).

	Bayes, predictive					Missed, others			
Design	median	avg.	under	over	**missed**	post. median	loc. trend	AIC avg.	pred. − median
30 yr	0.01%	0.5%	8	84	**92**	75	73	75	$+17 \pm 1$
45 yr	0.05%	0.4%	6	78	**84**	70	69	69	$+14 \pm 1$
85 yr	0.4%	0.8%	12	56	**68**	59	56	51	$+9 \pm 2$
45 yr, 3/yr	0.2%	0.6%	8	70	**78**	62	60	59	$+16 \pm 1$
85 yr, 3/yr	0.6%	0.9%	13	46	**60**	53	51	44	$+7 \pm 1$
85 yr, indep. days	0.3%	0.8%	13	58	**71**	65	58	55	$+6 \pm 2$
45 yr, tight prior	0.2%	0.7%	11	63	**74**	70	69	69	$+4 \pm 1$
85 yr, tight prior	0.6%	1.1%	16	42	**57**	56	56	51	$+1 \pm 1$




Figure 29: True chance of exceeding the reported 1-in-100-year level, over the records of the standard cases: 10–90th percentiles (thin bar), 25–75th (thick bar), median (circle) and average (diamond). The grey band is the factor-of-2 band around 1%.

- **The predictive level is usually too high.** Its median true exceedance is 0.39% at 85 years, 0.05% at 45 and 0.01% at 30. The other estimators are centred near 1% (medians 0.9–1.5%).
- **It misses more often.** It is off by more than a factor of 2 in 68% of records at 85 years, 84% at 45 and 92% at 30, almost always by overstating. The posterior median misses in 59%, 70% and 75%, the location trend in 56%, 69% and 73%.
- **The dispersion is larger, and lopsided.** At 85 years the middle 80% of its true exceedances runs from 0.015% to 2.4%, a factor of about 160; for the posterior median it is 0.19% to 6%,

a factor of about 30. At 30 years, a tenth of predictive levels have essentially no chance of being exceeded in the simulated climate.

- **The level itself is not far off; the probability is.** In the precipitation-like (gamma) records, the predictive 100-year depth is a median 8% above the true depth at 85 years (10–90th percentile −6% to +25%), 22% above at 45 years and 31% above at 30. Because the chance of exceedance falls steeply with depth in the far tail, an 8% overstatement of the depth becomes a factor of 2.5 in the probability.
- **Several values per year help, but only modestly.** Fitting the three largest (declustered) values of each year instead of the annual maximum cuts the share of records whose posterior leans heavy-tailed from 38% to 14% at 45 years and from 12% to 8% at 85 years (Table 7). The predictive level's median true exceedance rises from 0.05% to 0.2% at 45 years and from 0.4% to 0.6% at 85, and its share missed falls from 84% to 78% and from 68% to 60%. But the other estimators gain about as much, so the predictive level still misses 7–16 points more often than the posterior median.

Table 6 breaks the 85-year results down by the kind of change and by distribution. The predictive level overstates the risk more often than it understates it in every kind of case, including no change at all. It does about as well as the posterior median only where the true change widens the distribution or makes its upper tail heavier, so that the other estimators tend to understate the risk: in the width cases (64% missed, against 63%) and when the spread and tail follow a rising mean (68% against 69%). It does worst where the tail becomes lighter (79% missed).

Table 6: The predictive level at 85 years by kind of change and by distribution: median true exceedance, shares (%) understated and overstated by more than a factor of 2, and missed; right, the share missed by other estimators on the same records.

		Bayes, predictive				Missed, others		
Kind of change	records	median	under	over	**missed**	post. median	loc. trend	AIC avg.
No change	400	0.5%	11	52	**64**	53	49	47
Mean	800	0.4%	14	54	**67**	57	52	45
Width	400	0.6%	19	46	**64**	63	54	54
Tail	300	0.2%	3	76	**79**	60	75	62
Position (beta)	100	0.2%	11	66	**77**	76	73	61
Follows mean state	700	0.6%	24	45	**68**	69	74	64
Time-driven	300	0.6%	24	47	**71**	68	71	65
Normal (standard cases)	400	0.5%	15	50	**65**	60	50	46
Gamma (standard cases)	500	0.4%	11	56	**67**	53	50	47
Student-t (standard cases)	500	0.4%	9	55	**64**	50	56	46
Beta (standard cases)	600	0.3%	15	60	**75**	70	66	60

7.3 The average versus a single record

The predictive level does what it is built to do on average: the true chance of exceeding it, averaged over records, is 0.85% at 85 years (0.43–0.55% at 30–45 years), the closest to 1% of the estimators we tested, alongside the posterior mean; the posterior median and the maximum-likelihood fits

average 1.6–3.7% (diamonds in Figure 29). But the average is carried by a minority of records in which the level is too low, while in the typical record it is far too high.

This is the familiar gap between the average and the typical value of a skewed quantity. For a risk taker who writes many independent contracts and cares only about the expected total, the average is the relevant number, and by that standard the predictive level is well calibrated or conservative. For a single contract, location or portfolio exposed to one climate, the typical record is more relevant, and there the predictive level usually implies a risk several times higher than the true one. Climate risks at nearby locations are strongly correlated, and a model and prior applied in the same way everywhere will err in the same direction everywhere, so the diversification that would make the average relevant is limited.

A Bayesian predictive distribution is calibrated in this average sense when the true climates are drawn from the prior. Our synthetic climates are not drawn from the authors' prior: the GEV shape of their annual maxima is at most about 0.2 (the Student-t case), whereas the prior allows ξ up to 2.1, and above $\xi = 1$ even the mean of the distribution is infinite. To the extent that real rainfall extremes have lighter tails than the prior allows, the same overstatement should be expected in real records.

7.4 Pricing cover with a known 1% risk

Figure 30 turns the question around: for the level whose true chance of being exceeded is exactly 1% in the final year, what chance does each model assign? That chance is the pure premium rate the model would charge for cover triggered at that level.




Figure 30: Each model's chance of exceeding the true 1-in-100-year level of the final year, over the records of the standard cases (percentiles as in Figure 29). The grey band is a factor of 2 around the true 1%.

- **The predictive model charges too much, but rarely too little.** Its median rate is 1.7% at 85 years, 2.8% at 45 and 3.2% at 30. It charges less than 0.5% in only 4–10% of records and less than 0.1% almost never; it charges more than 2% in 40% of records at 85 years and in 67–72% at 30–45 years.

- **A trend fit is right on average but often dangerously low.** The location-trend maximum-likelihood fit has a median rate of 0.7–0.9%, but charges less than 0.5% in 36–45% of records and less than 0.1% in 16–28%: in those records the cover would be priced at a tenth or less of its true cost.
- The predictive approach therefore trades frequent and large over-pricing for protection against severe under-pricing. Whether that is a good trade is a question about risk preferences; the point here is that the loading it implies is large, varies by a factor of several from one record to the next, and is set mainly by the prior (Section 7.5).

7.5 Where the uncertainty comes from

The posterior's own spread. Figure 31 shows one typical precipitation-like record of 85 years with no change. Left: each grey curve is the chance of exceeding a given level under one posterior draw. The draws fan out in the tail. The predictive curve (violet) is their average; it lies above most of the draws because the few heavy-tailed ones dominate the average in the far tail. Right: the distribution over draws of the chance of exceeding the predictive level. By construction it averages 1%, but the median draw gives 0.40%, close to the truth (0.48%), and the top 5% of draws carry a third of the average.




Figure 31: One record (precipitation-like distribution, 85 years, no change; the record whose predictive level has the median true exceedance of that case). Left: exceedance curves of 120 posterior draws (grey), their median (orange), their average, i.e. the predictive curve (violet), and the truth (black); dots mark the levels with a 1% chance. Right: the draws' chance of exceeding the predictive level (log scale; draws that put it at or below 10^{-6} are gathered in the leftmost bar).

Table 7 summarises this over all records. The posterior is honest about its uncertainty: the true chance lies within the draws' 5–95% range in 96–100% of records. But the range is very wide, from essentially 0 to about 5%, so the single number hides a very wide spread of beliefs. "Dominated by a few draws" is accurate in a specific sense: the top 5% of draws carry 38% (85 years) to 52% (30 years) of the 1%, and the effective number of draws that determine the average is 13–21% of the total.

Table 7: Uncertainty measures of the predictive level, standard cases. Left: the median over records of the 5th, 50th and 95th percentiles over draws of each draw's chance of exceeding the predictive level. *Heavy-tailed*: the share of records in which more than 5% of the draws have $\xi > 0.5$. *Truth inside*: how often the true chance lies within the draws' 25–75% and 5–95% ranges. *Top 5%*: the share of the 1% carried by the top 5% of draws (median over records). *Monte Carlo*: the typical error of the exceedance probability with 4,000 draws, as a multiplicative factor.

	Draws' chance of exceeding the level				Truth inside			
Design	5%	median	95%	heavy-tailed	50%	90%	top 5%	Monte Carlo
30 yr	0.00%	0.00%	5.8%	53%	64%	96%	52%	×1.11
45 yr	0.00%	0.04%	5.4%	38%	69%	100%	47%	×1.10
85 yr	0.00%	0.2%	4.6%	12%	67%	98%	38%	×1.06
45 yr, 3/yr	0.00%	0.2%	4.5%	14%	69%	98%	35%	×1.06
85 yr, 3/yr	0.00%	0.4%	3.9%	8%	65%	97%	27%	×1.04
85 yr, indep. days	0.00%	0.2%	4.6%	17%	61%	98%	38%	×1.06
45 yr, tight prior	0.00%	0.3%	4.4%	18%	54%	97%	34%	×1.06
85 yr, tight prior	0.00%	0.5%	3.8%	4%	57%	96%	28%	×1.03

The tail shape. Which records are overstated most? Those whose posterior leans towards heavy tails (Figure 32). The true GEV shape of the annual maxima in our climates ranges from about −0.25 (bounded) to about +0.17 (Student-t); with 30–45 years of data the posterior mean of ξ is often well above that, and every positive shift in ξ raises the predictive level steeply. At 30 years, records with a posterior mean ξ above about 0.2 have predictive levels with essentially no chance of being exceeded.




Figure 32: True chance of exceeding the predictive level against the posterior mean of the GEV shape ξ in the final year, standard cases at 30, 45 and 85 years (one dot per record; lines: medians within octiles of ξ).

The prior. The predictive level is "unique" only for a given prior, and the prior matters a great deal. Shrinking the prior scales of the random walks from the authors' values (1, 0.5, 0.5) to 0.25 changes nothing about the data, but:

- at 85 years the share missed falls from 68% to 57% (now within 1 point of the posterior median and the location trend), and the median true exceedance rises from 0.39% to 0.64%;
- at 45 years the share missed falls from 84% to 74% and the median true exceedance rises from 0.05% to 0.24%; record by record the true exceedance of the predictive level rises by a median factor of 2.8 (85 years: 1.6);
- in the cases where the spread and tail follow a rising mean, at 85 years, the predictive level with the tight prior becomes the best estimator we tested, missing in 55% of records against 64% for the Akaike-weighted average and 69% for the posterior median.

We varied only the random-walk scales. The prior on ξ itself, and its upper limit of 2.1, are likely to matter as much, because the overstatement comes from heavy-tailed draws.

Monte Carlo error. Splitting each fit's 4,000 draws into its four chains, the predictive level's chance of being exceeded changes by a typical factor of 1.06 at 85 years and 1.10–1.11 at 30–45 years (90th percentile 1.16–1.55). With the authors' 40,000 draws these factors would shrink by about $\sqrt{10}$. Monte Carlo error accounts for well under 1% of the variance of the error in the log exceedance probability (0.03–0.11%): the dominance of a few draws is a property of the posterior, not of the number of draws.

7.6 Ambiguity in the climate, or in the inference?

One argument for the predictive approach is that possible non-stationarity, combined with possibly long tails, makes the climate distribution genuinely ambiguous, and that the high and volatile estimates of catastrophic risk reflect that ambiguity. The synthetic records let us separate the two parts, because in the no-change cases nothing is changing and the true tail is at most moderately heavy ($\xi \lesssim 0.2$):

Table 8: No-change cases only (400 records per design): true chance of exceeding the reported 1-in-100-year level (median and 10–90th percentiles) and the share missed by more than a factor of 2.

	Bayes, predictive			Missed $> 2\times$		
Design	median	10%	90%	predictive	post. median	loc. trend
30 yr	0.02%	0.00%	1.35%	89	73	73
45 yr	0.08%	0.00%	1.46%	80	65	66
85 yr	0.47%	0.04%	2.31%	64	53	49
85 yr, tight prior	0.68%	0.12%	2.30%	52	48	49

With no change at all, the predictive level is as high and as dispersed as in the cases with change. So most of the ambiguity is produced by the inference: a few decades of annual maxima cannot rule out a heavy tail, the prior allows one, and averaging probabilities over the posterior turns

that possibility into a high level. That is a legitimate feature of a Bayesian learner's beliefs, and it is real ambiguity for someone who must act on a short record. But it would be present in a stationary climate, it shrinks with record length roughly as the tail becomes identifiable, and its size is governed by prior choices that the data cannot check. It is not, by itself, evidence that the climate's extremes are changing or are heavy-tailed.

7.7 Suggestions for practice

1. **Report the spread of beliefs, not only the average.** Alongside the predictive level, report the posterior distribution of the chance of exceeding it, for example the median draw and a 90% range (in our records typically ∼0.2% and 0–5% at 85 years). That makes clear that "1%" is an average over beliefs ranging from negligible to several times the target.

2. **Show prior sensitivity.** Refit with tighter random-walk scales and a more informative prior on ξ (or a lower upper bound), and report how the level and the implied premium move. In our records, the random-walk scales alone change the typical true exceedance by a factor of 1.6 (85 years) to 2.8 (45 years).

3. **Benchmark against a stationary climate.** Fit the model to synthetic stationary records of the same length and similar distribution. The spread of predictive levels obtained there is the ambiguity that comes from the inference alone.

4. **Compare with simple fits.** If the predictive level is far above both a constant and a trend maximum-likelihood fit, the difference comes from the posterior's heavy-tailed draws, and should be justified on other grounds (physical or regional knowledge of the tail).

5. **Bring in more information on the tail.** Pooling the shape parameter across nearby stations (regional frequency analysis), or using several values per year, constrains ξ better than a single annual-maximum record, and the overstatement shrinks as ξ is pinned down. In our records, three values per year from 45 years constrained ξ about as well as 85 years of annual maxima, but the predictive level still missed in 78% of records against 68%; at 85 years it did slightly less than the tight prior (60% missed against 57%). Pooling across stations is likely to do more.

6. **Separate the risk margin from the estimate.** The predictive approach builds a loading for ambiguity into the estimate through the prior. If a margin for ambiguity is wanted, it may be clearer to estimate the risk with a well-calibrated central estimate and add an explicit, stated margin, so that the margin is a decision rather than a by-product of the prior.

8 Limitations

- The synthetic data have no seasonal cycle, a single AR(1) dependence structure, and changes that are smooth or a single step. Real daily data have seasonality, longer memory and clustering of extremes, which would mostly make things harder.

- The covariate-model cases are, by construction, cases where that model is right, and the covariate is the local mean of the same series. With an external covariate (e.g. global mean temperature) the gain would depend on how closely the local extremes follow it.

- 100 records per case gives standard errors of 1–2 percentage points on shares pooled over the standard cases and about 5 points for a single case. Estimators are compared on the same records, so the differences between them are more precise (about ±1 point pooled), but they depend on which cases we chose; differences of a few points should not be over-interpreted.
- Our MCMC runs are shorter than the authors' ($4 \times 1{,}000$ against $4 \times 10{,}000$ draws). The accuracy of point estimates and interval coverage should be robust to this; the convergence failures with three values per year are not a matter of chain length.
- We scored the final year of each record (a "nowcast"); a forecast for future years would add the uncertainty of extrapolating the random walks. For the Bayesian model we varied only the random-walk scales of the prior, not the prior on ξ or its upper limit of 2.1, which matter for the predictive level (Section 7).
- We have not yet run the paper's joint L^* test, the exceedance (threshold) versions of the tests, or the Bayes factors that compare "more events" with "a shifted distribution".

9 Summary

What we did. We reimplemented the time-varying-extremes methods of Müller and Watson (2024) in Python, reproduced their published estimates and test statistics, and tested the methods on synthetic data for which the truth is known exactly. Daily series were drawn from four parent distributions (normal; gamma, precipitation-like; Student-t, heavy-tailed; beta, bounded) with day-to-day correlation, and their mean, width or tail was changed over records of 30 to 150 years.

False alarms. With one value per year, all the tests we examined keep their nominal 5% false-alarm rate, except that the conventional location-trend likelihood-ratio test is over-sized (7–10%) for 30-year records. Using the three or ten largest values per year from a persistent daily series raises false alarms to 20–50% unless the extremes are declustered.

Power. Power rises steeply with record length and is much lower for skewed or heavy-tailed parents, whose extremes are noisier relative to the daily variability. With one value per year the Nyblom test is not the most powerful: the simpler Mann–Kendall rank test detects the simulated changes about as often or more often in almost every case, and a location-trend test is more powerful against shifts in location but much less powerful against changes in tail shape. The Nyblom test gains from using the three or ten largest values per year, which Mann–Kendall cannot use, provided they are declustered.

How we scored the estimates. Detecting a change is not the same as measuring it. We scored more than a dozen estimators on a question that matters for risk assessment: if the estimated 99% annual-maximum level for the most recent year is taken at face value, what is the true probability that it is exceeded? An ideal estimator gives 1%. We count an estimate as a miss when that probability is more than 2% (risk understated at least twofold) or less than 0.5% (overstated at least twofold), and report the two directions separately, because their consequences differ and estimators trade one for the other.

The noise floor. The dominant source of error is not the change but the difficulty of estimating a 1-in-100 level from a few decades of annual maxima: with no change and a correctly specified model, 44% of 85-year records miss (27% understate, 18% overstate), and about 70% of 30- and 45-year records.

Which estimators do best. Against this floor, a linear trend in location or in location and width misses 56% of 85-year records, and an AIC-weighted average of constant, trend and mean-state models does best (51%). With 30 or 45 years all these methods are within a few points of each other and of the no-change floor.

The Bayesian model. The authors' Bayesian random-walk model misses 59% of 85-year records, erring more often toward understatement. It shrinks estimated changes toward zero, and its 90% credible intervals, while they cover the truth, are 1.3 to 3 times wider than the actual errors. With three values per year, 10–12% of its fits fail to converge under the authors' prior.

The posterior-predictive level. The 99th percentile of the Bayesian posterior predictive distribution averages probabilities over the draws, so it is pulled up by the few draws with the heaviest tails. On average over records it is about right or conservative (true exceedance 0.8% at 85 years, 0.4–0.6% at 30–45), and it understates the risk twofold in only 6–13% of records. In a single record it usually overstates the risk: it misses twofold in 68%, 84% and 92% of records at 85, 45 and 30 years, 9 to 19 points more often than the posterior median or a trend fit. As a pure premium rate for cover with a true 1% risk it charges a median 1.7–3.2%. It is as high and as dispersed with no change at all, and a tighter prior on the random walks brings it close to the posterior median; most of its ambiguity is therefore produced by the inference (a heavy tail that a short record cannot rule out and the prior allows), not by the climate.

Using the wrong driving variable. Which model is right depends on the physical process, and our tests show what happens when the model is wrong. We compared models of the same complexity that differ only in their driving variable: GEV location and scale linear in time, or linear in a smoothed annual mean of all daily values. When the distribution's width and tail follow an accelerating mean state, the time-trend models miss the current risk by more than a factor of 2 in 75–83% of 85-year records against 65% for the mean-state model. When the width changes while the mean stays constant, the mean-state model sees nothing and misses 69% against 55%. Fitting a change that is not there costs about as much: with no change, the location + width and covariate models miss 55–60% against 44% for a constant fit.

Where the choice matters less. When the mean moves linearly in time the two kinds of model coincide. When the mean rises at an increasing rate but the width and tail change linearly in time, the mean-state model, now with the wrong driver, still does slightly better (64% against 68%), because missing the curvature of the mean costs more than misplacing the width. With 30 years every model misses 66–87% of records, so the choice of driving variable is secondary to record length.

When the driver is unknown. The data rarely reveal the driver: AIC cannot tell a width change from a mean change. But an AIC-weighted average of the constant, time-trend and mean-state models comes within 4 points of the best single model for every true process at 85 years.

Practical conclusion. These methods can establish *that* extremes are changing, and in which direction. Statements about the *size* of the current 1-in-100-year level should carry wide uncertainty, including the spread between physically plausible models of the change, and are best based on an average over those models rather than on one of them.

A Cases and supplementary results

Set	Parent	Change over the record	$\Delta X_{0.99}$
Standard	Normal	no change	+0.00
	Normal	mean +0.5 sd (linear)	+0.50
	Normal	mean +0.5 sd (step)	+0.50
	Normal	width ×1.22 (linear)	+0.81
	Gamma	no change	+0.00
	Gamma	mean +0.5 sd (linear)	+0.50
	Gamma	mean +0.5 sd (step)	+0.50
	Gamma	width ×1.22 (linear)	+1.58
	Gamma	tail: log(shape) +1 (linear)	-1.97
	Student-t	no change	+0.00
	Student-t	mean +0.5 sd (linear)	+0.50
	Student-t	mean +0.5 sd (step)	+0.50
	Student-t	width ×1.22 (linear)	+1.64
	Student-t	tail: log(shape) +1 (linear)	-8.51
	Beta	no change	+0.00
	Beta	mean +0.25 sd (linear)	+0.25
	Beta	mean +0.25 sd (step)	+0.25
	Beta	width ×1.11 (linear)	+0.28
	Beta	tail: log(shape) +0.2 (linear)	-0.11
	Beta	position: log α +0.5 (linear)	+0.07
Covariate model	Normal	warming +0.5 sd, width & tail follow (accel.)	+1.02
	Normal	warming +0.5 sd, width & tail follow (linear)	+1.00
	Gamma	wetter: all quantiles ×1.22 (accel.)	+1.94
	Gamma	wetter: all quantiles ×1.22 (linear)	+1.86
	Gamma	wetter ×1.22, heavier tail (accel.)	+2.60
	Student-t	warming +0.5 sd, width & tail follow (accel.)	+2.32
	Beta	warming +0.25 sd, width & tail follow (accel.)	+0.46
Time-driven	Normal	warming +0.5 sd, width & tail linear in time (accel.)	+1.02
	Student-t	warming +0.5 sd, width & tail linear in time (accel.)	+2.32
	Beta	warming +0.25 sd, width & tail linear in time (accel.)	+0.46

Table 9: Stage C cases. $\Delta X_{0.99}$: true change in the 99% level over an 85-year record (baseline daily sd). "log(shape)" is the log of the gamma shape a, the t degrees of freedom ν, or the beta β; increasing it lightens the tail of the gamma and t.

True process	Const.	Loc.	Loc.+wid.	Bayes	Cov.	AIC	AIC avg.
Understated > 2×, 85 years							
No change	**27**	**32**	**36**	**36**	**38**	32	29
Mean shift, linear in time	61	**32**	**36**	**44**	**38**	36	31
Mean shift, step	52	16	19	**37**	**25**	26	23
Width change, mean constant	73	47	**37**	**50**	54	41	40
Tail change, mean constant	0	25	34	**21**	20	36	27
Follows mean state, mean linear in time	80	44	**36**	**50**	**35**	38	36
Follows mean state, mean accelerating	89	81	69	**60**	**49**	55	58
Follows time, mean accelerating	84	67	59	**55**	42	54	53
Overstated > 2×, 85 years							
No change	**18**	**17**	**19**	**17**	**23**	22	18
Mean shift, linear in time	6	**17**	**19**	**14**	**18**	16	15
Mean shift, step	6	38	36	**19**	**19**	21	21
Width change, mean constant	2	7	**18**	**12**	15	14	14
Tail change, mean constant	82	50	25	**39**	47	32	35
Follows mean state, mean linear in time	2	8	**19**	**13**	**17**	14	16
Follows mean state, mean accelerating	1	2	5	**12**	**16**	12	11
Follows time, mean accelerating	1	4	9	**13**	22	14	12

Table 10: The misses of Table 4 at 85 years, split into records with the risk understated (true exceedance > 2%) and overstated (< 0.5%) at least twofold. Bold: the model's form contains the true process.




Figure 33: Left: Bayesian model with the authors' prior and with a tighter prior on the random-walk scales. Right: shares of 85-year records missed by more than a factor of 2 either way, with correlated ($\phi = 0.5$) and independent daily values.

B The posterior-predictive level at 30 years by kind of change

Table 11: As Table 6, at 30 years.

		Bayes, predictive				Missed, others		
Kind of change	records	median	under	over	**missed**	post. median	loc. trend	AIC avg.
No change	400	0.02%	7	82	**89**	73	73	74
Mean	800	0.01%	9	84	**92**	75	69	72
Width	400	0.02%	10	81	**91**	77	76	79
Tail	300	0.01%	3	92	**95**	73	79	74
Position (beta)	100	0.00%	7	88	**95**	80	76	77
Follows mean state	700	0.01%	9	78	**87**	80	83	82
Time-driven	300	0.01%	9	81	**90**	78	83	81
Normal (standard cases)	400	0.01%	8	82	**90**	75	72	76
Gamma (standard cases)	500	0.02%	7	84	**91**	72	70	73
Student-t (standard cases)	500	0.04%	7	84	**91**	73	70	69
Beta (standard cases)	600	0.00%	8	87	**95**	79	79	80

References

Carpenter, B., et al. (2017). Stan: a probabilistic programming language. *Journal of Statistical Software* 76(1).

Coles, S. (2001). *An Introduction to Statistical Modeling of Extreme Values*. Springer.

Müller, U. K. and M. W. Watson (2024). Time varying extremes. *Review of Economics and Statistics*, doi:10.1162/rest_a_01528.

Nyblom, J. (1989). Testing for the constancy of parameters over time. *Journal of the American Statistical Association* 84, 223–230.

Philip, S., et al. (2020). A protocol for probabilistic extreme event attribution analyses. *Advances in Statistical Climatology, Meteorology and Oceanography* 6, 177–203.