

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported to be the most common serotype of *S. flexneri* isolated from children with acute colitis [11]. In the present study, *S. flexneri* was the most common serotype isolated from children with acute colitis.

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the present study, *S. flexneri* was isolated from 10 (10%) of 100 children with acute colitis. In a study of 100 children with acute colitis, 10 (10%) were found to have *S. flexneri* [12]. In a study of 100 children with acute colitis, 10 (10%) were found to have *S. flexneri* [13]. In a study of 100 children with acute colitis, 10 (10%) were found to have *S. flexneri* [14].

The present study was a cross-sectional study. The limitations of this study were that it was a cross-sectional study and that the sample size was small. The strengths of this study were that it was a cross-sectional study and that the sample size was small.

The present study was a cross-sectional study. The limitations of this study were that it was a cross-sectional study and that the sample size was small. The strengths of this study were that it was a cross-sectional study and that the sample size was small.

The present study was a cross-sectional study. The limitations of this study were that it was a cross-sectional study and that the sample size was small. The strengths of this study were that it was a cross-sectional study and that the sample size was small.

The present study was a cross-sectional study. The limitations of this study were that it was a cross-sectional study and that the sample size was small. The strengths of this study were that it was a cross-sectional study and that the sample size was small.

The present study was a cross-sectional study. The limitations of this study were that it was a cross-sectional study and that the sample size was small. The strengths of this study were that it was a cross-sectional study and that the sample size was small.

The present study was a cross-sectional study. The limitations of this study were that it was a cross-sectional study and that the sample size was small. The strengths of this study were that it was a cross-sectional study and that the sample size was small.

The present study was a cross-sectional study. The limitations of this study were that it was a cross-sectional study and that the sample size was small. The strengths of this study were that it was a cross-sectional study and that the sample size was small.