

NOTIFIABLE DISEASES - Summary of reported cases per 100,000 population, United States, 1971-1980

Disease	1980	1979	1978	1977	1976	1975	1974	1973	1972	1971
Amebiasis	2.38	1.90	1.84	1.41	1.35	1.30	1.30	1.07	1.06	1.33
Anthrax	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Aseptic meningitis	3.61	4.05	3.01	2.24	1.84	2.10	1.53	2.33	2.23	2.51
Botulism, total	0.04	0.02	0.05	0.08	0.03	0.01	0.01	0.02	0.01	0.01
Foodborne	0.01	0.01	0.03	0.04	0.01	0.01	0.01	0.02	0.01	0.01
Infant	0.03	0.01	0.02	0.02	0.01	0.00	-	-	-	-
Brucellosis (undulant fever)	0.08	0.10	0.08	0.11	0.14	0.15	0.11	0.10	0.09	0.09
Chancroid	0.35	0.38	0.24	0.21	0.29	0.33	0.46	0.56	0.68	0.84
Chickenpox	96.69	102.93	80.42	97.63	96.06	78.11	72.20	97.68	87.34	...
Cholera	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Diphtheria	0.00	0.03	0.03	0.04	0.06	0.14	0.13	0.11	0.07	0.10
Encephalitis, primary	0.13	0.15	0.13	0.16	0.25	1.11	0.10	0.08	0.08	0.16
Indeterminate	0.48	0.54	0.49	0.50	0.52	0.80	0.45	0.68	0.43	0.58
Post childhood infections	0.02	0.04	0.04	0.06	0.08	0.11	0.10	0.17	0.12	0.21
Gonorrhea	443.27	459.49	468.25	466.83	470.47	472.91	432.12	404.92	371.62	328.11
Granuloma inguinale	0.02	0.03	0.03	0.03	0.03	0.03	0.02	0.03	0.04	0.04
Hepatitis A	12.84	13.82	13.53	14.40	15.51	16.82	19.54	24.18	25.97	28.90
Hepatitis B	8.39	7.02	6.89	7.78	7.14	6.30	5.15	4.03	4.52	4.74
Hepatitis, unspecified	6.25	4.79	4.02	3.99	3.57	3.44	3.95			
Legionellosis	0.19	0.26	0.35	0.17	0.11					
Leprosy	0.10	0.08	0.08	0.07	0.07	0.08	0.06	0.07	0.06	0.06
Leptospirosis	0.04	0.04	0.05	0.03	0.03	0.04	0.03	0.03	0.02	0.03
Lymphogranuloma venereum	0.09	0.11	0.13	0.16	0.17	0.17	0.19	0.20	0.37	0.34
Malaria	0.91	0.41	0.34	0.25	0.22	0.18	0.14	0.11	0.36	1.15
Measles (rubeola)	5.96	6.18	12.32	26.51	19.16	11.44	10.45	12.72	15.50	36.50
Meningococcal infections, total	1.25	1.24	1.16	0.84	0.75	0.69	0.64	0.66	0.64	1.10
Mumps	3.86	6.55	7.81	10.02	17.93	27.89	28.00	36.23	38.42	65.33
Pertussis (whooping cough)	0.76	0.74	0.95	1.02	0.47	0.82	1.16	0.84	1.58	1.47
Plague	0.01	0.01	0.01	0.01	0.01	0.01	0.00	0.00	0.00	0.00
Polio, total	0.00	0.02	0.01	0.01	0.01	0.00	0.00	0.00	0.01	0.01
Paralytic	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.00	0.01	0.01
Psittacosis	0.05	0.06	0.06	0.04	0.04	0.02	0.08	0.02	0.02	0.02
Rabies, human	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Rheumatic fever, acute	0.30	0.44	0.60	1.23	1.32	2.01	1.79	1.92	2.01	2.16
Rubella (German measles)	1.72	5.36	8.38	9.43	5.82	7.81	5.64	13.25	12.25	21.86
Rubella congenital syndrome**	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02
Salmonellosis, excluding typhoid fever	14.88	15.06	13.49	12.87	10.74	10.61	10.40	11.35	10.64	10.63
Shigellosis	8.41	9.16	8.95	7.42	6.15	7.78	10.69	10.79	9.70	7.83
Smallpox										
Syphilis, primary and secondary	12.01	11.38	10.00	9.50	11.14	12.09	12.11	11.93	11.83	11.64
Total all stages	30.39	30.68	30.00	30.10	33.69	38.00	39.95	42.03	44.15	46.99
Tetanus	0.04	0.04	0.04	0.04	0.03	0.05	0.05	0.05	0.06	0.06
Trichinosis	0.06	0.07	0.03	0.07	0.05	0.12	0.06	0.05	0.04	0.05
Tuberculosis	12.25	12.57	13.08	13.93	14.96	15.95	14.25	14.77	15.79	17.07
Tularemia	0.10	0.09	0.06	0.08	0.07	0.06	0.07	0.08	0.07	0.09
Typhoid fever (cases)	0.23	0.24	0.23	0.18	0.20	0.18	0.21	0.32	0.19	0.20
(Carriers)	0.03	0.03	0.03			NA				
Typhus fever, flea-borne (endemic, murine)	0.04	0.03	0.02	0.04	0.03	0.02	0.01	0.02	0.01	0.01
Typhus fever, tick-borne (Rocky Mountain spotted)	0.52	0.49	0.49	0.53	0.44	0.40	0.36	0.32	0.25	0.21
Yellow fever										

Note: Rates less than 0.01 after rounding are shown as 0.00.

Population data from those states where diseases were not notifiable (NN) were excluded from rate calculation. Civilian resident population used for chancroid, gonorrhea, granuloma inguinale, lymphogranuloma venereum, and syphilis.

*Not previously notifiable nationally.

**Per 1,000 live births.