

Lake Okeechobee System Operating Manual

**Iteration 2 Modeling -
Evaluation Technical workshop
Climate perspective**


Sanibel-Captiva Conservation Foundation

Conservancy of Southwest Florida

June 29, 2021




Paul Julian PhD 



Monthly observed Atlantic Multidecadal Oscillation (AMO) Index from 1856 to 2021 for the northern Atlantic (Enfield et al. 2001).


- Recent debate on AMO index suggest decadal or longer-term internal oscillatory signals that are distinguishable from climatic noise (see Steinman et al 2015; Mann et al 2020; Mann et al 2021).
- Only variability in the interannual range associated with the El Niño/Southern Oscillation is found to be distinguishable from the noise background.
 - Still an indicator of warm/cool phase shift.

Extreme High/Low




Percent of time where stages ≤ 10 or ≥ 17 Ft NGVD29 during the period of simulation (1965 - 2016).

Extreme High/Low




Percent of time where stages ≤ 10 or ≥ 17 Ft NGVD29 during the period of simulation (1965 - 2016) broken down into dry (1965-1994) and wet (1995-2016) phases.

Lake Stage Envelope




Percent of time within normal and recovery stage envelope during dry (1965-1994) and wet (1995-2016) phases for each alternative.

Lake Stage Duration Curves





Stage duration curves for the entire period of simulation (Jan 1, 1965 - Dec 31, 2016) within each phase for each alternative compared to FWO (NA25) and ECB (ECBr).




Difference in stage duration curves (SDC; Cool - Wet) between climate phases for the entire period of simulation (Jan 1, 1965 - Dec 31, 2016) for each alternative includes FWO and ECB.

Daily time-series of S79 discharges for each alternative with mean discharge for dry/cold (1965 – 1994) and wet/warm (1995 – 2016) AMO phase depicted (red and blue, respectively).





Boxplot of daily discharge comparing dry/cold (1965 – 1994) and wet/warm (1995 – 2016) AMO phase for each alternative for the entire period of simulation (Jan 1, 1965 - Dec 31, 2016).



Percent of time within each respective flow category for S-79 comparing dry/cold (1965 – 1994) and wet/warm (1995 – 2016) AMO phase for each alternative for the entire period of simulation (Jan 1, 1965 - Dec 31, 2016).

Acknowledgments



South Florida Water Management District ([DBHYDRO](#))

 US Army Corps of Engineers ([USACE LOSOM](#))

- Interagency Modeling Center

[HTML Slide deck](#) | [PDF Slide deck](#) | [RMarkdown Source](#) © Julian (2021)




[Analysis Script](#)

[Additional Supplemental Slides](#); [Workshop #1 Presentation](#)



References

- Enfield DB, Mestas-Nuñez AM, Trimble PJ (2001) The Atlantic Multidecadal Oscillation and its relation to rainfall and river flows in the continental U.S. *Geophysical Research Letters* 28:2077–2080. doi: 10.1029/2000GL012745
- Mann ME, Steinman BA, Brouillette DJ, Miller SK (2021) Multidecadal climate oscillations during the past millennium driven by volcanic forcing. *Science* 371:1014–1019. doi: 10.1126/science.abc5810
- Mann ME, Steinman BA, Miller SK (2020) Absence of internal multidecadal and interdecadal oscillations in climate model simulations. *Nature Communications* 11:49. doi: 10.1038/s41467-019-13823-w
- Steinman BA, Mann ME, Miller SK (2015) Atlantic and Pacific multidecadal oscillations and Northern Hemisphere temperatures. 347:5.



CY 1965 - 2016

