

1 Significant plate motion during the early magmatic
2 stage of North American Midcontinent Rift
3 development

Nicholas. L. Swanson-Hysell,^{1,2} Angus A. Vaughan,^{1,3} Monica R. Mustain,^{1,4}
and Kristofer E. Asp^{1,5}

¹Institute for Rock Magnetism,
Department of Earth Sciences, University of
Minnesota, Minneapolis, Minnesota, USA

²Department of Earth and Planetary
Science, University of California, Berkeley,
California, USA

³Department of Geology, Carleton
College, Northfield, Minnesota, USA

⁴Department of Geology and Geography,
Illinois State University, Normal, Illinois,
USA

⁵Department of Geological Sciences,
University of Minnesota, Duluth,
Minnesota, USA

Abstract. As the supercontinent Rodinia was assembling ca. 1.1 billion years ago, there was extensive magmatism on at least five Proterozoic continents including the development of the North American Midcontinent Rift. New paleomagnetic data from 84 lava flows of the Osler Volcanic Group of the Midcontinent Rift reveal that there was a significant and progressive decrease in inclination between the initiation of extrusive volcanism in the region (ca. 1110 Ma) and ca. 1105 ± 2 Ma (the “early stage” of rift development). Paleomagnetic poles can be calculated for the lower portion of the reversed Osler Group (41.0°N , 218.7°E , $A_{95}=4.8^\circ$, $N=30$) and the upper portion of the reversed Osler Group (42.5°N , 201.6°E , $A_{95}=3.7^\circ$, $N=59$; this pole can be assigned the age of ca. 1105 ± 2 Ma). This result is a positive test of the hypothesis that there was significant plate motion during the early stage of rift development. In addition to being a time of widespread volcanism on Laurentia and other continents, this interval of the late Mesoproterozoic was characterized by rapid paleogeographic change.

1. Introduction

Despite being active for more than 20 million years [*Davis and Green, 1997*] and resulting in the thinning of pre-rift crust to less than 10 km [*Cannon, 1992*], the 1.1 Ga Midcontinent Rift failed to dismember the Laurentian craton. This failure resulted in the preservation of a thick record of rift-related volcanic and sedimentary rocks that gives geoscientists insight into the development of this ancient rift. Most models for the development of the Midcontinent Rift attribute its origin to the upwelling and decompression melting of a mantle plume [*Shirey, 1997*]. On the basis of the great volume of generated magma and interpretation of geochemical data, it is argued that the early stage plateau flood basalts of the rift (ca. 1110-1105 Ma) and the main stage volcanics that erupted into the central basin (ca. 1100-1095 Ma) were both dominated by plume-sourced melts. This deep-plume origin for the rift needs to be considered in conjunction with paleogeographic change that has been inferred to have been ongoing throughout rift development. Fully constraining this paleogeographic change is essential for understanding rift development and for constraining late Mesoproterozoic paleogeographic reconstructions given the centrality of Laurentia's apparent polar wander path to such efforts.

It has long been noted that there is a significant difference in paleomagnetic inclination between the steep (dominantly reversed polarity) magnetizations of the oldest volcanics and intrusives of the Midcontinent Rift and the shallower (dominantly normal polarity) magnetizations from the younger main stage volcanics and intrusives [*Halls and Pesonen, 1982*]. This inclination change has been interpreted either as resulting from rapid plate motion [*Davis and Green, 1997*] or as being the result of large non-dipolar contributions

to the late Mesoproterozoic geomagnetic field that led to asymmetry across reversals
Pesonen and Nevanlinna [1981]. The interpretation of the record as recording stepwise
inclination change across reversals associated with a significant sustained departure from
a geocentric axial dipole (GAD) dominated field was challenged by the observation of a
progressive decrease in paleomagnetic inclination across multiple geomagnetic reversals up
through the stratigraphy of Midcontinent Rift lavas at Mamainse Point, Ontario [*Swanson-*
Hysell et al., 2009; ?]. This progressive decrease in inclination leads to the interpretation
of multiple symmetric reversals when the data are considered in stratigraphic context
and these results have been used to support the hypothesis that rapid paleogeographic
change was ongoing during Midcontinent Rift magmatism [??]. To date, Mamainse Point
is the only succession where a progressive decrease in inclination through an exposure
of rift stratigraphy has been reported. At Mamainse Point, much of the decrease in
paleomagnetic inclination occurs within the lowermost reversed polarity portion of the
stratigraphy that erupted during the early magmatic stage of rift development [?]. This
result sets up the prediction that other localities in the rift that span the same period
of time should also record such a decrease. This work tests that hypothesis with new
paleomagnetic data developed in stratigraphic context from the Osler Volcanic Group.

2. Geology and context of the Osler Volcanic Group

2.1. Osler Volcanic Group lithologies

The Osler Volcanic Group overlies the epicontinental sediments of the Mesoproterozoic
Sibley Group. The lowest 100 m of stratigraphy of the Osler Volcanic Group contains
rift-related sandstones and conglomerates [*Hollings et al.*, 2007a], which are followed by
a succession of relatively continuous tholeiitic basalt flows. We studied these flows along

the east shore of Simpson Island in the Lake Superior Archipelago. This location has well-preserved basalt flows and exposure of a significant thickness of the stratigraphy. This location occurs at a great enough distance (> 10 km) from the intrusive St. Ignace Island Complex so that the magnetizations of the flows are apparently unaffected by its emplacement. The St. Ignace Island Complex is dominated by felsic lithologies and was emplaced into the mafic flows of the Osler Volcanic Group near the end of Midcontinent Rift volcanism [*Hollings et al.*, 2007b].

Along the east shore of Simpson Island, where we measured stratigraphic sections (Fig. 1), the stratigraphy is dominated by basaltic lava flows with typical thicknesses of ca. 5 meters. For the 105 flows in the measured stratigraphic sections where there is sufficient exposure of the flow base, interior and top to determine thickness, the median flow thickness is 4.9 meters with a first quartile thickness of 2.0 meters and third quartile thickness of 9.8 meters. Minor interflow siltstone and conglomerate occur between some flows and provide constraints on paleo-horizontal that were utilized for structural corrections of paleomagnetic data (Fig. 1). The red-brown siltstone beds are generally centimeters to tens of centimeters thick while the conglomerate beds can be up to several meters in thickness. Clasts in the conglomerate are basaltic in composition and are primarily pebble-sized, but can range up to 10 cm in diameter. Within the stratigraphic sections, individual flows can be distinguished by their characteristic transition in texture moving up through the flow: pipe vesicles at the base of flows, to massive basalt, to massive basalt with some amygdules, to a highly amygdaloidal texture at the top of the flows. There is well-preserved pahoehoe flow-banding on some flow tops. Using our measurements of

bedding orientation, we estimate that slightly over 3000 meters of Osler Volcanic Group stratigraphy is exposed on Simpson Island (Fig. 1).

2.2. Angular unconformity and associated paleomagnetic reversal

Southwest of Simpson Island in the Nipigon Strait region there is exposure on Puff Island, which contains an angular unconformity marked by a conglomerate that separates underlying flows of reversed magnetic polarity from overlying flows of normal magnetic polarity [Halls, 1974]. This unconformity is higher in the stratigraphy than the top of the exposure at Simpson Island where all studied flows have reversed magnetic polarity. Only ca. 110 stratigraphic meters of normally magnetized flows are exposed south (i.e. stratigraphically above) of the unconformity on Puff Island before the sequence is covered by Lake Superior. This unconformity has been interpreted to be due to a period of quiescence of local volcanism [Halls, 1974]. Halls [1974] was the first to suggest that such quiescence may have been widespread throughout the Midcontinent Rift—an idea that is now incorporated into models of rift development that utilize interpretations of the distribution of U-Pb dates and has been termed the “latent stage” [Vervoort *et al.*, 2007]. As occurs in sequences across the Midcontinent Rift, the lower reversed flows studied paleomagnetically by Halls [1974] have a steeper inclination, in an absolute sense, than the younger normal flows above the unconformity. The missing time evidenced by the angular unconformity likely correlates with the missing time inferred from radiometric dates on units within the stratigraphy of the North Shore Volcanic Group and the Powder Mill Group [Davis and Green, 1997; Zartman *et al.*, 1997]. In these rift successions, the stratigraphic intervals where time is inferred to be missing are associated with a switch from reversed to normal magnetic polarity.

The paleomagnetic data that *Halls* [1974] developed from Osler Volcanic Group flows were in relatively close stratigraphic proximity below and above the unconformity (as the goal of the study was to confirm the presence of a geomagnetic reversal that had been inferred from aeromagnetic data; *Halls* [1972]). The aeromagnetic data demonstrated that the remainder of the flows below the unconformity (i.e. to the north) and stratigraphically below those studied by *Halls* [1974] are also of reversed magnetic polarity. On the basis of polarity and the geochronology discussed below, the interval has been correlated with the “early stage” of Midcontinent Rift development and the other basal sequences of lava flows within the rift. These early stage flows are interpreted to be plateau lavas that erupted over a broad geographic area prior to significant development of the main central rift graben that underlies present day Lake Superior [*Cannon*, 1992]. Due to being limited to the stratigraphy in close proximity to the angular unconformity, existing data have not permitted evaluation of whether or not there is a progressive change in paleomagnetic inclination through the reversed polarity flows as observed in the “early stage” volcanics at Mamainse Point [*Swanson-Hysell et al.*, 2009].

2.3. Age constraints on the Osler Volcanic Group

In some locales, a quartz-feldspar, porphyritic felsic unit occurs near the base of the Osler Group for which a U-Pb zircon date of 1107.5^{+4}_{-2} Ma has been reported [*Davis and Sutcliffe*, 1985]. This date was obtained from an outcrop of felsic porphyry on Black Bay Peninsula, ca. 40 km to the west of Simpson Island. This unit was tentatively interpreted as extrusive [*Davis and Sutcliffe*, 1985; *Lightfoot et al.*, 1991] and the date has been interpreted as constraining the time at which Osler Group volcanism commenced. New observations made of a quartz-feldspar porphyry unit on Simpson Island, mapped as

equivalent to the unit from which the date was obtained [*Giguere*, 1975], provide additional evidence for the inference made by *Giguere* [1975] that this unit is actually intrusive. On Simpson Island, the basal sedimentary units of the Osler Volcanic Group are overlain by the quartz-feldspar porphyry which itself underlies the basalt flows (Fig. 1). A thin (1-2 mm thick) veneer of basalt is variably present overlying the porphyry. The basalt veneer displays pahoehoe flow banding, which is unlikely to have developed if the flow was originally this thin, implying that the felsic unit intruded into the basalt, cutting into an originally thicker flow. An additional observation is that there are protrusions of porphyry surrounded by host basalt, providing further support for an intrusive relationship. If the Simpson Island intrusion is indeed equivalent to the Black Bay Peninsula unit, this evidence suggests that the 1107.5^{+4}_{-2} Ma date is a minimum age for the eruption of the first Osler basalt flows, rather than an absolute age for that point in the Osler Group stratigraphy. This unit should be a target for future geochronology.

A sequence of quartz-feldspar phyric rhyolite flows occurs near the top of the magnetically reversed portion of the stratigraphy at Agate Point (*Davis and Sutcliffe* [1985]; Fig. 3; stratigraphically higher than the highest flow on Simpson Island). *Davis and Green* [1997] obtained a U-Pb zircon date from the Agate Point Rhyolite of 1105 ± 2 Ma which, if the extrusive interpretation is correct, is a robust age for that point in the Osler Group stratigraphy.

3. Methods

We collected oriented samples for magnetic laboratory measurements during the course of measuring stratigraphic sections (Fig. 1). Each site consists of an individual lava flow from which we collected 6 to 10 small (2 cm diameter) rock cores with a hand-held drill.

These small core samples were oriented with a magnetic and sun compass, when possible, such that their spatial orientation is known. To minimize the visual impact of collecting these small cores along the pristine Lake Superior shoreline, we knocked out the portion of the outcrop from which they were collected.

At the Institute for Rock Magnetism, specimens were prepared from the samples and subjected to progressive thermal or alternating field (AF) demagnetization (Fig 2A). Initial results on sister specimens demonstrated the simplicity of the magnetizations and the similarity between results obtained through thermal and AF demagnetization (Fig 2A). Low-temperature remanence experiments were run on representative samples and loss of remanence across the Verwey transition demonstrates that the magnetic mineralogy is dominated by low-titanium magnetite (e.g. Fig 2B). These results revealed the ability of AF demagnetization to isolate the characteristic remanent magnetization (ChRM) held by magnetite with relatively small and variably present overprints being effectively removed by low-field AF steps. Given these results, the majority of flows were studied with AF demagnetization alone.

4. Results and Discussion

Flow means from the data generated for 84 flows of the Osler Volcanic Group are summarized in Figures 1 and 3 and the table in the supporting information. To consider whether the flows of the Osler Volcanic Group record progressive paleogeographic change, we take the approach of grouping and comparing data from the lower third of the stratigraphy (0 to 1041 meters; 30 flows), the middle third of the stratigraphy (1041 to 2083 meters; 20 flows) and the upper third of the stratigraphy (2083 to 3124 meters; 34 flows). To test whether these subsets of the data could have been drawn from a common mean,

we apply both the Watson V_w test with Monte Carlo simulation [Watson, 1983] and the bootstrap test for a common mean [Tauxe, 2010]. Full details associated with these statistical tests are provided in the supporting information. The results from these common mean statistical tests show that the directions from the lower third of the stratigraphy cannot be distinguished from those from the middle third, nor can the directions from the upper third of the stratigraphy be distinguished from those in the middle third. In contrast, directions from the lower third of the stratigraphy can be distinguished from those of the upper third at the 95% confidence level. This result can be seen visually in Figure 3 as the α_{95} ellipses associated with the directional means and the A_{95} ellipses associated with the pole means do not overlap. The statistically significant difference between the populations of flow means in the lower and upper third of the stratigraphy, combined with the result that the middle third data has a mean that is an intermediate direction between the lower and upper means, supports the hypothesis that progressive paleogeographic change was ongoing throughout the eruption of the Osler Volcanic Group with Laurentia moving to lower latitudes (see reconstruction Fig. 3). The paleomagnetic poles calculated and used in the paleogeographic reconstruction in Fig. 3 are: lower third Simpson Island Osler Group (40.9°N, 218.6°E, $A_{95}=4.8^\circ$, N=30); middle third Simpson Island Osler Group (42.7°N, 211.3°E, $A_{95}=8.2^\circ$, N=20) and the upper portion of the reversed Osler Group (41.6°N, 205.4°E, $A_{95}=4.8^\circ$, N=34). Stratigraphic subgroups of the Simpson Island data can be made in many different ways than this approach of dividing the stratigraphy into thirds. For example, comparing the 17 flows in the lowermost 500 meters against the 17 flows in the uppermost 500 meters, demonstrates that those populations are dramatically different such that the bootstrap test for a common mean shows

their x, y and z components to be distinct at the 99% confidence level (see supporting information for details). Comparing the 41 flows in the lower half of the stratigraphy to the 43 flows in the upper half of the stratigraphy also reveals a statistically significant, but relatively small, difference between the populations (see supporting information for details). We focus on the lower, middle and upper third grouping in our analysis making the judgement that such an analysis strikes the balance between considering the possibility of change through the stratigraphy while binning enough data over thick enough intervals to not to be significantly biased by under-averaging secular variation.

The paleomagnetic data developed by *Halls* [1974] come from the Osler Volcanic Group in the Nipigon Straits region. The data were obtained from the uppermost part of the stratigraphy below the angular unconformity on Puff Island that separates the flows of reversed polarity from younger flows of normal polarity (only a few of which are preserved before the sequence is submerged beneath Lake Superior). The *Halls* [1974] data of reversed polarity (N=25, <http://earthref.org/MAGIC/9518>) comes from a portion of the stratigraphy that should correlate with the upper third of the stratigraphy at Simpson Island. Watson and bootstrap tests for a common mean between the *Halls* [1974] data of reversed polarity and data from the upper third of the stratigraphy at Simpson Island are positive indicating that the populations of directions cannot be distinguished from one another—consistent with this stratigraphic correlation. In contrast, tests for a common mean between the reversed polarity *Halls* [1974] data and the lower third of the Simpson Island stratigraphy fail. This result indicates that the populations are statistically distinct, building on the result that there was significant paleogeographic change recorded by the Osler Volcanic Group. Given that the data from the upper third of the stratigraphy

on Simpson Island correlate stratigraphically and share a common mean with the Halls
(1974) data from the Nipigon Straits region, we can calculate a mean paleomagnetic pole
for the upper portion of the reversed Osler Group stratigraphy (42.5°N , 201.6°E , $A_{95}=3.7^{\circ}$,
 $N=59$) that can be assigned an approximate age of 1105 ± 2 Ma.

These new data from 84 flows of Osler Volcanic Group from the early stage of the Mid-
continent Rift bolster evidence from the succession at Mamainse Point that the decrease
in inclination through the history of the rift is a progressive change rather than a step-
wise change across reversals. The interpretation of a step-wise change of inclination across
reversals has been used to argue for reversal asymmetry at the time that was proposed
to result from significant deviation from an axial dipole geomagnetic field [*Pesonen and
Nevanlinna*, 1981]. The observed progressive change in inclination is more consistent with
the hypothesis that inclination decrease is a result of fast equatorward motion of Lau-
rentia [*Davis and Green*, 1997; *Swanson-Hysell et al.*, 2009]. The poles calculated from
the stratigraphic groupings of the Osler Group stratigraphy at Simpson Island fit the
progression along the path suggested by paleomagnetic poles from the lowermost polarity
zone at Mamainse Point, Ontario. These data sets combined indicate that there was sig-
nificant plate motion during early magmatic stage of North American Midcontinent Rift
development.

From the beginning of Midcontinent Rift extrusive volcanism in the early magmatic
stage to the voluminous volcanism associated with the main magmatic stage there was
ca. 25° of latitudinal motion of North America. The voluminous volcanism appears to
have been concentrated in the Lake Superior region both during the early magmatic stage
and during the eruption of the thick main stage volcanics in the central graben. Arguments

for a plume-origin for Midcontinent Rift volcanism have argued that a plume is necessary to explain isotopic signatures in lava flows (Nd isotopes, *Nicholson et al.* [1997]; Re-Os data, *Shirey* [1997]) and necessary as a heat source in generating the large volumes of basaltic magma associated with the rift [*Cannon*, 1992]. If a long-lived plume was in a fixed position relative to Earth's spin axis, large relative motion of Laurentia would make it unable to continue to be a source of melt to the rift. Furthermore, no evidence to date has revealed a hotspot track of progressively younger volcanics off-axis of the Midcontinent Rift. One possibility is that the North American plate and a deep-seated mantle plume traveled in unison to lower latitudes as a result of large-scale rapid true polar wander. The motion implied by the Keweenaw poles has been interpreted as a result of true polar wander [*Evans*, 2003; *Mitchell et al.*, 2012]. An interpretation of an active contribution from a narrow fixed plume throughout rift development implicitly favors this true polar wander hypothesis. The true polar wander explanation for Keweenaw paleogeographic change has posited that the entire swath from the early Midcontinent Rift poles down to the Grenville poles is the result of TPW about a fixed minimum inertia axis and can be fit with a great circle. With new paleomagnetic data from the Osler Volcanic Group, the progression of the apparent polar wander path through the early magmatic stage of the rift can be further resolved. The path that emerges from these additional data shows that there was a marked change in the trajectory of the path at the beginning of rift's main magmatic stage. Separate Euler poles are needed to explain the motion of the early+latent magmatic stages and the main+late magmatic stages. This result doesn't rule out the possibility that true polar wander was a significant contributor to the rapid paleogeographic change recorded by the Midcontinent Rift. However, it does demonstrate

that fitting a single great circle to all data from the beginning of the rift up through the apex of the Grenville loop (e.g. the ca. 1015 Ma Haliburton pole as done by *Mitchell et al.* [2012]) and explaining all of the paleogeographic changes as being a result of true polar wander about a single fixed minimum inertia axis is not a good fit to the data.

At the same time that the Midcontinent Rift was initiating in Laurentia the following igneous provinces were emplaced on four other cratons:

- the Umkondo large igneous province of the Kalahari Craton (many ID-TIMS U-Pb dates on zircon and baddeleyite between 1112 and 1108 Ma; *Hanson et al.* [2004])
- thick and extensive gabbro-norite (GN) dikes exposed in the southwest Angola portion of the Congo Craton (one of which has an ID-TIMS U-Pb date on baddeleyite of 1110.3 ± 2.5 Ma; *Ernst et al.* [2013])
- the Mahoba suite of dikes of the India Craton (one of which, the “Great Dike of Mahoba” has an laser ablation U-Pb date on zircon of 1112.7 ± 7.4 Ma; *Pradhan et al.* [2012])
- a putative ca. 1110 Ma large igneous province in the southwest portion of the Amazonia craton inferred from dates on two widely separated intrusions (ID-TIMS U-Pb date on baddeleyite from the Rincón del Tigre intrusion of 1110.4 ± 1.8 Ma and a sill within Aguepeí sediments with a ID-TIMS U-Pb date on baddeleyite of 1111.5 ± 1.5 Ma *Hamilton et al.* [2012])

This contemporaneous voluminous volcanism on five late Mesoproterozoic cratons is coincident with the onset of Laurentia’s rapid plate motion. This temporal correlation suggests that this time period was characterized by particularly vigorous mantle con-

vection, which could have contributed to large igneous province development and to the driving forces that resulted in rapid plate motion and/or true polar wander.

5. Conclusion

Lava flows of the Osler Volcanic Group below the Puff Island unconformity erupted during the early magmatic stage of Midcontinent Rift development during a time interval characterized by reversed magnetic polarity. New paleomagnetic data from 84 Osler Volcanic Group lava flows reveal a significant decrease in paleomagnetic inclination through the stratigraphy. These results support the hypothesis that the difference between the steep paleomagnetic inclinations characteristic of the early magmatic stage and the relatively shallower inclinations of the main magmatic stage throughout the rift are the result of rapid plate motion that was recorded by the Osler Volcanic Group.

Acknowledgments. The Ontario Ministry of Natural Resources granted permits for work to be conducted in the Lake Superior Archipelago Conservation Reserve. Pete Hollings of Lakehead University along with Dorothy Campbell and John Scott of the Ontario Geological Survey provided an introduction to the geology of the Lake Superior Archipelago. This work benefitted from discussions with Robert Cundari, Josh Feinberg and Henry Halls. Julie Bowles, Mike Jackson and Peat Solheid provided support for measurements at the Institute for Rock Magnetism. This research was supported by NSF EAR-1045635 to N.L.S.-H. and the NSF-funded Research Experiences for Undergraduates program in the Department of Earth Sciences at the University of Minnesota that supported A.A.V. and M.R.M.. This is IRM contribution #1312.

References

- Cannon, W. F. (1992), The Midcontinent rift in the Lake Superior region with emphasis on its geodynamic evolution, *Tectonophysics*, *213*(1-2), 41–48, doi:10.1016/0040-1951(92)90250-A.
- Davis, D., and J. Green (1997), Geochronology of the North American Midcontinent rift in western Lake Superior and implications for its geodynamic evolution, *Canadian Journal of Earth Science*, *34*, 476–488, doi:10.1139/e17-039.
- Davis, D., and R. Sutcliffe (1985), U-Pb ages from the Nipigon plate and northern Lake Superior, *Geological Society of America Bulletin*, *96*, 1572–1579, doi:10.1130/0016-7606(1985)96<1572:UAFTNP>2.0.CO;2.
- Ernst, R. E., E. Pereira, M. A. Hamilton, S. A. Pisarevsky, J. Rodriques, C. C. G. Tassinari, W. Teixeira, and V. Van-Dunem (2013), Mesoproterozoic intraplate magmatic ‘barcode’ record of the Angola portion of the Congo Craton: Newly dated magmatic events at 1505 and 1110 Ma and implications for Nuna (Columbia) supercontinent reconstructions, *Precambrian Research*, *230*(0), 103–118, doi:10.1016/j.precamres.2013.01.010.
- Evans, D. (2003), True polar wander and supercontinents, *Tectonophysics*, *362*, 303–320, doi:10.1016/S0040-1951(02)000642-X.
- Giguere, J. F. (1975), Geology of St. Ignace Island and adjacent islands, District of Thunder Bay, Canada, *Tech. rep.*, Ontario Geological Survey.
- Halls, H. (1974), A paleomagnetic reversal in the Osler Volcanic Group, northern Lake Superior, *Canadian Journal of Earth Science*, *11*, 1200–1207, doi:10.1139/e74-113.

- 326 Halls, H., and L. Pesonen (1982), Paleomagnetism of Keweenawan rocks, *Geological So-*
 327 *ciety of America Memoirs*, 156, 173–201, doi:10.1130/MEM156-p173.
- 328 Halls, H. C. (1972), Magnetic studies in northern Lake Superior, *Canadian Journal of*
 329 *Earth Sciences*, 9(11), 1349–1367, doi:10.1139/e72-123.
- 330 Hamilton, M. A., G. R. Sadowski, W. Teixeira, R. E. Ernst, and A. S. Ruiz (2012), Precise,
 331 matching U-Pb ages for the Rincon del Tigre mafic layered intrusion and Huanchaca
 332 gabbro sill, Bolivia: Evidence for a late Mesoproterozoic LIP in SW Amazonia?, in
 333 *Geological Association of Canada Annual Meeting Abstracts*, vol. 35.
- 334 Hanson, R., J. Crowley, S. Bowring, J. Ramezani, W. Gose, I. Dalziel, J. Pancake, E. Sei-
 335 del, T. Blenkinsop, and J. Mukwakwami (2004), Coeval large-scale magmatism in the
 336 Kalahari and Laurentian Cratons during Rodinia assembly, *Science*, 304, 1126–1129,
 337 doi:10.1126/science.1096329.
- 338 Hollings, P., P. Fralick, and B. Cousens (2007a), Early history of the Midcontinent Rift in-
 339 ferred from geochemistry and sedimentology of the Mesoproterozoic Osler Group, north-
 340 western Ontario, *Canadian Journal of Earth Sciences*, 44(3), 389–412, doi:10.1139/e06-
 341 084.
- 342 Hollings, P., M. Smyk, and L. Heaman (2007b), Preliminary investigations of the ~ 1 Ga
 343 St. Ignace Island Complex, northern Lake Superior, Ontario, *Geological Association of*
 344 *Canada/Mineralogical Association of Canada Program with Abstracts*, 32, 39.
- 345 Lightfoot, P. C., R. H. Sutcliffe, and W. Doherty (1991), Crustal contamination identi-
 346 fied in Keweenawan Osler group tholeiites, Ontario: A trace element perspective, *The*
 347 *Journal of Geology*, 99(5), 739–760, doi:10.1086/629536.

348 Mitchell, R. N., T. M. Kilian, and D. A. D. Evans (2012), Supercontinent cycles and
349 the calculation of absolute palaeolongitude in deep time, *Nature*, *482*(7384), 208–211,
350 doi:10.1038/nature10800.

351 Nicholson, S., S. Shirey, K. Schultz, and J. Green (1997), Rift-wide correlation of 1.1 Ga
352 Midcontinent rift system basalts: implications for multiple mantle sources during rift
353 development, *Canadian Journal of Earth Science*, *34*, 504–520, doi:10.1139/e17-041.

354 Pesonen, L., and H. Nevanlinna (1981), Late Precambrian Keweenawan asymmetric re-
355 versals, *Nature*, *294*, 436–439, doi:10.1038/294436a0.

356 Pradhan, V. R., J. G. Meert, M. K. Pandit, G. Kamenov, and M. E. A. Mon-
357 dal (2012), Paleomagnetic and geochronological studies of the mafic dyke swarms
358 of Bundelkhand craton, central India: Implications for the tectonic evolution and
359 paleogeographic reconstructions, *Precambrian Research*, *198–199*(0), 51–76, doi:
360 10.1016/j.precamres.2011.11.011.

361 Shirey, S. B. (1997), Re-Os isotopic compositions of Midcontinent rift system picrites:
362 implications for plume –lithosphere interaction and enriched mantle sources, *Canadian*
363 *Journal of Earth Sciences*, *34*(4), 489–503, doi:10.1139/e17-040.

364 Swanson-Hysell, N. L., A. C. Maloof, B. P. Weiss, and D. A. D. Evans (2009), No asym-
365 metry in geomagnetic reversals recorded by 1.1-billion-year-old Keweenawan basalts,
366 *Nature Geoscience*, *2*, 713–717, doi:10.1038/ngeo622.

367 Tauxe, L. (2010), *Essentials of Paleomagnetism*, University of California Press.

368 Vervoort, J., K. Wirth, B. Kennedy, T. Sandland, and K. Harpp (2007), The
369 magmatic evolution of the Midcontinent rift: New geochronologic and geochem-
370 ical evidence from felsic magmatism, *Precambrian Research*, *157*, 235–268, doi:

371 10.1016/j.precamres.2007.02.019.

372 Watson, G. S. (1983), Large sample theory of the langevin distribution, *Journal of Sta-*
373 *tistical Planning and Inference*, 8(3), 245–256, doi:10.1016/0378-3758(83)90043-5.

374 Zartman, R., S. Nicholson, W. Cannon, and G. Morey (1997), U-Th-Pb zircon ages of
375 some Keweenawan Supergroup rocks from the south shore of Lake Superior, *Canadian*
376 *Journal of Earth Science*, 34, 549–561, doi:10.1139/e17-044.




Figure 1. Geological map of Simpson Island in the Lake Superior Archipelago with studied stratigraphic sections and lava flows shown. Data from flows are color-coded by inclination. The lava flows of the Osler Volcanic group are tilted such that more southward flows are higher in the stratigraphy. The lower, middle and upper thirds divisions of the stratigraphy that are used in the text and in Figure 3 are shown. The inset map shows the location of the geological map.




Figure 2. A) Paleomagnetic data for basaltic lava flow sample SI5b-185.7. Alternating field and thermal demagnetization data are shown for sister specimens of this sample in vector component and equal area diagrams. These data reveal a dominantly single-component remanence with both demagnetization protocols isolating the same direction. B) Low temperature demagnetization data from a sample in the same lava flow wherein a saturating isothermal remanent magnetization (SIRM) was imposed at 20K after cooling in a zero field. Subsequent warming of this remanence led to significant demagnetization across temperatures characteristic of the Verwey transition indicating that the magnetic mineralogy of the sample is dominated by low-titanium magnetite.




Figure 3. Summary of paleomagnetic data from the Simpson Island exposure of the Osler Volcanic Group. The equal area plot shows the mean directions for each of the individual studied flows ($N=84$). All studied flows are of reverse polarity such that the plotted directions are in the upper hemisphere of the projection. Means are calculated and plotted on the equal area plot with α_{95} confidence ellipses for flows in the lower third (red; Dec=99.3, Inc=-68.0, $\alpha_{95}=3.4$, $N=30$), the middle third (yellow; Dec=105.9, Inc=-64.9, $\alpha_{95}=5.9$, $N=20$) and upper third (blue; Dec=107.6, Inc=-61.6, $\alpha_{95}=3.5$, $N=34$) of the stratigraphy. The mean paleolatitudes for these portions of the stratigraphy calculated from the Fisher means are also shown on the simplified composite stratigraphy with corresponding 2σ error bars. The paleogeographic reconstruction shows Laurentia's progressive equatorward motion as constrained by the poles from each of the stratigraphic groupings.