

26.1.1 Geometry: UGC NET CSE | December 2014 | Part 3 | Question: 16



Consider a triangle represented by $A(0, 0)$, $B(1, 1)$, $C(5, 2)$. The triangle is rotated by 45° about a point $P(-1, -1)$. The co ordinates of the new triangle obtained after rotation shall be _____.

- A. $A'(-1, \sqrt{2} - 1)$, $B'(-1, 2\sqrt{2} - 1)$, $C'(\frac{3}{2}\sqrt{2} - 1, \frac{9}{2}\sqrt{2} - 1)$
- B. $A'(\sqrt{2} - 1, -1)$, $B'(2\sqrt{2} - 1, -1)$, $C'(\frac{3}{2}\sqrt{2} - 1, \frac{9}{2}\sqrt{2} - 1)$
- C. $A'(-1, \sqrt{2} - 1)$, $B'(2\sqrt{2} - 1, -1)$, $C'(\frac{3}{2}\sqrt{2} - 1, \frac{9}{2}\sqrt{2} - 1)$
- D. $A'(-1, \sqrt{2} - 1)$, $B'(2\sqrt{2} - 1, -1)$, $C'(\frac{9}{2}\sqrt{2} - 1, \frac{3}{2}\sqrt{2} - 1)$

ugnetcse-dec2014-paper3 geometry non-gatecse

Answer key

26.2.1 Lines Curves: UGC NET CSE | December 2015 | Part 3 | Question: 32



The end points of a given line are $(0, 0)$ and $(6, 18)$. Compute each value of y as x steps from 0 to 3, by using equation of straight line:

- A. For $x = 0, y = 0; x = 1, y = 3; x = 2, y = 6; x = 3, y = 9$
- B. For $x = 0, y = 1; x = 1, y = 3; x = 2, y = 4; x = 3, y = 9$
- C. For $x = 0, y = 2; x = 1, y = 3; x = 2, y = 6; x = 3, y = 9$
- D. For $x = 0, y = 0; x = 1, y = 3; x = 2, y = 4; x = 3, y = 6$

ugnetcse-dec2015-paper3 geometry lines-curves

Answer key

Answer Keys

26.1.1

A

26.2.1

C